


What Climate Change Means for New Hampshire

New Hampshire's climate is changing. Most of the state has warmed two to three degrees (F) in the last century. Throughout the northeastern United States, spring is arriving earlier and bringing more precipitation, heavy rainstorms are more frequent, and summers are hotter and drier. Sea level is rising, and severe storms cause floods that damage property and infrastructure. In the coming decades, the changing climate is likely to increase flooding, harm ecosystems and winter recreation, disrupt farming, and increase some risks to human health.

Our climate is changing because the earth is warming. People have increased the amount of carbon dioxide in the air by 40 percent since the late 1700s. Other heat-trapping greenhouse gases are also increasing. These gases have warmed the surface and lower atmosphere of our planet about one degree during the last 50 years. Evaporation increases as the atmosphere warms, which increases humidity, average rainfall, and the frequency of heavy rainstorms in many places—but contributes to drought in others.

Greenhouse gases are also changing the world's oceans and ice cover. Carbon dioxide reacts with water to form carbonic acid, so the oceans are becoming more acidic. The surface of the ocean has warmed about one degree during the last 80 years. Warming is causing snow to melt earlier in spring, and mountain glaciers are retreating. Even the great ice sheets on Greenland and Antarctica are shrinking. Thus the sea is rising at an increasing rate.



Rising temperatures in the last century. New Hampshire has warmed twice as much as the rest of the contiguous 48 states. Source: EPA, *Climate Change Indicators in the United States*.

Increasing Temperature and Changing Precipitation Patterns

Rising temperatures and shifting rainfall patterns are likely to increase the intensity of both floods and droughts. Average annual precipitation in the Northeast increased 10 percent from 1895 to 2011, and precipitation from extremely heavy storms has increased 70 percent since 1958. During the next century, average annual precipitation and the frequency of heavy downpours are likely to keep rising. Average precipitation is likely to increase during winter and spring, but not change significantly during summer and fall. Rising temperatures will melt snow earlier in spring and increase evaporation, and thereby dry the soil during summer and fall. So flooding is likely to be worse during winter and spring, and droughts worse during summer and fall.



Torrential downpours flooded Alstead and neighboring communities in 2005, causing millions of dollars of damage. Credit: EPA.

Winter Recreation

Warmer winters may bring more rain and less snow to New Hampshire. A decline in snowfall would shorten the season during which the ground is covered with snow, which could harm recreational industries like skiing, snowboarding, and snowmobiling, and the local economies that depend on them.

Agriculture

Climate change may pose challenges for agriculture. Some farms may be harmed if more hot days and droughts reduce crop yields, or if more flooding and wetter springs delay their planting dates. Other farms may benefit from a longer growing season and the fertilizing effect of carbon dioxide.

Sea Level Rise, Wetland Loss, and Coastal Flooding

Rising sea level erodes wetlands and beaches and increases damage from coastal storms. Tidal wetlands are inherently vulnerable because of their low elevations, and shoreline development prevents them from migrating inland onto higher ground. Human activities such as filling wetlands have destroyed about one third of New England's coastal wetlands since the early 1800s. Wetlands provide habitat for many bird species, such as osprey and heron, as well as several fish species. Losing coastal wetlands would harm coastal ecosystems and remove an important line of defense against coastal flooding.

Coastal cities and towns will become more vulnerable to storms in the coming century as sea level rises, shorelines erode, and storm surges become higher. Storms can destroy coastal homes, wash out highways and rail lines, and damage essential communication, energy, and wastewater management infrastructure.



The Great Bay area has marshes and infrastructure that are at risk from sea level rise. © Ben Miller; used by permission.

Ecosystems

Changing the climate threatens ecosystems by disrupting relationships between species. Wildflowers and woody perennials are blooming—and migratory birds are arriving—sooner in spring. Not all species adjust in the same way, however, so the food that one species needs may no longer be available when that species arrives on its migration. Rising temperatures allow deer populations to increase, reducing forest underbrush, which makes some animals more vulnerable to predators.


Climate change can allow invasive species to expand their ranges. For example, the hemlock woolly adelgid has infested hemlock trees in southern New Hampshire. Infestation eventually kills almost all hemlock trees, which are replaced by black oaks,

black birch, and other hardwoods. Warmer temperatures are likely to enable the woolly adelgid to expand northward. The loss of hemlock trees would remove the primary habitat for the blue-headed vireo and Blackburnian warbler. It could also change stream temperatures and cause streams to run dry more often, harming brook trout and brown trout.

Human Health

Changes in temperature and precipitation could increase the incidence of acute and chronic respiratory conditions such as asthma. Higher temperatures can increase the formation of ground-level ozone (smog), a pollutant that can contribute to respiratory problems. Rising temperatures may also increase the length and severity of the pollen season for plants such as ragweed—which has already been observed in other regions. Certain people are especially vulnerable, including children, the elderly, the sick, and the poor.

The risk of some diseases carried by insects may also increase. The ticks that transmit Lyme disease are active when temperatures are above 45°F, so warmer winters could lengthen the season during which ticks can become infected or people can be exposed to the ticks. Higher temperatures would also make more of New England warm enough for the Asian tiger mosquito, a common carrier of West Nile virus. The number of cases may or may not increase, depending on what people do to control insect populations and avoid insect bites.



Increase in Lyme disease between 1996 and 2013. Each dark dot shows one case reported in 1996; light dots show 2013. The increased range shown here has been attributed to factors other than climate change. Nevertheless, additional warming will lengthen the season during which people are exposed to Lyme disease and may allow the disease to spread to colder areas. Source: CDC.