



PROTOCOL FOR TESTS ON DISTINCTNESS, UNIFORMITY AND STABILITY

Agapanthus L'Hér.

AFRICAN LILY

UPOV Code: AGAPA

Adopted on 19/03/2014

Entry into force on 19/03/2014

TABLE OF CONTENTS

CPVO-TP/266/1

| | | |
|-----|---|----|
| 1. | SUBJECT OF THE PROTOCOL AND REPORTING..... | 3 |
| 1.1 | Scope of the technical protocol..... | 3 |
| 1.2 | Entry Into Force..... | 3 |
| 1.3 | Reporting between Examination Office and CPVO and Liaison with Applicant..... | 3 |
| 2. | MATERIAL REQUIRED | 4 |
| 2.1 | Plant material requirements | 4 |
| 2.2 | Informing the applicant of plant material requirements..... | 4 |
| 2.3 | Informing about problems on the submission of material | 4 |
| 3. | METHOD OF EXAMINATION..... | 4 |
| 3.1 | Number of growing cycles..... | 4 |
| 3.2 | Testing Place | 4 |
| 3.3 | Conditions for Conducting the Examination..... | 4 |
| 3.4 | Test design..... | 4 |
| 3.5 | Additional tests | 5 |
| 3.6 | Constitution and maintenance of a variety collection | 5 |
| 4. | ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY | 5 |
| 4.1 | Distinctness | 5 |
| 4.2 | Uniformity | 6 |
| 4.3 | Stability..... | 6 |
| 5. | GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL..... | 7 |
| 6. | INTRODUCTION TO THE TABLE OF CHARACTERISTICS | 7 |
| 6.1 | Characteristics to be used | 7 |
| 6.2 | Example Varieties..... | 8 |
| 6.3 | Legend..... | 8 |
| 7. | TABLE OF CHARACTERISTICS..... | 9 |
| 8. | EXPLANATIONS ON THE TABLE OF CHARACTERISTICS..... | 17 |
| 8.1 | Explanations covering several characteristics | 17 |
| 8.2 | Explanations for individual characteristics..... | 17 |
| 9. | LITERATURE | 23 |
| 10. | TECHNICAL QUESTIONNAIRE | 24 |

1. SUBJECT OF THE PROTOCOL AND REPORTING

1.1 Scope of the technical protocol

This Technical Protocol applies to all varieties of *Agapanthus L'Hér.*

The protocol describes the technical procedures to be followed in order to meet the requirements of Council Regulation 2100/94 on Community Plant Variety Rights. The technical procedures have been agreed by the Administrative Council and are based on documents agreed by the International Union for the Protection of New Varieties of Plants (UPOV), such as the General Introduction to DUS (UPOV Document TG/1/3 http://www.upov.int/export/sites/upov/resource/en/tg_1_3.pdf), its associated TGP documents (<http://www.upov.int/tgp/en/>) and the relevant UPOV Test Guideline TG/266/1 Rev. dated 20/03/2013 (<http://www.upov.int/edocs/tgdocs/en/tg266.pdf>) for the conduct of tests for Distinctness, Uniformity and Stability.

1.2 Entry into Force

The present protocol enters into force on **19.03.2014**. Any on-going DUS examination of candidate varieties started before the aforesaid date will not be affected by the approval of the Technical Protocol. Technical examinations of candidate varieties are carried out according to the TP in force when the DUS test starts. The starting date of a DUS examination is considered to be the due date for submitting of plant material for the first test period.

In cases where the Office requests to take-over a DUS report for which the technical examination has either been finalized or which is in the process to be carried out at the moment of this request, such report can only be accepted if the technical examination has been carried out according to the CPVO TP which was in force at the moment when the technical examination started.

1.3 Reporting between Examination Office and CPVO and Liaison with Applicant

1.3.1 Reporting between Examination Office and CPVO

The Examination Office shall deliver to the CPVO a preliminary report ("the preliminary report") no later than two weeks after the date of the request for technical examination by the CPVO.

The Examination Office shall also deliver to the CPVO a report relating to each growing period ("the interim report") and, when the Examination Office considers the results of the technical examination to be adequate to evaluate the variety or the CPVO so requests, a report relating to the examination ("the final report").

The final report shall state the opinion of the Examination Office on the distinctness, uniformity and stability of the variety. Where it considers those criteria to be satisfied, or where the CPVO so requests, a description of the variety shall be added to the report. If a report is negative the Examination Office shall set out the detailed reasons for its findings.

The interim and the final reports shall be delivered to the CPVO as soon as possible and no later than on the deadlines as laid down in the designation agreement.

1.3.2 Informing on problems in the DUS test

If problems arise during the course of the test the CPVO should be informed immediately so that the information can be passed on to the applicant. Subject to prior permanent agreement, the applicant may be directly informed at the same time as the CPVO particularly if a visit to the trial is advisable.

1.3.3 Sample keeping in case of problems

If the technical examination has resulted in a negative report, the CPVO shall inform the Examination Office as soon as possible in case that a representative sample of any relevant testing material shall be kept.

2. MATERIAL REQUIRED

2.1 Plant material requirements

Information with respect to the agreed closing dates and submission requirements of plant material for the technical examination of varieties can be found on <http://cpvo.europa.eu/applications-and-examinations/technical-examinations/submission-of-plant-material-s2-publication> in the special issue S2 of the Official Gazette of the Office. General requirements on submission of samples are also to be found following the same link.

2.2 Informing the applicant of plant material requirements

The CPVO informs the applicant that

- he is responsible for ensuring compliance with any customs and plant health requirements.
- the plant material supplied should be visibly healthy, not lacking in vigour, nor affected by any important pest or disease.
- the plant material should not have undergone any treatment which would affect the expression of the characteristics of the variety, unless the competent authorities allow or request such treatment. If it has been treated, full details of the treatment must be given.

2.3 Informing about problems on the submission of material

The Examination Office shall report to the CPVO immediately in cases where the test material of the candidate variety has not arrived in time or in cases where the material submitted does not fulfil the conditions laid down in the request for material issued by the CPVO.

In cases where the examination office encounters difficulties to obtain plant material of reference varieties the CPVO should be informed.

3. METHOD OF EXAMINATION

3.1 Number of growing cycles

The minimum duration of tests should normally be a single growing cycle.

3.2 Testing Place

Tests are normally conducted at one place. In the case of tests conducted at more than one place, guidance is provided in TGP/9 "Examining Distinctness"

http://www.upov.int/export/sites/upov/en/publications/tgp/documents/tgp_9_1.pdf.

3.3 Conditions for Conducting the Examination

The tests should be carried out under conditions ensuring satisfactory growth for the expression of the relevant characteristics of the variety and for the conduct of the examination.

Because daylight varies, colour determinations made against a colour chart should be made either in a suitable cabinet providing artificial daylight or in the middle of the day in a room without direct sunlight. The spectral distribution of the illuminant for artificial daylight should conform with the CIE Standard of Preferred Daylight D 6500 and should fall within the tolerances set out in the British Standard 950, Part I. These determinations should be made with the plant part placed against a white background. The colour chart and version used should be specified in the variety description.

3.4 Test design

3.4.1 Each test should be designed to result in a total of at least 10 plants.

3.4.2 The design of the tests should be such that plants or parts of plants may be removed for measurement or counting without prejudice to the observations which must be made up to the end of the growing cycle.

3.5 Additional tests

In accordance with Article 83(3) of Council Regulation No. 2100/94 an applicant may claim either in the Technical Questionnaire or during the test that a candidate has a characteristic which would be helpful in establishing distinctness. If such a claim is made and is supported by reliable technical data, an additional test may be undertaken providing that a technically acceptable test procedure can be devised.

Additional tests will be undertaken, with the agreement of the President of CPVO, where distinctness is unlikely to be shown using the characters listed in the protocol.

3.6 Constitution and maintenance of a variety collection

The process for the constitution and the maintenance of a variety collection can be summarized as follows:

Step 1: Making an inventory of the varieties of common knowledge

Step 2: Establishing a collection ("variety collection") of varieties of common knowledge which are relevant for the examination of distinctness of candidate varieties

Step 3: Selecting the varieties from the variety collection which need to be included in the growing trial or other tests for the examination of distinctness of a particular candidate variety.

3.6.1 Forms of variety collection

The variety collection shall comprise variety descriptions and may comprise living plant material. The variety description shall be produced by the examination office unless special cooperation exists between examination offices and the CPVO. The descriptive and pictorial information produced by the examination office shall be held and maintained in a form of a database.

3.6.2 Living Plant Material

The examination office shall obtain living plant material of reference varieties as and when those varieties need to be included in growing trials or other tests.

3.6.3 Making an inventory of varieties of common knowledge for inclusion in the variety collection

The inventory shall include varieties protected under National and Community PBR and varieties in trade or in commercial registers.

In addition to the above, the inventory shall be extended to the appropriate to

- any commercial document in which varieties are marketed as propagating or harvested material, especially when there is no official registration system;
- any list including varieties which are publicly available within plant collections (varieties included in genetic resource collections, collection of old varieties, etc.);
- information provided by relevant plant experts;
- relevant example varieties referred to in the technical protocols for the examination of distinctness.

4. ASSESSMENT OF DISTINCTNESS, UNIFORMITY AND STABILITY

The prescribed procedure is to assess distinctness, uniformity and stability in a growing trial.

4.1 Distinctness

4.1.1 General recommendations

It is of particular importance for users of this Technical Protocol to consult the UPOV-General Introduction to DUS (link in chapter 1 of this document) and TGP 9 'Examining Distinctness' (http://www.upov.int/edocs/tgpdocs/en/tgp_9.pdf) prior to making decisions regarding distinctness. However, the following points are provided for elaboration or emphasis in this Technical Protocol.

4.1.2. Consistent differences

The differences observed between varieties may be so clear that more than one growing cycle is not necessary. In addition, in some circumstances, the influence of the environment is not such that more than a single growing cycle is required to provide assurance that the differences observed between varieties are sufficiently consistent. One means of ensuring that a difference in a characteristic, observed in a growing trial, is sufficiently consistent is to examine the characteristic in at least two independent growing cycles.

4.1.3 Clear differences

Determining whether a difference between two varieties is clear depends on many factors, and should consider, in particular, the type of expression of the characteristic being examined, i.e. whether it is expressed in a qualitative, quantitative, or pseudo-qualitative manner. Therefore, it is important that users of these Technical Protocols are familiar with the recommendations contained in the UPOV-General Introduction to DUS prior to making decisions regarding distinctness.

4.1.4 Number of plants/parts of plants to be examined

Unless otherwise indicated, for the purposes of distinctness, all observations on single plants should be made on 5 plants or parts taken from each of 5 plants and any other observations made on all plants in the test, disregarding any off-type plants.

4.1.5 Method of observation

The recommended method of observing the characteristic for the purposes of distinctness is indicated by the following key in the third column of the Table of Characteristics (see document TGP/9 "Examining Distinctness", Section 4 "Observation of characteristics"):

| | |
|-----|---|
| MG: | single measurement of a group of plants or parts of plants |
| MS: | measurement of a number of individual plants or parts of plants |
| VG: | visual assessment by a single observation of a group of plants or parts of plants |
| VS: | visual assessment by observation of individual plants or parts of plants |

Type of observation: visual (V) or measurement (M)

"Visual" observation (V) is an observation made on the basis of the expert's judgment. For the purposes of this document, "visual" observation refers to the sensory observations of the experts and, therefore, also includes smell, taste and touch. Visual observation includes observations where the expert uses reference points (e.g. diagrams, example varieties, side-by-side comparison) or non-linear charts (e.g. colour charts). Measurement (M) is an objective observation against a calibrated, linear scale e.g. using a ruler, weighing scales, colorimeter, dates, counts, etc.

Type of record: for a group of plants (G) or for single, individual plants (S)

For the purposes of distinctness, observations may be recorded as a single record for a group of plants or parts of plants (G), or may be recorded as records for a number of single, individual plants or parts of plants (S). In most cases, "G" provides a single record per variety and it is not possible or necessary to apply statistical methods in a plant-by-plant analysis for the assessment of distinctness."

In cases where more than one method of observing the characteristic is indicated in the Table of Characteristics (e.g. VG/MG), guidance on selecting an appropriate method is provided in document TGP/9, Section 4.2.

4.2 **Uniformity**

It is of particular importance for users of this Technical Protocol to consult the UPOV-General Introduction to DUS (link in chapter 1 of this document) and TGP 10 'Examining Uniformity' (http://www.upov.int/edocs/tgpdocs/en/tgp_10.pdf) prior to making decisions regarding uniformity. However, the following point is provided for elaboration or emphasis in this Technical Protocol.

Uniformity assessment by off-types

For the assessment of uniformity, a population standard of 1% and an acceptance probability of at least 95 % should be applied. In the case of a sample size of 10 plants, 1 off-type is allowed.

4.3 **Stability**

4.3.1 It is of particular importance for users of this Technical Protocol to consult the UPOV-General Introduction to DUS (link in chapter 1 of this document) and TGP 11 'Examining Stability' (http://www.upov.int/edocs/tgpdocs/en/tgp_11.pdf).

In practice, it is not usual to perform tests of stability that produce results as certain as those of the testing of distinctness and uniformity. However, experience has demonstrated that, for many types of variety, when a variety has been shown to be uniform, it can also be considered to be stable.

4.3.2 Where appropriate, or in cases of doubt, stability may be further examined by testing a new plant stock to ensure that it exhibits the same characteristics as those shown by the initial material supplied.

5. GROUPING OF VARIETIES AND ORGANIZATION OF THE GROWING TRIAL

- 5.1** The selection of varieties of common knowledge to be grown in the trial with the candidate varieties and the way in which these varieties are divided into groups to facilitate the assessment of distinctness are aided by the use of grouping characteristics.
- 5.2** Grouping characteristics are those in which the documented states of expression, even where produced at different locations, can be used, either individually or in combination with other such characteristics: (a) to select varieties of common knowledge that can be excluded from the growing trial used for examination of distinctness; and (b) to organize the growing trial so that similar varieties are grouped together.
- 5.3** The following have been agreed as useful grouping characteristics:
- a) Plant: type (characteristic 1)
 - b) Leaf: variegation (characteristic 7)
 - c) Inflorescence bract: opening (characteristic 14)
 - d) Inflorescence: number of flowers (characteristic 19)
 - e) Flower bud: main colour (characteristic 22) with the following groups:
 - Gr. 1: white
 - Gr. 2: yellow green
 - Gr. 3: violet
 - Gr. 4: violet blue
 - Gr. 5: blue
 - f) Flower: type (characteristic 29)
 - g) Anther: colour (characteristic 43)
- 5.4** If other characteristics than those from the TP are used for the selection of varieties to be included into the growing trial, the examination office shall inform the CPVO and seek the prior consent of the CPVO before using these characteristics.

6. INTRODUCTION TO THE TABLE OF CHARACTERISTICS

6.1 Characteristics to be used

The characteristics to be used in DUS tests and preparation of descriptions shall be those referred to in the table of characteristics. All the characteristics shall be used, providing that observation of a characteristic is not rendered impossible by the expression of any other characteristic, or the expression of a characteristic is prevented by the environmental conditions under which the test is conducted or by specific legislation on plant health. In the latter case, the CPVO should be informed.

The Administrative Council empowers the President, in accordance with Article 23 of Commission Regulation N°874/2009, to insert additional characteristics and their expressions in respect of a variety.

States of expression and corresponding notes

In the case of qualitative and pseudo-qualitative characteristics, all relevant states of expression are presented in the characteristic. However, in the case of quantitative characteristics with 5 or more states, an abbreviated scale may be used to minimize the size of the Table of Characteristics. For example, in the case of a quantitative characteristic with 9 states, the presentation of states of expression in the Test Guidelines may be abbreviated as follows:

| State | Note |
|--------|------|
| small | 3 |
| medium | 5 |
| large | 7 |

However, it should be noted that all of the following 9 states of expression exist to describe varieties and should be used as appropriate:

| State | Note |
|---------------------|------|
| very small | 1 |
| very small to small | 2 |
| small | 3 |
| small to medium | 4 |
| medium | 5 |
| medium to large | 6 |
| large | 7 |
| large to very large | 8 |
| very large | 9 |

6.2 Example Varieties

Where appropriate, example varieties are provided to clarify the states of expression of each characteristic.

6.3 Legend

For the CPVO N° column:

| | | |
|----------------|-----------------------------------|---------------------|
| G | Grouping characteristic | – see Chapter 5 |
| MG, MS, VG, VS | | – see Chapter 4.1.5 |
| QL | Qualitative characteristic | |
| QN | Quantitative characteristic | |
| PQ | Pseudo-qualitative characteristic | |

For the UPOV N° column:

The numbering of the characteristics is provided as a reference to the ad hoc UPOV guideline.

(*) UPOV Asterisked characteristic – Characteristics that are important for the international harmonization of variety descriptions.

| | |
|---------|---|
| (a)-(g) | See Explanations on the Table of Characteristics in Chapter 8.1 |
| (+) | See Explanations on the Table of Characteristics in Chapter 8.2 |

7. TABLE OF CHARACTERISTICS

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|-----------------------|-------------------------|----------------------|--|-----------------|-------------|
| 1. | 1. (*) | VG | Plant: type | | |
| (+) | | | deciduous | Deep Blue | 1 |
| QL G | | | evergreen | Cloudy Skies | 2 |
| 2. | 2. (*) | VG | Plant: density of foliage | | |
| QN | | (a) | sparse | Baby Pete | 3 |
| | | | medium | Snow Cloud | 5 |
| | | | dense | Snowstorm | 7 |
| 3. | 3. | MG | Plant: number of leaves per shoot | | |
| (+) | | (a) | few | Lapis | 1 |
| QN | | | medium | Regal Beauty | 2 |
| | | | many | Snow Cloud | 3 |
| 4. | 4. | VG/MS | Leaf: length | | |
| QN | | (a) | short | Tinkerbelle | 3 |
| | | (b) | medium | Everblue | 5 |
| | | | long | Deep Blue | 7 |
| 5. | 5. (*) | VG/MS | Leaf: width | | |
| QN | | (a) | narrow | Deep Blue | 3 |
| | | (b) | medium | Buddy Blue | 5 |
| | | | broad | Glen Avon | 7 |
| 6. | 6. | VG | Leaf: curvature | | |
| QN | | (a) | absent or slightly recurved | Tinkerbelle | 1 |
| | | (b) | moderately recurved | Aureovittatus | 2 |
| | | | strongly recurved | Summer Gold | 3 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|------------|----------------|---------------|---|-------------------------|------|
| 7. | 7. (*) | VG | Leaf: variegation | | |
| QL | | (a) | absent | Blue Velvet | 1 |
| G | | (c) | present | Tinkerbelle | 9 |
| 8. | 8. | VG | Leaf: disappearance of variegation with development | | |
| (+) | | | absent or weak | Tinkerbelle | 1 |
| QN | | | medium | | 2 |
| | | | strong | Lemon & Lime | 3 |
| 9. | 9. (*) | VG | Leaf: green colour of upper side (excluding variegation) | | |
| PQ | | (a) | yellow green | | 1 |
| | | (b) | light green | Sybil Martin | 2 |
| | | (d) | medium green | Snowstorm | 3 |
| | | | dark green | Blue Horizons | 4 |
| | | | grey green | Goldstrike | 5 |
| 10. | 10. (*) | VG | Leaf: colour of variegation of upper side | | |
| PQ | | (a) | white | Tinkerbelle | 1 |
| | | (c) | yellow white | Silver Star | 2 |
| | | (d) | yellow | Goldstrike, Summer Gold | 3 |
| | | | yellow green | Sybil Martin | 4 |
| 11. | 11. (*) | VG | Leaf: anthocyanin coloration at base | | |
| (+) | | (a) | absent | Blue Velvet | 1 |
| QL | | | present | Wiley J | 9 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|-------------|----------------|---------------|---|-------------------|------|
| 12. | 12. | VG | Inflorescence bract: length of tip relative to total length of bract | | |
| (+) | | (a) | very short | Wiley J | 1 |
| QN | | | short | Lilac Beauty | 3 |
| | | | medium | Nana Blue | 5 |
| | | | long | Smurfy Blue | 7 |
| | | | very long | Hartenbos White | 9 |
| 13. | 13. (*) | VG | Inflorescence bract: anthocyanin coloration | | |
| QN | | (a) | absent or weak | ATl blu | 1 |
| | | | medium | Peter Pan | 2 |
| | | | strong | Victoria Bay | 3 |
| 14. | 14. (*) | VG | Inflorescence bract: opening | | |
| (+) | | | one side | Wiley J | 1 |
| QL G | | | both sides | Double Diamond | 2 |
| 15. | 15. (*) | VG/MS | Peduncle: length | | |
| QN | | | very short | Double Diamond | 1 |
| | | | short | Princess Margaret | 3 |
| | | | medium | Shinkai | 5 |
| | | | long | Ivory Bells | 7 |
| | | | very long | Purple Cloud | 9 |
| 16. | 16. (*) | VG | Peduncle: thickness | | |
| QN | | (e) | thin | Everblue | 3 |
| | | | medium | Buddy Blue | 5 |
| | | | thick | Cloudy Skies | 7 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|------------|-------------------|---------------|---|---------------|------|
| 17. | 17. (*) | VG | Peduncle: shape in cross section | | |
| (+) | | (e) | medium elliptic | Blue Velvet | 1 |
| QN | | | broad elliptic | Wiley J | 2 |
| | | | circular | Peter Pan | 3 |
| 18. | 18. (*) | VG | Peduncle: anthocyanin coloration | | |
| QN | | (e) | absent or weak | Peter Pan | 1 |
| | | | medium | Midnight Blue | 2 |
| | | | strong | Black Beauty | 3 |
| 19. | 19. (*) | VG/MS | Inflorescence: number of flowers | | |
| (+) | | | very few | Peter Pan | 1 |
| QN | | | few | Bright Eyes | 3 |
| | | | medium | Blue Velvet | 5 |
| | | | many | Magnifico | 7 |
| G | | | very many | Maximus | 9 |
| 20. | 20. (*) | VG/MS | Inflorescence: diameter | | |
| (+) | | | very small | Franni | 1 |
| QN | | | small | Adonis | 3 |
| | | | medium | Atlas | 5 |
| | | | large | Colossus | 7 |
| | | | very large | Trudy | 9 |
| 21. | 21. (*) | VG | Inflorescence: shape in lateral view | | |
| (+) | | | elliptic | Tall Boy | 1 |
| PQ | | | circular | Pinchbeck | 2 |
| | | | narrow oblate | Deep Blue | 3 |
| | | | broad oblate | Loch Hope | 4 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|-----------------------|--------------------|--------------------------|--|-----------------|------|
| 22. | 22. (*) | VG | Flower bud: main colour | | |
| PQ G | | (d) (f) | RHS Colour Chart (indicate reference number) | | |
| 23. | 23. | VG | Flower bud: secondary colour | | |
| PQ | | (d) (f) | RHS Colour Chart (indicate reference number) | | |
| 24. | 24. (*) | VG | Flower bud: distribution of secondary colour | | |
| PQ | | (d) | none | Double Diamond | 1 |
| | | (f) | towards base | Cloudy Skies | 2 |
| | | | towards apex | White Beauty | 3 |
| 25. | 25. | VG/MS | Pedicle: length | | |
| QN | | (f) | short | Deep Blue | 3 |
| | | | medium | Blue Velvet | 5 |
| | | | long | Cloudy Skies | 7 |
| 26. | 26. | VG | Pedicle: anthocyanin coloration | | |
| QN | | (f) | absent or weak | Stéphanie Charm | 1 |
| | | | medium | Silver Jubilee | 2 |
| | | | strong | Black Beauty | 3 |
| 27. | 27. | VG | Pedicle: distribution of anthocyanin coloration | | |
| (+) | | (f) | only on middle third | Everblue | 1 |
| PQ | | | entire | Black Beauty | 2 |
| | | | only on upper and lower third | Victoria Bay | 3 |
| 28. | 28. (*) | VG | Flower: shape | | |
| (+) | | (g) | globose | Buddy Blue | 1 |
| PQ | | | tubular | Graskop | 2 |
| | | | funnel | Amethyst | 3 |
| | | | campanulate | Blue Stars | 4 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|-----------------------|--------------------|---------------|---|----------------|------|
| 29. | 29. (*) | VG | Flower: type | | |
| (+) | | (g) | single | Blue Velvet | 1 |
| QL G | | | semi-double | Double Diamond | 2 |
| 30. | 30. (*) | VG/MS | Perianth: length | | |
| (+) | | (g) | short | Lilliput | 3 |
| QN | | | medium | Blue Velvet | 5 |
| | | | long | Graskop | 7 |
| 31. | 31. (*) | VG/MS | Perianth: diameter | | |
| (+) | | (g) | small | Graskop | 3 |
| QN | | | medium | Amethyst | 5 |
| | | | large | Atlantic Ocean | 7 |
| 32. | 32. | VG | Perianth: overlapping of tepal lobes | | |
| (+) | | (g) | absent | Goliath | 1 |
| QN | | | incomplete | Amethyst | 2 |
| | | | complete | Graskop | 3 |
| 33. | 33. (*) | VG/MS | Perianth tube: length | | |
| (+) | | (g) | short | Peter Pan | 3 |
| QN | | | medium | Goliath | 5 |
| | | | long | Graskop | 7 |
| 34. | 34. (*) | VG | Perianth tube: main colour of outer side | | |
| PQ | | (d) (g) | RHS Colour Chart (indicate reference number) | | |
| 35. | 35. | VG | Tepal lobe: ratio length/width | | |
| (+) | | (g) | slightly elongated | Blue Globe | 1 |
| QN | | | moderately elongated | Elisabeth | 2 |
| | | | strongly elongated | Atlantic Ocean | 3 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|------------|--------------------|---------------|--|--------------|------|
| 36. | 36. (*) | VG | Tepal lobe: colour of marginal zone of inner side | | |
| PQ | | (g) | RHS Colour Chart (indicate reference number) | | |
| 37. | 37. (*) | VG | Tepal lobe: colour of midrib zone of inner side | | |
| PQ | | (g) | RHS Colour Chart (indicate reference number) | | |
| 38. | 38. | VG | Tepal lobe: transparency of midrib zone of inner side | | |
| QN | | (g) | absent or weak | Amethyst | 1 |
| | | | medium | Cloudy Skies | 2 |
| | | | strong | Windsor Grey | 3 |
| 39. | 39. | VG | Tepal lobe: undulation of margin | | |
| QN | | (g) | weak | Amethyst | 1 |
| | | | medium | Blue Heaven | 2 |
| | | | strong | Blue Stars | 3 |
| 40. | 40. (*) | VG | Flower: tepal-like staminodes and pistillodes | | |
| (+) | | (g) | absent | Blue Velvet | 1 |
| QL | | | present | Flore Pleno | 9 |
| 41. | 41. (*) | VG | Flower: extrusion of stamens | | |
| (+) | | (g) | absent or weak | Kama | 1 |
| QN | | | medium | Blue Velvet | 2 |
| | | | strong | Blue Stars | 3 |
| 42. | 42. (*) | VG | Flower: colour | | |
| PQ | | (g) | white | Blue Moon | 1 |
| | | | violet | Wiley J | 2 |
| | | | violet blue | Black Pantha | 3 |

| CPVO N° | UPOV N° | Stage, Method | Characteristics | Examples | Note |
|------------|--------------------------|---------------|---------------------------------------|------------------|------|
| 43. | 43. (*) | VG | Anther: colour | | |
| (+) | | | white | Lavender Haze | 1 |
| PQ | | | green | Blue Brush | 2 |
| | | | blue green | Glen Avon | 3 |
| | | | light yellow | Ossato Snow | 4 |
| | | | medium yellow | Polar Ice | 5 |
| | | | purple | Corinne | 6 |
| | | | brown | Umbellatus Albus | 7 |
| | | | blue grey | Sarah | 8 |
| G | | | black | Aberdeen | 9 |
| 44. | 44. (*) | VG | Style: colour | | |
| PQ | | (g) | white | Blue Moon | 1 |
| | | | violet | Wiley J | 2 |
| | | | violet blue | Glen Avon | 3 |
| 45. | 45. (*) | MG/VG | Time of beginning of flowering | | |
| (+) | | | early | ATl blu | 3 |
| QN | | | medium | Blue Velvet | 5 |
| | | | late | New Blue | 7 |

8. EXPLANATIONS ON THE TABLE OF CHARACTERISTICS


8.1 Explanations covering several characteristics

Unless otherwise indicated, all characteristics should be observed at the time when at least 50% of all flowers have opened.

Characteristics containing the following key in the third column of the Table of Characteristics should be examined as indicated below:

- (a) To be observed when the first flower bud starts to protrude from the inflorescence bract.
- (b) Observations should be made on fully developed leaves.
- (c) Observations on leaf variegation should be made on young leaves.
- (d) The main colour is the colour of the largest surface area. In cases where it is difficult to determine the largest surface area, the darkest colour is considered to be the main colour. The secondary colour is the colour of the second largest surface area.
- (e) To be observed in the middle third of the peduncle.
- (f) Observations should be made when the flower bud is fully expanded, just prior to reflexing of the tepals.
- (g) Observations should be made on fresh fully open flowers.

Diagram of flower parts:



8.2 Explanations for individual characteristics

Ad. 1: Plant: type

Observations should be made during winter and spring.

Ad. 3: Plant: number of leaves per shoot



Ad. 8: Leaf: disappearance of variegation with development

Observations should be made by comparing variegation of young leaves with variegation of mature leaves. Disappearance of variegation is present when mature leaves have less variegation than young leaves.

Ad. 11: Leaf: anthocyanin coloration at base



1
absent



9
present

Ad. 12: Inflorescence bract: length of tip relative to total length of bract



1
very short



3
short



5
medium



7
long



9
very long

Ad. 14: Inflorescence bract: opening

To be observed when most of the flower buds have protruded.



1
one side




2
both sides


Ad. 17: Peduncle: shape in cross section



1
medium elliptic



2
broad elliptic




3
circular


Ad. 19: Inflorescence: number of flowers

The total number of flowers should be assessed, including flower buds, open flowers and faded flowers.


Ad. 20: Inflorescence: diameter




Ad. 21: Inflorescence: shape in lateral view



Ad. 27: Pedicel: distribution of anthocyanin coloration




Ad. 28: Flower: shape



Ad. 29: Flower: type

Single type flowers have six tepals.
Semi-double type flowers have more than six tepals.

Ad. 30: Perianth: length



Ad. 30: Perianth: diameter

The maximum natural diameter should be assessed.

Ad. 32: Perianth: overlapping of tepals lobes



1
absent




2
incomplete



3
complete

Ad. 32: Perianth: overlapping of tepals lobes

Ad. 33: Perianth tube: length



Ad. 35: Tepal lobe: ratio length/width

The tepal lobe is the part of the tepal that does not form part of the perianth tube.

Ad. 40: Flower: tepal-like staminodes and pistillodes




1
absent



9
present

Ad. 41: Flower: extrusion of stamens

The extrusion of stamens is observed as the protrusion of the stamens in relation to the apex of the tepal lobes.



1
absent or weak



2
medium



3
strong

Ad. 43: Anther: colour

The colour of the anthers should be observed just before dehiscence.

Ad. 45: Time of beginning of flowering

Time of beginning of flowering is when 50% of plants have at least one flower fully open.

9. LITERATURE

Duncan, G.D., 1998: Kirstenbosch Gardening Series: Grow Agapanthus: a guide to the species, cultivation and propagation of the genus *Agapanthus*. National Botanical Institute, Kirstenbosch, Cape Town, ZA, 32 pp.

Germishuizen, G., Meyer, N.L., Steenkamp, Y., Keith, M., 2006: A checklist of South African plants. Southern African Botanical Diversity Network Report No. 41, SABONET, Pretoria, ZA

Hattatt, L., 2001: Encyclopedia of garden plants and flowers. Parragon, Bath, UK, 256 pp.

Leighton, F.M., 1965: The genus *Agapanthus* L'Heritier. Journal of South African Botany, Supplementary Volume No. IV, ZA, 50 pp.

Perry, F. (ed.), 1980: The Macdonald encyclopedia of plants & flowers. Macdonald General Books, London, UK

Snoeiijer, W., 2004: *Agapanthus*: a revision of the genus. Timber Press, Inc., Portland, Oregon, US, 320 pp.

Zonneveld, B.J.M., Duncan, G.D., 2003: Taxonomic implications of genome size and pollen colour and vitality for species of *Agapanthus* L'Heritier (Agapanthaceae). Plant Syst. Evol. 241: 115-123

10. TECHNICAL QUESTIONNAIRE

The Technical Questionnaire is available on the CPVO website under the following reference: CPVO-TQ/266/1