

A scoping review of social media in child, adolescents and young adults: research findings in depression, anxiety and other clinical challenges

Donald M. Hilty, Dorothy Stubbe, Alastair J. McKean, Pamela E. Hoffman, Isheeta Zalpuri, Myo T. Myint, Shashank V. Joshi, Murat Pakyurek and Su-Ting T. Li

Background

Social media and other technologies are reshaping communication and health.

Aims

This review addresses the relationship between social media use, behavioural health conditions and psychological well-being for youth aged <25 years.

Method

A scoping review of 11 literature databases from 2000 to 2020 explored research studies in youth in five areas: clinical depression and anxiety, quantitative use, social media mode, engagement and qualitative dimensions and health and well-being.

Results

Out of 2820 potential literature references, 140 met the inclusion criteria. The foci were clinical depression and anxiety disorders ($n = 78$), clinical challenges (e.g. suicidal ideation, cyberbullying) ($n = 34$) and psychological well-being ($n = 28$). Most studies focused on Facebook, Twitter, Instagram and YouTube. Few studies are longitudinal in design ($n = 26$), had comparison groups ($n = 27$), were randomised controlled trials ($n = 3$) or used structured assessments ($n = 4$). Few focused on different youth and sociodemographic populations, particularly for low-income, equity-seeking and deserving populations. Studies examined

association ($n = 120$; 85.7%), mediating ($n = 16$; 11.4%) and causal ($n = 4$; 2.9%) relationships. Prospective, longitudinal studies of depression and anxiety appear to indicate that shorter use (≤ 3 h/day) and purposeful engagement is associated with better mood and psychological well-being. Depression may predict social media use and reduce perception of support. Findings provide families, teachers and providers ways to engage youth.

Conclusions

Research opportunities include clinical outcomes from functional perspective on a health continuum, diverse youth and sociodemographic populations, methodology, intervention and privacy issues. More longitudinal studies, comparison designs and effectiveness approaches are also needed. Health systems face clinical, training and professional development challenges.

Keywords

Social media; adolescents; children; suicide; youth.

Copyright and usage

© The Author(s), 2023. Published by Cambridge University Press on behalf of the Royal College of Psychiatrists. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Children, adolescents and young adults under 25 years of age (i.e. youth) are raised in an increasingly digitalised society, with technology as an integral part of daily life; some researchers suggest 30 years of age as a limit of youth, but there is not consensus on this.¹ Social media is very attractive to youth as it is portable and offers ever-changing, immersive, diverse, individualised social engagement. The following social media platforms have launched since 2000: networks Facebook (2004), Twitter (2006) and LinkedIn (2002); media-sharing networks Instagram (2010), Snapchat (2011) and YouTube (2005); discussion forums Reddit (2005), Quora (2009) and Digg (2004); and bookmarking and content curation networks Pinterest (2010) and Flipboard (2010). Youth mostly use YouTube (81%) and Facebook (69%).¹ Instagram and Snapchat are also commonly used, with the latter as the most important social network for 44% of youth.

Youth are vulnerable in many ways, and may need supervision with social media because of their limited ability to self-regulate, vulnerability to peer pressure and susceptibility to sharing personal information.² Teenagers acknowledge social media's role in helping build their social connections and expose them to a diverse world, and cite concerns around the social pressure that it generates.³ Most (65%) parents worry about their children spending too much time in front of screens, and its impact on mental and physical health, safety, well-being, social development and family dynamics.⁴

The USA Children's Online Privacy Protection Act has effectively guided participants since 1998, if and when those aged ≤ 13 years adhere to parental/guardian permission.⁵

Current state

This review attempts to describe and consider improvements to the literature about social media use in youth and young adults, as there are many things that are still unknown despite past studies and reviews.^{3–16} How social media is used may make a difference in how it is experienced – from browsing through content to posting content to directed communication (e.g. conversational or liking content) – and if this is self-reported, methods are needed to monitor and verify. The positive and negative effects of social media related to clinical populations (i.e. normal versus problematic use) are not well described. Past studies and reviews are limited by the lack of consensus on definitions of terminology (e.g. normal versus problematic use, sexting, cyberbullying);^{3,4,6} the quality of social-media-specific assessment tools and the rigor of other tools applied to social media; quality of study designs (e.g. cross-sectional or short-term designs that limit evaluation of outcomes) and summarising data, with emphasis on the better designs. Prior reviews found that social media use is negatively correlated with well-being,^{7–12} but the linkage to depression and/or lower self-esteem is not clear.^{11–15} Many reviews reported both negative effects (low

mood or esteem, decreased offline prosocial activity, overuse, impulsivity) and positive effects (developing friends, feeling connected, social capital).¹⁶ Unfortunately, many prior reviews did not clarify the relationship between social media and behavioural health issues (i.e. associative, mediating versus causal relationships).^{8,12} Ideally, more data from across the world is needed, rather than studies from a few countries.

This scoping review explores the question ‘What is the nature of the relationship (i.e. association, mediation, causation and/or other) between social media use in children/adolescents/young adults, psychopathology and mental and/or behavioural health conditions or problems?’. This review is intended to assist providers in educating adolescent/young adult patients and their families in how to best interact with social media. The review has several aims.

- To summarise findings of the relationship (association, mediation, causation) between social media use in children/adolescents/young adults, psychopathology and mental and/or behavioural health conditions or problems.
- To explore the unique challenges, effects and benefits of social media use by youth, related to clinical populations for depression and anxiety (Supplementary Table 1 available at <https://doi.org/10.1192/bjo.2023.523>);^{17–90} clinical challenges like cyberbullying, sexting and suicide (Supplementary Table 2);^{91–123} and health behaviour and well-being (Supplementary Table 3).^{12,124–149}
- Based on the literature, to provide an approach for future clinical research and approaches for providers and health systems to social media in youth (Table 1).

Method

Approach

The literature search was conducted from January 2000 to December 2020. The philosophical approach to the search was done according to the original six-stage process¹⁵⁰ and updated modifications¹⁵¹ (purposeful research question, identifying relevant studies, selecting studies based on an iterative process, charting the data, analysis of findings and consultation from stakeholders). The Preferred Reporting Extension for Systematic Reviews and Meta-Analyses (PRISMA) for scoping reviews¹⁵² has additional suggestions for sources of information, the search and appraising data.

Research question

This review addresses the overarching question: ‘What is the nature of the relationship between social media use, psychopathology and mental and/or behavioural health conditions or problems?’ The population of interest is children, adolescents and young adults (aged ≤25 years). Secondary questions are as follows.

- What social media is commonly used, in what ways and for what purpose(s) (i.e. approach, interest, motivation)?
- In what ways is social media helpful, neutral or negative related to clinical populations for depression and anxiety, and specific problems like cyberbullying, sexting and suicide?
- What is the relationship (i.e. association, mediation, causation and/or other) between social media (e.g. Facebook, Twitter, Instagram) and behavioural health?
- What methods of assessment, triage and approaches, interventions and professional development can help providers, parents, teachers and others in the community to help?

Identifying relevant studies

Eleven databases were queried: PubMed/Medline, APA PsycNET, Cochrane Database of Systematic Reviews, EMBASE, PsycINFO, Web of Science and Scopus, Social Sciences Citation Index (SSCI), Centre for Reviews and Dissemination, Cochrane Central Register of Controlled Trials, Cumulative Index to Nursing and Allied Health Literature (CINAHL) and Google Scholar.

The search focused on youth (adolescent, child, children, high, junior, juvenile, middle, minor, secondary, teenager, youth) and social media use in five concept areas (Fig. 1): clinical depression and anxiety and problematic challenges (e.g. suicidal ideation, cyberbullying); quantitative data; social media mode; engagement and qualitative dimensions; and health and psychological well-being. Definitions were used based on consensus literature: bullying is a subset of aggressive behaviour that involves repeated and intentional attempts to damage/distress a weaker victim by a more powerful perpetrator;¹⁵³ and sexting is sending or receiving of sexually explicit pictures, videos, or text messages via smartphone, digital camera or computer.⁹⁶ Exclusion criteria included studies focusing on anorexia, attention-deficit hyperactivity disorder, physical or intellectual disabilities, genetics, substance use, gambling, sleep/insomnia, cognitive disorders and aggression/violence beyond cyberbullying and suicide) (Fig. 1).

Study selection

One author (D.M.H.) screened titles and abstracts of potential references, excluding duplicates and those that did not meet the search criteria. Two authors (D.M.H., D.S.) reviewed the full text of remaining abstracts to find those meeting inclusion criteria; additional studies that met inclusion criteria were added from references.

Data charting

A data-charting form was used to extract data, and notes were organised with a descriptive analytical method. The reviewers (D.M.H., D.S.) compared and consolidated information by using a modified content analysis with thematic components;¹⁵⁴ a third author (A.J.M.) moderated any disagreement and a fourth author (S.-T.T.L.) analysed consistency of the approach. The information was shared with selected experts, their input summarised and themes extracted.

Analysis, reporting and the meaning of findings

Results were organised into tables, with key concepts and components outlined and described, partially based on excerpts from published topics. The studies varied considerably, and therefore were challenging to compare. Qualitative steps to analyse disparate populations, methods and data of studies were used (Fig. 2).¹⁵⁴ Content, discourse and framework qualitative analysis techniques were used to analyse findings from papers and classify, summarise and tabulate the behavioural data; discourse and thematic analyses were used to search for themes and patterns; and framework analysis was used to sift through, chart and sort data in accordance with key issues and themes a series of steps (e.g. indexing, charting, mapping and interpretation).¹⁵⁴ Data in Supplementary Tables 1–3 are organised by study, sample size, population (e.g. country), objective and design, methods and measures, outcomes and clinical implications/challenges and training/research foci.

Expert opinions and feedback

Expert opinions were solicited to review preliminary findings and suggest additional steps for improvement. A list of relevant experts was compiled from (a) behavioural health organisations




Fig. 1 Search flow diagram for child, adolescent and young adult social media articles reviewed.

across professions internationally; (b) technology-related special interest groups of organisations (e.g. American Telemedicine Medical, Nursing and Informatics Associations); (c) educational and professional development organisations (e.g. Accreditation Council of Graduate Medical Education, American Academy of Child and Adolescent Psychiatry, American Academy of Pediatrics); (d) academic institutions and (e) researchers, authors, editors and editorial board members of journals related to social media.

Experts were invited by email ($N = 24$) and attended a live expert feedback session for discussion and feedback; completed a qualitative and quantitative five-item Likert scale survey ($n = 20$; 83.3%) and/or provided qualitative feedback via email ($n = 4$; 16.7%). The data charting and the search criteria plan were reviewed; their input did not suggest a search with additional terminology or otherwise change the scope. Input was summarised and themes were extracted to guide the organisation (e.g. headings in rows) and content (e.g. in




Fig. 2 Qualitative steps to analyse disparate study populations, methodology and data.

the columns) of Table 1 and Supplementary Tables 1–3, based on previous work using consensus and modified Delphi processes.¹⁵⁴ Results showed that the majority agreed or strongly agreed that the search strategy was effective using the research question ($n = 21$; 87.5%); it was systematic/thorough (20; 83.3%); and adequately scientific in methodology ($n = 18$; 75%); and '[The tables] are organised in a practical way to summarise social media study findings for providers, teachers and systems' ($n = 20$; 83.3%), once more specific outcomes were entered in the final column of each.

Results

Literature overview

Out of 2820 potential literature references, 112 duplicates and 2284 studies that were outside of the scope of this review were excluded (Fig. 1). Full-text review of 424 articles revealed that 128 met full inclusion criteria; 12 additional studies were found within those, for a total of 140 studies.^{12,17–95,97–149,155} The studies focused on clinical populations for depression and anxiety,⁷¹ clinical challenges (e.g. suicidal ideation, cyberbullying)²⁷ or psychological well-being.²¹ Studies were of children aged 12 years and younger ($n = 1$; 0.01%), adolescents (13–18 years) ($n = 72$; 54.1%) and young adults (19–25 years) ($n = 48$; 34.2%); the rest were aggregates of the above ($n = 18$; 12.9%). The overall mean age was 18.78 years. The most common social media studied were Facebook ($n = 62$), Twitter ($n = 20$), Instagram ($n = 11$), YouTube ($n = 6$) and MySpace ($n = 5$). Studies varied in identifying gender identity ($n = 63$; 45%), ethnicity and race ($n = 42$; 30%) or neither ($n = 35$; 25%).

Most studies were cross-sectional cohort studies using self-report questionnaires. Few studies were longitudinal in design,¹⁹ had comparison groups²⁰ or were randomised controlled trials.³ Few studies used clinician/provider-administered instruments^{2,20,30} or structured assessments.^{4,17,51,82,131} Timing or temporal dimensions are generally quite limited and studies span across acute disorders, subacute symptoms and trait/personality factors among a wide variety of ethnic, clinical and non-clinical populations. Broadly speaking, the studies looked at associations ($n = 120$; 85.7%),^{17–23,25–29,31–33,36–41,43–53,55–61,63–73,75,76,78–84,86–95,97–100,102,103,105,112,114–127,129–131,133–142,144,146,148,149,154,155} and mediating ($n = 16$; 11.4%)^{24,30,34,35,62,74,77,85,101,104,113,128,132,143,145} and causal ($n = 4$; 2.9%)^{42,54,104,147} relationships between social media and behavioural health issues.

Clinical populations, depression and anxiety

There were 78 studies of social media with outcomes in clinical populations and disorders (Supplementary Table 1). The mean age was 18.4 years (median 18 years) and included adolescents⁴² and young adults.²¹ The study populations were diverse in terms of ethnicity, but were predominately White, and 46 studies were $\geq 50\%$ female. The mean sample size was 8332.4 (median 310). The most common social media sites studied were Facebook ($n = 37$), Twitter ($n = 10$), Instagram ($n = 5$), MySpace ($n = 2$) and YouTube ($n = 2$); two were on screen time.^{13,33}

Cross-sectional and longitudinal studies^{12,19,22,23,25,27,28,31,35,41,42,47,48,51,65,67,76,79,87,89} of social media use and depression found that shorter periods of social media use (<3 h), particularly with purposeful or active engagement, are associated with better mood

and psychological well-being, whereas longer periods of social media use predict depression (and often anxiety) or poorer psychological function^{22,25,27,28,31,35,48,59,65,87} (particularly browsing^{30,89}), partly because of sleep disruptions.⁴⁷ Cross-sectional and longitudinal studies are consistent with one prospective study that suggests a threshold effect around 3 h that has negative impact for many, but not all, users: low use-stable (80% at 3–4 h/day/item), high use-decreasing (12.3% at 4–5 h/day/item) and low use-increasing (7.3% at 3 to nearly 5 h/day).³³ Two studies found that depression predicts social media use^{33,67} and reduces perception of support.⁵¹ Specifically, Twitter use may be associated with depressive thoughts and symptoms, but only for people with low initial levels of in-person social support, and conveying positive sentiment helped to reduce depressive thoughts and feelings irrespective of people's level of in-person social support.²³ Depressive signals observed in Tweets may predict future depression.⁷⁶ Instagram browsing was associated with increases in depressed mood in adolescents.⁴²

Type of media use is important, since hours spent on social media and internet use were more strongly associated with self-harm behaviours, depressive symptoms, low life satisfaction and low self-esteem than hours spent electronic gaming and watching television.¹² In addition, girls generally demonstrated stronger associations between screen media time and mental health indicators than boys (e.g. heavy internet users were 166% more likely to have clinically relevant levels of depressive symptoms than low users for girls, compared with 75% more likely for boys). A cross-sectional study showed that cortisol systemic output was positively associated with Facebook network size and negatively associated with Facebook peer interactions.⁵⁰

Studies of anxiety disorders are similar to findings in depression studies, with social anxiety symptoms mediated by spending more time on Facebook and passively using Facebook (i.e. viewing other's profiles without interacting).⁶² In a study with three focus groups of those with anxiety disorders, six themes emerged: seeking approval, fearing judgement, escalating interpersonal issues, wanting privacy, negotiating self and social identity and connecting and disconnecting.⁴¹ A qualitative study revealed three types of negative use, including 'oversharing' (frequent updates or too much personal information), 'stressed posting' (sharing negative updates) and encountering 'triggering posts'.⁴⁶ Both social anxiety and need for social assurance had a significant positive association with problematic use of Facebook^{41,54} or 'fear of missing out' (FOMO).^{24,39}

Clinical challenges like suicide, cyberbullying, sexting and other behaviours

The review found 34 studies on clinical challenges such as cyberbullying, sexting and posts on suicide (Supplementary Table 2). The primary populations were children ($n = 1$), adolescent ($n = 15$) and young adults ($n = 14$, with 3 for college students), with a mean age of 18 (median 17.9) years. The study populations were diverse in terms of ethnicity, but were predominately White and 15 studies were $\geq 50\%$ female. The mean sample size was 34934.5 (median 524). The most common social media types studied were Facebook ($n = 10$), Twitter ($n = 8$), Instagram ($n = 5$), YouTube ($n = 3$) and MySpace ($n = 2$).

Excessive social media use, depression, suicide and school burn-out appear strongly related.^{103,107,109,115} One longitudinal study found that, compared with matched non-suicide-related Twitter posts, suicide-related posts were characterised by a higher word count, increased use of first-person pronouns and more references to death.¹⁰³ In this study, emotional engagement, school burn-out and depression contributed to excessive social media use. Similarly, students with burn-out are at higher risk for depression

and excessive social media use. Excessive social media use leads to school burn-out and school burn-out leads to excessive social media use. Individuals who were suicidal felt significantly less belongingness and significantly higher burdensomeness; they also use a higher proportion of achievement-related words and appear protective. Studies have compared artificial intelligence/machine learning to self-report measures to evaluate risk of suicide,¹⁰⁷ para-suicidal events,¹⁰⁹ suicide-related Tweets¹¹² and other behaviors.¹¹⁵ Machine learning can easily differentiate people who are at high suicidal risk from those who are not (linguistic inquiry and word count, decision tree and cross-validation analyses).¹⁰⁷ Machine-learning algorithms accurately identify the clinically significant suicidal rate in 92% of cases (sensitivity: 53%, specificity: 97%, positive predictive value: 75%, negative predictive value: 93%); a higher proportion of achievement-related words appears protective. For a single point of performance for comparison, artificial intelligence/machine learning had roughly 10% false alarms, but correctly identified about 70% of those who will attempt suicide.¹⁰⁹

The relationship of depression, self-esteem and cyberbullying has been evaluated. A study of 8- to 13-year-olds evaluated whether cybervictimisation is prospectively related to negative self-cognitions and depressive symptoms beyond other types of victimisation.¹¹⁰ The majority of participants reported experiencing at least some degree of peer victimisation at either wave 1 or wave 2 (physical: 68.1%, relational: 89.8%, verbal: 87.9%, property related: 65.8%, cyber: 63.1%). Of note, 16.1% of participants obtained raw scores >75 on the Reynolds Adolescent Depression Scale – Version 2 (RADS-2), and 8.1% obtained scores >82 (signifying mild and moderate depression, respectively). Victimisation was correlated with negative cognition and depressive symptoms; it predicted depressive symptoms; age and gender were not predictors of cybervictimisation or depression. Depression is associated with problematic social media use and indirectly predicted cyberbullying perpetration (associations were weak). Another study found that problematic social media use is weakly correlated with depression ($r = 0.22$), gender ($r = -0.15$), age ($r = -0.13$) and self-esteem ($r = -0.11$).⁹⁵ Experiences of LGBTQ participants included both help for coping and cyberbullying leading to depression, stress and suicidal ideation.⁹⁷

Bystander responses to suicidal behaviour and cyberbullying are in sharp contrast. Only 33.6% of participants left a positive, supportive comment on at least one of two suicide posts. Content severity, experience with a loved one's suicide attempts and use of Facebook to meet people were predictive of providing positive comments.⁹⁴ Positive bystander responses (PBRs) were higher in cyberbullying than traditional bullying incidents.¹⁵⁴ Females exhibited more PBRs across both types of bullying. Bullying severity affected PBRs, in that PBRs increased across mild, moderate and severe incidents, consistent across traditional bullying and cyberbullying. PBRs related to cyberbullying included (a) seek help from a teacher or parent, (b) seek help from a peer or friend, (3) direct intervention and (d) providing comfort or emotional support.

Provider access to a patient's social media could assist in identifying suicidal ideation and/or acts, since patients fail to disclose risk factors to physicians; however, there are ethical and privacy concerns about searching a patient's social media platforms.¹⁰⁰

Health behaviour and well-being topics

There were 28 studies on health behaviour and well-being (Supplementary Table 3). The primary populations were adolescents,⁸ college students¹⁴ and young adults.⁶ The study populations were diverse in terms of ethnicity, but were predominately White and 19 studies were $\geq 50\%$ female. The mean sample size was 1558.8 (median 15.8). The most common social media types

studied were Facebook,¹⁵ Twitter ($n = 2$), Instagram ($n = 1$), YouTube ($n = 1$) and MySpace ($n = 1$). The most study population or disorder was depression (8) or anxiety (6).

Of the longitudinal studies, one found that a group deactivated from Facebook for 4 weeks showed small increases in well-being, but no changes in loneliness, compared with a usual use group.¹²⁴ Another study over 2 months examined internalising symptoms (e.g. depression, anxiety and loneliness) related to the content of their Facebook communication and the responses they received from peers.¹³⁵ The mean number of posts was 60.2 overall (88 for girls and 37 for boys). For girls, internalising symptoms predicted negative affect, somatic complaints and eliciting support; they also predicted receiving more peer comments expressing negative affect and peer responses offering support. A study over 9 months evaluated how social media activity affected individual social communication skill and self-esteem.¹⁴⁶ Active social media use (i.e. directed, person-to-person exchanges) increases bonding and bridging social capital and decreases loneliness; passive use does not.

Cross-sectional studies of teenagers examined psychological well-being and differences between girls and boys in use of technologies,¹² screen time^{27,125,126} and social networking services (SNS).¹⁴² The study found that adolescent girls spent more time on smartphones, social media, texting, general computer use and online, and boys spent more time gaming and on electronic devices in general.¹² Associations between moderate or heavy digital media use and low psychological well-being/mental health issues were generally larger for girls than for boys. For both girls and boys, heavy users (≥ 5 h) often rated twice as likely to experience well-being and mental health issues (e.g. risk factors for suicide) as low users. Also important was that the time 12th graders spent online doubled between 2006 and 2016; girls tend to spend more time in friendship dyads and boys in groups, and girls focus more on social relationships and popularity. A study of SNS and social self-concept, self-esteem and depressed mood found that the association between having an SNS and these negative indicators is more common with female youth; overall, frequency of SNS use is a positive predictor of social self-concept.¹⁴²

With regard to college students, studies examined the relationship of social media with well-being,¹²⁸ FOMO,¹³⁰ attachment, social capital¹³⁰ and social closeness based on activity.¹³⁹ Social media use is not associated with mental health problems, nor is emotional regulation; however, emotional regulation is associated with perceived stress and perceived stress is associated with mental health problems.¹²⁸ Social media use does not indirectly predict mental health problems as mediated by perceived stress or emotional regulation. Social media use may indicate challenges with mental health issues or be a way of dealing with difficult emotions. When attachment theory was used to explore individuals' attachment orientations towards Facebook use related to online and offline social capital, a secure attachment was positively associated with online bonding, bridging and all capital, and offline bridging capital; an avoidant attachment was negatively associated with online bonding capital.¹³⁸ Anxious-ambivalent attachment had a direct association with online bonding capital and an indirect effect on all capital through Facebook. Users in the study on social closeness spent 7.82 min consuming content and 3.13 min on participation.¹³⁹ Interacting with others on social media (e.g. commenting on updates) helps users feel closer to other people and this predicts positive emotional states after Facebook use. A study on FOMO involved two groups (10 min/day versus usual use), and both showed decreases in anxiety and FOMO; only the experimental group showed additional decreases in loneliness and depression.¹³⁰ Moderation helps with mood and loneliness, and reduces anxiety and FOMO.

In a study on giving up Facebook, pre- and post-evaluation of perceived stress and well-being was measured by salivary cortisol between 14.00 and 17.00 h; those using Facebook had lower cortisol levels, less perceived stress, decreased life satisfaction and lower social loneliness on the Social and Emotional Loneliness Scale for Adults.¹³¹ One study examined that a user's activities on Twitter estimate a depressive tendency, based on a medium positive correlation ($r = 0.45$) between the Zung Self-Rating Depression Scale and the model estimations of potentially meaningful words (≤ 20).¹⁴⁶ Although a total of 99 words had absolute values of correlation coefficients with Zung scores >0.4 , the highest scores were associated with the following words: even if, very, workplace, hopeless, disappear, too much, sickness, bad and hospital.

Implications for clinicians and researchers across clinical populations, problems and well-being

Findings of this scoping review inform approaches by providers, families and teachers when working with social media in children, adolescents and young adults (Table 1). To understand how technology affects the lives of adolescents and emergent adults, it is necessary to engage them in a conversation, share ideas and be available to help with problems. As many young people (and adults) may consider the internet their 'lifeline' to social engagement, consideration of the problematic aspects of internet use may be met with reluctance.^{6,12,96,156} Exploring beliefs, norms, values, cultural and language factors, and the meaning of technology to the individual, is integral to understanding and meeting the needs of each patient.^{16,23,24,132} For providers, the value of forming and maintaining a trusting, therapeutic alliance with youth cannot be overstated, as quality care depends on patient-provider engagement, open and honest communication and shared decision-making for treatment.^{11,96,157}

An accurate assessment or history is needed of online activities and associated health and risk factors. Internet use may be healthy or problematic, and this continuum may be explored with youth and parents via non-judgemental questioning to clarify the types and extent of technology used (Table 1).^{4,5,17,156} Assessment is enhanced with multiple informants: parents, significant others, schools, primary care providers and/or others that know the youth well.^{156,157} How they use their time, what they enjoy, how they want others to view them, awareness/use of privacy settings and proneness to risky behaviours is a snapshot of esteem and quality of relationships.¹⁵⁷⁻¹⁵⁹

Providers, families and others need an approach to promote healthy use of social media and prevent problematic social media behaviours. Data on the relationship of social media use and its impact on behaviour – association, mediation or causation – and clinical interventions are limited.^{4,5,9,14,158} Nonetheless, positive family/home life, good engagement, supervision and other approaches may reduce risk of risky or dangerous behaviour.^{4,24,38,156} A shared understanding is needed about healthy versus problematic use, how to monitor use and blending social media with alternative activities to meet emotional needs. Individual, peer/group and family education and therapy is often helpful. Motivational interviewing techniques may help co-construct a plan that meshes with values, with parent and provider input.^{3,24,156}

Discussion

This scoping review provides an update to past reviews on evaluation, interventions and outcomes of social media related to clinical populations (e.g. mood and anxiety disorders), clinical challenges

Table 1 Approach for providers to social media use by youth and young adults: clinical questions and protective factors

Areas/questions	Prompts or specifics	Follow-up questions	Protective factors
Normalise, when possible, the use of electronics and devices <ul style="list-style-type: none"> Which do you use? How much time per day? 	Teenagers spend an average of 6–8 h daily <ul style="list-style-type: none"> Options <ul style="list-style-type: none"> Mobile phone Computer/tablet Gaming systems (X-Box, Wii, PlayStation) Television Other 	<ul style="list-style-type: none"> Which do you use the most? Which is your favourite? Compared with the average teen, about how much screen time do you have daily? Do you use privacy settings? Have you lied about your age to gain access? 	<ul style="list-style-type: none"> Purposeful networking/ communication Use focused on new friendships or building current relationship Emotionally stable, restrained or high harm avoidance Approachable, high social skills Serene, satisfied with life and high self-esteem
Screen for social networking accounts/profile. Which is your favourite? Why? <ul style="list-style-type: none"> Are you 'friends' with your parents or siblings? Do your parents limit your time? Daytime Evening/night Do your parents have your passwords? 	Facebook <ul style="list-style-type: none"> Twitter Instagram YouTube Tumblr Google Plus Pinterest Email Snapchat Video-sharing platforms Chat rooms MySpace Other 	<ul style="list-style-type: none"> Do you use media to deal with stress? <ul style="list-style-type: none"> How often? Number of hours per day? Does media use cause you stress, and if so, how? What do you post? <ul style="list-style-type: none"> Photos, posts, phone, school, city, birthdate? Videos: what kind? Have you posted or received inappropriate photos? Have your posts caused problems for you, a friend or family member? How many friends do you have? <ul style="list-style-type: none"> Do you personally know all of the friends on your social media? Have you interacted with strangers online, and if so, do you use your real name? Have you had bad experiences? 	<ul style="list-style-type: none"> Open discussion of use and high level of supervision Those who may need more supervision <ul style="list-style-type: none"> Low levels of psychopathology <ul style="list-style-type: none"> Anxiety Depression Not psychotic Not autism spectrum disorder Not attention-deficit hyperactivity disorder Males Younger Lower-income family (i.e. may need more structure and support)
Screen for positive aspects of internet and media use	Which specific sites have been helpful to you?	<ul style="list-style-type: none"> Have you gotten health or mental health information online? Have you used self-help sites? Have you made friends or gotten closer to a friend online? Have you used apps for wellness? Have you ever helped a friend who felt suicidal online or via text/chat? 	<ul style="list-style-type: none"> Parallel positive family involvement/function <ul style="list-style-type: none"> Interested/involved Higher functioning and education Positive relationship role modelling Positive peer relations <ul style="list-style-type: none"> Opposed to bullying Supportive Friendly actions/ pranks
Screen for problematic use or risky behaviour	Online use disrupts your sleep <ul style="list-style-type: none"> Visiting sites with sensitive topics that you do not want others to know Meeting others online that you do not know Times you were involved with or seen bullying or harassment Sexting 	<ul style="list-style-type: none"> How often do you stay up at night on media? Have you visited sites regarding: <ul style="list-style-type: none"> Weapons Porn Suicide Anorexia Other? Have you interacted with strangers online, and if so, do you use your real name and identification info? Have you planned to meet up with someone you met on the internet? Have you ever been bullied or harassed online? Or have you bullied or harassed others online? Have you regretted posts or do not want friends/parents to see your posts? Have you ever sexted? 	<ul style="list-style-type: none"> Interest shown and support offered Structured periods of access Reward systems in place Regular parenting schedules Consistent availability for chatting and encouragement Lower stress High stimulation/activity Autonomy supported if earned

(e.g. suicide, cyberbullying) and health behaviour and psychological well-being in youth.^{11,12,14–16,150} This scoping review cast a much broader net and shows how substantial data can contribute to diagnosis, monitor symptoms and collect ecologically rich behavioural data as a foundation for future interventions. Of 140 studies reviewed, longitudinal design,¹⁹ comparison groups²⁰ and randomised controlled trials³ were uncommon, resulting in association ($n = 120$; 85.7%), mediating ($n = 16$; 11.4%) and causal ($n = 4$; 2.9%) relationships between social media and behavioural health issues. Specifically, the review found that social media use of >3 h appears to be associated with increased depression and anxiety,

and passive browsing of social media appears to be associated with depression/anxiety compared with purposeful, positive and active engagement; more research is needed to verify these findings. Girls/young women are more likely to be disproportionately affected by depression/anxiety with regards to social media, which is potentially mediated by the type of interaction, whereas boys/young men have more difficult experiences with gaming. However, positive social support inside/outside of social media is protective (Supplementary Tables 1–3). Some studies have overlooked the impact of equity, diversity and inclusion related to social media use, and care is needed so that technology does not

inadvertently contribute to inequity and other injustices. Any of the many dimensions of diversity or differences (e.g. culture, ethnicity, race, religion, sexual orientation, gender identity, language, nationality, immigration status, socioeconomic status, geography) could affect evaluation and intervention.

Research into social media is moving towards standardised methods, interventions and evaluation measures. Studies are limited or have not looked at key issues, such as (a) sociodemographics and health, digital and language literacy; (b) clinical population state or trait; (c) passive consumption, broadcasting and directed purposeful or active engagement/communication; (d) quality of assessment measures (e.g. standardised, clinician/provider-administered instruments or structured assessments rather than self-report questionnaires without confirmation, verification, observation and corroboration); (e) temporal dimensions of symptoms and assessment; and (f) longitudinal design and comparison groups. More information related to equity, diversity and inclusion for the populations using social media, their families and the clinicians involved with assessment and care is needed to evaluate the impact of differences, cultural safety and humility and potential interventions.¹⁶⁰ This could include, but is not limited to, culture, ethnicity, race, religion, sexual orientation, gender identity, language, nationality, immigration status, socioeconomic status, spirituality, disability status, education, clinical diagnoses and geography. Implementation/ effectiveness designs – with longitudinal, quality of life and other dimensions – are also suggested,¹⁵⁷ if well-anchored to health improvement.¹⁶¹ Data from existing empirical foundations, hierarchical evaluation systems and statistical analyses for multiple comparisons and un/adjusted analyses are needed.^{157,161,162}

Research into social media could be helped by other advances in artificial intelligence, informatics and cognitive computing methods. These advance data processing, stratify risk (e.g. suicide) and predict future negative outcomes with longitudinal correlation, predict biomarkers/digital phenotypes (e.g. depression during and after pregnancy) and allow patients or providers to intervene for mood^{65,76} and suicide.^{107,109,112,115,163} Challenging issues include unique populations (e.g. culture, youth, college), the trade-off of privacy versus suicide detection and comparing artificial intelligence approaches with traditional methods. Social media, like wearable sensors, is transforming care by moving from manual transfer of subjective self-reported information during a patient visit to an integrated, longitudinal, minimally intrusive and interactive sharing of data based on the ecology of a person in their natural setting.^{164,165} Artificial intelligence inferential techniques (i.e. applied or performing functions similar to human thinking and analysis) have high predictive power and are reusable; suicide hotlines and face-to-face evaluations are effective methods for suicide intervention, but depend on action by the person with suicidal ideation.

Providers, parents/families and healthcare systems are facing challenges with social media, partly related to how youth live and how their developing brains are shaped by peers and the pervasive influence of technology.¹⁵⁶ There are a range of behaviours across teenagers, adolescents and other age groups, and so a behaviour may be normal for one group and not for another; a behaviour may be healthy or problematic, depending on age. Families, teachers and providers can use data to engage youth with non-judgemental questioning about social media use, use preventive/risk factors for making decisions and, most importantly, stay as close as possible to their young loved ones who may be at risk for hurting themselves – while privacy is important on one hand, notification of families, clinicians and others who could help them may be helpful. Resources are also available from the American Academy of Pediatrics' Media and Communication Toolkit and Family Media

Use Plan,¹⁵⁸ and other agencies.¹⁶⁶ Competencies for social media, mobile health, wearable sensors and other asynchronous technologies^{157,159} include suggestions for training programmes (undergraduate/medical student, graduate/resident). These also address professional development of faculty and institutional change of health systems or academic centres to integrate video¹⁶⁷ and asynchronous technologies.¹⁵⁷

Scoping reviews appear more helpful than other types of reviews for evaluating the broad context, asking questions of the literature and generating questions, approaches, questions and methodologies for current and target states of research.¹⁶⁸ There are limitations to this scoping review. First, a small team conducted the study selection and review, with only one reviewer screening all titles and abstracts. Second, a modified content analysis with thematic analysis components was presented, rather than a quantitative/numerical analysis of the extent and nature of the studies. Similarly, we categorised data into clinical disorders, but a different framework that looks at health from a functional perspective may have been a better option, such as the health continuum (from poor health/illness/languishing to good health/positive health/flourishing). Third, a quality evaluation tool was not used, partly because the diversity of study methodologies, duration and data collection make a thorough integrated review challenging, using a systematic quality evaluation system or the equivalent of a quantitative meta-analysis. In addition, a measure of risk of bias was not used, and is suggested when applicable and possible. There is also an inherent bias in studies of youth populations published in peer-reviewed literature. Cross-sectional studies of associations with multiple factors in applied rather than controlled settings have limitations. Fourth, the review does not cover all of the potentially relevant psychological well-being, stress and related life dimensions of youth. Fifth, this study did not assess if age or other sociodemographic characteristics were associated with or predicted types of social media use; furthermore, future studies and reviews may take the literature further by distinguishing between populations aged ≤17 years and those aged 18–25 years, as well as not extending this to 30 years of age. Sixth, broader input for consensus across organisations could have been helpful, and a qualitative, small-group interview approach with experts, using a semi-structured guide, could have discovered more information. Seventh, the review falls short of covering all psychiatric disorders (e.g. bipolar disorder, schizophrenia, developmental and other childhood disorders). Eighth, the review has some specific findings, yet points out generalised themes and questions; it is not a conclusive data analysis like a systematic review. Lastly, it is important to recognise the digital divide in social media use across different youth and sociodemographic populations, particularly for low-income, equity-seeking and deserving populations and populations in Latin America, Asia, Africa and Oceania.

In conclusion, research is moving forward on evaluation, intervention, monitoring and outcomes of social media use in youth related to clinical disorders, challenges like suicide and cyberbullying, and psychological well-being. Families, teachers and providers can use current data to engage youth with non-judgemental questioning about social media use and be aware of preventive/risk factors. Longitudinal comparison designs, effectiveness approaches, artificial intelligence and biomarking/digital phenotyping may provide a foundation for future interventions to examine causal relationships between social media use and behavioural health. Research opportunities and challenges can be broadly organised into the following categories: clinical outcomes from a functional perspective on a health continuum; diverse youth and sociodemographic populations, with age stratification by consensus, if possible (e.g. early adulthood to age 25, 30 or 34 years); methodology, models and data analytic approaches; development of consensus by 'youth

experts' to provide input on the results and suggest youth-led and other intervention initiatives; study of human-computer-human interaction and privacy issues that inform policy. Whether effectiveness research on social media use can lead to better overall health outcomes and reduced disease burden is still unknown. Analysing large amounts of data will require close collaboration between partners from diverse areas of expertise, such as researchers, providers, statisticians, software developers and engineers. Health systems need to explore competencies for providers to place the person's/patient's needs first and embrace social media technology within healthcare reform, and this will require adjustment of clinical, training, professional development and administrative missions and workflow.

Donald M. Hilty , MD, MBA, Department of Psychiatry and Behavioral Sciences, University of California, Davis, California, USA; and Mental Health, Veterans Affairs Northern California Health Care System, California, USA; **Dorothy Stubbe**, MD, Child Study Center, Yale School of Medicine, Connecticut, USA; **Alastair J. McKean**, MD, Department of Psychiatry and Psychology, Mayo Clinic, Minnesota, USA; **Pamela E. Hoffman**, MD, Department of Psychiatry & Behavioral Science, Yale School of Medicine, Connecticut, USA; **Isheeta Zalpuri**, MD, Department of Psychiatry & Behavioral Science, Stanford University Medical Center, California, USA; **Myo T. Myint**, MD, Department of Psychiatry & Behavioral Science, Tulane University School of Medicine, Louisiana, USA; **Shashank V. Joshi**, MD, Department of Psychiatry & Behavioral Science, Stanford University Medical Center, California, USA; **Murat Pakyurek**, MD, Division of Child and Adolescent Psychiatry, University of California, Davis School of Medicine, California, USA; **Su-Ting T. Li**, MD, MPH, Department of Pediatrics, University of California, Davis School of Medicine, California, USA

Correspondence: Donald M. Hilty. Email: donald.hilty@uhsinc.com

First received 2 Apr 2022, final revision 8 Jun 2023, accepted 9 Jun 2023

Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjo.2023.523>

Data availability

The authors confirm that the data supporting the findings of this study are available within the article and its supplementary materials.

Acknowledgements

The authors thank the American Telemedicine Association and its Telemental Health Interest Group, as well as the Departments of Psychiatry and Behavioral Sciences and Pediatrics at the University of California, Davis School of Medicine, and the Veterans Affairs Northern California Health Care System Mental Health Service. They also thank the Expert Opinion Panel from behavioural health, technology, education and professional development, as well as researchers, authors, editors and editorial board members of journals. The help provided assisted with the recruitment of the expert panel, looked at the data and provided staff support for the authors.

Author contributions

All authors met the three criteria (substantial contribution, drafting, final approval) and agree to be accountable for the work.

Funding

None.

Declaration of interest

None.

References

- Anderson M, Jiang J. *Teens, Social Media and Technology*. Pew Research Center, 2018 (<https://www.pewresearch.org/internet/2018/05/31/teens-social-media-technology-2018/>).
- Madden M, Lenhart A, Cortes S, Gasser URS, Duggan M, Smith A, et al. *Teens, Social Media, and Privacy*. Pew Research Center, 2013 (<https://www.pewresearch.org/internet/2013/05/21/teens-social-media-and-privacy/>).
- Anderson EL, Steen E, Stavropoulos V. Internet use and problematic Internet use: a systematic review of longitudinal research trends in adolescence and emergent adulthood. *Int J Adolesc Youth* 2017; **22**(4): 430–54.
- O'Keeffe GS. Social media: challenges and concerns for families. *Pediatr Clin* 2016; **63**(5): 841–9.
- Federal Trade Commission. *The Children's Online Privacy Protection Rule*. Federal Trade Commission, 2013 (<https://www.ftc.gov/enforcement/rules/rulemaking-regulatory-reform-proceedings/childrens-online-privacy-protection-rule>).
- Domahidi E. The associations between online media use and users' perceived social resources: a meta-analysis. *J Comput Med Commun* 2018; **23**(4): 181–200.
- Hoare E, Milton K, Foster C, Allender S. The associations between sedentary behaviour and mental health among adolescents: a systematic review. *Int J Behav Nutr Phys Activity* 2016; **13**(1): 108.
- Seabrook EM, Kern ML, Rickard NS. Social networking sites, depression, and anxiety: a systematic review. *JMIR Ment Health* 2016; **3**(4): e50.
- Dickson K, Richardson M, Kwan I, MacDowall W, Burchett H, Stansfield C, et al. *Screen-Based Activities and Children and Young People's Mental Health: A Systematic Map of Reviews*. Department of Health Reviews Facility, 2018 (https://eppi.ioe.ac.uk/cms/Portals/0/PDF%20reviews%20and%20summaries/Systematic%20Map%20of%20Reviews%20on%20Screen-based%20activities_08.01.19.pdf?ver=2019-01-29-155200-517).
- Ogders CL, Jensen MR. Adolescent mental health in the digital age: facts, fears, and future directions. *J Child Psychol Psychiatry* 2020; **61**: 336–48.
- Orben A. Teenagers, screens and social media: a narrative review of reviews and key studies. *Soc Psychiatry Psychiatr Epidemiol* 2020; **55**(4): 407–14.
- Twenge JM, Farley E. Not all screen time is created equal: associations with mental health vary by activity and gender. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**(2): 207–17.
- Keles B, McCrae N, Grealish A. A systematic review: the influence of social media on depression, anxiety and psychological distress in adolescents. *Int J Adolesc Youth* 2019; **25**(1): 79–93.
- Kelly Y, Zilanawala A, Booker C, Sacker A. Social media use and adolescent mental health: findings from the UK Millennium Cohort Study. *EClinicalMedicine* 2019; **6**: 59–68.
- Vahedi Z, Zannella L. The association between self-reported depressive symptoms and the use of social networking sites (SNS): a meta-analysis. *Curr Psychol* 2021; **40**: 2174–89.
- Sharma MK, John N, Sahu M. Influence of social media on mental health: a systematic review. *Curr Opin Psychiatry* 2020; **33**(5): 467–75.
- Akkin Gürbüz HG, Albayrak ZS, Kadak MT. Online social network sites usage and impression management of adolescents and relationship with emotional and behavioral problems. *Psychiatry Behav Sci* 2020; **10**(3): 148–54.
- Barthorpe A, Winstone L, Mars B, Moran P. Is social media screen time really associated with poor adolescent mental health? A time use diary study. *J Affect Disord* 2020; **274**: 864–70.
- Coyne SM, Rogers AA, Zurcher JD, Stockdale L, Booth M. Does time spent using social media impact mental health?: an eight year longitudinal study. *Comput Hum Behav* 2020; **104**: 106160.
- Nereim CD, Bickham DS, Rich MO. Social media and adolescent mental health: who you are and what you do matter. *J Adolesc Health* 2020; **66**: 118–9.
- Twenge JM. Why increases in adolescent depression may be linked to the technological environment. *Curr Opin Psychol* 2020; **32**: 89–94.
- Brunborg G, Andreas JB. Increase in time spent on social media is associated with modest increase in depression, conduct problems, and episodic heavy drinking. *J Adolesc* 2019; **74**: 201–9.
- Cole DA, Nick EA, Varga G, Smith D, Zerkowitz RL, Ford MA, et al. Are aspects of twitter use associated with reduced depressive symptoms? The moderating role of in-person social support. *Cyberpsychol Behav Soc Netw* 2019; **22**(11): 692–9.
- Dempsey AE, O'Brien KD, Tiampiy MF, Elhai JD. Fear of missing out (FoMO) and rumination mediate relations between social anxiety and problematic Facebook use. *Addict Behav Rep* 2019; **9**: 100150.
- Jensen M, George M, Russell M, Odgers C. Young adolescents' digital technology use and mental health symptoms: little evidence of longitudinal or daily linkages. *Clin Psychol Sci* 2019; **7**(6): 1416–33.
- Ophir Y, Asterhan CS, Schwarz BB. The digital footprints of adolescent depression, social rejection and victimization of bullying on Facebook. *Comput Hum Behav* 2019; **91**: 62–71.
- Orben A, Dienlin T, Przybylski AK. Social media's enduring effect on adolescent life satisfaction. *Proc Natl Acad Sci USA* 2019; **116**(21): 10226–8.
- Riehm KE, Feder KA, Tormohlen KN, Crum RM, Young AS, Green K, et al. Associations between time spent using social media and internalizing and externalizing problems among US youth. *JAMA Psychiatry* 2019; **76**(12): 1266–73.

- 29 Xie W, Karan K. Predicting Facebook addiction and state anxiety without Facebook by gender, trait anxiety, Facebook intensity, and different Facebook activities. *J Behav Addict* 2019; **8**(1): 79–87.
- 30 Yuen EK, Koterba EA, Stasio MJ, Patrick RB. The effects of Facebook on mood in emerging adults. *Psychol Pop Media Cult* 2019; **8**: 198–206.
- 31 Bayer J, Ellison N, Schoenebeck S, Brady E, Falk EB. Facebook in context(s): measuring emotional responses across time and space. *New Media Soc* 2018; **20**(3): 1047–67.
- 32 Berryman C, Ferguson C, Negy C. Social media use and mental health among young adults. *Psychiatr Q* 2018; **89**: 307–14.
- 33 Houghton S, Lawrence D, Hunter SC, Rosenberg M, Zadow C, Wood L, et al. Reciprocal relationships between trajectories of depressive symptoms and screen media use during adolescence. *J Youth Adolesc* 2018; **47**(11): 2453–67.
- 34 Niu GF, Luo YJ, Sun XJ, Zhou ZK, Yu F, Yang SL, et al. Qzone use and depression among Chinese adolescents: a moderated mediation model. *J Affect Disord* 2018; **231**: 58–62.
- 35 Reinecke L, Meier A, Beutel ME, Schemer C, Stark B, Wölfing K, et al. The relationship between trait procrastination, internet use, and psychological functioning: results from a community sample of German adolescents. *Front Psychol* 2018; **9**: 913.
- 36 Twenge JM, Joiner TE, Rogers ML, Martin GN. Increases in depressive symptoms, suicide-related outcomes, and suicide rates among U.S. adolescents after 2010 and links to increased new media screen time. *Clin Psychol Sci* 2018; **6**: 3–17.
- 37 Akkin Gürbüz HG, Demir T, Gökalp T, Kadak MT, Poyraz BC. Use of social network sites among depressed adolescents. *Behav Inform Technol* 2017; **36**: 517–23.
- 38 Bányai F, Zsila Á, Király O, Maraz A, Elekes Z, Griffiths MD, et al. Problematic social media use: results from a large-scale nationally representative adolescent sample. *PLoS One* 2017; **12**: e0169839.
- 39 Barry CT, Sidoti CL, Briggs SM, Reiter SR, Lindsey RA. Adolescent social media use and mental health from adolescent and parent perspectives. *J Adolesc* 2017; **61**: 1–11.
- 40 Brunborg G, Burdzovic AJ, Kvaavik E. Social media use and episodic heavy drinking among adolescents. *Psychol Rep* 2017; **120**(3): 475–90.
- 41 Calancie O, Ewing L, Narducci LD, Horgan S. Exploring how social networking sites impact youth with anxiety: a qualitative study of Facebook stressors among adolescents with an anxiety disorder diagnosis. *Cyberpsychol* 2017; **11**(4): 2.
- 42 Frison E, Eggermont S. Browsing, posting, and liking on Instagram: the reciprocal relationships between different types of Instagram use and adolescents' depressed mood. *Cyberpsychol Behav Soc Netw* 2017; **20**(10): 603–9.
- 43 Kokkinos CM, Sariapanidis I. A lifestyle exposure perspective of victimization through Facebook among university students. Do individual differences matter? *Comput Hum Behav* 2017; **74**: 235–45.
- 44 Lee WJ, Ho SS, Lwin HMO. Extending the social cognitive model—Examining the external and personal antecedents of social network sites use among Singaporean adolescents. *Comput Hum Behav* 2017; **67**: 240–51.
- 45 Oberst U, Wegmann E, Stodt B, Brand M, Chamarro A. Negative consequences from heavy social networking in adolescents: the mediating role of fear of missing out. *J Adolesc* 2017; **55**: 51–60.
- 46 Radovic A, Gmelin T, Stein BD, Miller E. Depressed adolescents' positive and negative use of social media. *J Adolesc* 2017; **55**: 5–15.
- 47 Vernon L, Modecki KL, Barber BL. Tracking effects of problematic social networking on adolescent psychopathology: the mediating role of sleep disruptions. *J Clin Child Adolesc Psychol* 2017; **46**(2): 269–83.
- 48 Frison E, Eggermont S. Gender and Facebook motives as predictors of specific types of Facebook use: a latent growth curve analysis in adolescence. *J Adolesc* 2016; **52**: 182–90.
- 49 Frison E, Subrahmanyam K, Eggermont S. The short-term longitudinal and reciprocal relations between peer victimization on Facebook and adolescents' well-being. *J Youth Adolesc* 2016; **45**(9): 1755–71.
- 50 Morin-Major JK, Marin MF, Durand N, Wan N, Juster RP, Lupien SJ. Facebook behaviors associated with diurnal cortisol in adolescents: is befriending stressful? *Psychoneuroendocrinology* 2016; **63**: 238–46.
- 51 Park J, Lee DS, Shablack H, Verduyn P, Deldin P, Ybarra O, et al. When perceptions defy reality: the relationships between depression and actual and perceived Facebook social support. *J Affect Disord* 2016; **200**: 37–44.
- 52 Banjanin N, Banjanin N, Dimitrijevic I, Pantic I. Relationship between internet use and depression: focus on physiological mood oscillations, social networking and online addictive behavior. *Comput Hum Behav* 2015; **43**: 308–12.
- 53 Coppersmith G, Dredze M, Harman C, Hollingshead K. From ADHD to SAD: analyzing the language of mental health on Twitter through self-reported diagnoses. *Second Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality (North American Chapter of the Association for Computational Linguistics, 5 June 2015)*. Association for Computational Linguistics, 2015.
- 54 Lee-Won RJ, Herzog L, Park SG. Hooked on Facebook: the role of social anxiety and need for social assurance in problematic use of Facebook. *Cyberpsychol Behav Soc Netw* 2015; **18**(10): 567–74.
- 55 Lup K, Trub L, Rosenthal L. Instagram #instasad?: exploring associations among Instagram use, depressive symptoms, negative social comparison, and strangers followed. *Cyberpsychol Behav Soc Netw* 2015; **18**(5): 247–52.
- 56 McCloskey W, Iwanicki S, Lauterbach D, Giammittorio DM, Maxwell K. Are Facebook friends helpful? Development of a Facebook-based measure of social support and examination of relationships among depression, quality of life, and social support. *Cyberpsychol Behav Soc Netw* 2015; **18**(9): 499–505.
- 57 Mitchell M, Hollingshead K, Coppersmith G. Quantifying the language of schizophrenia in social media. *Second Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality (North American Chapter of the Association for Computational Linguistics, 5 June 2015)*. Association for Computational Linguistics, 2015.
- 58 Moberg FB, Anestis MD. A preliminary examination of the relationship between social networking interactions, internet use, and thwarted belongingness. *Crisis* 2015; **36**(3): 187–93.
- 59 Nesi J, Prinstein MJ. Using social media for social comparison and feedback-seeking: gender and popularity moderate associations with depressive symptoms. *J Abnorm Child Psychol* 2015; **43**: 1427–38.
- 60 Park S, Kim I, Lee SW, Yoo J, Jeong B, Cha M. Manifestation of depression and loneliness on social networks: a case study of young adults on Facebook. *18th ACM Conference on Computer Supported Cooperative Work & Social Computing (Vancouver, Canada; 14–18 March 2015)*. Association for Computing Machinery, 2015.
- 61 Preotiu-Pietro D, Eichstaedt J, Park G. The role of personality, age, and gender in tweeting about mental illness. *Second Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality (North American Chapter of the Association for Computational Linguistics, 5 June 2015)*. Association for Computational Linguistics, 2015.
- 62 Shaw AM, Timpano KR, Tran TB, Joormann J. Correlates of Facebook usage patterns: the relationship between passive Facebook use, social anxiety symptoms, and brooding. *Comput Hum Behav* 2015; **48**: 575–80.
- 63 Tandoc EC Jr., Ferrucci P, Duffy M. Facebook use, envy, and depression among college students: is Facebooking depressing? *Comput Hum Behav* 2015; **43**: 139–46.
- 64 Davidson TC, Farquhar LK. Correlates of social anxiety, religion, and Facebook. *J Media Religion* 2014; **13**: 208–25.
- 65 De Choudhury M, Counts S, Horvitz E, Hoff A. Characterizing and predicting postpartum depression from shared Facebook data. *17th ACM Conference on Computer Supported Cooperative Work & Social Computing (Baltimore, USA; 15–19 February 2015)*. Association for Computing Machinery, 2014.
- 66 Farquhar LK, Davidson TC. Facebook frets: the role of social media use in predicting social and Facebook-specific anxiety. *J Alab Acad Sci* 2014; **85**(1): 8–24.
- 67 Gámez-Gaudix M. Depressive symptoms and problematic use among adolescents: analysis of the longitudinal relationships from the cognitive-behavioral model. *Cyberpsychol Behav Soc Netw* 2014; **17**: 714–9.
- 68 Király O, Griffiths M, Urbán R, Farkas J, Kökönyei G, Elekes Z, et al. Problematic internet use and problematic online gaming are not the same: findings from a large nationally representative adolescent sample. *Cyberpsychol Behav Soc Netw* 2014; **17**: 749–54.
- 69 Labrague L. Facebook use and adolescents' emotional states of depression, anxiety, and stress. *Health Sci J* 2014; **8**(1): 80–9.
- 70 Lee S. How do people compare themselves with others on social network sites?: the case of Facebook. *Comput Hum Behav* 2014; **32**: 253–60.
- 71 Neira CJB, Barber BL. Social networking site use: linked to adolescents' social self-concept, self-esteem, and depressed mood. *Austral J Psychol* 2014; **66**(1): 56–64.
- 72 Schwartz HA, Eichstaedt J, Kern ML, Park G. Towards assessing changes in degree of depression through Facebook. *Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality (North American Chapter of the Association for Computational Linguistics, 27 June 2014)*. Association for Computational Linguistics, 2014.
- 73 Simoncic TE, Kuhlman KR, Vargas I, Houchins S, Lopez-Duran NL. Facebook use and depressive symptomatology: investigating the role of neuroticism and extraversion in youth. *Comput Hum Behav* 2014; **40**: 1–5.
- 74 Steers MN, Wickham RE, Acitelli LK. Seeing everyone else's highlight reels: how Facebook usage is linked to depressive symptoms. *J Soc Clin Psychol* 2014; **33**(8): 701–31.
- 75 Tsitsika AK, Tzavela EC, Janikian M, Ólafsson K, Iordache A, Schoenmakers TM, et al. Online social networking in adolescence: patterns of use in six European

- countries and links with psychosocial functioning. *J Adolesc Health* 2014; **55**(1): 141–7.
- 76 De Choudhury M, Counts S, Horvitz E. Predicting postpartum changes in emotion and behavior via social media. *CHI '13: CHI Conference on Human Factors in Computing Systems (Special Interest Group on Computer-Human Interaction, 27 April – 2 May 2013)*. Association for Computing Machinery, 2013.
- 77 Feinstein BA, Hershenberg R, Bhatia V, Latack J. Negative social comparison on Facebook and depressive symptoms: rumination as a mechanism. *Psychol Pop Media Cult* 2013; **2**(3): 161–70.
- 78 Jelenchick LA, Eickhoff JC, Moreno MA. Facebook depression? social networking site use and depression in older adolescents. *J Adolesc Health* 2013; **52**(1): 128–30.
- 79 Koc M, Gulyagci S. Facebook addiction among Turkish college students: the role of psychological health, demographic, and usage characteristics. *Cyberpsychol Behav Soc Netw* 2013; **16**(4): 279–84.
- 80 Kross E, Verduyn P, Demiralp E, Park J, Lee DS, Lin N, et al. Facebook use predicts declines in subjective well-being in young adults. *PLoS One* 2013; **8**(8): e69841.
- 81 Landoll RR, La Greca AM, Lai BS. Aversive peer experiences on social networking sites: development of the social networking-peer experiences questionnaire (SN-PEQ). *J Res Adolesc* 2013; **23**(4): 10.1111/jora.12022.
- 82 Park S, Lee SW, Kwak J, Cha M, Jeong B. Activities on Facebook reveal the depressive state of users. *J Med Internet Res* 2013; **15**(10): e217.
- 83 Wright EJ, White KM, Obst PL. Facebook false self-presentation behaviors and negative mental health. *Cyberpsychol Behav Soc Netw* 2018; **21**(1): 40–9.
- 84 Dumitrache SD, Mitrofan L, Petrov Z. Self-image and depressive tendencies among adolescent Facebook users. *Rev Psychol* 2012; **58**: 285–95.
- 85 Locatelli S, Kluge K, Bryant F. Facebook use and the tendency to ruminate among college students: testing mediational hypotheses. *J Educ Comput Res* 2012; **46**: 377–94.
- 86 Pantic I, Damjanovic A, Todorovic J, Topalovic D, Bojovic-Jovic D, Ristic S, et al. Association between online social networking and depression in high school students: behavioral physiology viewpoint. *Psychiatr Danub* 2012; **24**(1): 90–3.
- 87 Selfhout M, Branje SJT, Delsing M, ter Bogt TF, Meeus WH. Different types of internet use, depression, and social anxiety: the role of perceived friendship quality. *J Adolesc* 2009; **32**: 819–33.
- 88 Hwang JM, Cheong PH, Feeley TH. Being young and feeling blue in Taiwan: examining adolescent depressive mood and online and offline activities. *New Media Soc* 2009; **11**(7): 1101–21.
- 89 van den Eijnden RJJM, Meerkerk GJ, Vermulst AA, Spijkerman R, Engels RC. Online communication, compulsive internet use, and psychosocial well-being among adolescents: a longitudinal study. *Devel Psychol* 2008; **44**(3): 655–65.
- 90 Ybarra ML, Alexander C, Mitchell KJ. Depressive symptomology, youth internet use, and online interactions: a national survey. *J Adolesc Health* 2005; **36**: 9–18.
- 91 Khasawneh A, Chalil Madathil K, Dixon E, Wiśniewski P, Zinzow H, Roth R. Examining the self-harm and suicide contagion effects of the Blue Whale challenge on YouTube and Twitter: qualitative study. *JMIR Ment Health* 2020; **7**(6): e15973.
- 92 Mori K, Haruno M. Differential ability of network and natural language information on social media to predict interpersonal and mental health traits. *J Pers* 2020; **89**(2): 228–43.
- 93 Sindermann C, Elhai JD, Montag C. Predicting tendencies towards the disordered use of Facebook's social media platforms: on the role of personality, impulsivity, and social anxiety. *Psychiatry Res* 2020; **285**: 112793.
- 94 Corbitt-Hall DJ, Gauthier JM, Troop-Gordon W. Suicidality disclosed online: using a simulated Facebook task to identify predictors of support giving to friends at risk of self-harm. *Suicide Life Threat Behav* 2019; **49**(2): 598–613.
- 95 Kircaburun K, Kokkinos CM, Demetrovics Z, Király O, Griffiths Ko, Çolak TS. Problematic online behaviors among adolescents and emerging adults: associations between cyberbullying perpetration, problematic social media use, and psychosocial factors. *Int J Mental Health Addiction* 2019; **17**(4): 891–908.
- 96 Maheu M, Drude K, Hertlein K, Hilty DM. An interdisciplinary framework for telebehavioral health competencies. *J Tech Behav Sci* 2018; **3**(2): 108–40. correction 3(2): 107.
- 97 Escobar-Viera CG, Whitfield DL, Wessel CB, Shensa A, Sidani JE, Brown AL, et al. For better or for worse? A systematic review of the evidence on social media use and depression among lesbian, gay, and bisexual minorities. *JMIR Ment Health* 2018; **5**(3): e10496.
- 98 O'Reilly M, Dogra N, Hughes J, Reilly P, George R, Whiteman N. Potential of social media in promoting mental health in adolescents. *Health Promot Int* 2018; **34**: 981–91.
- 99 O'Reilly M, Dogra N, Whiteman N, Hughes J, Eruyer S, Reilly P. Is social media bad for mental health and wellbeing? Exploring the perspectives of adolescents. *Clin Child Psychol Psychiatry* 2018; **23**: 601–13.
- 100 Pourmand A, Roberson J, Caggiula A, Monsalve N, Rahimi M, Torres-Llenza V. Social media and suicide: a review of technology-based epidemiology and risk assessment. *Telemed J E Health* 2019; **25**(10): 880–8.
- 101 Wang P, Wang X, Wu Y, Biao L. Social networking sites addiction and adolescent depression: a moderated mediation model of rumination and self-esteem. *Pers Individ Differ* 2018; **127**: 162–7.
- 102 Chen H. Antecedents of positive self-disclosure online: an empirical study of US college students' Facebook usage. *Psychol Res Behav Manag* 2017; **10**: 147–53.
- 103 O'Dea B, Larsen ME, Batterham PJ, Caleare AL, Christensen H. A linguistic analysis of suicide-related Twitter posts. *Crisis* 2017; **38**(5): 319–29.
- 104 Salmela-Aro K, Upadaya K, Hakkarainen K, Lonka K, Alho K. The dark side of internet use: two longitudinal studies of excessive internet use, depressive symptoms, school burnout and engagement among Finnish early and late adolescents. *J Youth Adolesc* 2017; **46**(2): 343–57.
- 105 Van Rooij AJ, Ferguson CJ, Van de Mheen D, Schoenmakers TM. Time to abandon internet addiction? Predicting problematic Internet, game, and social media use from psychosocial well-being and application use. *Clin Neuropsychiatry* 2017; **14**(1): 113–21.
- 106 Yan H, Zhang R, Oniffrey TM, Chen G, Wang Y, Wu Y, et al. Associations among screen time and unhealthy behaviors, academic performance, and well-being in Chinese adolescents. *Int J Environ Res Public Health* 2017; **14**(6): 596.
- 107 Braithwaite SR, Giraud-Carrier C, West J, Barnes MD, Hanson CL. Validating machine learning algorithms for Twitter data against established measures of suicidality. *JMIR Ment Health* 2016; **3**(2): e21.
- 108 Cole DA, Zerkowicz RL, Nick E, Martin NC, Roeder KM, Sinclair-McBride K, et al. Longitudinal and incremental relation of cybervictimization to negative self-cognitions and depressive symptoms in young adolescents. *J Abnorm Child Psychol* 2016; **44**(7): 1321–32.
- 109 Coppersmith G, Ngo K, Leary R, Wood A. Exploratory analysis of social media prior to a suicide attempt. *3rd Workshop on Computational Linguistics and Clinical Psychology: From Linguistic Signal to Clinical Reality (North American Chapter of the Association for Computational Linguistics, 16 June 2016)*. Association for Computational Linguistics, 2016.
- 110 Burnap P, Colombo W, Scourfield J. Machine classification and analysis of suicide-related communication on Twitter. 26th ACM Conference on Hypertext & Social Media (Middle East Technical University Northern Cyprus Campus, 1–4 September 2015). Association for Computing Machinery, 2015.
- 111 Muench F, Hayes M, Kuerbis A, Shao S. The independent relationship between trouble controlling Facebook use, time spent on the site and distress. *J Behav Addict* 2015; **4**(3): 163–9.
- 112 O'Dea B, Wan S, Batterham PJ, Caleare AL, Paris C, Christensen H. Detecting suicidality on Twitter. *Internet Interv* 2015; **2**(2): 183–8.
- 113 Sampasa-Kanyinga H, Hamilton HA. Use of social networking sites and risk of cyberbullying victimization: a population-level study of adolescents. *Cyberpsychol Behav Soc Netw* 2015; **18**(12): 704–10.
- 114 Sampasa-Kanyinga H, Lewis RF. Frequent use of social networking sites is associated with poor psychological functioning among children and adolescents. *Cyberpsychol Behav Soc Netw* 2015; **18**(7): 380–5.
- 115 Zhang L, Huang X, Liu T, Li A, Chen Z, Zhu T. Using linguistic features to estimate suicide probability of Chinese microblog users. In *Human Centered Computing. HCC 2014. Lecture Notes in Computer Science. Volume 8944* (eds Q Zu, B Hu, N Gu, S Seng): 549–59. Springer, 2015.
- 116 La Sala L, Skues J, Grant S. Personality traits and Facebook use: the combined/interactive effect of extraversion, neuroticism and conscientiousness. *Soc Netw* 2014; **3**(5): 211–9.
- 117 Rodríguez Puentes AP, Fernández Parra A. Relationship between the time spent on internet social networking and mental health in Colombian adolescents. *Act Colom Psicol* 2014; **17**(1): 131–40.
- 118 Tseng F-Y, Yang H-J. Internet use and web communication networks, sources of social support, and forms of suicidal and nonsuicidal self-injury among adolescents: different patterns between genders. *Suic Life Threat Behav* 2015; **45**(2): 178–91.
- 119 Jashinsky J, Burton SH, Hanson CL, West J, Giraud-Carrier J, Barnes MD, et al. Tracking suicide risk factors through Twitter in the US. *Crisis* 2014; **35**(1): 51–9.
- 120 Romer D, Bagdasarov Z, More E. Older versus newer media and the well-being of United States youth: results from a national longitudinal panel. *J Adol Health* 2013; **52**(5): 613–9.

- 121 Dunlop SM, More E, Romer D. Where do youth learn about suicides on the internet, and what influence does this have on suicidal ideation? *J Child Psychol Psychiatry* 2011; **52**(10): 1073–80.
- 122 Szwedo DE, Mikami AY, Allen JP. Qualities of peer relations on social networking websites: predictions from negative mother-teen interactions. *J Res Adolesc* 2011; **21**(3): 595–607.
- 123 Mitrofan O, Paul M, Spencer N. Is aggression in children with behavioural and emotional difficulties associated with television viewing and video game playing? A systematic review. *Child Care Health Dev* 2009; **35**(1): 5–15.
- 124 Alcott H, Braghieri L, Eichmeyer S, Gentzkow M. *The Welfare Effects of Social Media. Working Paper 25514*. National Bureau Of Economic Research, 2019 (<http://www.nber.org/papers/w25514>).
- 125 Orben A, Przybylski AK. Screens, teens, and psychological well-being: evidence from three time-use-diary studies. *Psychol Sci* 2019; **30**(5): 682–96.
- 126 Orben A, Przybylski AK. The association between adolescent well-being and digital technology use. *Nat Hum Behav* 2019; **3**(2): 173–82.
- 127 Pope ZC, Barr-Anderson DJ, Lewis BA, Pereira MA, Gao Z. Use of wearable technology and social media to improve physical activity and dietary behaviors among college students: a 12-week randomized pilot study. *Int J Environ Res Public Health* 2019; **16**(19): 3579.
- 128 Rasmussen E, Punyanunt-Carter N, LaFreniere J, Norman MS. The serially mediated relationship between emerging adults' social media use and mental well-being. *Comput Hum Behav* 2020; **102**: 206–13.
- 129 Coyne SM, Padilla-Walker LM, Holmgren HG. A six-year longitudinal study of texting trajectories during adolescence. *Child Dev* 2018; **89**(1): 58–65.
- 130 Hunt MG, Marx R, Lison C, Young J. No more FOMO: limiting social media decreases loneliness and depression. *J Soc Clin Psychol* 2018; **37**(10): 751–68.
- 131 Vanman EJ, Baker R, Tobin SJ. The burden of online friends: the effects of giving up Facebook on stress and well-being. *J Soc Psychol* 2018; **158**(4): 496–508.
- 132 Mun IB, Kim H. Influence of false self-presentation on mental health and deleting behavior on Instagram: the mediating role of perceived popularity. *Front Psychol* 2021; **12**: 660484.
- 133 Frith E, Loprinzi P. Can Facebook reduce perceived anxiety among college students? randomized controlled exercise trial using the transtheoretical model of behavior change. *JMIR Ment Health* 2017; **4**(4): e50.
- 134 Sun WH, Wong CKH, Wong WCW. A peer-led, social media-delivered, safer sex intervention for Chinese college students: randomized controlled trial. *JMIR* 2017; **19**(8): e284.
- 135 Ehenreich SE, Underwood MK. Adolescents' internalizing symptoms as predictors of the content of their Facebook communication and responses received from peers. *Transl Issues Psychol Sci* 2016; **2**(3): 227–37.
- 136 Marder B, Joinson A, Shankar A, Thirlway K. Strength matters: self-presentation to the strongest audience rather than lowest common denominator when faced with multiple audiences in social network sites. *Comput Hum Behav* 2016; **61**: 56–62.
- 137 Gunnell KE, Flament MF, Buchholz A, Henderson KA, Obeid N, Schubert N, et al. Examining the bidirectional relationship between physical activity, screen time, and symptoms of anxiety and depression over time during adolescence. *Prev Med* 2016; **88**: 147–52.
- 138 Lin JH. The role of attachment style in Facebook use and social capital: evidence from university students and a national sample. *Cyberpsychol Behav Soc Netw* 2015; **18**(3): 173–80.
- 139 Neubaum G, Kramer NC. My friends right next to me: a laboratory investigation on predictors and consequences of experiencing social closeness on social networking sites. *Cyberpsychology Behav Soc Netw* 2015; **18**: 443–9.
- 140 Rae JR, Lonborg SD. Do motivations for using Facebook moderate the association between Facebook use and psychological well-being? *Front Psychol* 2015; **6**: 771.
- 141 Verduyn P, Lee DS, Park J, Shablack H, Orvell A, Bayer J, et al. Passive Facebook usage undermines affective well-being: experimental and longitudinal evidence. *J Exp Psychol Gen* 2015; **144**(2): 480–8.
- 142 Blomfield-Neira CJ, Barber BL. Social networking site use: linked to adolescents' social self-concept, self-esteem, and depressed mood. *Austral J Psychol* 2014; **66**(1): 56–64.
- 143 Coyne SM, Padilla-Walker LM, Harper J, Stockdale L A friend request from dear old dad: associations between parent-child social networking and adolescent outcomes. *Cyberpsychol Behav Soc Netw* 2014; **17**(1): 8–13.
- 144 Farquhar LK, Davidson TC. Tolerance on Facebook: exploring network diversity and social distance. *J New Media and Culture* 2015; **10**(1). Available from: <http://ibiblio.org/nmediac/summer2015/facebook.html>.
- 145 Vogel EA, Rose JP, Roberts LR, Eckles K. Social comparison, social media, and self-esteem. *Psychol Pop Media Cult* 2014; **3**: 206–22.
- 146 Tsugawa S, Mogi Y, Kikuchi Y, Kishino F. *On estimating depressive tendencies of Twitter users utilizing their tweet data. 2013 IEEE Virtual Reality Conference (Florida, USA, 16–23 March 2013)*. Institute of Electrical and Electronics Engineers, 2013.
- 147 Burke M, Kraut R, Marlow C. Social capital on Facebook: differentiating uses and users. *CHI '11: CHI Conference on Human Factors in Computing Systems (Special Interest Group on Computer-Human Interaction, 7–12 May 2011)*. Association for Computing Machinery, 2011.
- 148 Burke M, Marlow C, Lento T. Social network activity and social well-being. *CHI '10: CHI Conference on Human Factors in Computing Systems (Special Interest Group on Computer-Human Interaction, 10–15 April 2010)*. Association for Computing Machinery, 2010.
- 149 Ellison NB, Steinfield C, Lampe C. The benefits of Facebook friends: social capital and college students' use of online social network sites. *J Mediat Commun* 2007; **12**(4): 1143–68.
- 150 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Meth* 2005; **8**(1): 19–32.
- 151 Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci* 2010; **20**(5): 69.
- 152 Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018; **169**(7): 467–73.
- 153 Macaulay PJR, Boulton MJ, Betts LR. Comparing early adolescents' positive bystander responses to cyberbullying and traditional bullying: the impact of severity and gender. *J Tech Behav Sci* 2019; **4**: 253–61.
- 154 Crowe M, Inder M, Porter R. Conducting qualitative research in mental health: thematic and content analyses. *Aust N Z J Psychiatry* 2015; **49**(7): 616–23.
- 155 Strasburger VC, Zimmerman H, Temple JR, Madigan S. Teenagers, sexting, and the law. *Pediatrics* 2019; **143**(5): e20183183.
- 156 Joshi SV, Stubbe D, Li ST, Hilty DM. The use of technology by youth: implications for psychiatric educators. *Acad Psychiatry* 2019; **43**(1): 101–9.
- 157 Hilty DM, Torous J, Parish M, Chan SR, Xiong G, Scher L, et al. A literature review comparing clinicians' approaches and skills to in-person, synchronous and asynchronous care: moving toward asynchronous competencies to ensure quality care. *Telemed J E Health* 2021; **27**(4): 356–73.
- 158 American Academy of Pediatrics. *Family Media Use Plan*. American Academy of Pediatrics, 2022 (<https://www.healthychildren.org/English/media/Pages/default.aspx>).
- 159 Zalpuri I, Liu H, Stubbe D, Hadsu J, Hilty DM. Social media and networking competencies for psychiatric education: skills, teaching methods, and implications. *Acad Psychiatry* 2018; **42**(6): 808–17.
- 160 Hilty DM, Crawford A, Teshima J, Nasatir-Hilty SE, Luo J, Chisler LSM, et al. Mobile health and cultural competencies as a foundation for telehealth care: scoping review. *J Technol Behav Sci* 2021; **6**: 197–230.
- 161 Armstrong CM, McGee-Vincent P, Juhasz K, Owen J, Avery T, Jaworski B, et al. *VA Mobile Health Practice Guide* (1st edn). US Department of Veterans Affairs, 2021 (https://www.researchgate.net/publication/351563585_VA_Mobile_Health_Practice_Guide).
- 162 Chancellor S, De Choudhury M. Methods in predictive techniques for mental health status on social media: a critical review. *NPJ Digit Med* 2020; **3**: 43.
- 163 McIntyre RS, Cha DS, Jerrell JM, Swardfager W, Kim RD, Costa LG, et al. Advancing biomarker research: utilizing 'Big Data' approaches for the characterization and prevention of bipolar disorder. *Bipolar Disord* 2014; **16**(5): 531–47.
- 164 Garcia-Ceja E, Riegler M, Nordgreen T, Jakobsen P, Oedegaard KJ, Tørresen J, et al. Mental health monitoring with multimodal sensing and machine learning: a survey. *Pervas Mob Comput* 2018; **51**: 1–26.
- 165 Hilty DM, Armstrong CM, Luxton DD, Gentry MT, Luxton DD, Krupinski EA. Sensor, wearable and remote patient monitoring competencies for clinical care and training: scoping review. *J Tech Behav Sci* 2021; **6**(2): 252–77.
- 166 US Department of Health and Human Services. United States Stop Bullying (<https://www.stopbullying.gov>).
- 167 Hilty DM, Unutzer J, Ko DK, Luo J, Worley LLM, Yager J. Approaches for departments, schools and health systems to better implement technologies used for clinical care and education. *Acad Psychiatry* 2019; **43**(6): 611–6.
- 168 Vidal C, Lhaksampa T, Miller L, Platt R. Social media use and depression in adolescents: a scoping review. *Int Rev Psychiatry* 2020; **32**(3): 235–53.

