

# Brain and Spinal Cord Tumors in Children Causes, Risk Factors, and Prevention

Learn about the risk factors for brain and spinal cord tumors in children.

## Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for brain and spinal cord tumors in children.

- [Risk Factors for Brain and Spinal Cord Tumors in Children](#)
- [What Causes Brain and Spinal Cord Tumors in Children?](#)

## Prevention

Other than exposure to radiation, there are no known lifestyle-related or environmental causes of risk factors for brain and spinal cord tumors in children, so at this time there is no way to protect against most of these cancers.

- [Can Brain and Spinal Cord Tumors in Children Be Prevented?](#)
- 

# Risk Factors for Brain and Spinal Cord

# Tumors in Children

- [Radiation exposure](#)
- [Inherited and genetic conditions](#)
- [Factors with uncertain, controversial, or unproven effects on brain tumor risk](#)

A risk factor is anything that affects a person's chance of getting a disease such as a brain or spinal cord tumor. Different types of cancer have different risk factors.

Lifestyle-related risk factors such as diet, body weight, physical activity, and tobacco use play a major role in many adult cancers. But these factors usually take many years to influence cancer risk, and they are not thought to play much of a role in childhood cancers, including brain tumors.

Very few risk factors have been found for brain and spinal cord tumors. There is no clear [cause](#) for most of these tumors.

## Radiation exposure

The only well-established environmental risk factor for brain tumors is radiation exposure to the head, which most often comes from the treatment of other conditions.

For example, before the risks of radiation were well known (more than 50 years ago), children with ringworm of the scalp (a fungal infection) often received low-dose radiation therapy. This was later found to increase their risk of some types of brain tumors as they got older.

Today, most radiation-induced brain tumors are caused by radiation given to the head to treat other cancers, such as leukemia. These brain tumors usually develop around 10 to 15 years after getting radiation therapy.

Radiation-induced tumors are still fairly rare, but because of the increased risk (as well as the other possible side effects), radiation therapy is only given to the head after carefully weighing the possible benefits and risks. For most patients with cancer in or near the brain, the benefits of getting radiation therapy as part of their treatment far outweigh the small risk of developing a brain tumor years later.

The possible risk from fetal or childhood exposure to imaging tests that use radiation, such as x-rays or CT scans, is not known for sure. These tests use much lower levels of radiation than those used in radiation treatments, so if there is any increase in risk, it is

likely to be very small. But to be safe, most doctors recommend that pregnant women and children not get these tests unless they are absolutely needed.

## Inherited and genetic conditions

Rarely, children have inherited abnormal **genes** from a parent that put them at increased risk for certain types of brain tumors. In other cases, these abnormal genes are not inherited but occur as a result of changes (mutations) in the gene before birth.

People with inherited tumor syndromes often have many tumors that start when they are young. Some of the better known syndromes include:

### Neurofibromatosis type 1 (von Recklinghausen disease)

This is the most common syndrome linked to brain or spinal cord tumors. It is often inherited from a parent, but it can also start in some children whose parents don't have it. Children with this syndrome may have optic **gliomas** or other gliomas of the brain or spinal cord, or **neurofibromas** (benign tumors of peripheral nerves). Changes in the *NF1* gene cause this disorder.

### Neurofibromatosis type 2

This condition is less common than von Recklinghausen disease. It can also either be inherited or may start in children without a family history. It is associated with cranial or spinal nerve **schwannomas**, especially vestibular schwannomas (acoustic neuromas), which almost always occur on both sides of the head. It is also linked to an increased risk of **meningiomas**, as well as spinal cord **gliomas** or **ependymomas**. Changes in the *NF2* gene are nearly always responsible for neurofibromatosis type 2.

### Tuberous sclerosis

Children with this condition may develop **subependymal giant cell astrocytomas (SEGAs)**, as well as other **benign tumors** of the brain, skin, heart, kidneys, or other organs. This condition is caused by changes in either the *TSC1* or the *TSC2* gene.

### Von Hippel-Lindau disease

Children with this disease tend to develop **blood vessel tumors (hemangioblastomas)** of the cerebellum, spinal cord, or retina, as well as tumors in the kidney, pancreas, and some other parts of the body. It is caused by changes in the *VHL*

gene.

### **Li-Fraumeni syndrome**

People with this syndrome have an increased risk of **gliomas**, as well as breast cancer, soft tissue sarcomas, leukemia, and some other types of cancer. It is caused by changes in the *TP53* gene.

### **Other syndromes**

Other inherited conditions linked with increased risks of certain types of brain and spinal cord tumors include:

- Gorlin syndrome (basal cell nevus syndrome)
- Turcot syndrome
- Cowden syndrome
- Hereditary retinoblastoma
- Rubinstein-Taybi syndrome

Some families may have genetic disorders that are not well recognized or that could even be unique to a particular family.

## **Factors with uncertain, controversial, or unproven effects on brain tumor risk**

### **Cell phone use**

Cell phones give off radiofrequency (RF) rays, a form of electromagnetic energy on the spectrum between FM radio waves and those used in microwave ovens, radar, and satellite stations. Cell phones do not give off ionizing radiation, the type that can cause cancer by damaging the DNA inside cells. Still, there have been concerns that the phones, whose antennae are built-in and therefore are placed close to the head when being used, might somehow raise the risk of brain tumors.

Some studies have suggested a possible increased risk of brain tumors or of vestibular schwannomas (acoustic neuromas) in adults with cell phone use, but most of the larger studies done so far have not found an increased risk, either overall or among specific types of tumors. Still, there are very few studies of long-term use (10 years or more), and cell phones haven't been around long enough to determine the possible risks of

lifetime use. The same is true of any possible higher risks in children, who are increasingly using cell phones. Cell phone technology also continues to change, and it's not clear how this might affect any risk.

These risks are being studied, but it will likely be many years before firm conclusions can be made. In the meantime, for people concerned about the possible risks, there are ways to lower their (and their children's) exposure, such as using the phone's speaker or an earpiece to move the phone itself away from the head when used. For more information, see [Cellular Phones](#)<sup>1</sup>.

## Other factors

Exposure to [aspartame](#)<sup>2</sup> (a sugar substitute), exposure to [electromagnetic fields](#)<sup>3</sup> from power lines and other sources, and infection with certain viruses have been suggested as possible risk factors, but most researchers agree that there is no convincing evidence to link these factors to brain tumors. Research on these and other potential risk factors continues.

## Hyperlinks

1. [www.cancer.org/cancer/risk-prevention/radiation-exposure/cellular-phones.html](http://www.cancer.org/cancer/risk-prevention/radiation-exposure/cellular-phones.html)
2. [www.cancer.org/cancer/risk-prevention/chemicals/aspartame.html](http://www.cancer.org/cancer/risk-prevention/chemicals/aspartame.html)
3. [www.cancer.org/cancer/risk-prevention/radiation-exposure/extremely-low-frequency-radiation.html](http://www.cancer.org/cancer/risk-prevention/radiation-exposure/extremely-low-frequency-radiation.html)

## References

Dorsey JF, Hollander AB, Alonso-Basanta M, et al. Chapter 66: Cancer of the Central Nervous System. In: Abeloff MD, Armitage JO, Niederhuber JE, Kastan MB, McKenna WG, eds. *Abeloff's Clinical Oncology*. 5th ed. Philadelphia, Pa: Elsevier; 2014.

Lau C, Teo WY. Epidemiology of central nervous system tumors in children. UpToDate. 2018. Accessed at [www.uptodate.com/contents/epidemiology-of-central-nervous-system-tumors-in-children](http://www.uptodate.com/contents/epidemiology-of-central-nervous-system-tumors-in-children) on April 23, 2018.

Williams D, Parsons IF, Pollack DA. Chapter 26A: Gliomas, Ependymomas, and Other Nonembryonal Tumors of the Central Nervous System. In: Pizzo PA, Poplack DG, eds. *Principles and Practice of Pediatric Oncology*. 7th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2016.

Last Revised: June 20, 2018

# What Causes Brain and Spinal Cord Tumors in Children?

- [Inherited gene changes](#)
- [Acquired gene changes](#)

The cause of most brain and spinal cord tumors is not fully understood, and there are very few known [risk factors for these tumors](#). But researchers have found some of the changes that occur in normal brain cells that may lead them to form tumors.

Normal human cells grow and function based mainly on the information contained in each cell's DNA. Brain and spinal cord tumors, like other tumors, are usually caused by changes (mutations) in the DNA inside cells. DNA is the chemical that makes up our genes, which control how our cells function. We usually look like our parents because they are the source of our DNA. But DNA affects more than how we look.

Some genes control when our cells grow, divide into new cells, and die:

- Certain genes that help cells grow, divide, and stay alive are called **oncogenes**.
- Genes that help keep cell division under control, or cause cells to die at the right time, are called **tumor suppressor genes**.

Cancers can be caused by DNA changes that turn on oncogenes or turn off tumor suppressor genes. These gene changes can be inherited from a parent (as is sometimes the case with childhood cancers), but more often they are acquired during a person's lifetime.

## Inherited gene changes

Researchers have found the gene changes that cause some rare [inherited syndromes](#) (like neurofibromatosis, tuberous sclerosis, Li-Fraumeni syndrome, and von Hippel-Lindau disease) and increase the risk of developing some brain and spinal cord tumors. For example, the Li-Fraumeni syndrome is caused by changes in the *TP53* tumor suppressor gene. Normally, this gene prevents cells with damaged DNA from growing. Changes in this gene increase the risk of developing brain tumors (particularly gliomas), as well as some other cancers.

## Acquired gene changes

Most often, it's not known why children *without* inherited syndromes develop brain or spinal cord tumors. Most exposures that cause cancer, such as tobacco smoke, somehow damage DNA. But the brain is relatively protected from most cancer-causing chemicals that we might breathe in or eat. What's more, children are less likely to have been exposed to many of these chemicals.

Several different gene changes usually occur in normal cells before they become cancerous. There are many kinds of brain tumors, each of which may have different sets of gene changes. A number of gene changes have been found in different brain tumor types, but there are probably many others that have not yet been found.

Researchers now understand some of the gene changes that occur in different types of brain tumors, but it's still not clear what causes these changes. Some gene changes might be inherited, but most brain and spinal cord tumors in children are not the result of known inherited syndromes. Most gene changes are probably just random events that sometimes happen inside a cell, without having an outside cause.

Other than [radiation](#)<sup>1</sup>, there are no known lifestyle-related or environmental factors clearly linked to childhood brain tumors, so it's important to remember that there is nothing these children or their parents could have done to prevent these cancers.

## Hyperlinks

1. [www.cancer.org/cancer/risk-prevention/radiation-exposure/x-rays-gamma-rays.html](http://www.cancer.org/cancer/risk-prevention/radiation-exposure/x-rays-gamma-rays.html)

## References

Dorsey JF, Hollander AB, Alonso-Basanta M, et al. Chapter 66: Cancer of the Central Nervous System. In: Abeloff MD, Armitage JO, Niederhuber JE, Kastan MB, McKenna WG, eds. *Abeloff's Clinical Oncology*. 5th ed. Philadelphia, Pa: Elsevier; 2014.

Lau C, Teo WY. Epidemiology of central nervous system tumors in children. UpToDate. 2018. Accessed at [www.uptodate.com/contents/epidemiology-of-central-nervous-system-tumors-in-children](http://www.uptodate.com/contents/epidemiology-of-central-nervous-system-tumors-in-children) on April 23, 2018.

Williams D, Parsons IF, Pollack DA. Chapter 26A: Gliomas, Ependymomas, and Other Nonembryonal Tumors of the Central Nervous System. In: Pizzo PA, Poplack DG, eds. *Principles and Practice of Pediatric Oncology*. 7th ed. Philadelphia, Pa: Lippincott

Williams & Wilkins; 2016.

Last Revised: June 20, 2018

## Can Brain and Spinal Cord Tumors in Children Be Prevented?

- [Limiting radiation exposure to the head](#)

Adults can lower their risk of certain cancers with lifestyle changes (such as staying at a healthy weight or quitting smoking), but at this time there are no known ways to prevent most cancers in children.

Other than exposure to [radiation](#)<sup>1</sup>, there are no known lifestyle-related or environmental [risk factors for brain and spinal cord tumors in children](#), so at this time there is no way to protect against most of these cancers.

### Limiting radiation exposure to the head

For most children with other types of cancer in or near the head, radiation therapy may be given if the doctors feel the benefits outweigh the small risk of developing a brain tumor years later. Still, when it is needed, doctors try to limit the dose of radiation as much as possible.

X-rays or CT scans done before birth or during childhood use much lower levels of radiation than those used for cancer treatment. If there is any increase in risk from these tests, it is likely to be very small, but to be safe, most doctors recommend that pregnant women and children not get these tests unless they are absolutely needed.

### Hyperlinks

1. [www.cancer.org/cancer/risk-prevention/radiation-exposure/x-rays-gamma-rays.html](http://www.cancer.org/cancer/risk-prevention/radiation-exposure/x-rays-gamma-rays.html)



## References

Chang SM, Mehta MP, Vogelbaum MA, Taylor MD, Ahluwalia MS. Chapter 97: Neoplasms of the central nervous system. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 10th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2015.

Dorsey JF, Hollander AB, Alonso-Basanta M, et al. Chapter 66: Cancer of the central nervous system. In: Abeloff MD, Armitage JO, Niederhuber JE, Kastan MB, McKenna WG, eds. *Abeloff's Clinical Oncology*. 5th ed. Philadelphia, Pa: Elsevier; 2014.

Last Revised: June 20, 2018

## Written by

The American Cancer Society medical and editorial content team  
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy ([www.cancer.org/about-us/policies/content-usage.html](http://www.cancer.org/about-us/policies/content-usage.html)).

**cancer.org | 1.800.227.2345**