

Infectious Diseases and Clinical Xenotransplantation

Jay A. Fishman, Nicolas J. Mueller



In support of improving patient care, this activity has been planned and implemented by Medscape, LLC and Emerging Infectious Diseases. Medscape, LLC is jointly accredited with commendation by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medscape, LLC designates this Journal-based CME activity for a maximum of 1.00 **AMA PRA Category 1 Credit(s)**[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1.0 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

All other clinicians completing this activity will be issued a certificate of participation. To participate in this journal CME activity: (1) review the learning objectives and author disclosures; (2) study the education content; (3) take the post-test with a 75% minimum passing score and complete the evaluation at <http://www.medscape.org/journal/eid>; and (4) view/print certificate. For CME questions, see page XXX.

NOTE: It is Medscape's policy to avoid the use of Brand names in accredited activities. However, in an effort to be as clear as possible, trade names are used in this activity to distinguish between the mixtures and different tests. It is not meant to promote any particular product.

Release date: June 20, 2024; Expiration date: June 20, 2025

Learning Objectives

Upon completion of this activity, participants will be able to:

- Assess how swine are prepared as source animals for xenotransplantation
- Distinguish the most common potentially harmful organisms associated with xenotransplantation
- Compare viruses that may infect human and swine cells vs swine cells only
- Evaluate therapeutic interventions to reduce the risk for infectious complications of xenotransplantation

CME Editor

Dana C. Dolan, BS, Technical Writer/Editor, Emerging Infectious Diseases. *Disclosure: Dana C. Dolan, BS, has no relevant financial relationships.*

CME Author

Charles P. Vega, MD, Health Sciences Clinical Professor of Family Medicine, University of California, Irvine School of Medicine, Irvine, California. *Charles P. Vega, MD, has the following relevant financial relationships: consultant or advisor for Boehringer Ingelheim; GlaxoSmithKline.*

Authors

Jay A. Fishman, MD; Nicolas J. Mueller.

Author affiliations: Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA (J.A. Fishman); University Hospital Zurich, University of Zurich, Switzerland (N.J. Mueller)

DOI: <http://doi.org/10.3201/eid3007.240273>

Xenotransplantation, transplantation into humans of vascularized organs or viable cells from nonhuman species, is a potential solution to shortages of transplantable human organs. Among challenges to application of clinical xenotransplantation are unknown risks of transmission of animal microbes to immunosuppressed recipients or the community. Experience in allotransplantation and in preclinical models suggests that viral infections are the greatest concern. Worldwide, the distribution of swine pathogens is heterogeneous and cannot be fully controlled by international agricultural regulations. It is possible to screen source animals for potential human pathogens before procuring organs in a manner not possible within the time available for surveillance testing in allotransplantation. Infection control measures require microbiological assays for surveillance of source animals and xenograft recipients and research into zoonotic potential of porcine organisms. Available data suggest that infectious risks of xenotransplantation are manageable and that clinical trials can advance with appropriate protocols for microbiological monitoring of source animals and recipients.

Xenotransplantation, the implantation of vascularized organs or viable cells from nonhuman species into humans, is under development to address the shortage of human organs for transplantation. Clinical xenotransplantation from swine has become more practical through advances in molecular biology (e.g., CRISPR manipulations) that have

enabled the breeding of swine with advantageous immunologic traits coupled with newer immunosuppressive regimens (Figure; Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/30/7/24-0273-App1.pdf>). Recent porcine cardiac and renal transplants survived for about 2 months in hosts with multiple comorbid conditions and who were not candidates for allotransplantation. Decedent xenografts of hearts and kidneys have been used to demonstrate fundamental functions and immune responses of porcine xenografts in human hosts. Prior experience with xenogeneic (pig, bovine) heart valves, tendons, and skin have generally been fixed or sterilized tissues not carrying viable cells. Regulatory guidelines exist for the clinical use of genetically modified animals but incompletely address microbiologic standards (1–5). Experience in allotransplantation indicates that the risk for xenosis or xenozoonosis (transmission of infection from animals to humans from viable cells of organs or cellular transplantation) is determined by epidemiologic exposure of source animals and human recipients, the net state of immunosuppression, and the underlying factors contributing to infectious risk, including the type, intensity and duration of immunosuppression (6,7). In human allotransplantation, immunosuppression is largely standardized, the pattern of infections is predictable, and prophylactic regimens are standardized (6,7). Some infections are considered




Figure. Advances in genetic engineering have led to the breeding of pigs with advantages in infection, immunology, coagulation, size, and inflammation. Breeding of source animals in biosecure facilities enables screening for potential pathogens. B4GalNT2, glycosyltransferase; CD46, human membrane cofactor protein; CD47, block SIRPα tyrosine phosphorylation; CD55, human decay-accelerating factor; CMAH, cytidine monophosphate-N-acetylneuraminic acid hydroxylase; EPCR, endothelial cell protein C receptor; GGAT1, α-1,3-glycosyltransferase; HO1, heme oxygenase-1; HA20, human A20; PERV, porcine endogenous retrovirus; THBD, human thrombomodulin gene.

routine (human cytomegalovirus [CMV], Epstein-Barr virus [EBV]); unexpected infections reflect excess immunosuppression, unusual exposures in the hospital or community, or donor organ-derived infections. Unexpected donor-derived infections are uncommon despite the urgency of screening, given time limitations for organ implantation (8,9). Data from microbiologic screening studies are often not available until after implantation.

Immunosuppressed xenograft recipients have potential exposure to microbes carried by xenografts as well as to community-derived exposures and reactivation of latent infections in the human host (10). The likelihood of infection caused by any specific organism is unknown, particularly without clinical trials or validated assays for pig-specific pathogens. A challenge and benefit of xenotransplantation is the ability to develop herds of animals free of potential pathogens; thus, developing serologic and molecular assays for use in swine herds and to monitor for infection in recipients is an important component of safety in clinical xenotransplantation.

Developing Swine as Source Animals for Clinical Xenotransplantation

Consensus does not exist around optimal screening paradigms for source animals and for monitoring of human recipients (Appendix Tables 2, 3). Effective prophylactic strategies require gaining clinical experience, identifying important pathogens, and studying antimicrobial efficacy for organisms from pigs (10,11). Veterinary guidance for pig health tracks common pathogens and requires screening of animal care providers and animals for infectious exposures. Animals determined specific pathogen-free (SPF) are generally screened for drug-resistant organisms, have limited routine use of antimicrobial agents, are vaccinated extensively, and receive sterile feed in biosecure facilities. Herds of swine for xenotransplantation are maintained in biosecure facilities and monitored routinely to exclude potential human pathogens; this practice is termed designated pathogen-free (DPF) status, a term adopted by regulatory authorities (12,13). This list of potential pathogens is based on experience in allotransplantation and pre-clinical xenotransplantation; it includes organisms that cause infection in immunocompromised persons (e.g., *Toxoplasma gondii*) or that are like those causing infection in transplant recipients (Appendix Table 2). Some porcine viruses have known zoonotic potential, including hepatitis E virus, influenza A virus (IAV), Japanese encephalitis virus, and Nipah virus. Pathogens known to be infectious in both

pigs and humans (e.g., hepatitis E, influenza) merit monitoring and exclusion from breeding herds (14). Depending on the sensitivity of the assays used, organisms excluded from the breeding herd should not pose a threat to xenograft recipients. Pigs are rescreened at the time of organ procurement for xenotransplantation for known pathogens (Appendix Table 3). They are also screened by histopathology, by metagenomic sequencing for unknown organisms, and by routine blood and tissue cultures for bacteria or fungi; not all data are available at the time of procurement.

Immunosuppression for xenotransplantation in nonhuman primates and in recent human xenocardiac and xenokidney recipients has included costimulatory blockade by CD40 or CD154 monoclonal antibodies, mycophenylate mofetil, and T-cell (anti-thymocyte globulin), B-cell (CD20), and complement inhibition or depletion with perfusion solutions containing anti-inflammatory agents (15–18). Similar regimens in humans are associated with increased risk for infections from certain viruses (CMV, EBV, BK polyomavirus), *Pneumocystis* spp., *Toxoplasma* spp., and encapsulated bacteria (e.g., *Neisseria meningitidis* A and B).

Groenendaal et al. compiled a list of all known organisms infecting swine from the literature and sorted these based on pathogenicity for pigs and similarity to organisms causing infection in immunocompromised human hosts (19). The report identified 254 potential pathogens in pigs in the United States: 108 viruses, 75 bacteria, 59 parasites, 11 fungi and 1 prion; it did not include organisms potentially introduced during the procurement and transportation of organs intended for transplantation (19). The list includes organisms important to routine pig health status, pig pathogens, and potential human pathogens; clear distinction is not yet feasible given limited clinical experience (19). Most (~130) pathogens are routinely excluded from biosecure pig-breeding facilities. The list supports development of risk mitigation strategies including requirements for biosecure breeding facilities, pathogen monitoring and exclusion, pharmaceutical treatment or vaccination, and genome editing; however, screening and monitoring for infection remained difficult.

Some viruses identified in preclinical pig-to-primate xenotransplantation appear to be species specific and do not infect human cells; those viruses include porcine cytomegalovirus, PCMV, and porcine circovirus. The viruses proliferate in xenografts in immunosuppressed nonhuman primates (NHP) and may cause graft dysfunction, graft rejection,

coagulopathy, or other syndromes (20–23). In baboon recipients of porcine heart and kidney xenografts, PCMV-infected pig cells and viral DNA are found in circulation despite ganciclovir prophylaxis. PCMV-infected porcine endothelial cells become activated to produce tissue factor, leading to systemic consumptive coagulopathy and accelerated renal xenograft rejection (12,20,21,23). PCMV can be excluded from pig colonies by caesarean delivery, early weaning, and biosecure isolation but is easily reintroduced (24–26). Those interventions inconsistently reduce transmission of porcine lymphotropic herpesvirus (PLHV) from sow to piglet (25,27). In a human recipient of a porcine cardiac xenograft, PCMV was detected by unbiased plasma microbial cell-free DNA testing despite negative molecular testing of nasal swab and buffy coat before organ procurement and ganciclovir prophylaxis (28,29). These observations demonstrate the value of pig screening using serologic testing, nucleic acid testing (NAT), and other advanced techniques. Four species of porcine circovirus, 1–4, cause infection in swine; diverse clinical associations have been made between PCVs in swine. PCV2 and PCV3 disseminate with shed cells from cardiac xenografts in baboons; transmission to primate cells has not been demonstrated. No transmission of PCV was identified in a seropositive pig-to-human cardiac recipient (28).

The porcine endogenous retroviruses (PERV) have the theoretical capacity to integrate into the hosts' germline DNA causing insertional effects (13,30). PERV receptors for PERV-A and -B are ubiquitous in human cells (HuPAR-1 and HuPAR-2) (31). PERV-A and PERV-B can infect transformed human cells. PERV-C infects only pig cells. Receptor homologs in porcine cells are active while those in baboon appear inactive; baboons do not have PERV infection develop after xenotransplantation. Recombinant PERV-AC is a naturally occurring recombinant between PERV-A and -C and infects with greater efficiency than PERV-A into transformed human cells via the PERV-A receptor domain. PERV infection of humans exposed to porcine cells has not been reported. Various PERV mitigation strategies for source pigs include selective breeding of PERV-C-free pigs (which does not preclude recipient infection due to PERV-A or PERV-B), or genetic inactivation of the polymerase gene of PERV proviral elements using CRISPR-Cas9 (32).

Shifting Epidemiology of Organisms of Swine

Infectious disease management is a central component of the pork industry. Biosecurity precautions

vary across breeding facilities; one precaution is the exclusion of birds, rodents, and amphibians. Strict regulations exist for the international movement of pigs, feed, and pork products. The intensification of animal agriculture, applying technological advances to allow increased density of animal rearing, has accounted for emergence of new zoonoses resulting from various practices including crowded housing, use of antibiotics, deforestation, and inadequate waste management and contributes to global warming (33). The spread of animal microbes to humans has increased with contacts between humans and wild or domesticated animal hosts in agriculture and markets (33–35). For example, the spread of HIV, SARS-CoV-2, Middle East respiratory syndrome virus, swine influenza virus, hepatitis E virus, and Japanese encephalitis virus was the result of contacts between animal reservoirs and humans (36–38). The need for constant review of pathogens that require surveillance in swine raised for clinical xenotransplantation is demonstrated by porcine circovirus type 4, which was not reported in the literature until 2019 but had been identified in epidemiologic studies of swine for more than a decade (39). Global warming and intensified pig farming in previous bat habitats may have contributed to the spread of Nipah virus to swine and farmers in Malaysia. Epidemiologically restricted pathogens of swine are likely to spread to new areas with global warming, economic development, and international travel and trade. These may include many parasites, bacteria such as *Burkholderia* species and viruses such as Nipah, PCV4, lymphocytic choriomeningitis, and Japanese encephalitis. Worldwide, porcine organisms of concern with zoonotic potential are increasing; among those, use of antimicrobial agents is associated with increasing antimicrobial resistance. They include bacteria (*Salmonella*, *S. suis*, *S. aureus*, *Campylobacter*, *Mycobacteria*, *Brucella*, *Leptospira*, *E. coli*), parasites (*Trichinella*, *Toxoplasma*, *Trypanosoma*), and viruses (influenza, Nipah, Japanese encephalitis, Menangle) (19). Those pathogens merit surveillance in pig herds as their epidemiologic footprint expands. At the same time, biosecurity facilities have improved through experience and necessity in genetic manipulation and oocyte implantation, which may mitigate some of the challenges of maintenance of DPF status.

Microbiological Testing in the Human Recipient

The key elements of infection control are exclusion of potential pathogens from breeding herds (DPF status) and monitoring in xenograft recipients and

clinical staff (Appendix Table 3) (11,30,40,41). Although bacteria, fungi, and parasites can generally be identified in veterinary or clinical microbiologic labs by culture-based techniques, viruses require both serologic and NAT. Multiplexed PCRs against multiple viral targets have been reported for use in pigs (42). Pathogen-directed viral assays are not yet validated in humans; some assays may not be able to distinguish between similar porcine and human pathogens (43). Porcine retroviruses such as PERV AC have some unstable target sequences or variable tissue tropism and may require functional assays (e.g., reverse transcribed retrovirus on productively infected target cells), full sequence analysis, or in situ hybridization (44).

The availability of archived biospecimens from source pigs and recipients, and from persons with significant exposures to donor swine and recipients, will enable researchers to evaluate infections and possible donor-derived transmissions. Metagenomic or next-generation sequencing (NGS) approaches rely on available pathogen sequence data for analyzing sequences derived from animals or in preclinical or clinical recipients. As genetic databases for genomic and microbial sequences grow, retrospective analysis of stored clinical samples is feasible. Because infections are common in immunosuppressed allotransplant recipients, such techniques are also helpful for evaluating infectious syndromes for which a specific diagnosis cannot be established. NGS approaches are pathogen agnostic and may also detect colonizing species or replication-incompetent sequences of unclear clinical significance (40). Using a NGS approach is of particular interest for pathogen discovery in the context of xenotransplantation where knowledge of potential porcine pathogens is limited (45); the technology was instrumental in the discovery of several new viruses, some associated with human disease (46). Those data will also address concerns regarding potential spread of xenogeneic organisms to the general population.

Prevention and Surveillance of Infection in the Xenograft Recipient

After xenotransplantation, recipient surveillance must consider both swine and human pathogens (Appendix Table 3). Standard allotransplantation prophylactic regimens can be used for perioperative bacterial infections, herpesviruses, molds, *Toxoplasma gondii*, and *Pneumocystis jirovecii*. Novel immunosuppression regimens may alter the spectrum of opportunistic infections. Testing should be guided by knowledge of microbes not excluded from the

breeding herd (e.g., PERV and PCMV status). Surveillance will require use of laboratory-developed assays or off-label use of available tests for more extensive pathogen discovery (e.g., NGS). Recent porcine-to-human cardiac and renal xenotransplants successfully used NGS for posttransplant surveillance (28). Biopsies used to monitor graft rejection should include microbial analysis using cultures, NGS, immunohistology, and electron microscopy for viral infections. Clinical trials should consider standard protocols for management of fevers or infectious syndromes in addition to routine screening during early periods. Treating graft rejection or infectious syndromes requires increased testing.

Porcine Antiviral Therapy Prophylaxis and Treatment

Strategies for prevention and treatment of potential viral infections in xenotransplantation, as for allotransplantation, include understanding of the antiviral susceptibilities of porcine viruses. Data on antiviral therapy for porcine viruses are limited (41). PCMV does not infect human cells but can provoke graft dysfunction and coagulopathy and will merit prophylaxis and therapy. PCMV has reduced susceptibility to acyclovir, ganciclovir, and foscarnet; ganciclovir prophylaxis at full treatment doses is inconsistently effective in vivo in immunosuppressed NHP xenograft recipients. Consistent with homology with human herpesvirus 6, the nephrotoxic agent cidofovir is more effective at therapeutic concentrations in vitro (22,47). Porcine lymphotropic herpesviruses (PLHV) 1, 2, and 3 have been associated with a lymphoproliferative disorder after experimental hematopoietic stem cell transplantation in pigs; the viruses are not known to be pathogens in NHP or in humans, and no effective antiviral drugs exist. PLHV was not activated after xenotransplantation of various organs from swine infected with PLHV into nonhuman primates (20)

Regarding PERV, transmission was not identified in decedent recipients of renal xenografts for ≤ 72 hours or in recipients of PERV-C negative cardiac xenografts for ≤ 60 days; chimerism of cells infected with PERV-A or -B is expected. Retroviral transmission to xenograft recipients remains a concern (28,48,49). Antiretroviral drugs used to treat HIV-1, including reverse transcription inhibitors zidovudine, tenofovir, and adefovir, as well as the integrase inhibitors raltegravir and dolutegravir, can inhibit PERV. Nonnucleoside reverse transcriptase inhibitors (nevirapine) and protease inhibitors lack inhibitory activity for PERV. Should PERV therapy

or postexposure prophylaxis be required, combination antiretroviral therapy using integrase inhibitors and active nucleoside reverse transcriptase inhibitors would be recommended.

There are no specific treatments known for circoviruses PCV1-4; however, swine vaccination is available. Caesarean delivery and colostrum deprivation with use of NAT can prevent PCV transmission to piglets.

Infection Control in Clinical Xenotransplantation

As part of protocol development and the informed consent process, prospective xenograft recipients require education about infectious risks of xenotransplantation to themselves and potentially to social and sexual partners, of which data are limited. In the absence of PERV risk, standard universal precautions for xenograft recipients should be adequate to protect hospital staff and social contacts. No infections have been reported among veterinary staff, scientists, or surgeons participating in preclinical xenotransplant studies. As for any surgical procedure, the risk for exposure is greatest for operating room staff handling pig organs and fluids or via splash or needlestick injury. Standard surgical infection control practices should prevent such exposures. Given the unknowns, archiving baseline leukocyte and plasma samples could enable future investigations should infectious syndromes emerge in xenograft recipients. Additional samples can be obtained for documented exposures to bodily fluids or with undiagnosed infectious syndromes in xenograft recipients or surgical teams. General hospital care workers for xenotransplant recipients should not have risks of exposure beyond those prevented by universal precautions. Infection and infectious syndromes are common in immunosuppressed transplant recipients; recipients should follow isolation precautions based on the primary syndrome (e.g., for diarrhea or pneumonitis).

Occupational health service staff should be aware of xenotransplantation protocols for blood or body fluid exposure from source animals or xenotransplant recipients. In such situations, knowing the infectious status of the source pig and the recipient is invaluable. If the donor animal is PERV negative, postexposure retroviral prophylaxis should not be required. For PERV-positive donors, prophylaxis after needlestick exposure to porcine tissues recommends use of a reverse transcription inhibitor and integrase inhibitor. Testing should include NAT for swine-specific pathogens, as well as standard tests for HIV, hepatitis C and hepatitis B. Repeat NAT

testing should be performed at regular intervals (e.g., 1, 3, and 6 months) after a blood or body fluid exposure. Plans for passive surveillance and active testing and treatment will be required for clinical trials; those plans should be developed in conjunction with Infection Control and Occupational Health groups. Informed consent may be required for acquiring and storing blood samples from clinical care providers.

Because clinical experience is limited, infectious risks to close contacts of the xenotransplant recipient are not defined. The clinical trial design and consent process should address the benefits and feasibility of posttransplant surveillance of close contacts to inform blood sample archiving in advance of the procedure in the event of blood or body fluid exposure. As part of pretransplant education, the recipient and close contacts should be instructed to refrain from blood donation and unprotected sexual contacts; household members may be counseled to avoid sharing items that could be contaminated with blood. Education on potential risks to recipients, healthcare providers, and the general public includes ethical considerations for unknown hazards. The actual risk for infectious spread to the public is unknown; most potential pathogens are species specific. Active PERV can be excluded; recombination events should not occur without viral replication but cannot be completely excluded. With careful screening of source animals and monitoring of recipients for unknown as well as known microbes, the risk for xenogeneic spread to the public is very limited. Data from clinical trials will refine our understanding of disease transmission via xenotransplantation and will inform education for potential recipients and the public.

Conclusions

The risk for transmission of infection due to novel pathogens in association with xenotransplantation is unknown. Microbiological screening of source animals may reduce infectious risk; however, unknown porcine pathogens with capacity to infect humans may exist and are unlikely to be identified in the absence of clinical trials. The effect of the activation of PCMV in 1 cardiac recipient demonstrated the importance of herd screening for xenotransplants (28). Studies in deceased recipients of kidneys and hearts have provided information on metabolic and immunologic aspects (e.g., role of innate immunity), but they have reported limited immunosuppression and are of limited durations (<2 months) and so are less informative regarding infectious risks (50). Infection control measures include storage of baseline blood samples from the

xenograft donor, persons involved in procurement and transplantation of pig organs, and serial monitoring of the recipient and close contacts for known and possible unknown pathogens. Assays, including metagenomics, for potential pig pathogens need to be developed and validated. Transparency is essential in microbiologic investigations performed in clinical xenotransplantation trials.

J.A.F. is a consultant to United Therapeutics, eGenesis, Makana, Elion, Jura, Well Medical, Kamada, CLD Inc, OM1, and Vertex Inc.

About the Author

Dr. Fishman is professor of medicine at Harvard Medical School, director of the Transplant Infectious Diseases Program at Massachusetts General Hospital, associate director of the MGH Transplant Center, and president-elect of the International Xenotransplantation Society. His primary research interests include managing infectious risk in allotransplantation and xenotransplantation and studied porcine cytomegalovirus and porcine endogenous retrovirus in preclinical models. Dr. Mueller is a transplant infectious diseases physician and head of the transplant center at the University Hospital Zurich and chairman of the Scientific Committee of the Swiss Transplant Cohort Study. His initial research focused on viral infections in animal models of allo- and xenotransplantation; more recently, his primary research focus has been the role of latent herpes virus infections in the immunocompromised host.

References

- Hering BJ, Cozzi E, Spizzo T, Cowan PJ, Rayat GR, Cooper DK, et al. First update of the International Xenotransplantation Association consensus statement on conditions for undertaking clinical trials of porcine islet products in type 1 diabetes – executive summary. *Xenotransplantation*. 2016;23:3-13. <https://doi.org/10.1111/xen.12231>
- Food and Drug Administration. Source animal, product, preclinical, and clinical issues concerning the use of xenotransplantation products in humans; guidance for industry. 2016 [cited 2024 May 24]. <https://www.fda.gov/downloads/biologicsbloodvaccines/guidancecompliance-regulatoryinformation/guidances/xenotransplantation/ucm533036.pdf>
- Food and Drug Administration. PHS guideline on infectious disease issues in xenotransplantation. 66 F.R. 8120. 2001 [cited 2024 May 24]. <https://www.federalregister.gov/documents/2001/01/29/01-2419/phs-guideline-on-infectious-disease-issues-in-xenotransplantation-availability>
- World Health Organization. Second WHO global consultation on regulatory requirements for xenotransplantation clinical trials. 2001 [cited 2024 May 24]. http://www.who.int/transplantation/xeno/report2nd_global_consultation_txt.pdf
- Hawthorne WJ, Cowan PJ, Bühler LH, Yi S, Bottino R, Pierson RN III, et al. Third WHO global consultation on regulatory requirements for xenotransplantation clinical trials, Changsha, Hunan, China, December 12–14, 2018. *Xenotransplantation*. 2019;26:e12513. <https://doi.org/10.1111/xen.12513>
- Fishman JA. Infection in solid-organ transplant recipients. *N Engl J Med*. 2007;357:2601–14. <https://doi.org/10.1056/NEJMra064928>
- Fishman JA. Infection in organ transplantation. *Am J Transplant*. 2017;17:856–79. <https://doi.org/10.1111/ajt.14208>
- Fishman JA, Grossi PA. Donor-derived infection—the challenge for transplant safety. *Nat Rev Nephrol*. 2014;10:663–72. <https://doi.org/10.1038/nrneph.2014.159>
- Fishman JA. Xenosis and xenotransplantation: addressing the infectious risks posed by an emerging technology. *Kidney Int Suppl*. 1997;58:S41–5.
- Fishman JA. Risks of infectious disease in xenotransplantation. *N Engl J Med*. 2022;387:2258–67. <https://doi.org/10.1056/NEJMra2207462>
- Nellore A, Walker J, Kahn MJ, Fishman JA. Moving xenotransplantation from bench to bedside: managing infectious risk. *Transpl Infect Dis*. 2022:e13909. <https://doi.org/10.1111/tid.13909>
- Fishman JA. Infection in xenotransplantation. *J Card Surg*. 2001;16:363–73. <https://doi.org/10.1111/j.1540-8191.2001.tb00536.x>
- Fishman JA, Patience C. Xenotransplantation: infectious risk revisited. *Am J Transplant*. 2004;4:1383–90. <https://doi.org/10.1111/j.1600-6143.2004.00542.x>
- Denner J. Hepatitis E virus (HEV) – the future. *Viruses*. 2019;11:251. <https://doi.org/10.3390/v11030251>
- Längin M, Mayr T, Reichart B, Michel S, Buchholz S, Guethoff S, et al. Consistent success in life-supporting porcine cardiac xenotransplantation. *Nature*. 2018;564:430–3. <https://doi.org/10.1038/s41586-018-0765-z>
- Mohiuddin MM, Singh AK, Corcoran PC, Thomas ML III, Clark T, Lewis BG, et al. Chimeric 2C10R4 anti-CD40 antibody therapy is critical for long-term survival of GTKO. hCD46.hTBM pig-to-primate cardiac xenograft. *Nat Commun*. 2016;7:11138. <https://doi.org/10.1038/ncomms11138>
- Ma D, Hirose T, Lassiter G, Sasaki H, Rosales I, Coe TM, et al. Kidney transplantation from triple-knockout pigs expressing multiple human proteins in cynomolgus macaques. *Am J Transplant*. 2022;22:46–57. <https://doi.org/10.1111/ajt.16780>
- Firl DJ, Markmann JF. Measuring success in pig to non-human-primate renal xenotransplantation: Systematic review and comparative outcomes analysis of 1051 life-sustaining NHP renal allo- and xeno-transplants. *Am J Transplant*. 2022;22:1527–36. <https://doi.org/10.1111/ajt.16994>
- Groenendaal H, Costard S, Ballard R, Bienhoff S, Challen DC, Dominguez BJ, et al. Expert opinion on the identification, risk assessment, and mitigation of microorganisms and parasites relevant to xenotransplantation products from pigs. *Xenotransplantation*. 2023;30:e12815. <https://doi.org/10.1111/xen.12815>
- Mueller NJ, Barth RN, Yamamoto S, Kitamura H, Patience C, Yamada K, et al. Activation of cytomegalovirus in pig-to-primate organ xenotransplantation. *J Virol*. 2002;76:4734–40. <https://doi.org/10.1128/JVI.76.10.4734-4740.2002>
- Gollackner B, Mueller NJ, Houser S, Qawi I, Soizic D, Knosalla C, et al. Porcine cytomegalovirus and coagulopathy in pig-to-primate xenotransplantation. *Transplantation*. 2003;75:1841–7. <https://doi.org/10.1097/01.TP.0000065806.90840.C1>

22. Mueller NJ, Sulling K, Gollackner B, Yamamoto S, Knosalla C, Wilkinson RA, et al. Reduced efficacy of ganciclovir against porcine and baboon cytomegalovirus in pig-to-baboon xenotransplantation. *Am J Transplant*. 2003; 3:1057–64. <https://doi.org/10.1034/j.1600-6143.2003.00192.x>
23. Mueller NJ, Kuwaki K, Dor FJ, Knosalla C, Gollackner B, Wilkinson RA, et al. Reduction of consumptive coagulopathy using porcine cytomegalovirus-free cardiac porcine grafts in pig-to-primate xenotransplantation. *Transplantation*. 2004;78:1449–53. <https://doi.org/10.1097/01.TP.0000141361.68446.1F>
24. Clark DA, Fryer JF, Tucker AW, McArdle PD, Hughes AE, Emery VC, et al. Porcine cytomegalovirus in pigs being bred for xenograft organs: progress towards control. *Xenotransplantation*. 2003;10:142–8. <https://doi.org/10.1034/j.1399-3089.2003.01128.x>
25. Mueller NJ, Kuwaki K, Knosalla C, Dor FJ, Gollackner B, Wilkinson RA, et al. Early weaning of piglets fails to exclude porcine lymphotropic herpesvirus. *Xenotransplantation*. 2005;12:59–62. <https://doi.org/10.1111/j.1399-3089.2004.00196.x>
26. Yamada K, Tasaki M, Sekijima M, Wilkinson RA, Villani V, Moran SG, et al. Porcine cytomegalovirus infection is associated with early rejection of kidney grafts in a pig to baboon xenotransplantation model. *Transplantation*. 2014;98:411–8. <https://doi.org/10.1097/TP.0000000000000232>
27. Denner J. Porcine lymphotropic herpesviruses (PLHVs) and xenotransplantation. *Viruses*. 2021;13:1072. <https://doi.org/10.3390/v13061072>
28. Griffith BP, Goerlich CE, Singh AK, Rothblatt M, Lau CL, Shah A, et al. Genetically modified porcine-to-human cardiac xenotransplantation. *N Engl J Med*. 2022;387:35–44. <https://doi.org/10.1056/NEJMoa2201422>
29. Fishman JA. Next-generation sequencing for identifying unknown pathogens in sentinel immunocompromised hosts. *Emerg Infect Dis*. 2023;29:431–2. <https://doi.org/10.3201/eid2902.221829>
30. Fishman JA. Infectious disease risks in xenotransplantation. *Am J Transplant*. 2018;18:1857–64. <https://doi.org/10.1111/ajt.14725>
31. Ericsson TA, Takeuchi Y, Templin C, Quinn G, Farhadian SF, Wood JC, et al. Identification of receptors for pig endogenous retrovirus. *Proc Natl Acad Sci U S A*. 2003;100:6759–64. <https://doi.org/10.1073/pnas.1138025100>
32. Niu D, Wei HJ, Lin L, George H, Wang T, Lee IH, et al. Inactivation of porcine endogenous retrovirus in pigs using CRISPR-Cas9. *Science*. 2017;357:1303–7. <https://doi.org/10.1126/science.aan4187>
33. Hayek MN. The infectious disease trap of animal agriculture. *Sci Adv*. 2022;8:eadd6681. <https://doi.org/10.1126/sciadv.add6681>
34. Linder A. Animal markets and zoonotic disease in the United States. 2023 [cited 2024 May 24]. <https://animal.law.harvard.edu/wp-content/uploads/Animal-Markets-and-Zoonotic-Disease-in-the-United-States.pdf>
35. VanderWaal K, Deen J. Global trends in infectious diseases of swine. *Proc Natl Acad Sci U S A*. 2018;115:11495–500. <https://doi.org/10.1073/pnas.1806068115>
36. Pavio N, Meng XJ, Renou C. Zoonotic hepatitis E: animal reservoirs and emerging risks. *Vet Res*. 2010;41:46. <https://doi.org/10.1051/vetres/2010018>
37. Lopez-Moreno G, Davies P, Yang M, Culhane MR, Corzo CA, Li C, et al. Evidence of influenza A infection and risk of transmission between pigs and farmworkers. *Zoonoses Public Health*. 2022;69:560–71. <https://doi.org/10.1111/zph.12948>
38. McLean RK, Graham SP. The pig as an amplifying host for new and emerging zoonotic viruses. *One Health*. 2022;14:100384. <https://doi.org/10.1016/j.onehlt.2022.100384>
39. Zhai SL, Lu SS, Wei WK, Lv DH, Wen XH, Zhai Q, et al. Reservoirs of porcine circoviruses: a mini review. *Front Vet Sci*. 2019;6:319. <https://doi.org/10.3389/fvets.2019.00319>
40. Fishman JA. Prevention of infection in xenotransplantation: designated pathogen-free swine in the safety equation. *Xenotransplantation*. 2020;27:e12595. <https://doi.org/10.1111/xen.12595>
41. Mehta SA, Saharia KK, Nellore A, Blumberg EA, Fishman JA. Infection and clinical xenotransplantation: guidance from the Infectious Disease Community of Practice of the American Society of Transplantation. *Am J Transplant*. 2023;23:309–15. <https://doi.org/10.1016/j.ajt.2022.12.013>
42. Hartline CB, Conner RL, James SH, Potter J, Gray E, Estrada J, et al. Xenotransplantation panel for the detection of infectious agents in pigs. *Xenotransplantation*. 2018;25:e12427. <https://doi.org/10.1111/xen.12427>
43. Fiebig U, Holzer A, Ivanusic D, Plotzki E, Hengel H, Neipel F, et al. Antibody cross-reactivity between porcine cytomegalovirus (PCMV) and human herpesvirus-6 (HHV-6). *Viruses*. 2017;9:317. <https://doi.org/10.3390/v9110317>
44. Kono K, Kataoka K, Yuan Y, Yusa K, Uchida K, Sato Y. A highly sensitive method for the detection of recombinant PERV-A/C env RNA using next generation sequencing technologies. *Sci Rep*. 2020;10:21935. <https://doi.org/10.1038/s41598-020-78890-2>
45. Santiago-Rodriguez TM, Hollister EB. Unraveling the viral dark matter through viral metagenomics. *Front Immunol*. 2022;13:1005107. <https://doi.org/10.3389/fimmu.2022.1005107>
46. Li L, Kapoor A, Slikas B, Bamidele OS, Wang C, Shaikat S, et al. Multiple diverse circoviruses infect farm animals and are commonly found in human and chimpanzee feces. *J Virol*. 2010;84:1674–82. <https://doi.org/10.1128/JVI.02109-09>
47. Fryer JF, Griffiths PD, Fishman JA, Emery VC, Clark DA. Quantitation of porcine cytomegalovirus in pig tissues by PCR. *J Clin Microbiol*. 2001;39:1155–6. <https://doi.org/10.1128/JCM.39.3.1155-1156.2001>
48. Porrett PM, Orandi BJ, Kumar V, Houpp J, Anderson D, Cozette Killian A, et al. First clinical-grade porcine kidney xenotransplant using a human decedent model. *Am J Transplant*. 2022;22:1037–53. <https://doi.org/10.1111/ajt.16930>
49. Montgomery RA, Stern JM, Lonze BE, Tatapudi VS, Mangiola M, Wu M, et al. Results of two cases of pig-to-human kidney xenotransplantation. *N Engl J Med*. 2022;386:1889–98. <https://doi.org/10.1056/NEJMoa2120238>
50. Loupy A, Goutaudier V, Giarraputo A, Mezine F, Morgand E, Robin B, et al. Immune response after pig-to-human kidney xenotransplantation: a multimodal phenotyping study. *Lancet*. 2023;402:1158–69. [https://doi.org/10.1016/S0140-6736\(23\)01349-1](https://doi.org/10.1016/S0140-6736(23)01349-1)

Address for correspondence: Jay Fishman, MGH Transplant Center, 55 Fruit St, Boston, MA 02114 USA; email: fishman.jay@mgh.harvard.edu

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Infectious Diseases and Clinical Xenotransplantation

Appendix

Appendix Table 1. Potential targets for genetic manipulation in swine xenograft donors

Target type	Target	Potential gene targets for xenotransplantation
Pig breed	—	Various pig breeds
Endogenous retrovirus	Virus	Porcine endogenous retrovirus (PERV A, B, C, AC)
Inactivation		
Knockout	Carbohydrate antigens	GGAT1 (α -1,3-glycosyltransferase)
	Carbohydrate antigens	B4GalNT2 (glycosyltransferase)
	Carbohydrate antigens	CMAH (cytidine monophosphate-N-acetylneuraminic acid hydroxylase)
	Organ growth	Growth hormone receptor
Added human transgenes	Complement regulation	CD46 (hMCP, human membrane cofactor protein)
	Complement regulation	CD55 (hDAF human decay-accelerating factor)
	Coagulation	THBD (human thrombomodulin gene)
	Coagulation	EPCR (Endothelial cell protein C receptor)
	Innate immunity	CD47 (Block SIRP α tyrosine phosphorylation)
	Inflammation, apoptosis	HO1 (Heme Oxygenase-1)
	Inflammation, apoptosis	HA20 (Human A20)

Appendix Table 2. Risk categories for potential pathogens in recipients of porcine xenografts (1,2)*

Organism hosts	Examples	Microbiologic assays available?	Monitor in breeding colony?
Pathogens of immunologically normal humans and swine	Influenza viruses* (3–6); Hepatitis E virus* (7–9); <i>Mycobacterium tuberculosis</i> *; rabies*; many bacterial and parasitic species (e.g., <i>Ascaris</i> , <i>Toxocara</i> , <i>Pasteurella multocida</i> ; <i>Mycoplasma</i> spp.)	Yes	Yes
Known pathogens of immunosuppressed human transplant recipients	<i>Toxoplasma gondii</i> *; <i>Strongyloides</i> spp.*; <i>Aspergillus</i> sp.; <i>Cryptococcus</i> spp; <i>Cryptosporidium</i> spp.	Yes	Based on risk with organism
Porcine organisms similar to common pathogens of immunosuppressed human transplant recipients	Porcine adenovirus; porcine parvovirus 1; porcine respiratory coronavirus; parainfluenza virus 3	Few	Require validation of assays in human blood or tissues.
Unique swine pathogens (may replicate only in pig cells)	Porcine cytomegalovirus (PCMV)* (14–18); Porcine circovirus (PCV 1–4) (19–24); porcine lymphotropic herpesvirus (PLHV 1,2); porcine endogenous retrovirus* (PERV A, B, C, AC) (1,25)	Some	Herpesviruses generally species-specific. Risk requires clinical study
Organisms routinely tested for health status of swine	Porcine enterovirus spp; <i>Lawsonia Intracellularis</i> ; porcine epidemic diarrhea virus; transmissible gastroenteritis virus; porcine delta coronavirus; <i>Brucella suis</i> ; porcine reproductive and respiratory syndrome virus; porcine epidemic diarrhea virus; <i>Pseudorabies</i> virus	Yes	Yes

Organism hosts	Examples	Microbiologic assays available?	Monitor in breeding colony?
Porcine organisms largely geographically restricted (4,26,27) (examples)	<i>Burkholderia pseudomallei</i> ; <i>Clonorchis sinensis</i> ; <i>Echinococcus</i> spp; <i>Schistosoma</i> spp; African swine fever (ASF) virus; Menangle virus; Nipah virus (28,29); porcine circovirus type 4 (PCV4)	Some	Monitor for future geographic spread

*Consider exclusion of infected animals carrying these species. PERV may be excluded genetically (e.g., CRISPR-cas9) ; PERV-C negative animals carry potentially infectious PERV-A and B.

Appendix Table 3. Considerations in routine testing of xenograft recipients*

Virus	Testing method
Porcine endogenous retrovirus (PERV) A, B, C, AC (if present in source animal) *	Qualitative and quantitative (QNAT) nucleic acid testing (NAT); antibody- based tests (serology, ELISA, Western Bylot) ⁺
Porcine lymphotropic herpesvirus type 2 (PLHV-1–2)	QNAT ⁺
Porcine circovirus (PCV 1–4)	QNAT
Porcine cytomegalovirus (PCMV)	NAT ⁺ ; serology
Human cytomegalovirus (HCMV) – per risk status	QNAT, serology
Human Epstein-Barr virus (EBV) – per risk status	QNAT, serology
BK polyomavirus (kidney recipients) – per protocol	QNAT
Pig cell chimerism in circulation (PBMC)	QNAT ⁺ (e.g., P-MHC class I gene; p-mtCOII gene) in recipient PBMC DNA.
Unknown pathogens	Metagenomics or next generation sequencing (10–13)

* Additional testing is needed for individuals with infectious syndromes. QNAT: quantitative nucleic acid test; P-MHC: porcine major histocompatibility complex; p-mtCOII: pig mitochondrial cytochrome c oxidase subunit II gene

⁺ Quantitative NAT for PERV and other viruses must be normalized against chimerism studies to correct for the number of circulating pig cells in blood samples.

References

1. Fishman JA. Infectious disease risks in xenotransplantation. *Am J Transplant*. 2018;18:1857–64. [PubMed https://doi.org/10.1111/ajt.14725](https://doi.org/10.1111/ajt.14725)
2. Fishman JA. Risks of Infectious Disease in Xenotransplantation. *N Engl J Med*. 2022;387:2258–67. [PubMed https://doi.org/10.1056/NEJMra2207462](https://doi.org/10.1056/NEJMra2207462)
3. Campitelli L, Donatelli I, Foni E, Castrucci MR, Fabiani C, Kawaoka Y, et al. Continued evolution of H1N1 and H3N2 influenza viruses in pigs in Italy. *Virology*. 1997;232:310–8. [PubMed https://doi.org/10.1006/viro.1997.8514](https://doi.org/10.1006/viro.1997.8514)
4. Lopez-Moreno G, Davies P, Yang M, Culhane MR, Corzo CA, Li C, et al. Evidence of influenza A infection and risk of transmission between pigs and farmworkers. *Zoonoses Public Health*. 2022;69:560–71. [PubMed https://doi.org/10.1111/zph.12948](https://doi.org/10.1111/zph.12948)
5. Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology of influenza A viruses. *Microbiol Rev*. 1992;56:152–79. [PubMed https://doi.org/10.1128/mr.56.1.152-179.1992](https://doi.org/10.1128/mr.56.1.152-179.1992)

6. Zhou NN, Senne DA, Landgraf JS, Swenson SL, Erickson G, Rossow K, et al. Genetic reassortment of avian, swine, and human influenza A viruses in American pigs. *J Virol.* 1999;73:8851–6. [PubMed https://doi.org/10.1128/JVI.73.10.8851-8856.1999](https://doi.org/10.1128/JVI.73.10.8851-8856.1999)
7. Fang SY, Han H. Hepatitis E viral infection in solid organ transplant patients. *Curr Opin Organ Transplant.* 2017;22:351–5. [PubMed https://doi.org/10.1097/MOT.0000000000000432](https://doi.org/10.1097/MOT.0000000000000432)
8. Meng XJ. Swine hepatitis E virus: cross-species infection and risk in xenotransplantation. *Curr Top Microbiol Immunol.* 2003;278:185–216. [PubMed https://doi.org/10.1007/978-3-642-55541-1_7](https://doi.org/10.1007/978-3-642-55541-1_7)
9. Wang B, Meng XJ. Hepatitis E virus: host tropism and zoonotic infection. *Curr Opin Microbiol.* 2021;59:8–15. [PubMed https://doi.org/10.1016/j.mib.2020.07.004](https://doi.org/10.1016/j.mib.2020.07.004)
10. Casto AM, Fredricks DN, Hill JA. Diagnosis of infectious diseases in immunocompromised hosts using metagenomic next generation sequencing-based diagnostics. *Blood Rev.* 2022;53:100906. [PubMed https://doi.org/10.1016/j.blre.2021.100906](https://doi.org/10.1016/j.blre.2021.100906)
11. Greninger AL. The challenge of diagnostic metagenomics. *Expert Rev Mol Diagn.* 2018;18:605–15. [PubMed https://doi.org/10.1080/14737159.2018.1487292](https://doi.org/10.1080/14737159.2018.1487292)
12. Santiago-Rodriguez TM, Hollister EB. Unraveling the viral dark matter through viral metagenomics. *Front Immunol.* 2022;13:1005107. [PubMed https://doi.org/10.3389/fimmu.2022.1005107](https://doi.org/10.3389/fimmu.2022.1005107)
13. Simner PJ, Miller S, Carroll KC. Understanding the promises and hurdles of metagenomic next-generation sequencing as a diagnostic tool for infectious diseases. *Clin Infect Dis.* 2018;66:778–88. [PubMed https://doi.org/10.1093/cid/cix881](https://doi.org/10.1093/cid/cix881)
14. Mueller NJ, Barth RN, Yamamoto S, et al. Activation of cytomegalovirus in pig-to-primate organ xenotransplantation. *J Virol.* 2002;76:4734–40.
15. Gollackner B, Mueller NJ, Houser S, Qawi I, Soizic D, Knosalla C, et al. Porcine cytomegalovirus and coagulopathy in pig-to-primate xenotransplantation. *Transplantation.* 2003;75:1841–7. [PubMed https://doi.org/10.1097/01.TP.0000065806.90840.C1](https://doi.org/10.1097/01.TP.0000065806.90840.C1)
16. Yamada K, Tasaki M, Sekijima M, Wilkinson RA, Villani V, Moran SG, et al. Porcine cytomegalovirus infection is associated with early rejection of kidney grafts in a pig to baboon xenotransplantation model. *Transplantation.* 2014;98:411–8. [PubMed https://doi.org/10.1097/TP.0000000000000232](https://doi.org/10.1097/TP.0000000000000232)

17. Sekijima M, Waki S, Sahara H, Tasaki M, Wilkinson RA, Villani V, et al. Results of life-supporting galactosyltransferase knockout kidneys in cynomolgus monkeys using two different sources of galactosyltransferase knockout Swine. Transplantation. 2014;98:419–26. [PubMed](#) <https://doi.org/10.1097/TP.0000000000000314>
18. Denner J, Längin M, Reichart B, Krüger L, Fiebig U, Mokolke M, et al. Impact of porcine cytomegalovirus on long-term orthotopic cardiac xenotransplant survival. Sci Rep. 2020;10:17531. [PubMed](#) <https://doi.org/10.1038/s41598-020-73150-9>
19. Allan GM, McNeilly F, Kennedy S, et al. Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. J Vet Diagn Invest. 1998;10:3–10. <https://doi.org/10.1177/104063879801000102>
20. Denner J, Mankertz A. Porcine circoviruses and xenotransplantation. Viruses. 2017;9:83. [PubMed](#) <https://doi.org/10.3390/v9040083>
21. Klaumann F, Correa-Fiz F, Franzo G, Sibila M, Núñez JI, Segalés J. Current knowledge on *Porcine circovirus 3* (PCV-3): a novel virus with a yet unknown impact on the swine industry. Front Vet Sci. 2018;5:315. [PubMed](#) <https://doi.org/10.3389/fvets.2018.00315>
22. Krüger L, Längin M, Reichart B, Fiebig U, Kristiansen Y, Prinz C, et al. Transmission of porcine circovirus 3 (PCV3) by xenotransplantation of pig hearts into baboons. Viruses. 2019;11:650. [PubMed](#) <https://doi.org/10.3390/v11070650>
23. Zhai SL, Lu SS, Wei WK, Lv DH, Wen XH, Zhai Q, et al. Reservoirs of porcine circoviruses: a mini review. Front Vet Sci. 2019;6:319. [PubMed](#) <https://doi.org/10.3389/fvets.2019.00319>
24. Zhang H-H, Hu W-Q, Li J-Y, Liu TN, Zhou JY, Opriessnig T, et al. Novel circovirus species identified in farmed pigs designated as Porcine circovirus 4, Hunan province, China. Transbound Emerg Dis. 2020;67:1057–61. [PubMed](#) <https://doi.org/10.1111/tbed.13446>
25. Fishman JA, Patience C. Xenotransplantation: infectious risk revisited. Am J Transplant. 2004;4:1383–90. [PubMed](#) <https://doi.org/10.1111/j.1600-6143.2004.00542.x>
26. McLean RK, Graham SP. The pig as an amplifying host for new and emerging zoonotic viruses. One Health. 2022;14:100384. [PubMed](#) <https://doi.org/10.1016/j.onehlt.2022.100384>
27. Pavio N, Meng XJ, Renou C. Zoonotic hepatitis E: animal reservoirs and emerging risks. Vet Res. 2010;41:46. [PubMed](#) <https://doi.org/10.1051/vetres/2010018>

28. Drew TW. The emergence and evolution of swine viral diseases: to what extent have husbandry systems and global trade contributed to their distribution and diversity? *Rev Sci Tech*. 2011;30:95–106. [PubMed https://doi.org/10.20506/rst.30.1.2020](https://doi.org/10.20506/rst.30.1.2020)
29. VanderWaal K, Deen J. Global trends in infectious diseases of swine. *Proc Natl Acad Sci U S A*. 2018;115:11495–500. [PubMed https://doi.org/10.1073/pnas.1806068115](https://doi.org/10.1073/pnas.1806068115)