

Multiplex Dual-Target Reverse Transcription PCR for Subtyping Avian Influenza A(H5) Virus

Malaya K. Sahoo, Ingrid E.A. Morante, ChunHong Huang, Daniel Solis, Fumiko Yamamoto, Uzoamaka C. Ohiri, Daniel Romero, Benjamin A. Pinsky

An increased risk for human infection with avian influenza A(H5N1) viruses is of concern. We developed an internally controlled, dual-target reverse transcription PCR for influenza A(H5) subtyping. This test could be used to detect influenza A(H5) in clinical samples.

Highly pathogenic avian influenza (HPAI) viruses cause high mortality rates in wild birds and farmed poultry, and their potential for adaptation to humans remains a major pandemic threat (1). In 2021, a notable increase in avian influenza activity occurred, driven by the emergence of influenza A(H5N1) clade 2.3.4.4b viruses (2,3). Suspected mammal-to-mammal transmission during outbreaks among seals and minks led to concerns regarding the escalation of risk to humans (4–7). In 2024, the virus was identified in dairy cows in the United States (8), and exposure to infected cows has resulted in transmission to humans (9). To prepare for possible additional human cases, we developed an internally controlled, dual-target H5 subtyping quantitative reverse transcription PCR (qRT-PCR) based on primer–probe sequences from the World Health Organization (10). We evaluated this qRT-PCR by using synthesized nucleic acids, cultured avian and human virus isolates, and clinical upper respiratory specimens collected from patients with influenza A virus infection.

The Study

We identified previously published primer–probe sets from the World Health Organization that target different regions of the influenza A(H5) hemagglutinin

(HA) gene for combination into a dual-target H5 subtyping qRT-PCR (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/30/8/24-0785-App1.pdf>) (10). We introduced minor sequence changes to account for recent clade 2.3.4.4b diversity, and simultaneously, to limit primer–probe complexity (Table 1). We evaluated these primers and probes against complete North American H5 clade 2.3.4.4b HA gene sequences in the GISAID database (<https://www.gisaid.org>) during January 1, 2022–May 29, 2024, and found that 99.8% (5,975/5,987) of sequences aligned with a maximum of 1 mismatch to primer–probe set 1 and 97.7% (5,972/5,990) aligned with a maximum of 1 mismatch to primer–probe set 2 (Appendix Tables 2, 3). We observed no primer–probe mismatches compared with the influenza A H5 sequences obtained from persons with H5 infections in the United States (4 persons as of May 29, 2024) (Appendix Tables 4, 5). We combined the primer–probe sets in multiplex with primers–probes for the influenza A matrix (M) gene for pan-influenza A detection and with primers–probes for RNase P as an internal control (11).

We determined the 95% lower limit of detection (LLOD) by using single-stranded DNA (ssDNA) encoding the H5 target sequences from HPAI virus clade 1 (GenBank accession no. JQ966928, A/concador/Guangdong/139/2003 [H5N1]) (12) and clade 2.3.4.4b (GenBank accession no. OP499866, A/red-tailed hawk/Kansas/W22-198/2022 [H5N1]) viruses (Appendix Table 6). The primer–probe target regions of this clade 2.3.4.4b H5 sequence are identical to the recent H5 sequences from dairy cows and humans. We combined the 2 clade 1 ssDNA targets at equal copy numbers; we did the same with the clade 2.3.4.4b ssDNA targets. We also combined these H5 clade mixes at equal copy numbers with ssDNA encoding the pan-influenza A M gene target. We then diluted the clade 1 ssDNA mix and clade 2.3.4.4b ssDNA mix to 2, 1, and 0.5 copies/μL in pooled influenza

Author affiliations: Stanford Health Care Clinical Virology Laboratory, Stanford, California, USA (M.K. Sahoo, I.E.A. Morante, U.C. Ohiri, D. Romero, B.A. Pinsky); Stanford University School of Medicine, Stanford (C. Huang, D. Solis, F. Yamamoto, B.A. Pinsky)

DOI: <https://doi.org/10.3201/eid3008.240785>

Table 1. Primers and probes used in study of multiplex dual-target reverse transcription PCR for subtyping avian influenza A(H5) virus*

Target	Name	Sequence, 5' → 3'	Final concentration
A(H5)	FluA_H5_v4_1F	TACCAGATACTGTCAATTTATTCAAC	400 nM
	FluA_H5_v4_1R	GTAACGACCCATTGGAGCACATCC	400 nM
	FluA_H5_v4_1Prb.FAM	FAM-CTGGCAATC/ZEN/ATGRTRGCTGGTCT-3IABkFQ	200 nM
	FluA_H5_v4_2F	TGGGTACCATCATAGCAATGAGCA	400 nM
	FluA_H5_v4_2R	AACTCCCTCCAAGTGCCTCAA	400 nM
	FluA_H5_v4_2Prb.FAM	FAM-TGGGTACGC/ZEN/TGCGGACAAAGAATCCA-3IABkFQ	200 nM
A (M)	Pan-FluA-F	GACCRATCCTGTACCTCTGAC	400 nM
	Pan-FluA-R	AGGGCATTYTGACAAAKCGTCTA	400 nM
	Pan-FluA-prb_Q705	Q705-TGCAGTCTCGTCACTGGGCACG-BHQ-3	200 nM
Human RNase P	RNase P Fwd	AGATTTGGACCTGCCGAGCG	100 nM
	RNase P Rev	GAGCGCTGTCTCCACAAGT	100 nM
	RNase P Probe CF560	CF560-TTCTGACCTGAAGGCTCTGCGCG-BHQ-1	50 nM

*H5 probes were purchased from Integrated DNA Technologies (<https://www.idtdna.com>). The probes contain the internal quencher, ZEN, in addition to the 3' Iowa BHQ FQ (3IABkFQ) (proprietary to Integrated DNA Technologies). Influenza A (M) and human RNase P probes were purchased from Biosearch Technologies (<https://www.biosearchtech.com>). Bold type indicates mixed bases. BHQ, Black Hole Quencher; CF560, CalFluor 560; FAM, 6-carboxyfluorescein; M, matrix gene; Q705, Quasar 705.

A-negative eluates extracted from clinical upper respiratory specimens. We then performed the H5 subtyping qRT-PCR (Appendix). We tested 20 replicates at each level for both mixes. The 95% LLOD for the H5 target was 2.5 copies/μL (95% CI 1.8–5.3 copies/μL) for the clade 1 ssDNA mix and <0.5 copies/μL (we were unable to calculate 95% CI) for the clade 2.3.4.4b ssDNA mix.

We also evaluated the H5 subtyping multiplex qRT-PCR by using genomic RNA from a clade Eurasian non-goose/GuangDong (Gs/Gd) isolate, Kilbourne F181, A/duck/Singapore/645/1997 (H5N3). We assigned this genomic RNA an estimated concentration in copies/μL by using the clade 2.3.4.4b ssDNA as standard curve, then diluted it to 2, 1, and 0.5 copies/μL in pooled influenza A-negative eluates. We tested 20 replicates at each level. The 95% LLOD for the H5 target was 0.6 copies/μL (95% CI 0.5–0.7 copies/μL). In addition, we tested 10-fold serial dilutions from 4.6 to 0.6 log₁₀ copies/mL of the Kilbourne F181 genomic RNA in duplicate (Appendix Figure), which indicated 0.97 reaction efficiency for the H5 target.

We then evaluated the H5 subtyping qRT-PCR by using genomic RNA from additional H5 viruses, including 5 low pathogenicity avian influenza non-

Gs/Gd isolates from the United States (Table 2). We tested each eluate in duplicate. Although we detected H5 RNA in all reactions, the difference in mean cycle threshold (Ct) values between the H5 and pan-influenza A (M gene) targets varied widely (range 0.6–9.9 cycles). Testing the H5 primer-probe sets individually revealed that the US non-Gs/Gd genomic RNAs were detected only by primer-probe set 1. The A/mallard/Minnesota/16-041335-3/2016 (H5N2) eluate showed the greatest Ct difference, consistent with the highest number of H5 primer-probe mismatches in primer-probe set 1 (9 mismatches) (Appendix Tables 5, 6).

To determine specificity, we tested genomic RNA from cultured human and avian virus isolates encoding non-H5 HA genes (Table 3). We detected the pan-influenza A target in all eluates; median Ct value was 17.3 (range 13.6–20.7). We did not detect the H5 target in any eluates.

We performed further specificity experiments by using residual upper respiratory swab samples in viral transport media submitted for clinical respiratory virus testing. We conducted this work with Stanford Institutional Review Board approval (protocol no. 68234); individual consent was waived. We evaluated 100 samples collected

Table 2. Avian influenza A(H5) virus genomic RNA used to evaluate inclusivity in study of multiplex dual-target reverse transcription PCR for subtyping avian influenza A(H5) virus*

Genome	Clade	GenBank accession no.	H5 Ct	M Ct	Ct difference
Kilbourne F181, A/duck/Singapore/645/1997 (H5N3)	EA-non-Gs/Gd	NA	18.3	17.7	0.6
A/quail/California/14-012546-1/2014 (H5N8)	Am-non-Gs/Gd	NA	22.9	19.0	3.9
A/mallard/Minnesota/16-041335-3/2016 (H5N2)	Am-non-Gs/Gd	MH546659	32.1	22.3	9.9
A/emperor goose/Alaska/17-004479-1/2016 (H5N2)	Am-non-Gs/Gd	MH546451	26.8	23.3	3.6
A/glaucous-winged gull/Alaska/16-041335-19/2016 (H5N2)	Am-non-Gs/Gd	MH546475	23.3	21.8	1.5
A/northern pintail/Alaska/16-041335-5/2016 (H5N2)	Am-non-Gs/Gd	MH546883	20.9	19.1	1.9

*Kilbourne F181 genomic RNA was obtained from BEI Resources (<https://www.beiresources.org>). Am-non-Gs/GD A(H5) genomic RNA was obtained from the US Department of Agriculture. The genomic RNA from these influenza A(H5) viruses was diluted 1:10 buffer AVE plus carrier RNA (QIAGEN, <https://www.qiagen.com>) and tested in duplicate. Ct values are means. Am-non-Gs/Gd, non-goose/GuangDong from the United States; Ct, cycle threshold; EA-non-Gs/Gd, Eurasian non-goose/GuangDong; M, matrix gene; NA, not available.

Table 3. Non-H5 human and avian influenza A virus genomic RNA used to evaluate H5 specificity in study of multiplex dual-target reverse transcription PCR for subtyping avian influenza A(H5) virus*

HA gene	Genome	BEI catalog no.
H1	A/Brisbane/59/2007 (H1N1)	NR-20080
H1 pdm09	A/California/04/2009 (H1N1)pdm09	NR-14689
H2	A/duck/Germany/1215/1973 (H2N3)	NR-2762
H3	A/Brisbane/10/2007 (H3N2)	NR-20081
H4	A/duck/Czechoslovakia/1956 (H4N6)	NR-43012
H6	A/shearwater/Australia/1/1973 (H6N5)	NR-43014
H8	A/turkey/Ontario/6118/1968 (H8N4)	NR-43015
H9	A/turkey/Wisconsin/1/1966 (H9N2)	NR-43016
H10	A/chicken/Germany/N/1949 (H10N7)	NR-2765
H11	A/duck/Memphis/546/1974 (H11N9)	NR-43017
H12	A/duck/Alberta/60/1976 (H12N5)	NR-43018
H13	A/gull/Maryland/704/1977 (H13N6)	NR-43019
H14	A/mallard/Astrakhan/263/1982 (H14N5)	NR-43020
H15	A/shearwater/Australia/2576/1979 (H15N9)	NR-43021
H16	A/shorebird/Delaware/172/2006 (H16N3)	NR-43022

*All genomic RNA obtained from BEI Resources (<https://www.beiresources.org>). The genomic RNA from these influenza A (H5) viruses was diluted 1:10 in buffer AVE plus carrier RNA (QIAGEN, <https://www.qiagen.com>). HA, hemagglutinin; pdm09, pandemic 2009.

during the 2022–2023 respiratory virus season that were positive for influenza A by using routine clinical methods (98 by using Hologic Panther Fusion [Hologic, <https://www.hologic.com>] and 2 by using Cepheid GeneXpert [Cepheid, <https://www.cephid.com>]), including 29 known co-infections (13 respiratory syncytial virus [RSV], 7 rhinovirus, 3 adenovirus, 3 human metapneumovirus, 1 parainfluenza-3, 1 parainfluenza 4, and 1 with both RSV and rhinovirus). In addition, we tested 55 upper respiratory samples collected in April 2024 that were negative for influenza A, influenza B, RSV, parainfluenza virus 1–4, rhinovirus, adenovirus, and human metapneumovirus on the Panther Fusion by using the H5 subtyping qRT-PCR. We extracted total nucleic acids from 400 μ L and eluted them in 60 μ L buffer AVE on the BioRobot EZ1 (Qiagen) by using the EZ1 virus mini kit 2.0, according to the manufacturer's recommendations. We detected the pan-influenza A target in 97.0% (97/100) of the influenza A-positive samples; median Ct value was 24.1 (range 12.6–37.7). All 3 of the samples that were not detected in the pan-influenza A channel of the H5 multiplex had detectable RNase P (Ct values of 22.5, 18.7, and 20.8), indicating adequate extraction and the absence of inhibitors. The original influenza A Ct values (Panther Fusion) for those 3 samples were 41.4, 38.0, and 38.2. Upon repeat testing, influenza A RNA was detected only in the sample with the lowest original Ct value for influenza A (C_t 34.8) by the H5 multiplex. None of those influenza A-positive samples had detectable H5 RNA. Of the respiratory virus panel negative samples, all had detectable RNase P; median Ct value was 23.9 (range 17.7–29.8). None of the panel negative samples had detectable H5 RNA.

Conclusions

HPAI H5 viruses, particularly clade 2.3.4.4b viruses, pose an emerging pandemic threat. This internally controlled dual-target influenza A(H5) qRT-PCR demonstrated high analytical performance and could be used to directly test samples from patients with suspected HPAI A(H5) virus infection or as a secondary test to subtype known influenza A-positive samples after routine respiratory virus testing. The dual-target design reduces the likelihood of false negatives, as evidenced by detection of low pathogenicity avian influenza H5 viruses with substantial target mismatch. However, continuous sequence surveillance and updating of the primer-probe sets will be required to ensure that this assay accounts for ongoing viral evolution (13).

Acknowledgments

We thank Erica Spackman for providing genomic RNA from US avian influenza A(H5) non-goose/GuangDong isolates.

We obtained additional influenza A genomic RNA through the Biodefense and Emerging Infections Research Resources Repository, National Institute for Allergy and Infectious Diseases, National Institutes of Health.

About the Author

Dr. Sahoo is senior bioinformatic staff scientist in the Stanford Health Care Clinical Virology Laboratory. His research involves the design and evaluation of nucleic acid amplification and sequencing assays for the diagnosis and characterization of infectious diseases.

References

1. Morens DM, Park J, Taubenberger JK. Many potential pathways to future pandemic influenza. *Sci Transl Med*. 2023; 15:eadj2379. <https://doi.org/10.1126/scitranslmed.adj2379>

2. Sun Y, Zhang T, Zhao X, Qian J, Jiang M, Jia M, et al. High activity levels of avian influenza upwards 2018–2022: a global epidemiological overview of fowl and human infections. *One Health*. 2023;16:100511. <https://doi.org/10.1016/j.onehlt.2023.100511>
3. Szablewski CM, Iwamoto C, Olsen SJ, Greene CM, Duca LM, Davis CT, et al. Reported global avian influenza detections among humans and animals during 2013–2022: comprehensive review and analysis of available surveillance data. *JMIR Public Health Surveill*. 2023;9:e46383. <https://doi.org/10.2196/46383>
4. Kniss K, Sumner KM, Tastad KJ, Lewis NM, Jansen L, Julian D, et al. Risk for infection in humans after exposure to birds infected with highly pathogenic avian influenza A(H5N1) virus, United States, 2022. *Emerg Infect Dis*. 2023;29:1215–9. <https://doi.org/10.3201/eid2906.230103>
5. Gilbertson B, Subbarao K. Mammalian infections with highly pathogenic avian influenza viruses renew concerns of pandemic potential. *J Exp Med*. 2023;220:e20230447. <https://doi.org/10.1084/jem.20230447>
6. Puryear W, Sawatzki K, Hill N, Foss A, Stone JJ, Doughty L, et al. Highly pathogenic avian influenza A(H5N1) virus outbreak in New England seals, United States. *Emerg Infect Dis*. 2023;29:786–91. <https://doi.org/10.3201/eid2904.221538>
7. Agüero M, Monne I, Sánchez A, Zecchin B, Fusaro A, Ruano MJ, et al. Highly pathogenic avian influenza A(H5N1) virus infection in farmed minks, Spain, October 2022. *Euro Surveill*. 2023;28:2300001. <https://doi.org/10.2807/1560-7917.ES.2023.28.3.2300001>
8. Burrough ER, Magstadt DR, Petersen B, Timmermans SJ, Gauger PC, Zhang J, et al. Highly pathogenic avian influenza A(H5N1) clade 2.3.4.4b virus infection in domestic dairy cattle and cats, United States, 2024. *Emerg Infect Dis*. 2024;30:1335–43. <https://doi.org/10.3201/eid3007.240508>
9. Uyeki TM, Milton S, Abdul Hamid C, Reinoso Webb C, Presley SM, Shetty V, et al. Highly pathogenic avian influenza A(H5N1) virus infection in a dairy farm worker. *N Engl J Med*. 2024;390:2028–9. <https://doi.org/10.1056/NEJMc2405371>
10. World Health Organization. WHO information for molecular diagnosis of influenza virus – update. 2021 Feb 7 [cited 2023 Jun 25]. <https://www.who.int/teams/global-influenza-programme/laboratory-network/quality-assurance/eqa-project/information-for-molecular-diagnosis-of-influenza-virus>
11. Huang C, Sahoo MK, Verghese M, Sibai M, Solis D, Mfuh KO, et al. Interepidemic respiratory syncytial virus during the COVID-19 pandemic. *Microbiol Spectr*. 2022;10:e0094722. <https://doi.org/10.1128/spectrum.00947-22>
12. Jiao P, Yuan R, Song Y, Wei L, Ren T, Liao M, et al. Full genome sequence of a recombinant H5N1 influenza virus from a condor in southern China. *J Virol*. 2012;86:7722–3. <https://doi.org/10.1128/JVI.01043-12>
13. Kandeil A, Patton C, Jones JC, Jeevan T, Harrington WN, Trifkovic S, et al. Rapid evolution of A(H5N1) influenza viruses after intercontinental spread to North America. *Nat Commun*. 2023;14:3082. <https://doi.org/10.1038/s41467-023-38415-7>

Address for correspondence: Benjamin A. Pinsky, Stanford University School of Medicine, 3375 Hillview Ave, Palo Alto, CA 94304, USA; email bpinsky@stanford.edu

EID Podcast

Heartland Virus from Lone Star Ticks, Georgia, USA, 2019

Heartland virus is an emerging infectious disease that is not well understood. A report of a human case and exposure of white-tailed deer to Heartland virus in Georgia prompted the sampling of questing ticks during 2018–2019. With the confirmation that Heartland virus is actively circulating in locally infected ticks in Georgia, clinicians should be alerted to the presence of this emerging tickborne virus.

In this EID podcast, Dr. Gonzalo Vazquez-Prokopec, an associate professor of environmental sciences at Emory University in Atlanta, discusses the presence of Heartland virus in lone star ticks in Georgia.

Visit our website to listen: <https://go.usa.gov/xy6UH> **EMERGING INFECTIOUS DISEASES®**

Article DOI: <https://doi.org/10.3201/eid3008.240785>

EID cannot ensure accessibility for supplementary materials supplied by authors. Readers who have difficulty accessing supplementary content should contact the authors for assistance.

Multiplex Dual-Target Reverse Transcription PCR for Subtyping Avian Influenza A(H5) Virus

Appendix

Methods

We performed reverse transcription quantitative PCR (RT-qPCR) by using the Invitrogen Superscript III Platinum One-Step qRT-PCR kit (Thermo Fisher Scientific, <https://www.thermofisher.com>) on the Rotor-Gene Q real-time PCR instrument (QIAGEN, <https://www.qiagen.com>). Each 25- μ L reaction contained 12.5 μ L of 2X buffer, 0.5 μ L of enzyme mix, 2 μ L of primer–probe mix, and 10 μ L of nucleic acids. Cycling conditions were as follows: hold at 52°C for 15 min, 94°C for 2 min, then 45 cycles of 94°C for 15 sec, 55°C for 40 sec, and 68°C for 20 sec. We analyzed the fluorescence signal in the green (H5), crimson (pan-influenza A), and yellow (RNase P) channels. We set thresholds at 0.1 normalized fluorescence units for all targets. We interpreted any exponential amplification curve crossing the green channel as subtype H5. The 95% lower limit of detection was determined in R (<https://www.r-project.org>) by using probit regression.

Appendix Table 1. Original Influenza A(H5) Primer and Probe Sequences

Name	Sequence (5'→3')	Source	Notes
H5.2344-1673F	TACCAAATAYTGTCAATTTATTCAAC	School of Public Health, Hong Kong University, Hong Kong SAR, China	A (H5) clade 2.3.4.4
H5.2344-1749R	GTAAYGACCCRTTRGARCACATCC		
H5.2344-1718Prb FAM	CTGGCAATCAT DRT GGCTGGTCT		
H5-1012F	TGGGTACCACCATAGCAATGAGCA	Centre for Health Protection, Hong Kong SAR, China	A (H5)
H5-1155R	AATCCCTTCCAACGGCCTCAAAC		
H5-1042P-P2_CFO560	TGGGTACGCTGCAGACAAAGAATCCA		

SAR, Special Administrative Region. Mixed bases are in bold

Appendix Table 2. Alignment of Primer/Probe Set 1 to North American Influenza A(H5) Hemagglutinin (HA) Sequences available from GISAID January 1, 2022 to May 29, 2024.

Reference	Forward Primer	Probe	Reverse Primer*	No.	%
Set 1	TACCAGATACTGTCAATTTATTCAAC	CTGGCAATCATG TR RGCTGGTCT	GGATGTGCTCCAATGGGTCGTTAC		
	TACCAGATACTGTCAATTTATTCAAC	CTGGCAATCATG TR RGCTGGTCT	GGATGTGCTCCAATGGGTCGTTAC	5529	92.4
C.....	82	1.4
T.....	58	1.0
G.....G.....	44	0.7
G.....	39	0.7
T.....	36	0.6
A.....	25	0.4
T.....	24	0.4
A.....A.....	21	0.4
A.....	13	0.2
T.....	11	0.2
A.....	8	0.1
T.....	8	0.1
A.....	8	0.1
C.....	7	0.1
G.....	6	0.1
G.....C.....	6	0.1
G.....	5	0.1
T.....	5	0.1
T.....	4	0.1
T.....	4	0.1
A.....	4	0.1
T.....	4	0.1
	T.....	4	0.1
T.....T.....	3	0.1

*The reverse primer sequence is presented as its reverse complement.

Sequences accessed May 29, 2024. Aligned sequences: 5987. GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference.

Primer and probe alignments are in-phase. Top 25 most abundant sequences are listed. Mixed bases are in bold.

Appendix Table 3. Alignment of Primer/Probe Set 2 to North American Influenza A(H5) Hemagglutinin (HA) Sequences available from GISAID January 1, 2022 to May 29, 2024.

Reference	Forward Primer	Probe	Reverse Primer*	No.	%
Set 2	TGGGTACCATCATAGCAATGAGCA	TGGGTACGCTGCGGACAAAGAATCCA	TTTGAGGCAGTTGGAAGGGAGTT	5085	84.9
	209	3.5
T.....A.....	93	1.6
	..A.....	74	1.2
T.....	48	0.8
C.....C.....	45	0.8
A.....	43	0.7
T.....	41	0.7
A.....	32	0.5
T.....	26	0.4
A.....	17	0.3
T.....G.....	15	0.3
	..T.....	15	0.3
T.....	14	0.2
A.....	13	0.2
C.....	12	0.2
T.....	12	0.2
A.....G.....	10	0.2
C.....	9	0.2
A.....	9	0.2
A.....	9	0.2
T.....	8	0.1
G.....	8	0.1
A.....	8	0.1
G.....	8	0.1

*The reverse primer sequence is presented as its reverse complement.
 Sequences accessed May 29, 2024. Aligned sequences: 5990. GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference.
 Primer and probe alignments are in-phase. Top 25 most abundant sequences are listed.

Appendix Table 4. Alignment of Primer/Probe Set 1 to Avian Influenza A(H5) Sequences from Human Cases and Dairy Cows.

Reference	Forward Primer	Probe	Reverse Primer*
Set 1	TACCAGATACTGTCAATTTATTCAAC	CTGGCAATCATGATGGCTGGTCT	GGATGTGCTCCAATGGGTCGTTAC
Human: A/Texas/37/2024
Human: A/Michigan/90/2024
Human: A/Michigan/91/2024
Human: A/Colorado/18/2022
A/dairy cow/Texas/24_009367-003/2024G.....
A/dairy cow/Texas/24_009367-005/2024G.....
A/dairy cow/Texas/24_009367-009/2024G.....
A/dairy cow/Texas/24_009367-010/2024G.....
A/dairy cow/Texas/24_009367-013/2024G.....
A/dairy cow/Texas/24_009367-014/2024G.....
Other 190 sequences from dairy cows

*The reverse primer sequence is presented as its reverse complement.
 GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference. Mixed bases are in bold. Primer and probe alignments are in-phase.
 Note: The 6 dairy cow sequences showing a mismatch with primer/probe set 1 do not contain any mismatches with primer/probe set 2.

Appendix Table 5. Alignment of Primer/Probe Set 2 to Avian Influenza A(H5) Sequences from Human Cases and Dairy Cows.

Reference	Forward Primer	Probe	Reverse Primer*
Set 2	TGGGTACCATCATAGCAATGAGCA	TGGGTACGCTGCCGACAAAGAATCCA	TTTGAGGCAGTTGGAAGGGAGTT
Human: A/Texas/37/2024
Human: A/Michigan/90/2024
Human: A/Michigan/91/2024
Human: A/Colorado/18/2022
A/dairy cow/Texas/24_009110-003/2024A.....
A/dairy cow/South Dakota/24_010354-001/2024A.....
A/dairy cow/Texas/24_009776-002/2024T.....
Other 193 sequences from dairy cows

*The reverse primer sequence is presented as its reverse complement.
 GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference. Primer and probe alignments are in-phase.
 Note: The 3 dairy cow sequences showing a mismatch with primer/probe set 2 do not contain any mismatches with primer/probe set 1.

Appendix Table 6. Single-Stranded DNA Sequences

Name	H5 Target Sequence (5'→3')	GenBank Reference	Location
ssD_FluA_H5_T1C1	GTATGACTACCCGCAGTATTCAGAAGAAGCAAGACTAAAAAGAGAGGAAATAAGTGGAGTAAAATTGGAATCAATAGGAACT <u>TACCAA</u> AACTGTCAATTTATTCTACAGTGGCGAGTCCCTAGCACTGGCAATCATGGTAGCTGGTCTATCTTTATGGATG <u>TGCTCCAATGGGTCGTTACAATGCAGAAAT</u>	JQ966928	1534–1725
ssD_FluA_H5_T2C1	ATGGTTGGTATGGGTACCACCATAGCAATGAACAGGGGAGTGGGTACGCTGCAGACAAAGAATCCACTCAAAGGCAAT AGATGGAGTACCAATAAGGTCAACTCGATCATTGACAAAATGAACACTCAGTTT <u>GAGGCCGTTGGAAGGGA</u> TTTAATA ACTTA	JQ966928:	1122–1285
ssD_FluA_H5_T1C2	AGTAGGAACTTACCAGATACTGTCAATTTATTCAACAGCGGCAAGTCCCTAGCACTGGCAATCATGATGGCTGGTCTAT CTTTATGGATGTGCTCCAATGGGTCGTTACAGTGCAGAAAT	OP499866	1591–1710
ssD_FluA_H5_T2C2	ATGGTTGGTATGGGTACCATCATAGCAATGAGCAGGGGAGTGGGTACGCTGCGGACAAAGAATCCACCCAAAAGGCAAT AGATGGAGTTACCAATAAGGTCAACTCAATCATTGACAAAATGAACACTCAATTTGAGGCAGTTGGAAGGGAGTTAATAA CTTA	OP499866	1107–1270

Primer and probe targets are underlined. Mismatched bases are in bold. Single-stranded DNA was synthesized by Integrated DNA Technologies.

Appendix Table 7. Alignment of Primer/Probe Set 1 to Avian Influenza A (H5) American-non Goose GuangDong Sequences

Reference	Forward Primer	Probe	Reverse Primer*
Set 1	TACCAGATACTGTCAATTTATTCAAC	CTGGCAATCATGATGGCTGGTCT	GGATGTGCTCCAATGGGTCGTTAC
A/mallard/Minnesota/16–041335–3/2016 (H5N2)	. . T T . A C . . CG A T A . . G . .
A/emperor goose/Alaska/17–004479–1/2016 (H5N2)	. . T A C . . C A A A . . G . .
A/glaucous-winged gull/Alaska/16–041335–19/2016 (H5N2)	. . T A C A A A . . G . .
A/northern pintail/Alaska/16–041335–5/2016 (H5N2)	. . T A C A A A . . G . .

*The reverse primer sequence is presented as its reverse complement.


GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference. Mixed bases are in bold.

Appendix Table 8. Alignment of Primer/Probe Set 2 to Avian Influenza A(H5) American-non Goose GuangDong Sequences

Reference	Forward Primer	Probe	Reverse Primer*
Set 2	TGGGTACCATCATAGCAATGAGCA	TGGGTACGCTGCGGACAAAGAATCCA	TTTGAGGCAGTTGGAAGGGAGTT
A/mallard/Minnesota/16–041335–3/2016 (H5N2)	C A . . T A T C . . A . . T G . AA . . A . .
A/emperor goose/Alaska/17–004479–1/2016 (H5N2)	C . . A A . . T A T C . . A . . T AA . . A . .
A/glaucous-winged gull/Alaska/16–041335–19/2016 (H5N2)	C . . A A . . T A T C . . A . . C G . AA . . A . .
A/northern pintail/Alaska/16–041335–5/2016 (H5N2)	C . . A A . . T A T C . . A . . C G . AA . . A . .

*The reverse primer sequence is presented as its reverse complement.

GenBank: PP577943/GISAID: EPI3171488 (A/Texas/37/2024) is used as reference. Primer and probe alignments are in-phase.



Appendix Figure. Ten-fold serial dilutions of Kilbourne F181, A/duck/Singapore/645/1997 (H5N3) (BEI Resources Catalog # NR-9682) genomic RNA from 4.6 to 0.6 log₁₀ copies/mL were tested in duplicate using the H5 subtyping multiplex RT-qPCR. Filled circles, individual replicates. Solid line, simple linear regression (GraphPad Prism 10 for macOS, version 10.2.3).