Table S2. Prevalence of lifetime DSM-IV disorders among suicidality in developing countries

	Among total sample						Among total sample						Among Ideators						Among ideators with a lifetime plan						Among ideators without a lifetime plan					
	% with disorder among						% with disorder among						% with disorder among						% with disorder among						% with disorder among					
	Ideation			No ideation			Attempt			No attempt			Plan			No plan			Attempt			No attempt			Attempt			No attempt		
	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE	N[2]	%[3]	SE
I. Anxiety Disorders																														
Panic Disorder[4]	85	2.0	0.3	442	0.7	0.1	40	3.1	0.6	532	0.8	0.1	48	3.1	0.5	81	2.8	0.4	35	4.0	0.8	20	3.1	0.6	5	1.3	0.6	70	2.8	0.4
General Anxiety Disorder[4]	129	3.2	0.4	826	1.4	0.1	63	5.0	0.7	1013	1.7	0.1	64	4.8	0.7	159	5.7	0.6	41	5.5	1.0	64	11.7	1.7	22	4.0	1.0	123	5.3	0.7
Specific Phobia[4]	571	13.7	0.7	2689	5.1	0.1	248	19.3	1.4	3051	5.5	0.1	282	18.8	1.5	326	12.4	0.8	188	21.4	1.9	102	17.4	1.9	60	15.0	2.2	260	11.7	0.8
Social Phobia[4]	250	6.2	0.5	839	1.6	0.1	105	8.7	1.0	1021	1.9	0.1	119	8.7	1.0	158	6.0	0.5	76	9.3	1.3	56	10.4	1.7	29	7.7	1.6	126	5.6	0.6
Obsessive-Compulsive Disorder[9]	102	3.3	0.5	513	1.3	0.1	62	5.8	0.8	624	1.5	0.1	66	5.0	0.7	93	5.1	0.7	48	6.8	1.1	39	6.8	1.4	14	3.7	1.3	72	5.0	0.8
Post-Traumatic Disorder[5]	81	2.8	0.4	447	1.5	0.1	27	3.3	0.8	548	1.7	0.1	36	3.4	0.8	90	4.9	0.6	22	3.6	1.0	24	5.2	1.6	5	2.6	1.5	77	4.9	0.7
Separation Anxiety Disorder[5]	205	6.3	0.5	774	2.6	0.1	93	8.4	1.1	940	2.9	0.1	100	7.2	0.8	160	8.8	0.9	58	7.7	1.3	52	8.5	1.4	35	9.8	2.2	114	7.8	1.0
Agoraphobia[4]	119	2.6	0.3	636	1.2	0.1	55	3.7	0.6	733	1.3	0.1	58	3.7	0.6	92	3.1	0.4	38	3.9	0.7	25	4.7	1.2	17	3.4	1.0	72	2.9	0.4
Any Anxiety Disorder[5]	1020	27.1	0.9	4661	12.0	0.2	439	36.3	1.8	5450	13.0	0.2	512	35.3	1.7	696	30.4	1.3	323	39.9	2.6	242	40.3	2.7	116	29.3	2.9	547	29.2	1.4
II. Mood Disorders																														
Major Depression[4]	560	12.9	0.6	3846	6.8	0.2	243	18.2	1.3	4679	7.9	0.2	306	20.4	1.3	666	24.9	1.0	187	21.6	1.7	272	43.0	2.6	56	11.5	1.7	561	24.6	1.1
Dysthymia[4]	118	2.8	0.3	567	0.9	0.1	60	4.6	0.8	690	1.1	0.1	63	4.0	0.6	106	4.1	0.5	45	5.3	1.0	40	6.4	1.1	15	3.3	1.0	83	3.9	0.5
Bipolar[4]	64	1.7	0.3	350	0.8	0.1	30	2.4	0.5	450	0.9	0.1	44	3.3	0.7	74	2.9	0.4	27	3.3	0.7	38	7.2	1.4	3	0.7	0.4	62	2.8	0.5
Any Mood[5]	607	13.9	0.6	4162	7.5	0.2	260	19.4	1.3	5052	8.6	0.2	333	22.0	1.3	706	26.4	0.9	203	23.3	1.8	293	46.5	2.6	57	11.6	1.7	597	26.1	1.0
III. Impulse-Control Disorders																														
ODD[7]	79	3.0	0.5	146	0.5	0.1	47	5.0	1.0	179	0.6	0.1	46	4.6	0.9	36	2.2	0.4	35	5.3	1.3	10	3.5	1.3	12	4.2	1.4	23	1.8	0.5
Conduct[7]	86	3.2	0.5	137	0.5	0.1	53	5.9	1.1	174	0.6	0.1	48	4.8	0.9	42	2.6	0.6	38	6.8	1.4	10	2.3	0.9	15	4.2	1.3	27	2.3	0.6
ADD[7]	80	2.5	0.4	148	0.5	0.1	40	3.7	0.8	188	0.6	0.1	44	3.0	0.5	36	2.1	0.5	29	3.5	0.8	15	2.4	0.7	11	4.2	1.5	25	1.8	0.5
IED[4]	250	6.5	0.5	917	2.0	0.1	105	8.8	1.1	1148	2.3	0.1	128	9.8	1.0	210	8.2	0.7	74	9.7	1.3	72	12.6	1.7	31	7.1	1.5	159	7.5	0.7
Any Impulse[7]	358	13.2	1.0	968	3.8	0.2	172	19.0	1.8	1218	4.3	0.2	196	19.7	1.5	235	13.7	1.1	125	20.8	2.3	78	18.8	2.6	47	15.3	2.4	172	12.7	1.2
IV. Substance Use Disorders																														
Alcohol Abuse or Dependence[4]	267	7.9	0.6	2438	5.6	0.2	106	10.5	1.1	2757	6.0	0.2	137	11.1	1.2	282	13.1	0.9	79	10.6	1.4	103	20.4	2.2	27	10.3	2.3	216	11.8	1.0
Drug Abuse or Dependence[8]	90	3.1	0.4	332	1.1	0.1	52	4.9	0.7	421	1.4	0.1	52	4.6	0.7	76	5.3	0.7	44	5.8	0.9	31	8.8	2.5	8	3.0	1.1	58	5.0	0.8
Any Substance Disorder[8]	292	9.7	0.7	2107	6.7	0.2	128	13.3	1.3	2433	7.3	0.2	161	13.6	1.4	281	15.9	1.2	99	14.0	1.6	116	25.0	3.1	29	11.8	2.6	210	14.5	1.3
Any Disorder[10]	1611	42.9	1.1	8703	21.8	0.3	666	54.6	2.0	10112	23.6	0.3	804	55.5	1.7	1219	53.0	1.5	489	59.6	2.4	434	69.6	2.7	177	44.6	3.2	975	51.5	1.7
V. Number of Disorders																														
Exactly 1[10]	852	23.3	0.9	5408	14.4	0.3	311	25.7	1.6	6034	15.0	0.3	385	25.6	1.4	546	24.9	1.4	225	27.0	1.9	178	25.1	2.5	86	23.3	2.9	448	24.9	1.5
Exactly 2[10]	402	10.6	0.8	1980	4.6	0.2	163	13.5	1.5	2380	5.1	0.2	213	15.8	1.4	339	14.0	0.9	123	15.9	2.0	120	20.2	1.9	40	8.7	1.6	280	13.7	1.0
Exactly 3[10]	180	4.7	0.5	795	1.8	0.1	82	6.5	0.9	985	2.1	0.1	92	5.9	0.7	175	7.5	0.8	55	7.0	1.2	63	10.1	1.6	27	5.6	1.3	127	7.1	0.9
Exactly 4[10]	88	2.2	0.3	311	0.6	0.1	52	4.3	0.7	403	0.8	0.1	55	4.2	0.7	75	3.7	0.6	41	4.7	0.9	35	6.1	1.4	11	3.6	1.3	57	3.3	0.5
Exactly 5[10]	44	1.1	0.2	127	0.2	0.0	28	1.9	0.4	175	0.3	0.0	30	2.3	0.5	39	1.3	0.3	22	2.4	0.6	20	4.3	1.5	6	1.0	0.5	28	1.1	0.3
6 or more disorders[10]	45	1.0	0.2	82	0.2	0.0	30	2.6	0.6	135	0.3	0.0	29	1.8	0.4	45	1.7	0.3	23	2.7	0.7	18	3.9	1.4	7	2.4	1.0	35	1.4	0.3
(N)[1]	(4090)			(52803)			(1223)			(55670)			(1443)			(2647)			(835)			(608)			(388)			(2259)		

[1] Number of cases with the outcome variable

[2] Numerator N. Number of cases with the LT disorder among cases with the outcome variable. Columns with "Yes" in the headers represents cases with the dx among those with the outcome, and columns with "No" represents among those without the outcome. Part II disorders assessed in part II sample, and disorders with age restrictions restricted to appropriate age range

[3] % represents the percentage of people with the DSM-IV disorder among the cases with the outcome variable indicated in the column header. For example: the first cell is the % of those with Panic disorders among those with attempts

[4] assessed in part I sample

[5] assessed in part II sample

[6] assessed in part II sample. Restricted to age groups 18-44 for Lebanon and India, 18-39 for China and Nigeria, and not restricted for other countries.

[7] assessed in part II sample and restricted to age groups 18-44 for Mexico, Colombia, Bulgaria, India, 18-39 for China, Nigeria, Ukraine, and not restricted for Brazil and Romania.

[8] assessed in part II sample for all data except Colombia and Mexico, where part I weight is used. A weight variable is created that takes the part I weight values for Colombia and Mexico and part II for other countries.

[9] OCD was only assessed in Brazil, Bulgaria, India, Romania, and Shenzhen as there was a problem with the variable in other countries. Assessed in part I sample for Shenzhen and India while in part II sample for Brazil, Bulgaria, and Romania.

[10] assessed in part II sample for all data except for Shenzhen, where part I weight is used.