



Figure: Prevalence of Vaping Brand Used Most Often by Adolescents Who Had Vaped in the Past 30 Days for 8th, 10th, and 12th Grade Students Combined, by Year (95% Confidence Intervals in Parentheses)

* Change in prevalence from previous year is statistically significant at $p < .05$

** Change in prevalence from previous year is statistically significant at $p < .01$

Note: Adolescents who responded “Other” were asked to write in the brand name that they used.

Of those who did so, the % who wrote “Puff Bar” or a spelling variant thereof was 35% in 2020 and 26% in 2021. The prevalence of Puff Bar is therefore estimated as 9% in 2020 (35% of the 27% who reported “Other”) and 12% in 2021 (26% of the 48% who reported “Other”).

Table: Vaping Brand Use Most Often by Adolescents Who Had Vaped in the Past 30 Days:
Prevalence and 95% Confidence Intervals for for 8th, 10th, and 12th Grade Students
Combined, by Year

	----- 2020 -----		----- 2021 -----	
	Prevalence	95% C.I.	Prevalence	95% C.I.
Juul	43.4	(38.2-48.6)	24.9**	(22.0-28.0)
Smok	13.0	(9.3-17.9)	8.4*	(7.0-10.1)
Suorin	6.3	(4-9.8)	2.2**	(1.5-3.3)
Vuse	7.1	(4.9-10.2)	15.3**	(12.6-18.3)
Stig	3.2	(1.9-5.4)	1.3**	(0.9-2.1)
Other	27.0	(23.0-31.5)	47.9**	(44.7-51.1)

* Change in prevalence from previous year is statistically significant at p<.05

** Change in prevalence from previous year is statistically significant at p<.01

Note: Adolescents who responded “Other” were asked to write in the brand name that they used. Of those who did so, the % who wrote “Puff Bar” or a spelling variant thereof was 35% in 2020 and 26% in 2021. The prevalence of Puff Bar is therefore estimated as 9% in 2020 (35% of the 27% who reported “Other”) and 12% in 2021 (26% of the 48% who reported “Other”).