


MONITORING *the* FUTURE

NATIONAL SURVEY RESULTS
ON DRUG USE
1975-2013

2013
Volume 2

College Students & Adults Ages 19–55

Lloyd D. Johnston
Patrick M. O'Malley
Jerald G. Bachman
John E. Schulenberg
Richard A. Miech



MONITORING THE FUTURE
NATIONAL SURVEY RESULTS ON DRUG USE, 1975–2013

Volume 2

College Students and Adults Ages 19–55

by

Lloyd D. Johnston, Ph.D.
Patrick M. O'Malley, Ph.D.
Jerald G. Bachman, Ph.D.
John E. Schulenberg, Ph.D.
Richard A. Miech, Ph.D.

The University of Michigan
Institute for Social Research

Sponsored by:
The National Institute on Drug Abuse
National Institutes of Health

This publication was written by the principal investigators and staff of the Monitoring the Future project, at the Institute for Social Research, the University of Michigan, under Research Grants No. R01 DA 01411 and R01 DA 016575 from the National Institute on Drug Abuse.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the NIH.

Public Domain Notice

All material appearing in this publication is in the public domain and may be reproduced or copied, whether in print or non-print media including derivatives, without permission from the authors. If you plan to modify the material, please contact MTF at mtfinformation@umich.edu for verification of accuracy. Citation of the source is appreciated, including at least the following: Monitoring the Future, Institute for Social Research, University of Michigan.



<http://creativecommons.org/licenses/by/4.0/>

Recommended Citation

Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E. & Miech, R. A. (2014). *Monitoring the Future national survey results on drug use, 1975–2013: Volume 2, College students and adults ages 19–55*. Ann Arbor: Institute for Social Research, The University of Michigan.

Institute for Social Research
The University of Michigan
Ann Arbor, Michigan
Published August 2014

ABBREVIATED CONTENTS*

Click on any item below (in [blue](#)) to go directly to that page.

Detailed Contents	iv
List of Tables	vii
List of Figures.....	viii
Chapter 1 Introduction	1
Chapter 2 Key Findings: An Overview and Integration across Five Populations	6
Chapter 3 Study Design and Procedures	67
Chapter 4 Prevalence of Drug Use in Early and Middle Adulthood	87
Chapter 5 Trends in Drug Use in Early and Middle Adulthood	140
Chapter 6 Attitudes and Beliefs about Drugs among Young Adults	236
Chapter 7 The Social Context	292
Chapter 8 Prevalence of Drug Use among College Students	364
Chapter 9 Trends in Drug Use among College Students	376

*See next page for Detailed Contents.

DETAILED CONTENTS

Click on any item below (in blue) to go directly to that page.

Chapter 1	Introduction	1
	Surveys of Young Adults and Those Ages 35, 40, 45, 50, and 55	2
	Surveys of College Students and Noncollege Peers	3
	General Purposes of the Research.....	4
Chapter 2	Key Findings: An Overview and Integration across Five Populations	6
	Trends in Drug Use—The Advent of Cohort Effects	6
	College–Noncollege Differences in Illicit Drug Use.....	28
	Male–Female Differences in Substance Use	29
	Trends in Alcohol Use	30
	College–Noncollege Differences in Alcohol Use.....	31
	Male–Female Differences in Alcohol Use.....	32
	Trends in Cigarette Smoking	32
	Age- and Cohort-Related Differences in Cigarette Smoking	35
	College–Noncollege Differences in Cigarette Smoking.....	36
	Male–Female Differences in Cigarette Smoking.....	36
	Racial/Ethnic Comparisons.....	37
	Drug Use in Eighth Grade.....	39
	Drug Use by Age 50	41
	Summary and Conclusions	42
Chapter 3	Study Design and Procedures	67
	Research Design and Procedures for the Twelfth-Grade Surveys.....	67
	The Population under Study.....	68
	The Omission of Dropouts.....	68
	Sampling Procedures and Sample Weights	68
	Questionnaire Administration.....	69
	Questionnaire Format.....	69
	Research Design and Procedures for the Eighth- and Tenth-Grade Surveys	70
	Mode of Administration.....	70
	Questionnaire Forms and Sample Proportions	71
	Research Design and Procedures for the Twelfth-Grade Follow-up Surveys	72
	Follow-Up Procedures	72
	Follow-Up Questionnaire Format	73
	Representativeness and Sample Accuracy.....	74
	School Participation	74
	Student Participation.....	76

DETAILED CONTENTS (continued)

Sampling Accuracy of the Estimates	76
Panel Retention	77
The Problem of Panel Attrition.....	77
Response Rates	77
The Impact of Panel Attrition on Research Results.....	79
Effects on Relational Analyses	80
Validity of Measures of Self-Reported Drug Use.....	81
Consistency and Measurement of Trends	83
Chapter 4 Prevalence of Drug Use in Early and Middle Adulthood	87
Replicability of Findings.....	89
A Note on Adjusted Lifetime Prevalence Estimates.....	89
Prevalence of Drug Use across Age Groups.....	90
Prevalence Comparisons for Subgroups of Young Adults	99
Gender Differences	100
Regional Differences	101
Population Density Differences	102
Chapter 5 Trends in Drug Use in Early and Middle Adulthood	140
Recent Trends in Drug Use among Young Adults Ages 19-28.....	141
Longer-term Trends in Early and Middle Adulthood	142
Trends for Important Subgroups of Young Adults	154
Gender Differences in Trends.....	154
Regional Differences in Trends	161
Population Density Differences in Trends	167
Chapter 6 Attitudes and Beliefs about Drugs among Young Adults	236
Perceived Harmfulness of Drugs	236
Personal Disapproval of Drug Use	244
Cohort Differences and Their Implications for Prevention and Theory	248
Chapter 7 The Social Context	292
Peer Norms among Young Adults (Ages 18–30)	293
Current Perceptions of Close Friends’ Attitudes	293

DETAILED CONTENTS (continued)

Trends in Peer Norms	294
Adults' Exposure to Drug Use through Friends and Others (Ages 18–55)	297
Exposure to Drug Use (Ages 18-30).....	297
Trends in Exposure to Drug Use (Ages 18 to 55).....	300
Perceived Availability of Drugs among Adults (Ages 18–55)	304
Perceived Availability.....	304
Trends in Perceived Availability	305
Chapter 8 Prevalence of Drug Use among College Students	364
Prevalence of Drug Use among College Students versus Their Noncollege Peers	365
Gender Differences in Prevalence of Use among College Students.....	368
Chapter 9 Trends in Drug Use among College Students	376
Trends in Prevalence, 1980–2013: College Students versus Those Not Enrolled in College	377
Gender Differences in Trends among College Students.....	387

LIST OF TABLES

Click on any item below (in [blue](#)) to go directly to that page.

Table 2-1.	Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	48
Table 2-2.	Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	53
Table 2-3.	Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	59
Table 2-4.	Trends in 30-Day Prevalence of Daily Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28)	63
Table 3-1.	Sample Sizes and Response Rates	84
Table 4-1.	Prevalence of Use of Various Types of Drugs by Gender among Respondents of Modal Ages 19–30, 2013	105
Table 4-2.	Lifetime Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2013	108
Table 4-3.	Annual Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2013	111
Table 4-4.	Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2013	114
Table 4-5.	Thirty-Day Prevalence of Daily Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2013	117
Table 5-1.	Trends in Lifetime Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	175
Table 5-2.	Trends in Annual Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	177
Table 5-3.	Trends in 30-Day Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	179
Table 5-4.	Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs among Respondents of Modal Ages 19–28	181

LIST OF TABLES (continued)

Table 5-5.	Trends in Annual and 30-Day Prevalence of an Illicit Drug Use Index among Respondents of Modal Ages 19–28, Total and by Gender	183
Table 6-1.	Trends in Harmfulness as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	250
Table 6-2.	Trends in Proportions Disapproving of Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	256
Table 7-1.	Trends in Proportions of Respondents Reporting Their Close Friends Disapproving of Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	310
Table 7-2.	Trends in Friends’ Use of Drugs as Estimated by Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55	312
Table 7-3.	Trends in Exposure to Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	328
Table 7-4.	Trends in Availability of Drugs as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55	334
Table 8-1.	Lifetime Prevalence of Use for Various Types of Drugs, 2013: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School, by Gender.....	371
Table 8-2.	Annual Prevalence of Use for Various Types of Drugs, 2013: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School, by Gender.....	372
Table 8-3.	Thirty-Day Prevalence of Use for Various Types of Drugs, 2013: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School, by Gender.....	373
Table 8-4.	Thirty-Day Prevalence of Daily Use for Various Types of Drugs, 2013: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School, by Gender.....	374
Table 9-1.	Trends in Lifetime Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	390

LIST OF TABLES (continued)

Table 9-2.	Trends in Annual Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	392
Table 9-3.	Trends in 30-Day Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	394
Table 9-4.	Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs among College Students 1 to 4 Years beyond High School	396
Table 9-5.	Trends in Lifetime, Annual, and 30-Day Prevalence of an Illicit Drug Use Index among College Students 1 to 4 Years beyond High School, by Gender.....	398

LIST OF FIGURES

Click on any item below (in [blue](#)) to go directly to that page.

Figure 2-1.	Trends in Annual Prevalence of an Illicit Drug Use Index across 5 Populations.....	66
Figure 3-1.	Schools included in 1 Year’s Data Collection: 8th, 10th and 12th Grades.....	85
Figure 3-2.	School Participation Rates	86
Figure 4-1.	Any Illicit Drug: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	118
Figure 4-2.	Any Illicit Drug other than Marijuana: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	119
Figure 4-3.	Marijuana: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	120
Figure 4-4.	Amphetamines: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	121
Figure 4-5.	Methamphetamine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	122
Figure 4-6.	Crystal Methamphetamine (Ice): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	123
Figure 4-7.	Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	124
Figure 4-8.	Crack Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	125
Figure 4-9.	Other Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	126
Figure 4-10.	Hallucinogens: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	127
Figure 4-11.	LSD: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	128

LIST OF FIGURES (continued)

Figure 4-12.	Hallucinogens other than LSD: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	129
Figure 4-13.	Inhalants: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	130
Figure 4-14.	Sedatives (Barbiturates): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013	131
Figure 4-15.	Narcotics other than Heroin: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	132
Figure 4-16.	Tranquilizers: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	133
Figure 4-17.	Ecstasy (MDMA): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	134
Figure 4-18.	Steroids: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30, by Age Group, 2013.....	135
Figure 4-19.	Heroin: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	136
Figure 4-20a.	Alcohol: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	137
Figure 4-20b.	Alcohol: 2-Week Prevalence of 5 or More Drinks in a Row and 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 55, by Age Group, 2013	138
Figure 4-21.	Cigarettes: Annual, 30-Day, Daily, and Half-Pack-a-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group, 2013.....	139

LIST OF FIGURES (continued)

Figure 5-1.	Any Illicit Drug: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	184
Figure 5-2.	Any Illicit Drug other than Marijuana: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group.....	186
Figure 5-3a.	Marijuana: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group.....	188
Figure 5-3b.	Marijuana: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group.....	190
Figure 5-3c.	Marijuana: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 55, by Age Group.....	192
Figure 5-4.	Inhalants: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32, by Age Group	194
Figure 5-5.	Hallucinogens: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50, by Age Group.....	196
Figure 5-6.	LSD: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 35, by Age Group	198
Figure 5-7.	Hallucinogens other than LSD: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 35, by Age Group.....	200
Figure 5-8.	Ecstasy (MDMA): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32, by Age Group	202
Figure 5-9.	Cocaine: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	204
Figure 5-10.	Crack Cocaine: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	206
Figure 5-11.	Heroin: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	208
Figure 5-12.	Narcotics other than Heroin: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group.....	210

LIST OF FIGURES (continued)

Figure 5-13.	Amphetamines: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	212
Figure 5-14.	Crystal Methamphetamine (Ice): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32, by Age Group	214
Figure 5-15.	Sedatives (Barbiturates): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	216
Figure 5-16.	Tranquilizers: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	218
Figure 5-17.	Steroids: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32, by Age Group	220
Figure 5-18a.	Alcohol: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	222
Figure 5-18b.	Alcohol: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	224
Figure 5-18c.	Alcohol: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 55, by Age Group	226
Figure 5-18d.	Alcohol: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row among Respondents of Modal Ages 18 through 55, by Age Group	228
Figure 5-19a.	Cigarettes: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 55, by Age Group	230
Figure 5-19b.	Cigarettes: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 55, by Age Group	232
Figure 5-19c.	Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among Respondents of Modal Ages 18 through 55, by Age Group	234
Figure 6-1.	Trends in Harmfulness of Marijuana Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice	261
Figure 6-2.	Trends in Harmfulness of Marijuana Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use	262

LIST OF FIGURES (continued)

Figure 6-3.	Trends in Harmfulness of Marijuana Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	263
Figure 6-4.	Trends in Harmfulness of LSD Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	264
Figure 6-5.	Trends in Harmfulness of LSD Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	265
Figure 6-6.	Trends in Harmfulness of PCP Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	266
Figure 6-7.	Trends in Harmfulness of Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	267
Figure 6-8.	Trends in Harmfulness of Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use	268
Figure 6-9.	Trends in Harmfulness of Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	269
Figure 6-10.	Trends in Harmfulness of Crack Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	270
Figure 6-11.	Trends in Harmfulness of Crack Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use	271
Figure 6-12.	Trends in Harmfulness of Crack Cocaine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	272
Figure 6-13.	Trends in Harmfulness of Cocaine Powder Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	273
Figure 6-14.	Trends in Harmfulness of Cocaine Powder Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use.....	274
Figure 6-15.	Trends in Harmfulness of Cocaine Powder Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	275

LIST OF FIGURES (continued)

Figure 6-16.	Trends in Harmfulness of Ecstasy (MDMA) Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice	276
Figure 6-17.	Trends in Harmfulness of Ecstasy (MDMA) Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use.....	277
Figure 6-18.	Trends in Harmfulness of Heroin Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	278
Figure 6-19.	Trends in Harmfulness of Heroin Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Occasional Use	279
Figure 6-20.	Trends in Harmfulness of Heroin Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use	280
Figure 6-21.	Trends in Harmfulness of Amphetamine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice	281
Figure 6-22.	Trends in Harmfulness of Amphetamine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use.....	282
Figure 6-23.	Trends in Harmfulness of Crystal Methamphetamine (Ice) Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice	283
Figure 6-24.	Trends in Harmfulness of Amphetamine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice.....	284
Figure 6-25.	Trends in Harmfulness of Amphetamine Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use.....	285
Figure 6-26.	Trends in Harmfulness of Alcohol Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Trying Once or Twice	286

LIST OF FIGURES (continued)

Figure 6-27.	Trends in Harmfulness of Alcohol Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Having One or Two Drinks per Day	287
Figure 6-28.	Trends in Harmfulness of Alcohol Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Having Four or Five Drinks per Day	288
Figure 6-29.	Trends in Harmfulness of Binge Drinking as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Having Five or More Drinks Once or Twice Each Weekend.....	289
Figure 6-30.	Trends in Harmfulness of Tobacco Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Smoking One or More Packs of Cigarettes per Day	290
Figure 6-31.	Trends in Harmfulness of Smokeless Tobacco Use as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; Regular Use.....	291
Figure 7-1.	Trends in Exposure to Use of Any Illicit Drugs among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	340
Figure 7-2.	Trends in Exposure to Use of Any Illicit Drugs among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	341
Figure 7-3.	Trends in Exposure to Use of Any Illicit Drug other than Marijuana among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	342
Figure 7-4.	Trends in Exposure to Use of Any Illicit Drug other than Marijuana among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	343
Figure 7-5.	Trends in Exposure to Use of Marijuana among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	344
Figure 7-6.	Trends in Exposure to Use of Marijuana among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	345

LIST OF FIGURES (continued)

Figure 7-7.	Trends in Exposure to Use of LSD among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	346
Figure 7-8.	Trends in Exposure to Use of LSD among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	347
Figure 7-9.	Trends in Exposure to Use of Hallucinogens other than LSD among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	348
Figure 7-10.	Trends in Exposure to Use of Hallucinogens other than LSD among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	349
Figure 7-11.	Trends in Exposure to Use of Cocaine among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	350
Figure 7-12.	Trends in Exposure to Use of Cocaine among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	351
Figure 7-13.	Trends in Exposure to Use of Heroin among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	352
Figure 7-14.	Trends in Exposure to Use of Heroin among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	353
Figure 7-15.	Trends in Exposure to Use of Narcotics other than Heroin among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	354
Figure 7-16.	Trends in Exposure to Use of Narcotics other than Heroin among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	355
Figure 7-17.	Trends in Exposure to Use of Amphetamines among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	356
Figure 7-18.	Trends in Exposure to Use of Amphetamines among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	357

LIST OF FIGURES (continued)

Figure 7-19.	Trends in Exposure to Use of Sedatives (Barbiturates) among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	358
Figure 7-20.	Trends in Exposure to Use of Sedatives (Barbiturates) among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed	359
Figure 7-21.	Trends in Exposure to Use of Tranquilizers among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure	360
Figure 7-22.	Trends in Exposure to Use of Tranquilizers among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	361
Figure 7-23.	Trends in Exposure to Use of Alcohol among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Any Exposure.....	362
Figure 7-24.	Trends in Exposure to Use of Alcohol among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30; % Saying Often Exposed.....	363
Figure 9-1.	Any Illicit Drug: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	401
Figure 9-2.	Any Illicit Drug other than Marijuana: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	402
Figure 9-3a.	Marijuana: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	403
Figure 9-3b.	Marijuana: Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	404
Figure 9-4.	Inhalants: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	405

LIST OF FIGURES (continued)

Figure 9-5.	Hallucinogens: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	406
Figure 9-6.	LSD: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	407
Figure 9-7.	Hallucinogens other than LSD: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	408
Figure 9-8.	Ecstasy (MDMA): Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	409
Figure 9-9.	Cocaine: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	410
Figure 9-10a.	Narcotics other than Heroin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	411
Figure 9-10b.	Vicodin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School.....	412
Figure 9-10c.	OxyContin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School.....	413
Figure 9-11.	Amphetamines: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	414
Figure 9-12.	Sedatives (Barbiturates): Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	415
Figure 9-13.	Tranquilizers: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	416

LIST OF FIGURES (continued)

Figure 9-14a.	Alcohol: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	417
Figure 9-14b.	Alcohol: Trends in 30-day Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	418
Figure 9-14c.	Alcohol: Trends in 30-day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	419
Figure 9-14d.	Alcohol: Trends in 2-Week Prevalence of 5 or More Drinks in a Row among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	420
Figure 9-15a.	Cigarettes: Trends in 30-Day Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	421
Figure 9-15b.	Cigarettes: Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	422
Figure 9-15c.	Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	423

Chapter 1

INTRODUCTION

Monitoring the Future (MTF), now in its 39th year, is a research program conducted at the University of Michigan’s Institute for Social Research under a series of investigator-initiated research grants from the National Institute on Drug Abuse—one of the National Institutes of Health. The study comprises several ongoing series of annual surveys of nationally representative samples of 8th- and 10th-grade students (begun in 1991), 12th-grade students (begun in 1975), and high school graduates into adulthood (begun in 1976). The current monograph reports the results of the repeated cross-sectional surveys of high school graduates since 1976 as we follow them into their adult years. Several segments of the general adult population are represented in these follow-up surveys:

- American college students
- Their age peers who are not attending college, sometimes called the “forgotten half”
- All young adult high school graduates of modal ages 19 to 30, which we refer to as the “young adult” sample
- High school graduates at the specific later modal ages of 35, 40, 45, 50, and 55

In this volume, historical and developmental changes in substance abuse and related attitudes and beliefs occurring at these age strata receive particular emphasis.

The follow-up surveys have been conducted by mail on representative subsamples of the previous participants from each high school senior class. This volume presents data from the 1977 through 2013 follow-up surveys of the graduating high school classes of 1976 through 2012, as these respondents have progressed beyond high school and into adulthood. Data for the oldest respondents now extend through age 55 (the class of 1976) and are , included for the first time in the 2013 survey.

Other monographs in this series include the *Overview of Key Findings*,¹ which presents early results from the secondary school surveys; *Volume I*,² which provides an in-depth look at the secondary school survey results; and the *HIV/AIDS* monograph, drawn from the follow-up surveys of 21- to 30-year-olds, which focuses

¹Johnston, L. D., O’Malley, P. M., Miech, R. A., Bachman, J. G., & Schulenberg, J. E. (2014). *Monitoring the Future national survey results on drug use: Overview of key findings, 2013*. Ann Arbor, MI: Institute for Social Research, The University of Michigan, 84 pp. Available at <http://monitoringthefuture.org/pubs/monographs/mtf-overview2013.pdf>

²Johnston, L. D., O’Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Miech, R.A. (2014). *Monitoring the Future national survey results on drug use, 1975–2013: Volume I, Secondary school students*. Ann Arbor, MI: Institute for Social Research, The University of Michigan, 630 pp. Available at http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2013.pdf

on risk and protective behaviors among young adults related to the transmission of HIV/AIDS.³ This year's *Overview* and *Volume I* are currently available; the *HIV/AIDS* monograph will be published in the fall of 2014

To enable the present volume to stand alone, we have repeated chapters 2 and 3 from *Volume I*, providing key findings from the five populations under study as well as the study design and procedures. Readers already familiar with *Volume I* may wish to skip over these two chapters.

SURVEYS OF YOUNG ADULTS AND THOSE AGES 35, 40, 45, 50, AND 55

The young adult sample consists of representative samples from each graduating class from 2000 to 2012, all surveyed in 2013 and corresponding to modal ages 19 through 30. College students are included as part of this young adult sample. The MTF study design calls for annual follow-up surveys of each high school class cohort through modal age 30, though each individual participates in a follow-up survey only every two years. (High school seniors are considered to be modal age 18.) Chapter 4 presents results on the prevalence of drug use for this age group as well as middle adulthood through age 55. Chapter 5 presents the trends of adult drug use and covers young adult ages 19 through 28. The use of a somewhat shortened age range for young adults (age 28 rather than age 30) allows the trend results to begin two years earlier historically, providing data from 1986 rather than 1988; further, using the 19 through 28 modal age range does not substantially affect the young adult data, typically collected for ages 19 through 30.

Starting at modal age 35, surveys are conducted at five-year intervals. In 2013 the graduating classes of 1976, 1981, 1986, 1991, and 1996 were sent the age-55, age-50, age-45, age-40, and age-35 questionnaires, respectively. Panel data into middle adulthood on nationally representative samples of the population, as well as data on successive class cohorts, are extremely rare. These rare data make possible (1) analyses aimed at differentiating period-, age-, and cohort-related change; (2) analyses demonstrating long-term connections between use of various substances and many important potential outcomes (including eventual abuse and dependence, adverse health outcomes, and functioning in work and family roles); (3) tracking across the life course substance use involvement and how such involvement is affected by social roles and role transitions into and out of social environments; and (4) determining some of the factors in adolescence and early adulthood that are predictive of later substance use, abuse, and dependence.

In this volume, we have reweighted respondent data to adjust for the effects of panel attrition on measures such as drug consumption using a statistical technique called poststratification, explained in chapter 3 in the section on Panel Retention. We are less able to adjust for the absence of high school dropouts who were not included in

³Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., Patrick, M.E. & Miech R. A. (2013). HIV/AIDS: Risk & protective behaviors among American young adults, 2004–2012 Ann Arbor: Institute for Social Research, The University of Michigan, 123 pp. Available at http://www.monitoringthefuture.org/pubs/monographs/mtf-hiv-aids_2012.pdf

the original 12th-grade sample. Because nearly all college students have completed high school, the omission of dropouts should have almost no effect on the college student population estimates, but this omission does affect the estimates for entire age groups. Therefore, the reader is advised that the omission of, on average, about 8% to 15% of each cohort who drop out of high school likely make drug use estimates given here for the various age bands somewhat low for the age group as a whole. (Note, however, that the high school drop-out rate continues to decline, so this has become a bit less of a problem for more recent cohorts). The proportional effect of missing dropouts may be greatest for some of the most dangerous drugs such as heroin, crack, and methamphetamine, as well as for cigarettes—the use of which is highly correlated with educational aspirations and attainment. Nevertheless, even with some underreporting of usage rates, the year-to-year trends observed should be little affected by the limitations in sample coverage.

SURVEYS OF COLLEGE STUDENTS AND NONCOLLEGE PEERS

As defined here, the college student population comprises all full-time students, one to four years post high school, enrolled in a two- or four-year college in March during the year of the survey. More is said about this sample definition in chapters 3 and 8. Results on the *prevalence* of drug use in 2013 among college students and also among their noncollege peers are reported in chapter 8, and results on *trends* in substance use among college students and their noncollege peers are reported in chapter 9, covering the 34-year interval since 1980.

The MTF follow-up samples have provided excellent coverage of the U.S. college student population for more than three decades (1980–2013). College students tend to be a difficult population to study for a variety of reasons. For a number of years they were generally not well covered in household surveys, which tended to exclude dormitories, fraternities, and sororities. Further, institution-based samples of college students must be quite large in order to attain accurate national representation because of the great heterogeneity in colleges and universities and in the types of student populations they serve. Obtaining good samples and high response rates within many institutions also poses difficulties, because the cooperation of each institution must be obtained, as well as reasonable samples of the student body. The current study, which in essence draws the college sample prospectively in senior year of high school, has considerable advantages for generating a broadly representative sample of college students who emerge from each graduating cohort; moreover, it does so at very low cost. In addition, the “before, during, and after college” design permits examination of the many changes associated with the college experience. Finally, the MTF design generates comparable panel data on high school graduates who are *not* attending college, an important segment of the young adult population not only in its own right, but also as a comparison group for college students. This is a particularly valuable and rare feature of this research design.

GENERAL PURPOSES OF THE RESEARCH

MTF's research purposes are extensive and can be outlined here only briefly.⁴ One major purpose is to serve an epidemiological social indicator function, to accurately characterize the levels and trends in certain behaviors, attitudes, beliefs, and environmental conditions in the various populations covered. Social indicators can have important agenda-setting functions for society, drawing attention to new threats to the public health, and estimating the extent of those threats as well as determining where they are concentrated in the population. They are especially useful for gauging progress toward national goals and indicating the impacts of major historical events, including social or policy changes. Another purpose of the study is to develop knowledge that increases our understanding of how and why historical changes in these behaviors, attitudes, beliefs, and environmental conditions are taking place. Such work is usually considered to be social epidemiology. These two broad purposes are addressed in the current series of volumes.

Additional etiologic purposes of MTF include helping to discover what types of young people are at greatest risk for developing various patterns of drug abuse, gaining a better understanding of the lifestyles and value orientations associated with various patterns of drug use and monitoring how these are shifting over time, examining the immediate and more general aspects of the social environment that are associated with drug use and abuse, and discovering how drug use is affected by *major transitions* into and out of social environments (such as military service, civilian employment, college, unemployment) or social statuses (married, pregnant, parent, divorced, remarried). We also seek to examine the life course of various drug-using behaviors during this period of development (including progression to dependence), to distinguish such age effects from cohort and period effects that are influencing drug use and attitudes about drug use, to discover the effects of social legislation and changing regulations on various types of substance use, and to understand the changing connotations of drug use and changing patterns of multiple drug use among youth.

We believe that differentiating among period, age, and cohort effects on use of various types of substances has been a particularly important contribution of the project. The MTF cohort-sequential research design is especially well suited to discern changes with age common to all cohorts (age effects), differences among cohorts that tend to persist across time (cohort effects), and changes common to most or all ages in a given historical period (period effects).

⁴For a more complete listing and discussion of MTF's many objectives, see Johnston, L. D., O'Malley, P. M., Schulenberg, J. E., & Bachman,

J. G. (2006). The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2006 (Monitoring the Future Occasional Paper No. 65). Ann Arbor, MI: Institute for Social Research. Available online at <http://www.monitoringthefuture.org/pubs/occpapers/occ65.pdf>.

Knowing which type of change is occurring is important for at least three reasons. First, it can help to discover what types of causes account for the change. For example, age effects are often explained by maturation as well as by environmental and role transitions associated with age, as this study has demonstrated.⁵ Second, the type of change can indicate when in the life course the causes may have had their impact; in the case of cohort effects, it may well have been in earlier historical periods than when the change is actually documented. For example, we know from MTF data on age of initiation that the decline in cigarette smoking observed among 12th graders in the late 1970s actually reflected a cohort effect that emerged when those teens were younger, in the early 1970s, which was shortly after cigarette advertising was removed from radio and television. The third reason that knowing the type of change is important is that it can help in predicting future change more accurately. For example, the study has shown that perceived risk often is a leading indicator of change and that cohort effects help to predict forthcoming changes at later ages. Needless to say, predicting change is extremely valuable to the policy, prevention, and treatment communities. This volume features well-established age effects, some important cohort effects that emerged in the 1990s, and recent period effects.

One important purpose of MTF, related to but distinct from the ones described so far, is to study HIV/AIDS-related behaviors. This purpose is addressed in a separate monograph.⁶ In 2004, for the first time, questions were included on the prevalence and interconnectedness of risk and risk-reduction behaviors related to the spread of the human immunodeficiency virus (HIV), which causes acquired immunodeficiency syndrome (AIDS). The questions include, in addition to questions about drug involvement in general, questions about injection drug use, needle sharing in particular, as well as number of sexual partners, gender(s) of those partners, use of condoms, blood donation, getting tested for HIV/AIDS, and securing the results of such tests. Beginning in 2004 these questions were included in two of the six forms in the follow-up surveys of 21- to 30-year-olds. The questions were added to an additional form beginning in 2007, to the age-35 questionnaire in 2008, and then (having shown no deleterious effects on response rate at age 35) to the age-40 questionnaire in 2010.

Readers interested in publications dealing with any of these areas are invited to visit the MTF website at www.monitoringthefuture.org. For additional information, please e-mail us at MTFinfo@umich.edu.

⁵Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). The education-drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency. New York: Lawrence Erlbaum Associates/Taylor & Francis; Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs. Mahwah, NJ: Lawrence Erlbaum Associates.

⁶ Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., Patrick, M. E., & Miech, R. A. (2013). *HIV/AIDS: Risk & protective behaviors among American young adults, 2004-2012*. Ann Arbor: Institute for Social Research, The University of Michigan, 123 pp. Available at http://www.monitoringthefuture.org/pubs/monographs/mtf-hiv-aids_2012.pdf

Chapter 2

KEY FINDINGS AN OVERVIEW AND INTEGRATION ACROSS FIVE POPULATIONS

Monitoring the Future, now having completed its 39th year of data collection, has become one of the nation's most relied-upon scientific sources of valid information on trends in use of licit and illicit psychoactive drugs by American adolescents, college students, young adults, and adults up to age 55. During the last nearly four decades, the study has tracked and reported on the use of an ever-growing array of such substances in these populations.

This annual series of monographs is the primary mechanism through which the epidemiological findings from MTF are reported. Findings from the inception of the study in 1975 through 2013 are included—the results of 39 national in-school surveys and 37 national follow-up surveys.

MTF has conducted in-school surveys of nationally representative samples of (a) 12th-grade students each year since 1975 and (b) 8th- and 10th-grade students each year since 1991. In addition, beginning with the class of 1976, the project has conducted follow-up mail surveys on representative subsamples of the respondents from each previously participating 12th-grade class. These follow-up surveys now continue well into adulthood, currently up to age 55.

A number of important findings are summarized in this chapter to provide the reader with an overview of the key epidemiological results from the study. Because so many populations, drugs, and prevalence intervals are discussed here, a single integrative set of tables (Tables 2-1 through 2-4) show the 1991–2013 trends for all drugs on five populations: 8th-grade students, 10th-grade students, 12th-grade students, full-time college students modal ages 19–22, and all young adults modal ages 19–28 who are high school graduates. (Note that the young adult group includes the college student population.) *Volume II* also contains data on older age groups based on the longer term followup surveys, specifically ages 35, 40, 45, 50, and 55. (Data from the 55-year-olds first became available in 2013.)

TRENDS IN DRUG USE—THE ADVENT OF COHORT EFFECTS

Early in the 1990s, we reported an increase in use of several illicit drugs among secondary school students, and some important changes among the students in terms of certain key attitudes and beliefs related to drug use. In the volume reporting 1992 survey results, we noted the beginning of such reversals in both use and attitudes among 8th graders, the youngest respondents surveyed in this study, and also a reversal in attitudes among 12th graders. Specifically, the proportions seeing great

risk in using drugs began to decline, as did the proportions saying they disapproved of use. As we suggested then, those reversals indeed presaged “an end to the improvements in the drug situation that the nation may be taking for granted.” The use of illicit drugs rose sharply in all three grade levels after 1992, in what we refer to as the “relapse phase” in the larger epidemic of illicit drug use, as negative attitudes and beliefs about drug use continued to erode. This pattern continued into the mid-1990s, and beyond that for some drugs.

Then in 1997, for the first time in six years, the overall rate of *illicit drug* use finally showed a decline among 8th graders. Although marijuana use continued to rise that year among 10th and 12th graders, their use of several other drugs leveled off, and relevant attitudes and beliefs also began to reverse in many cases. In 1998, illicit drug use continued a gradual decline among 8th graders and also started to decline at 10th and 12th grades. In 1999 and 2000, the decline continued for 8th graders, while use held fairly level among 10th and 12th graders. In 2002 and 2003, use by 8th and 10th graders decreased significantly, and use by 12th graders finally began to drop; declines then continued for all three grades in 2004 and for several years thereafter. But in 2008, illicit drug use increased once again among 8th and 12th graders, followed by some increase in 8th and 10th grades in 2009, signaling an end to the immediately preceding period of decline. In 2010 the overall rate of illicit drug use increased for all grades, although only the increase among 8th graders was significant. In 2011 the increase continued among 10th and 12th graders and declined some at 8th grade. We expected a continued increase into 2012, in part because of the ongoing trend toward increased use and in part because some states have been moving to legalize the medical use (and simply the recreational use) of marijuana, which may serve to normalize use of marijuana, the most widely used of all illicit substances. Instead, in 2012, we found nonsignificant declines in illicit drug use in all three grades (when rates for all three grades were combined, there was also a nonsignificant decline). In 2013 an index of use of any illicit drug showed increases in all three grades and all prevalence periods (lifetime, annual, and 30-day). The only statistically significant increase was for annual use among 8th graders; but the three grades combined showed a significant increase in use for lifetime and annual use.

As will be illustrated below in the discussion of specific drugs, the increase in use of many drugs during the 1990s among secondary school students, combined with fairly level rates of use among college students and young adults, resulted in some unusual reversals in the usage rates by age (see Figure 2-1). In the early years of the epidemic, illicit drug use rates were clearly higher in the college-age group (and eventually the young adults) than they were among secondary school students. But by the late 1990s, the highest rates of active use (i.e., use within the prior year or prior 30 days) were found in the late secondary school years. In fact, in 1996 and 1997 both 10th and 12th graders actually had higher annual prevalence rates for illicit drug use (i.e., higher percentages reporting any use within the prior year) than either college students or all young adults. This changed somewhat after 2001, as the earlier, heavier using cohorts of adolescents began to comprise the college student and young adult populations, while at the same time use among the incoming secondary school students was declining.

- As can be seen by the divergence of trends for the different age groups, something other than a simple secular trend in drug use was taking place; important cohort differences were emerging. (A cohort refers to a group of people born at the same time or, in this case, of the same graduating class. A secular trend is a trend across time that occurs across multiple cohorts and multiple age groups.)
- In 2013, the rank order by age group for annual prevalence of using *any illicit drug* was 12th graders (40%), college students (39%), 19- to 28-year-olds (36%), 10th graders (32%), and 8th graders (15%). With respect to using *any illicit drug other than marijuana* in the past 12 months, there was less variability: college students (19%), 19- to 28-year-olds (18%), 12th graders (17%), 10th graders(11%), and 8th graders (6%).
- Bath Salts, so-called because they are sold over the counter as apparently innocuous products like bath salts but really contain strong stimulants, have been given much media attention in the past few years; however, there has been very little scientific information about the prevalence of their use. We believe that the 2012 MTF survey provided the first national survey data on their use. Fortunately, we found the annual prevalence rates in 2012 to be very low, at 0.8%, 0.6%, and 1.3% in grades 8, 10, and 12, respectively. In 2013 the prevalence rates are 1.0%, 0.9%, and 0.9% in grades 8, 10, and 12, respectively.
- From the early 1990s until 1997, *marijuana* use rose sharply among secondary school students, as did their use of a number of *other illicit drugs*, though more gradually. As previously stated, we have called this period a “relapse phase” in the longer term epidemic. An increase in marijuana use also began to occur among U.S. college students, largely reflecting “generational replacement” (i.e., a cohort effect), wherein earlier cohorts were replaced in the college population by more recent ones who were more drug-experienced before they left high school. This resurgence in illicit drug use spread *up* the age spectrum in a reversal of the way the epidemic spread several decades earlier. In the 1960s the epidemic began on the nation’s college campuses, and then diffused downward in age to high school students and eventually to middle school students. This time the increases began in middle schools and radiated up the age spectrum. The graduating class cohorts in the middle and late 1990s carried with them the pattern of heavier drug use that emerged while they were in secondary school in the early 1990s.

Increases during the 1990s in use of *any illicit drug* (including use of *marijuana* and use of *other illicit drugs* treated as a class) were substantially larger, in both proportional and absolute terms, in the three secondary school grades than in either the college or young adult populations. Among college students and young adults, the annual prevalence of use of *any illicit drug* held remarkably stable from 1991 through 1997, at the same time use rose appreciably among adolescents (see Figure 2-1). We predicted that, as

generational replacement continued to occur, we would likely see some increase in use of illicit drugs by the young adults. As would be expected given their younger age range (19–22), the increase happened sooner and more sharply among the college students than among the young adults in general (age range 19–28). Peak rates (since 1990) in annual prevalence of **any illicit drug** were reached in 1996 among 8th graders, in 1997 among 10th and 12th graders, in 2001 among college students (before leveling for some years), and in 2004 (before leveling) in the young adult segment. Similarly, the more recent declines in use among secondary students have thus far shown up only modestly and briefly among college students, and hardly at all among young adults (see Figure 2-1). In the last few years, including 2013, all five populations have shown some increase in the use of illicit drugs, largely as a result of increases in marijuana use; this appears to reflect a secular trend or historical period effect, a change seen across all ages in the same time period.

Again, the earlier diverging trends across the different age strata clearly show that changes during the 1990s reflected some important cohort effects rather than broad secular trends that would have appeared simultaneously in all of the age groups. During all of the previous years of the study, the use of most drugs moved in parallel across most age groups, indicating that secular change was prevailing then.

- Similar to the use patterns for illicit drugs, the trend for **cigarette** smoking evidenced a generational replacement effect during the 1990s in that college students showed a sharp increase in smoking beginning in 1995, as the heavier smoking cohorts of secondary school students from the early to mid-1990s entered college. This has been a more typical pattern of change for cigarettes, however, since differences in cigarette smoking rates among class cohorts tend to remain through the life course and also tend to account for much of the overall change in use observed at any given age.

In the early 1990s, cigarette smoking among 8th and 10th graders rose by about 50%—a particularly sharp and worrisome rise (based on 30-day prevalence rates shown in Table 2-3, and daily and half-pack rates shown in Table 2-4); MTF was the first study to draw national attention to this momentous development. Smoking also rose among 12th graders, beginning a year later.

The increase in current smoking ended among 8th and 10th graders in 1996, among 12th graders in 1997, and among college students in 1999. The nation then entered a period of appreciable decline in smoking rates that first began among 8th graders in 1997 and then began radiating up the age spectrum as those cohorts aged. (The 8th-grade 30-day prevalence rate fell by about three fourths, from 21% in 1996 to 4.5% in 2013.) Among the college and the young adult strata, the declines have been less sharp so far, but they are continuing. The 30-day smoking prevalence rate for college students in 2013 (14%) was down more than half from the recent peak of 31% in 1999, with

the decline accelerating after 2005 as the cohort effect worked its way up the age bands. Smoking among the young adult subgroup, on the other hand, has dropped by only about one third (to 20% by 2013) since its recent peak rate of 31% in 1998. The decline in smoking rates among secondary school students had been decelerating in all three grades in recent years; there was some evidence in 2010 that the decline had halted among 8th and 10th graders, and that a turnaround might be occurring. Fortunately, all three grades showed further declines in 2011, 2012, and 2013, including a significant drops in all three grades over that interval. (The recent decline may be due at least in part to a 2009 increase in federal taxes on tobacco products.) In 2013 neither of the older age groups—college students and young adults—showed any further decline in 30-day smoking rates.

- During the 1990s, the annual prevalence of *marijuana* use tripled among 8th graders (from 6% in 1991 to 18% in 1996), more than doubled among 10th graders (from 15% in 1992 to 35% in 1997), and nearly doubled among 12th graders (from 22% in 1992 to 39% in 1997). Among college students, however, the increase in marijuana use, presumably due to a generational replacement effect, was much more gradual. Annual prevalence of use rose by about one third, from 27% in 1991 to 36% in 1998. Marijuana use began to decline in 1997 among 8th graders and then did the same in 1998 among 10th and 12th graders. The rate of decline was rather modest, however, perhaps due in part to effects of the public debates over medical use of marijuana during that period. In 2001, use remained level in all three grades, but between 2001 and 2004 all three grades showed significant declines in their annual prevalence of marijuana use, with the proportional decline greatest among 8th graders. Eighth graders exhibited the most steady long-term decline since their recent peak, which occurred in 1996, although the decline halted in 2008, after a decline of more than four-tenths. After 2007 use began to increase among 8th graders (see Figure 5-4a in Chapter 5). Declines had been occurring in the upper grades after 1997, but mostly after about 2001, with their annual prevalence rates having fallen from recent peaks by 31% and 18% (roughly between 1997 and 2008) for 10th and 12th graders, respectively. All three grades have shown an increase in annual prevalence in recent years—8th graders since 2007 and 10th and 12th graders since about 2008, although the increases have been uneven. In 2013 use in the two lower grades continued to rise while use among the 12th graders leveled. Annual marijuana use among college students and young adults has been rising steadily since 2010. In 2013, increases in marijuana use occurred in all of the populations measured except 12th graders, who remained unchanged (see Table 2-2). Earlier, use declined modestly among college students from recent peak levels of 36% in 2001 to 30% in 2006. What seems clear is that the long decline in marijuana use, which extended over roughly a 10 year period, ended a few years back among secondary school students and has now ended among college students and young adults. We noted in 2010 that if a new cohort effect emerges, then within a few years we are likely to see an increase in marijuana use among college students and young adults generally. In 2011 both groups showed

some increase in marijuana use—for young adults a significant increase of 2.3 percentage points. We believe that some of this increase in use may reflect a secular trend. If the debate over medical marijuana use (and decriminalization and legalization) is reducing the perceived risk of that drug, the effect could well be occurring across various age groups, thus creating a secular trend rather than a cohort effect.

- ***Current daily marijuana*** use in all of these groups rose substantially after 1992, reaching peak levels in a somewhat staggered fashion as that just described (see Table 2-4 and Figure 5-4a in Chapter 5). Daily use began a slow decline after 1999 among 8th graders until 2007, after 2001 until 2009 among 10th graders, and after 2003 until 2010 among 12th graders, consistent with a cohort effect pattern. Use at all three grade levels was fairly level after 2004. In 2010 daily use at all three grade levels increased significantly and it increased further in grades 10 and 12 in 2011 and 2012, while holding steady in 8th grade. There was no further change in 2013 for grades 8 and 12, though daily use continued to rise in grade 10. The 2013 daily prevalence rates in grades 8, 10, and 12, respectively, are 1.1%, 4.0%, and 6.5%. *In other words, about one in every fifteen high school seniors is a current daily marijuana user.* College student and young adult rates of daily use have increased very gradually since 2007, from 3.5% to 5.1% in 2013 among college students and from 5.0% to 6.2% over that same interval among young adults. In general, prevalence of daily marijuana use was slow to decline, when annual and 30-day prevalence figures were dropping. Although the rates today are low relative to the peaks reported in the late 1970s, the 6.6% figure for 12th graders in 2011 was the highest observed in some 30 years. The fact that daily marijuana use was rising through 2011 in all three grades serves as a reminder that a relapse in the epidemic of marijuana use, as occurred in the early 1990s, could still occur. The role of the many debates on legalizing marijuana for medical use, the actual legalization for recreational use in some states, and the experiences those states have with the new laws likely will have an impact on present and future secular trends in use.

The amount of perceived risk associated with using marijuana fell during the earlier period of increased use in the late 1970s, and fell again during the more recent resurgence of use in the 1990s. Indeed, perceived risk among 12th graders began to decline a year *before* use began to rise in the upturn of the 1990s, making perceived risk a leading indicator of change in use. (The same may have happened in 8th grade, as well, but we do not have data starting early enough to check that possibility.) The decline in perceived risk halted after 1997 for 8th and 10th graders, and annual prevalence began to decline a year or two later. Perceived risk also declined prior to the recent rebound in marijuana use. Again, perceived risk has been a leading indicator of change in use, as it has proven to be for a number of drugs. As discussed in *Volume I*, chapter 8, these attitudes, as well as the behaviors that they predict, show evidence of cohort effects over the past decade and a half. Perceived risk of

trying marijuana has been declining in recent years and dropped for 8th, 10th, and 12th graders in 2013.

Personal disapproval of marijuana use slipped considerably among 8th graders between 1991 and 1996 and among 10th and 12th graders between 1992 and 1997, as use rose considerably. For example, the proportions of 8th, 10th, and 12th graders who said they disapproved of trying marijuana once or twice fell by 17, 21, and 19 percentage points, respectively, during their respective intervals of decline. Subsequently, disapproval began to rise among 8th graders after roughly 1997 and continued through 2007, while it began to rise in the upper grades in 2002 and also continued through 2007 among 10th and 12th graders, as use declined gradually. Since 2007 or 2008 there has been some reversal on this attitude as well as in use, with disapproval falling steadily in the upper grades and less consistently in grade 8 (see Figure 8-1b in Chapter 8). In 2013 there were significant drops in disapproval for various levels of use across all three grades.

- ***Synthetic marijuana***, so named because it contains synthetic versions of some of the cannabinoids found in marijuana, is a recent and important addition to the smorgasbord of drugs available to American young people. These designer chemicals are sprayed onto herbal materials that are then sold in small packets under such brand names as Spice and K-2. They have been readily available as over-the-counter drugs on the Internet, in head shops and gas stations, etc. While many of the most widely used chemicals were scheduled by the Drug Enforcement Administration in March of 2011, making their sale no longer legal, purveyors of these products have skirted the restrictions by making small changes in the chemical composition of the cannabinoids used. Use of these products was first measured in MTF in 2011 in a tripwire question for 12th graders, asking about their frequency of use in the prior 12 months (see Table 2-2). Annual prevalence was found to be 11.4%, making synthetic marijuana the second most widely used class of illicit drug after marijuana. In spite of the DEA's scheduling of the most common ingredients, use among 12th graders remained unchanged in 2012, with 11.3% annual prevalence. Eighth and 10th graders were also asked about use of these drugs in 2012, and their annual prevalence rates were 4.4% and 8.8%, respectively, making synthetic marijuana the second most widely used illicit drug among 10th graders, as well, and the third among 8th graders behind marijuana and inhalants. In 2013 use dropped appreciably in all five populations, including statistically significant drops among 12th graders, college students, and young adults. Efforts by the DEA and various states to make their sale illegal may well be making an impact. There is a relatively low level of perceived risk for trying synthetic marijuana once or twice, despite growing evidence of serious problems resulting from the use of these drugs.
- Among 12th graders, the proportions using ***any illicit drug other than marijuana*** in the past twelve months rose from a low of 15% in 1992 to a high of 21% in 1999 (see Table 2-2); these levels were substantially below the

34% peak rate reached two decades earlier, in 1981. All of the younger groups showed significant increases between 1992 and 1997, with use beginning to increase in 1992 among 8th graders, in 1993 among 10th and 12th graders, and in 1995 among college students—reflecting strong evidence of a cohort effect. Use peaked in 1996 among 8th and 10th graders, in 1997 among 12th graders, around 2004 for college students and in 2008 for young adults. Since 1996 the 8th graders have shown a gradual but considerable decline of one half in their use of illicit drugs other than marijuana, treated as a class (13.1% annual prevalence in 1996 to 5.5% in 2012, with no change in 2013). The decline among 10th graders paused from 1998 to 2001 with a net decline of about a third in annual prevalence from 18.4% in 1996 to 11.3% in 2008; use leveled again for several years and then declined further in 2011. It now stands at 10.9%. Twelfth-grade use also showed some decline beginning after 2001 (21.6%) and stands just 4.3 percentage points lower (17.3%) in 2013. College students so far have shown little change with a recent high of 18.6% in 2004 to 19.0% in 2013. Use among young adults also has remained at about the same level of annual use, between 17% and 19%, since 2003.

- Between 1989 and 1992 we noted an increase among 12th graders, college students, and young adults in their use of **LSD**, a drug quite popular in the late 1960s and early 1970s. In 1992 the newly added populations (8th and 10th graders) were also showing an increase in LSD use; for several more years, modest increases persisted in all five populations. Use of LSD peaked in 1995 among college students and young adults and in 1996 among 8th, 10th, and 12th graders, after which LSD use gradually declined in all five populations until 2005 for 8th, 10th, and 12th graders. Overall, the pattern for LSD use seems more consistent with secular change than a cohort effect. The different age groups moved in parallel for the most part, likely in response to historical events in the environment, including a sharp reduction in LSD availability after 2001.

The proportion of 12th graders seeing great risk associated with trying LSD declined by 4.3-percentage-points between 1991 and 1992, just prior to the significant increase in LSD use in 1993. The decline in perceived risk continued through 1997, halted in 1998, and has resumed since 2009. The proportion of 12th graders disapproving of LSD use began to decline in 1992, and continued to decline through 1996.

Because LSD was one of the earliest drugs to be popularly used in the American drug epidemic, young people in the 1990s may have been relatively unaware of the risks of use. They had less opportunity to learn vicariously about the consequences of use by observing others around them or to learn from intense media coverage of the issue, which occurred some years earlier. We were concerned that this type of generational forgetting of the dangers of a drug, which occurs as a result of generational replacement, could set the stage for a whole new epidemic of use. In fact, perceived harmfulness of LSD began to decline after 1991 among 12th graders. Perceived risk and

disapproval among 8th and 10th graders, first measured in 1993, both showed declines until 1997 or 1998, after which they leveled among 10th graders but then declined considerably more among 8th graders. In 2004, twelfth graders' personal disapproval of trying LSD increased significantly, with little change since. Because the decline in use in the last few years has generally *not* been accompanied by expected changes in these attitudes and beliefs, we suspected that some displacement by another drug might have been taking place, at least through 2001. The most logical candidate is *ecstasy (MDMA)*, which, like LSD, is used for its hallucinogenic effects; ecstasy was popular in the club and rave scenes, and was very much on the rise through 2001. After 2001, a sharp decline in the reported availability of LSD in all five populations (which corresponded to the closing of a major LSD lab by the Drug Enforcement Administration) very likely played a major role in the sharp decline in use of LSD among all groups. However, we want to caution that 8th graders' attitudes, in particular, are changing such as to make them receptive to LSD use some time in the future, should a plentiful supply re-emerge. Fortunately, 8th and 10th graders report declining availability in 2012. In 2013 there was some drop in perceived availability for 8th graders and 12th graders.

- Questions about the use of *ecstasy (MDMA)* have been included in the follow-up surveys of college students and young adults since 1989; however, because of our concern about stimulating interest in an attractive-sounding and little-known drug, these questions were not added to the secondary school surveys until 1996. From 1989 to 1994, the annual prevalence rates tended to be quite low in the older age groups for whom we had data, but in 1995 these rates increased—from 0.5% in 1994 to 2.4% in 1995 among college students, and from 0.7% to 1.6% over the same time span among young adults generally.

When usage data were first gathered on secondary school students in 1996, the 10th and 12th graders actually showed higher rates of annual use (both 4.6%) than the college students (2.8%). Ecstasy use then fell steadily in all three grades between 1996 and 1998, though it did not fall in the older age groups (see Table 2-2). But between 1998 and 2001, use rose sharply in all five populations. In fact, annual prevalence more than doubled in that three-year period among 12th graders, college students, and young adults, and nearly doubled in the lower grades. In 2000 even the 8th graders showed a significant increase in use. Ecstasy use for all five age groups declined slightly in 2002, but significantly only for 10th graders; declined again in 2003, with significant drops for all groups except the college students; and showed some decline again in 2004, with the largest decreases among college students and young adults. This pattern suggests that both cohort effects and a secular trend were at work. Once again, this decline in use among 12th graders was predicted by an increase in perceived risk in 2001—an increase that continued through 2005. Among college students the annual prevalence fell by half in 2004 alone, and all five groups are at rates that are still much lower than their recent peaks in 2001. Since 2005 or 2006, there was some rebound in use among all five populations, including a significant increase in the lower

grades in 2010. Except for the significant decline in use among 8th graders, there was little systematic change in 2011. In 2012 there were significant declines in annual prevalence in all three grades. There was little or no change in annual ecstasy use for any of the five groups in 2013.

Ecstasy use among all five populations has been moving fairly synchronously since 1999, which suggests a secular trend (some change in events in the social environment) that affected everyone. An important change during this period was the increasing availability of information on the adverse effects of ecstasy use via stories in the popular media, dissemination of the scientific evidence by the National Institute on Drug Abuse, and an anti-ecstasy media campaign by the Partnership for a Drug-Free America and the Office of National Drug Control Policy, initiated in 2002.

Availability of ecstasy increased dramatically through 2001, as reported by 12th graders and substantiated by law enforcement data on ecstasy seizures. Of the 12th graders surveyed in 1991, only 22% thought they could get ecstasy fairly easily, but a decade later (in 2001) 62% thought that they could. After 2001, however, the perceived availability of ecstasy began decreasing in all three grades, possibly due in part to the steep decline in the number of users who serve as supply points for others. The decreases continued through 2012 in the lower grades. In 12th grade, the decline in perceived availability continued through 2009, then leveled. (See Figure 8-6 in *Volume I*, chapter 8 for a graphic presentation of the trends in ecstasy use, availability, and perceived risk for 12th graders.) However, perhaps the most important change that has been taking place since 2005 is a continual decline in perceived risk for ecstasy use among 8th, 10th, and 12th graders, possibly as a result of generational forgetting. In our 2009 MTF report, we suggested that this decline in perceived risk was leaving high school students increasingly vulnerable to a possible rebound in use of ecstasy; indeed, there was some evidence that just such a rebound was occurring, at least through 2010 or 2011, but there was no further increase in 2012 and 2013.

- Between 1982 and 1992, annual prevalence rates for *amphetamine* use (other than use that was ordered by a physician) among 12th graders fell by nearly two thirds, from 20.3% to 7.1%. Rates among college students fell even more over the same interval, from 21.1% to 3.6%. During the relapse phase in the drug epidemic in the 1990s, annual amphetamine use increased by about half among 8th and 10th graders between 1991 and 1996, and also increased among 12th graders and college students between 1992 and 1996. After 1996 the age groups diverged, with amphetamine use declining gradually and substantially among 8th graders—where use is now a fraction of what it was in 1996—but continuing to rise among 12th graders (and eventually 10th graders), college students, and young adults until about 2002. The declines continued in the upper grades through about 2008 but through 2013 for 8th graders. Since 2009, annual prevalence has increased significantly among 12th graders (from 6.6% to 8.7% in 2013), perhaps as a result of more

students using amphetamines to help their academic work. Among students in college, amphetamine use rose even more sharply from 5.7% in 2008 to 10.6% in 2013, likely for the same reason. Young adults, who include the college students, showed less of an increase over the same interval, from 5.3% in 2008 to 7.8% in 2013. The pattern of cross-age-group change suggests a cohort effect at work for amphetamine use. Since the late 1990s there has been a greater difference between use among 8th graders and use by older students, suggesting that an age effect has emerged, possibly due to the older students becoming more likely to use amphetamines to aid their academic performance. (“To help me study” was the highest endorsed reason 12th graders gave for amphetamine use in 2012 and the third highest in 2013.)

Among 12th graders, the increase in nonmedical use of amphetamines (and a concurrent decrease in disapproval) began in 1993; this followed a sharp drop in perceived risk a year earlier (which, as we have noted for a number of drugs, often serves as a leading indicator). Following a period of decline, perceived risk among 12th graders increased gradually from 1995 through 2009.⁶

- Use of the amphetamine *Ritalin* outside of medical supervision showed a distinct increase around 1997—with annual prevalence among 12th graders going from 0.1% in 1992 to 2.8% in 1997—and then stayed level for a few years (see appendix E in Volume I, Table E-2⁷). Because of its increasing importance, a differently structured question was introduced for Ritalin use in 2001 (2002 in the follow-ups of college students and young adults). This new question, which we prefer to the original, does not use a prior branching question and produced somewhat higher prevalence rates. Results from the new question suggest an ongoing, gradual decline in Ritalin use, which continued into 2009 in all five populations. The decline continued further in 2012 among 8th graders, and until 2013 among 10th graders, Ritalin use leveled around 2010 among 12th graders. College students, but not all young adults, showed a nonsignificant increase in use in 2013.
- Another amphetamine used in the treatment of the symptoms of attention deficit hyperactivity disorder (ADHD) is *Adderall*. A new question on its use was introduced in 2009; annual prevalence rates in 2009 through 2013 were higher than those for Ritalin in all five populations. This suggests that Adderall may have to some degree replaced the use of Ritalin and may help to account for the declines that we have been observing for the latter drug. Annual prevalence of Adderall changed rather little between 2009 and 2013 in

⁶ In 2011 the question on perceived risk was modified to include Adderall and Ritalin as examples, which seems to have lowered the level of perceived risk (pep pills and bennies were deleted from the list of examples that same year).

⁷ As discussed in appendix E of *Volume I*, the absolute prevalence rates for Ritalin are probably higher than the statistics indicate, but the trend story is likely quite accurate. See Table 2-2 for more accurate estimates of the absolute annual prevalence rates in recent years; these estimates are based on a new question that does not require the respondent to indicate some amphetamine use before being branched to a question about Ritalin use.

8th and 10th grades, although the rates seem to be drifting down. In 12th grade, however, annual prevalence has risen from 5.4% in 2009 to 7.6% in 2012 and leveled in 2013. The absolute prevalence rates are fairly high, particularly among 12th graders (7.4%) and college students (10.7% in 2013).

- ***Methamphetamine*** questions were introduced in 1999 because of rising concern about use of this drug; but a decline in use has been observed among all five populations in the years since then, through about 2012. In 2013 use in all five populations was at very low rates of annual prevalence—particularly among college students (0.4%). These substantial declines occurred during a period in which there were many stories in the media suggesting that methamphetamine use was a *growing* problem—an example of the importance of having accurate epidemiological data available against which to test conventional wisdom.
- Measures on the use of ***crystal methamphetamine (ice)*** (a crystallized form of methamphetamine that can be smoked, much like crack) have been included in MTF since 1990. The use of crystal methamphetamine increased between the early and late 1990s among the three populations asked about their use: 12th graders, college students, and young adults. However, use never reached very high levels. The estimates are less stable than usual due to the relatively small samples asked about this drug, but it appears that among 12th graders crystal methamphetamine use held fairly steady from 1999 through 2005 (when it was 2.3%); since then it has declined by about half, to 1.1% in 2013. Use rose somewhat among college students and other young adults until 2005, before dropping substantially since then. After their peak levels were reached in 2005, college students and young adults showed substantial drops in annual prevalence to 0.0% by 2013 for college students and to 0.8% for young adults generally (see Table 2-2).
- ***Inhalants*** are defined as fumes or gases that are inhaled to get high, and they include common household substances such as glues, aerosols, butane, and solvents of various types. Among 12th graders there was a long-term gradual increase in the use of inhalants (unadjusted for nitrite inhalants) from 1976 to 1987, followed by a leveling for a few years and then a further increase in the early 1990s. This troublesome increase in inhalant use also occurred among students in the lower grades, and was followed by a reversal in all 3 grades after 1995. After reaching a low point by 2002 or 2003 in grades 8, 10, and 12, use of inhalants increased some in all grades, but then declined in all grades. Use is now at the lowest point in the history of the study for 8th and 10th graders and near the lowest point for 12th graders. Perceived risk for inhalant use among 8th and 10th graders declined fairly steadily after 2001, quite possibly as a result of generational forgetting of the dangers of these drugs; by 2013 the percent of 8th and 10th graders seeing great risk in trying inhalants fell by 12 and 7 percentage points. A new anti-inhalant campaign could well be effective in offsetting this decline in perceived risk in recent years, much as a similar campaign appeared to do in the mid-1990s.

- ***Amyl and butyl nitrites***, one class of inhalants, became somewhat popular in the late 1970s, but their use has been almost eliminated in the years since then. The annual prevalence rate among 12th-grade students was 6.5% in 1979 but only 0.9% in 2009. (Because of this decrease in use, and to allow for the addition of other questions, the questions on nitrite use were dropped from the study in 2010.) When nitrites were included in the definition of inhalants, they masked the increase that was occurring in the use of other inhalants, because their use was declining at the same time that the use of the other inhalants was increasing (see Figure 5-4c in *Volume I*).
- ***Crack cocaine*** use spread rapidly from the early to mid-1980s. Still, among 12th graders, the use of crack remained relatively low during this period (3.9% annual prevalence in 1987). Clearly, crack had quickly attained a reputation as a dangerous drug, and by the time of our first measurement of perceived risk in 1987, it was seen as the most dangerous of all drugs. Annual prevalence dropped sharply in the next few years, reaching 1.5% by 1991, where it remained through 1993. Perceived risk began a long and substantial decline after 1990—again serving as a driver and leading indicator of use. (The decline in perceived risk in this period may be an example of generational forgetting.) Annual prevalence among 12th graders rose gradually after 1993, from 1.5% to 2.7% by 1999. It finally declined slightly in 2000 and then held level through 2007. Since then, some additional decline has occurred. In 2013 annual prevalence for crack cocaine was at 1.1%.

Among 8th and 10th graders, crack use rose gradually in the 1990s: from 0.7% in 1991 to 2.1% by 1998 among 8th graders, and from 0.9% in 1992 to 2.5% in 1998 among 10th graders. And, as just discussed, use among 12th graders peaked in 1999 at 2.7% and among young adults at 1.4%. Since those peak years, crack use has declined appreciably—by more than half among 8th, 10th, and 12th graders—yet it held fairly steady among college students and young adults, at least until 2007, when use among college students finally began to decline. The 2013 prevalence rates for this drug are relatively low—between 0.3% and 1.1% in all five groups. Twelfth graders have the highest prevalence rate. Annual crack prevalence among the college-bound has generally been considerably lower than among those not bound for college. After a significant drop in use among the college-bound, the rates in 2013 are 0.7% for college-bound and 2.1% for noncollege-bound.

We believe that the particularly intense and early media coverage of the hazards of crack cocaine likely had the effect of capping an epidemic early by deterring many would-be users and motivating many experimenters to desist use. As has been mentioned, when we first measured crack use in 1987, it had the highest level of perceived risk of any illicit drug. Also, it did not turn out to be “instantly addicting” upon first-time use, as had been widely reported. In some earlier years, 1994 and 1995 for example, 3% of 12th graders reported ever trying crack; however, only about 2% used in the prior 12 months and only about 1.0% used in the prior 30 days. It thus appears that, among the

small numbers of 12th graders who have ever tried crack, the majority of those who tried it did not establish a pattern of continued use, let alone develop an addiction.

Perceived risk and disapproval associated with crack dropped in all three grade levels in 1993, foretelling the rise in use that occurred in all three grades between 1994 and 1999 in the case of the 12th graders. Because more than a decade had passed since the 1986 media frenzy over crack and its dangers, it is quite possible that generational forgetting of the risks of this drug contributed to the declines in perceived risk and disapproval. Indeed, perceived risk of crack use eroded steadily at all grade levels from 1991 (or 1992 for 12th graders) through 2000. There was not much systematic change in risk or disapproval of crack after that, though disapproval did rise some in all grades and perceived risk has increased some among the 12th graders since 2009. For 10th and 12th graders perceived risk of trying crack rose after 2007.

- Use of *cocaine*⁸ in general began to decline a year earlier than crack, probably because crack was still in the process of diffusing to new parts of the country, being still quite new. Between 1986 and 1987 the annual prevalence rate for cocaine dropped dramatically, by about one fifth in all three populations being studied at that time—12th graders, college students, and young adults. The decline occurred when young people finally began to view experimental and occasional use—the type of use in which they thought they would be most likely to engage—as more dangerous. This change was probably influenced by the extensive media campaigns that began in the preceding year, but also almost surely by the highly publicized cocaine-related deaths in 1986 of sports stars Len Bias and Don Rogers. By 1992 the annual prevalence of cocaine use had fallen by about two thirds among the three populations for which long-term data are available (12th graders, college students, and young adults).

During the resurgence of illicit drug use in the 1990s, however, cocaine use in all five populations increased once again, both beginning and ending in a staggered pattern by age, consistent with a cohort effect. Use rose among 8th graders from 1991 to 1998, among 10th and 12th graders from 1992 to 1999, among college students from 1994 to 2004, and among young adults from 1996 through 2004. As with crack, all five populations showed some decline in cocaine use in 2008 through 2011 with little change in 2012 and 2013. Annual prevalence rates in 2013 were 1.0%, 1.9%, 2.6%, 2.7%, and 3.9% for the five populations, respectively. For a few years (1996–1999) 12th graders had higher prevalence rates than did the young adults; but because of the staggered declines in use, young adults have had the highest prevalence rates in all years since then (see Table 2-4).

⁸ Unless otherwise specified, all references to cocaine concern the use of cocaine in any form, including crack.

The story regarding attitudes and beliefs about cocaine use is informative. Having risen substantially after 1986, the perceived risk of using cocaine showed some (nonsignificant) decline in 1992 among 12th graders. In 1993, perceived risk for cocaine powder fell sharply in all grades and disapproval began to decline in all grades, though not as sharply as perceived risk. During this time cocaine use was making a comeback. The decline in perceived risk had virtually ended by 1995 among 8th graders, by 1998 among 10th graders, and by 2001 among 12th graders, suggesting a cohort effect at work in this important *belief*, which tends to drive use. Tenth graders' perceived risk for trying cocaine powder rose further after 2007, and 12th graders' disapproval of trying cocaine also has increased in recent years.

The perceived availability of cocaine among 12th graders rose steadily from 1983 to 1989, suggesting that availability played *no* role in the substantial downturn in use that occurred after 1986. After 1989, however, perceived availability fell some among 12th graders—which may be explained in part by the greatly reduced proportions of 12th graders who said they have any friends who use, because friendship circles are an important part of the supply system. After 1995, availability began a long and substantial drop among 8th graders, as it did after 1998 among 10th graders and after 2006 among 12th graders. Twelfth graders' perceived availability rose some in 2013 for the first time since 2007.

- Use of **PCP**, measured and reported only for 12th graders and young adults, fell sharply among 12th graders between 1979 and 1982, from an annual prevalence of 7.0% to 2.2%. It reached a low point of 1.2% in 1988, rose some in the 1990s during the relapse period in the drug epidemic, reaching 2.6% by 1996, and then declined to 0.7% in 2013. For young adults, the annual prevalence rate has fluctuated between 0.1% and 0.6%, but has remained quite low in recent years, standing at 0.2% in 2013.
- Looking at the long-term trends, we see that the annual prevalence of **heroin** use among 12th graders fell by half between 1975 (1.0%) and 1979 (0.5%), then stabilized for 15 years, through 1994. Heroin use was also stable in the early 1990s among the other four populations covered here (see Table 2-2). Then, in 1994 for 8th graders and in 1995 for all other groups, use suddenly increased, with rates doubling or tripling in one or two years for 12th graders, college students, and young adults, and then remaining at the new higher levels among all five populations for the rest of the decade. After the period 1999 to 2001, heroin use fell back to lower levels than were observed in the mid- to late-1990s. Most of that decline was in heroin use without a needle, which we believe was largely responsible for the increase in use in the first half of the 1990s. In sum, all age groups except for the young adults had annual prevalence rates of heroin use in 2013 that were well below recent peaks (by roughly one half to two thirds). Young adults have remained at peak rates (0.5–0.6% in 2008–2013), perhaps due in part to a cohort effect working its way up the age spectrum. Twelfth graders did show a significant increase

to 0.7% annual prevalence in 2010 for heroin use with a needle, though there was no evidence of such an increase in any of the other four populations, which left us cautious about that finding. However, the 2011 rate provided some confirmation that an increase did occur—annual prevalence was at 0.6%, which, except for 2010, was higher than any rate reported since 1995 when this question was first asked. There is little evidence of any ongoing trend at present—indeed, the 12th graders’ annual prevalence for heroin use with a needle was down to 0.4% in 2013, suggesting that if there was an increase in use, it was short-lived. Four of the five populations show annual prevalence rates of either 0.5% or 0.6% in 2013; college students are the exception, at 0.3%.

Two factors very likely contributed to the upturn in heroin use in the 1990s. One is a long-term decline in the perceived risk of harm, probably due to generational forgetting, because it had been a long time since the country had experienced a heroin epidemic along with accompanying publicity about its casualties. The second factor, not unrelated to the first, is that in the 1990s the greatly increased purity of heroin allowed it to be used by means other than injection. This may have lowered an important psychological barrier for some potential users, making heroin use less aversive and seemingly less addictive and less risky in general, because avoiding injection reduces the likelihood of transmission of HIV, hepatitis, or other serious blood-borne diseases. By introducing some new questions on heroin use in 1995, we were able to show that significant proportions of past-year users in all five populations were indeed taking heroin by means other than injection at that point (see Table 2-2 and chapter 4 in both *Volume I* and *Volume II* for details).

The risk perceived to be associated with heroin fell for more than a decade after the study began, with 60% of the 1975 twelfth graders seeing a great risk of trying heroin once or twice, and only 46% of the 1986 twelfth graders saying the same. Between 1986 and 1991, perceived risk rose some, from 46% to 55%, undoubtedly reflecting the newly recognized threat of HIV infection associated with heroin injection. After 1991, however, perceived risk began to fall once again (to 51% by 1995), this time perhaps reflecting the fact that the newer heroin available on the street could be administered by methods other than injection. Between 1996 and 1998, perceived risk among 12th graders rose—possibly as the result of an anti-heroin campaign launched by the Partnership for a Drug-Free America in June 1996, as well as the visibility of heroin-related deaths of some celebrities in the entertainment and fashion design worlds (what we call the “unfortunate role models”). The perceived risk of trying heroin decreased among 12th graders in 1999, however, foretelling a significant increase in their use of the drug in 2000. In 2001, as the perceived risk of trying heroin increased slightly, 12th-grade use declined significantly. In recent years there has been little systematic change in the perceived risk nor in the very high levels of disapproval of heroin use.

Questions about the degree of risk perceived to be associated with heroin use were introduced into the questionnaires for 8th and 10th graders in 1995. The questions asked specifically and only about use “without using a needle” because we thought this was the form of heroin use of greatest concern at that point. (Similar questions were asked of 12th graders, as well, in one of the six questionnaire forms used in 12th grade.) In general, perceived risk for **heroin use without a needle** began rising after 1995, leveled for a while, and then began rising further. Perceived risk held fairly steady among 8th and 10th graders since it was first measured.

- The use of **narcotics other than heroin** is reported only for 12th graders and older populations because we believe that younger students are not accurately discriminating among the drugs that should be included or excluded from this general class. Use declined gradually over most of the first half of the study in these three older groups. Twelfth graders had an annual prevalence rate in 1977 of 6.4%, which fell to 3.3% by 1992. But after about 1992 or 1993, all of the older age groups showed continuing increases for a decade or more, through 2003 or 2004, before stabilizing. An updating of the list of examples given in the question stem in 2002 (to include Vicodin and OxyContin) led to an increase in reported prevalence. After a considerable increase in use from 1992 through 2001, during the relapse phase of the general epidemic and going beyond it, the use of narcotics other than heroin remained relatively constant at high levels through 2010; it then showed some decline in 2011 and 2012 among all three groups (see Table 2-2). In 2013 use continued to decline among 12th graders and young adults but leveled among college students.

The specific drugs in this class are listed in Table E-4 in appendix E of *Volume I*. Among these, **Vicodin**, **codeine**, **OxyContin**, and **Percocet** are commonly mentioned by 12th graders in recent years. In 2013 **hydrocodone** was added to the list of specific narcotics other than heroin, and was the most frequently mentioned.

- In 2002, specific questions were added for **Vicodin** and **OxyContin**, and the observed prevalence rates suggest that these two drugs likely help to account for the upturn in use of the general class of narcotics other than heroin. In 2003, Vicodin had attained surprisingly high prevalence rates in the five populations under study here—an annual prevalence of 2.8% in 8th grade, 7.2% in 10th grade, 10.5% in 12th grade, 7.5% among college students, and 8.6% among young adults. In 2013 the rates were down for all age groups: 1.4%, 4.6%, 5.3%, 4.4%, and 6.2%, respectively. OxyContin started with lower annual prevalence rates than those for Vicodin across all age groups in 2002 but given the highly addictive nature of this narcotic drug, the rates were not inconsequential. Annual prevalence for OxyContin increased in 2003 with slight further increases and leveling through 2011. In 2012 it dropped somewhat in all five populations to annual prevalence rates below the 2003 levels: 1.6%, 3.0%, 4.3%, 1.2%, and 2.3% for 8th, 10th, and 12th grades, college students, and young adults. In 2013 all 5 populations except 12th

graders showed increases. OxyContin use showed significant increases in 2009 among college students and young adults; but these were more than offset by significant decreases in 2010, suggesting that the 2009 values were overestimates (attributable to sampling error). Since 2010 OxyContin use has generally been in decline, while Vicodin use has declined among all groups except college students, where use has held steady. Because OxyContin has received considerable adverse publicity in recent years, it is possible that perceived risk (which we did not measure for this drug until 2012) increased. But because its use appears to have originated in several fairly delimited geographic areas, it seems likely that OxyContin was diffusing to new communities for some time, which may have delayed the turnaround in its use. We believe a similar process happened earlier when crack use and ecstasy use were rising. Questions on perceived risk of Vicodin and OxyContin were added to the 8th and 10th grade questionnaires in 2012; perceived risk is relatively low (and fell significantly in 2013) in both grades.

- Annual prevalence of *tranquilizer* use among 12th graders saw a long and substantial decline from 11% in 1977 to 2.8% in 1992. After 1992, use increased significantly among 12th graders as did most drugs, reaching 7.7% in 2002 (but the question was revised slightly in 2001 to include Xanax as an example of a tranquilizer, so a small portion of the increase may be an artifact). Since then, annual prevalence has leveled or even dropped a bit (4.6% in 2013). Reported tranquilizer use also increased modestly among 8th graders, from 1.8% in 1991 to 3.3% in 1996, before declining to 2.6% in 1998. It remained between 2.4% and 2.8% until 2011, when it declined significantly to 2.0%. It was at 1.8% in 2013. As with a number of other drugs, the downturn in use began considerably earlier among 8th graders compared to their older counterparts. Among 10th graders, annual prevalence remained stable between 1991 and 1994 at around 3.3%, and then increased significantly to 7.3% by 2001 (possibly including some artifact, as noted above). Since 2001 tranquilizer use has declined very gradually in all three grades. After a period of stability, college student use showed an increase between 1994 and 2003 (to 6.9%), more than tripling in that period. Since then there has been a gradual decline there as well, to 3.4% by 2012 followed by a nonsignificant increase to 4.4% in 2013. For the young adult sample, after a long period of decline, annual prevalence more than doubled between 1997 and 2002 to 7.0%, with little change thereafter (5.4% in 2013). Thus, while there was a considerable increase in use in all five populations, which reflected in part a cohort effect that first began in the early 1990s among 8th graders, that increase is clearly over and there has been some downward correction in recent years. Most of the reported tranquilizer use in recent years has involved *Valium*, *Xanax*, and more recently *Klonopin* (see Table E-3 in appendix E of *Volume I*).
- The long-term gradual decline in *sedative (barbiturate)* use among 12th graders, which has been observed since the start of the study in 1975, halted in 1992. (Data are not included here for 8th and 10th graders, again because we

believe that these students have more problems with proper classification of the relevant drugs.) Use among 12th graders then rose considerably during the relapse phase in the drug epidemic, from 2.8% in 1992 to 6.7% by 2002—but still well below the peak rate of 10.7% in 1975; use has shown a modest improvement since 2002 (4.8% in 2013). The 2013 annual prevalence of this class of drugs was lower among young adults (3.4% and college students (2.7%) than among 12th graders (4.8%). Use among college students began to rise a few years later than it did among 12th graders, again likely reflecting a cohort effect, but by 2011 it was at its lowest point since 1998. There followed a small increase in 2012 and 2013. Among young adults, sedative (barbiturate) use increased since the early 1990s, rising from 1.6% in 1992 to 4.4% in 2004. It stands at 3.4% in 2013, after declining some in recent years.

- ***Methaqualone***, although another sedative drug, has shown a trend pattern quite different from barbiturates. Methaqualone use rose among 12th graders from 1975 to 1981, when annual prevalence reached 7.6%. Its use then fell sharply, declining to 0.2% by 1993 before rising some during the general drug resurgence in the 1990s, although only to 1.1% by 1996. Prevalence rates have shown little consistent change since then, with use standing at 0.4% in 2012. (The question was dropped in 2013 to make room for other questions.) Use also fell in the 1980s among young adults and college students, who had annual prevalence rates by 1989—the last year they were asked about this drug—of only 0.3% and 0.2%, respectively. In the late 1980s, shrinking availability may well have played a role in the decline, as legal manufacture and distribution of methaqualone ceased. Because of very low usage rates, only 12th graders were asked about use of this drug for some years, before it was dropped. Methaqualone is one of the very few illegal drugs, the use of which has dropped to relatively negligible levels during the life of MTF. PCP is another.
- Clearly use of most of the several classes of ***psychotherapeutic drugs***—sedatives (barbiturates), tranquilizers, and narcotics other than heroin—has become a larger part of the nation’s drug abuse problem. While the rise in use appears to have halted, most rates remain relatively high. During much of the 1990s and into the 2000s, we were seeing a virtually uninterrupted increase among 12th graders, college students, and young adults in the use of all of these drugs, which had fallen from favor from the mid-1970s through the early 1990s. These drugs continued to rise, even after the increase in use of most illegal drugs ended in the late 1990s and began to reverse.
- For many years, five classes of illicitly used drugs—***marijuana, amphetamines, cocaine, LSD, and inhalants***—had an impact on appreciable proportions of young Americans in their late teens and 20s. In 2013, twelfth graders showed annual prevalence rates for these drugs of 36.4%, 8.7%, 2.6%, 2.2%, and 2.5%, respectively, reflecting declines in most of them, especially LSD. Among college students in 2013, the comparable annual prevalence

rates were 35.5%, 10.6%, 2.7%, 2.6%, and 0.5%; for all young adults the rates were 32.2%, 7.8%, 3.9%, 2.0%, and 0.5%. Because **LSD** use has fallen so precipitously since 2001 in all five populations, it no longer ranks as one of the major drugs of abuse, whereas **narcotics other than heroin** have become quite important due to the long-term rise in use that began in the 1990s. These narcotics now have annual prevalence rates of 5–7% among 12th graders, college students, and young adults. **Tranquilizers** have also become more important due to a similar rise in use, with prevalence rates in 2013 of about 4–5% across the same three populations, as have **sedatives (barbiturates)**, with rates of 4.8%, 2.7%, and 3.4%, respectively. The increase in use of these prescription-type drugs, combined with the decline in use of many illegal drugs, means that the misuse of prescription-type drugs clearly became a more important part of the nation’s drug problem.

- **Ecstasy (MDMA)** joined this set of long-established, more prevalent drugs for a period of time, but annual prevalence rates for ecstasy dropped considerably between 2000 and 2009, making ecstasy less prevalent than a number of other illicit drugs. In 2012 annual use declined significantly for all three grades. The 2013 rates are roughly half the peak rates observed in 2001 for 8th, 10th, and 12th graders, college students and young adults.
- In 8th grade **inhalants** rank second only to marijuana among the illicitly used drugs in terms of thirty-day, annual and lifetime prevalence. Because the use of inhalants reflects a form of illicit psychoactive drug use, and because of its importance among the younger adolescents, an additional index of “any illicit drug use including inhalants” was introduced in Tables 2-1 through 2-3. The inclusion of inhalants makes relatively little difference in the illicit drug index prevalence rates for the older age groups, but considerable difference for the younger ones. For example, in 2013 the proportion of 8th graders reporting any illicit drug use in their lifetime, exclusive of inhalants, was 20%, whereas including inhalants raised the figure to 26%.

Several drugs have been added to MTF’s coverage in recent years, and they are all discussed in *Volumes I and II*. These include **ketamine**, **GHB**, and **Rohypnol**, which are so-called “club drugs” (in addition to LSD and ecstasy). In general, these drugs have low prevalence rates that have declined over the past several years among 8th, 10th, and 12th graders. For that reason GHB and ketamine were dropped from the 8th and 10th grade surveys in 2012. For 12th graders, the 2013 annual prevalence rate was 1.4% for **ketamine** and 1.0% for **GHB**. Annual prevalence of **Rohypnol** was 0.4% for 8th graders, 0.6% for 10th graders, and 0.9% for 12th graders in 2013.

- The two narcotic drugs added to MTF’s coverage in 2002—**OxyContin** and **Vicodin**—show considerably higher prevalence rates, as noted earlier.
- In 2009 a question on past-year use without a doctor’s orders of **Adderall**, an amphetamine used to treat ADHD, was added to the MTF study for all three

grades and for the follow-up respondents. The 2013 annual prevalence rates were 1.8%, 4.4%, 7.4%, 10.7%, and 7.0% for 8th graders, 10th graders, 12th graders, college students, and young adults, respectively. The high rate of use among college students likely stems from its being used to stay awake and alert while studying for exams and doing assigned course work. Adderall use has not shown a clear trend in most of the five populations in the past five years, with the exception that it has been rising among 12th graders, from 5.4% in 2009 to 7.4% in 2013, though the increase did not continue in 2013. This increase could reflect a greater use of this drug among high school seniors attempting to enhance their academic performance, much as college students appear to have been doing for at least several years.

- Questions on use of *Provigil* (a prescription stay-awake drug used for narcolepsy, shift work, etc.) were added to the 12th-grade and follow-up questionnaires in 2009. In 2011 rates of Provigil use in the past year by 12th graders, college students, and young adults were 1.5%, 0.2%, and 0.3%, respectively, suggesting that this drug had not made serious inroads in terms of non-medically supervised use. Given the low use, questions on Provigil were dropped from the study in 2012.
- Questions on use of *salvia* (a plant-based psychoactive drug with dissociative effects, which is currently legal in most states) also were added to the 12th-grade and follow-up questionnaires in 2009 and were added to the 8th and 10th grade questionnaires in 2010. Unlike Provigil, the rates for annual prevalence of salvia were not inconsequential; in 2011, the rates were 1.6% among 8th graders, 3.9% among 10th graders, 5.9% among 12th graders, 3.1% for college students, and 2.2% for young adults (see Table 2-2). But by 2013 the rates for salvia use had declined in all five populations, suggesting that the popularity of this drug has peaked. Still, 3.4% of the 12th graders report some past-year use in 2013, but the college and young adult populations have rates at or below 1%.
- *Anabolic steroid* use occurs predominantly among males. In 2013 the annual prevalence rates for males in 8th, 10th, and 12th grades were 0.7%, 1.3%, and 2.2%, compared with 0.4%, 0.5%, and 0.7% for females. Between 1991 and 1998, the overall annual prevalence rate was fairly stable among 8th and 10th graders, ranging between 0.9% and 1.2%. In 1999, however, use jumped from 1.2% to 1.7% in both grades. Almost all of that increase occurred among males, from 1.6% in 1998 to 2.5% in 1999 in 8th grade and from 1.9% to 2.8% in 10th grade. Thus, rates among males increased by about half in a single year, which corresponded to stories in the news media about the use of androstenedione, a steroid precursor, by baseball home-run king Mark McGwire. Since then, among all 8th graders, anabolic steroid use has declined by almost two thirds to 0.6% in 2012 and 2013. Among 10th graders, use continued to increase, reaching 2.2% in 2002, suggesting a cohort effect, but then declined by more than half to 0.8% by 2012 and 2013. Among 12th graders, annual prevalence rose significantly to 2.4% in 2001, but then

decreased to 1.5% in 2013. Use generally has been much lower among college students and young adults, and declined to 0.2–0.4% annual prevalence in 2011 and 2012 in these groups. In 2013 the prevalence rates rose nonsignificantly, to 0.8% for college students and 0.5% for young adults.

- Two other substances used primarily by males to develop physique and physical strength were added to the question set in 2001. One is **androstenedione**, a precursor to anabolic steroids and available over the counter until early 2005. Among males, where use has tended to be more concentrated, the 2013 annual prevalence rates were 0.9%, 1.4%, and 1.0% in grades 8, 10, and 12, respectively. Among females, the rates were 0.6%, 0.4%, and 0.3%. As discussed in chapter 10 of *Volume I*, the proportion of young males who report past-year use of **androstenedione** and/or **steroids** was appreciable. In 2001, when the “andro” question was introduced, the annual prevalence rate for androstenedione and/or steroids was 8.0% for 12th-grade boys. The rate has fallen considerably in all three grades since then; among 12th-grade boys it was 2.9% in 2013.
- **Creatine** is another substance taken to enhance physique; it is not classified as a drug but rather as a type of protein supplement. Because we believed its use was often combined with the use of steroids and androstenedione, we introduced a question on it in 2001 and found prevalence of use to be very high. Among males, who again are the primary users, the 2013 annual prevalence for creatine was 3.3%, 10.9%, and 17.7% in grades 8, 10, and 12. In other words, one in every six 12th-grade boys used creatine in the prior year. For girls, the rates were far lower at 0.9%, 0.8%, and 1.2%, respectively.
- Beginning in 1982, MTF included a set of questions about the use of *nonprescription* stimulants, including stay-awake pills, diet pills, and the so-called “look-alikes” (see chapter 10 of *Volume I* for more detailed findings). One important finding shown in that chapter (see Table 10-3) is that the use of each of these over-the-counter substances is correlated positively with the respondent’s use of illicit drugs. *In other words, there is a more general propensity of some youth to use or not use psychoactive substances, regardless of their legal status*⁹.

The annual prevalence among 12th graders of over-the-counter **stay-awake pills**, which usually contain caffeine as their active ingredient, nearly doubled between 1982 and 1988, increasing from 12% to 26%. After 1988 this statistic fell considerably reaching 3.2% by 2010, the lowest level ever reported, where it remained in 2013.

⁹ For a more extended discussion and documentation of this point, see Johnston, L.D. (2003). Alcohol and illicit drugs: The role of risk perceptions. In Dan Romer (Ed.), *Reducing adolescent risk: Toward an integrated approach* (pp. 56-74). Thousand Oaks, CA: Sage. Available at <http://www.monitoringthefuture.org/pubs/chapters/ldj2003.pdf>.

- The *look-alike stimulants* have also shown considerable falloff since we first measured their use in 1982. Among 12th graders, annual prevalence decreased from 10.8% in 1982 to 5.2% in 1991. Their use rose only slightly during the relapse phase of the illicit drug epidemic in the 1990s, reaching 6.8% in 1995—roughly where it stayed through 2001. Since then the use of look-alikes decreased to 1.7% by 2010, the lowest level ever reported, which is also where it stands in 2013.
- Among 12th graders, annual prevalence rates for over-the-counter *diet pills* have fluctuated widely over the life of the study. Annual prevalence declined from 21% in 1983 to 8% a decade later, increased to 15% by 2002, then declined significantly to 4.3% by 2010, the lowest point since the questions were added in 1982. Use of this class of drugs in 2013 was up only slightly, to 5.3%. Among 12th-grade girls in 2013 substantial proportions were using over-the-counter diet pills—11.6% had tried diet pills by the end of senior year, 7.3% used them in the past year, and 3.3% used them in just the past 30 days.
- One additional type of over-the-counter drug was added to the 8th-, 10th-, and 12th-grade questionnaires in 2006—*dextromethorphan*, a cough suppressant found in many cough and cold medications. Respondents were asked, “How often have you taken cough or cold medicines to get high?” The proportions indicating such use in the prior 12 months were 4%, 5%, and 7% in grades 8, 10, and 12 in 2006—not inconsequential proportions. In 2013, the rates were similar (3%, 4%, and 5%). The 12th graders did show a significant 1.2 percentage-point decline in 2011 but there was little change after that.

College–Noncollege Differences in Illicit Drug Use

- For analytic purposes, “college students” are defined here as those respondents one to four years past high school who are actively enrolled full-time in a two- or four-year college in March of the year of the survey. For nearly all categories of illicit drugs, college students show *lower* rates of use than their age-mates not in college. However, for a few categories of drugs—including *any illicit drug*, *marijuana*, and *hallucinogens*—college students show annual usage rates that are about average for all high school graduates their age. (College students are about average on the index of any illicit drug use because they have average rates of marijuana use, which largely drives the index.)
- Although college-bound 12th graders have generally had below-average rates of use on *all* of the *illicit drugs* while they were in high school, these students’ eventual use of some illicit drugs attained equivalence with, or even exceeded, the rates of their age-mates who do not attend college. As MTF results have shown, this college effect of “catching up” is largely explainable in terms of differential rates of leaving the parental home after high school graduation and of getting married. College students are more likely than their age peers to

have left the parental home, and they tend to defer marriage, leaving them comparatively less constrained.¹⁰

- In general, the substantial decline in *illicit substance use* among American college students after 1980 paralleled that of their age peers not in college. Further, for the 12-year period 1980 to 1992, all young adult high school graduates through age 28, as well as college students taken separately, showed trends that were highly parallel, for the most part, to trends among 12th graders (see chapter 9 of *Volume II*). However, after 1992 a number of drugs showed an increase in use among 12th graders (as well as 8th and 10th graders), but *not* among college students and young adults for some period of time.

This divergence, combined with the fact that the upturn began first among 8th graders (in 1992), suggests that cohort effects were emerging for illicit drug use, as discussed earlier. Indeed, as those heavier using cohorts of 12th graders entered the college years, we saw a lagged increase in the use of several drugs in college. For example, annual prevalence reached a low point among 12th graders in 1992 for a number of drugs (e.g., *cocaine, amphetamines, sedatives, tranquilizers, narcotics other than heroin, and any illicit drug other than marijuana*) before rising thereafter. Among college students, those same drugs reached a low two years later in 1994, and then began to rise gradually. Then, in 1998, as *marijuana* use already was declining in secondary school, we saw a sharp increase in its use among college students. Consistent with our earlier predictions, the evidence for cohort effects resulting from generational replacement is quite substantial.

Male–Female Differences in Substance Use

- Regarding gender differences in the three older populations (12th graders, college students, and young adults), males are more likely to use most *illicit drugs*, and the differences tend to be largest at the higher frequency levels. For example, 2013 *daily marijuana* use rates among 12th graders are 8.9% for males versus 3.8% for females. Among all young adults (ages 19 to 30) the rates are very similar at 8.5% for males versus 3.8% for females, as they are among college students at 8.7% for males versus 2.9% for females.
- The 8th- and 10th-grade samples evidence fewer and smaller gender differences in the use of drugs than the older populations—perhaps because girls tend to date and then emulate older boys, who are in age groups considerably more likely to use drugs. While the rate of prior-year *marijuana* use is slightly higher for males, the rate for the use of *any illicit drug other*

¹⁰ Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates.

than marijuana generally has tended to be slightly higher for females. There is little or no gender difference in 2013 among 8th graders in their use of *LSD, MDMA, cocaine, crack, other cocaine, heroin, OxyContin, Vicodin, Ritalin, Adderall, methamphetamine, Bath Salts, Rohypnol, or over-the-counter cough and cold medications* taken to get high. The use of *inhalants, alcohol, flavored alcoholic beverages, and being drunk* is actually slightly higher among females in 8th grade. By 10th grade use among boys catches up and in some cases surpasses usage rates among girls on many of these drugs.

TRENDS IN ALCOHOL USE

- Several findings about *alcohol* use in these age groups are noteworthy. First, despite the fact that it is illegal for virtually all secondary school students and most college students to purchase alcoholic beverages, they have had a substantial amount of experience with alcohol. Alcohol has been tried by 28% of 8th graders, 52% of 10th graders, 68% of 12th graders, 78% of college students, and 86% of young adults (19 to 28 years old). Current use (use in past 30 days) is also widespread. Most important, perhaps, is the prevalence of *occasions of heavy drinking*—five or more drinks in a row at least once in the prior two-week period—which was reported by 5% of 8th graders, 14% of 10th graders, 22% of 12th graders, 35% of college students, and 35% of young adults who were surveyed in 2013. Such occasional heavy or “binge” drinking peaks in the early 20s and recedes with age after that, reflected by the 31% rate found among 29- to 30-year-olds.

Alcohol use did not increase as use of other illicit drugs decreased among 12th graders from the late 1970s to the early 1990s, although it was common to hear such a “displacement hypothesis” asserted. MTF demonstrates that the opposite seems to be true. After 1980, when illicit drug use was declining, the *monthly prevalence of alcohol use* among 12th graders also declined gradually, but substantially, from 72% in 1980 to 51% in 1992. *Daily alcohol use* declined by half over the same interval, from a peak of 6.9% in 1979 to 3.4% in 1992; the prevalence of drinking *five or more drinks in a row* during the prior two-week interval fell from 41% in 1983 to 28% in 1993—nearly a one-third decline. When illicit drug use rose again in the 1990s, alcohol use (particularly binge drinking) rose some as well—albeit not as sharply as marijuana use. In the late 1990s, as illicit drug use leveled in secondary schools and began a gradual decline, similar trends were observed for alcohol. Therefore, long-term evidence indicates that alcohol use moves much more in concert with illicit drug use than counter to it. From 2007 through 2011, however, alcohol use continued its long term decline, reaching historic lows in the life of the study, whereas marijuana use was rising gradually. In 2012 the story became more complicated, with marijuana use still rising among college students, leveling among 12th graders and actually declining a bit among 8th and 10th graders. Thirty-day alcohol use rose in all groups except the 8th

graders in 2012 (who had a significant decline of 1.7 percentage points); the increase of 4.2 percentage points among college students was significant. In 2013 annual marijuana use increased in all groups except 12th grade, and the increase among young adults was significant. In contrast, 30-day alcohol use dropped for all groups including significant drops for 12th graders and college students.

College–Noncollege Differences in Alcohol Use

- Trends in *alcohol* use among college students are quite different than those for 12th graders or noncollege respondents of the same age as the college students (see Figure 9-14 in *Volume II*). From 1980 to 1993, college students showed considerably less drop-off in monthly prevalence of *alcohol* use (82% to 70%) than did 12th graders (72% to 51%), and also less decline in *occasions of heavy drinking* (from 44% to 40%) than either 12th graders (41% to 28%) or their noncollege age-mates (41% to 34%). Because both the noncollege 19- to 22-year-olds and high school students were showing greater declines, the college students stood out as having maintained a high rate of episodic heavy (or binge) drinking. Since 1993, this behavior has not changed a great deal among college students—their rate of binge drinking in 2013 was 35%, down modestly from their 1993 (and 2008) rate of 40%. The rate among noncollege age-mates was 31% in 2013 (and 30% in 2012)—down from 34% in 1993. The 12th graders’ rate, after increasing to 32% in 1998, dropped to 25% by 2006 where it remained through 2009; it then declined significantly to 22% in 2011—a new low—where it remained in 2013. College students continue to stand out as having a relatively high rate of binge drinking, though at 35% it is still substantially below where it was in 1993 and 2008.

College-bound 12th graders are consistently less likely than their noncollege-bound counterparts to report occasions of heavy drinking, yet the higher rates of such drinking among college students compared to noncollege peers indicate that these 12th graders catch up to and pass their peers in binge drinking after high school graduation. As stated above, we have shown that this differential change after high school is largely attributable to the fact that college students are more likely to leave the parental home and less likely to get married in the four years after high school graduation than their age mates. An MTF journal article also shows that membership in a fraternity or sorority is associated with a greater than average increase in heavy episodic drinking and marijuana use in college.¹¹

- Since 1980, college students have generally had *daily drinking* rates that were slightly lower than their age peers, suggesting that they were more likely to confine their drinking to weekends, when they tend to drink a lot. The rate of

¹¹ McCabe, S. E., Schulenberg, J. E., Johnston, L. D., O’Malley, P. M., Bachman, J. G., & Kloska, D. D. (2005). Selection and socialization effects of fraternities and sororities on U.S. college student substance use: A multi-cohort national longitudinal study. *Addiction, 100*, 512–524.

daily drinking among the *noncollege* group fell from 8.3% in 1980 to 3.2% in 1994, rose to 5.8% by 2000, and dropped some to 4.1% in 2013. Daily drinking by the *college* group also dropped in approximately the same time period, from 6.5% in 1980 to 3.0% in 1995, then increased to 5.0% in 2002; since then it has hovered between 3.6% and 4.8%. By 2013 the two groups have similar rates of daily drinking, which is a change in a long-standing difference. We will have to see if this new parity continues in the future.

Male–Female Differences in Alcohol Use

- Given that the physiological impacts of five drinks are considerably greater for the typical young female versus the typical young male, it is not surprising that we find substantial gender differences in the prevalence of having ***five or more drinks in a row***. Among 12th graders, the rates in 2013 are 18% for females versus 26% for males. This difference has generally been diminishing since MTF began; in 1975 there was a 23-percentage-point difference, versus an 8-point difference in 2013. The proportions indicating in 2013 that they have ***been drunk*** in the prior 30 days are only slightly closer at 23% and 29% for females and males, respectively.
- Among college students and young adults generally, there are also substantial gender differences in alcohol use, with college males drinking the most. In 2013, for example, 43% of college males reported having ***five or more drinks in a row*** over the previous two weeks versus 30% of college females. Since MTF began, this gender difference has narrowed gradually, with the rate declining somewhat for males and increasing somewhat for females.
- College males report considerably higher rates of ***daily drinking*** than college females (5.6% vs. 2.3% in 2013). A similar gender difference also exists in the noncollege group (5.6% vs. 2.7% in 2013).

TRENDS IN CIGARETTE SMOKING

A number of very important findings about ***cigarette smoking*** among American adolescents and young adults have emerged during the life of the study, and we believe that one of the study's more important contributions to the long-term health of the nation has been to document and call public attention to these trends. Despite the demonstrated health risks associated with smoking, young people have continued to establish regular cigarette habits during late adolescence in sizeable proportions, and, during the first half of the 1990s, in rapidly growing proportions. In fact, since MTF began in 1975, cigarettes have consistently remained the class of abusable substances most frequently used on a daily basis by high school students.

- During most of the 1980s, when smoking rates were falling steadily among adults, we reported that smoking among adolescents was *not* declining. Then the situation went from bad to worse. Among 8th and 10th graders, the

current (past 30-day) smoking rate increased by about half between 1991 (when their use was first measured) and 1996; among 12th graders, the current smoking rate rose by nearly one third between 1992 and 1997. MTF played an important role in bringing these disturbing increases in adolescent smoking to public attention during those years, which was the historical period in which major social action was initiated in the White House, the Food & Drug Administration, the Congress, and eventually the state attorneys general, culminating in the 1998 Tobacco Master Settlement agreement between the tobacco industry and the states.

Fortunately—and largely as a result of that settlement, we believe—there have been some important declines in current smoking since 1996 among 8th and 10th graders, and since 1997 among 12th graders. In fact, the declines more than offset the increases observed earlier in the 1990s. In 2009, 7% of 8th graders (down from 14% in 1991 and 21% in 1996) reported smoking one or more cigarettes in the prior 30 days—a decline of two thirds from the recent peak rate. Some 13% of 10th graders were current smokers in 2009 (down from 21% in 1991 and 30% in 1996), representing a drop of nearly six tenths from the recent peak rate. And in 2010, 19% of 12th graders were current smokers (versus 28% in 1991 and 37% in 1997), representing a drop of roughly half from the recent peak. In recent years these declines decelerated, however, and in 2010 they stopped among 8th and 10th graders. Fortunately, there was some further decline in 2011, 2012, and 2013 in all three grades under study. Monthly prevalence of use for all three grades is now at the lowest point in the history of the study.

Several of the important attitudinal changes that accompanied these declines in use ended some years ago (around 2007), leading us to conclude that further improvement in smoking rates will likely have to come from changes in the environment—for example, enacting such policies as tobacco tax increases, further reducing the places in which smoking is permitted, and providing effective quit-smoking programs. In 2009, federal taxes on tobacco products were in fact raised, which may well have contributed to the resumption of declines in use starting in 2011. Despite these very important improvements in the past decade and a half, about one sixth (16%) of young Americans are current smokers by the time they complete high school. Other research consistently shows that smoking rates are substantially higher among those who drop out before graduating, so the estimates here, based on high school seniors, are low for the age cohort as a whole.¹²

Among college students, the peak rate in current smoking (31%) was not reached until 1999—reflecting a cohort effect—after which it declined

¹² For a recent analysis showing much higher smoking rates among 8th graders who later dropped out before completing high school, see Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

moderately to 24% in 2005. In 2006 a significant decline brought it down to 19%. By 2012 current smoking among college students stood at 13%, having fallen more than half since 1999. In 2013 the rate increased slightly to 14%, but this could simply indicate a sampling error effect. Young adults 19 to 28 years old have shown more modest change in rates of current smoking between 2001 (30%) and 2013 (20%)—a decline of one third including a significant decrease in 2012.

- The dangers that survey participants perceive to be associated with *pack-a-day smoking* differ greatly by grade level, and seem to be unrealistically low at all grade levels. Currently, about three quarters of 12th graders (78%) think that pack-a-day smokers run a great risk of harming themselves physically or in other ways, but only 62% of the 8th graders think the same. All three grades showed a decrease in perceived risk between 1993 and 1995, as use was rising rapidly, but a slightly larger and offsetting increase in perceived risk occurred between 1995 and 2000, presaging the subsequent downturn in smoking. After 2000 there was a slight upward drift in perceived risk at all three grade levels, but it leveled off after 2004 in the lower grades and after 2006 at 12th grade. After that the upward drift resumed in all three grades. In 2013 there was a slight drop in perceived risk among 8th and 10th grades and a leveling at 12th grade.
- Disapproval of *cigarette smoking* was in decline for a considerable period: from 1991 through 1996 among 8th and 10th graders, and from 1992 to 1996 among 12th graders. Since then there was a fairly steady increase in disapproval of cigarette smoking in all three grades—at least until 2007 or 2008, when the increase halted. In 2011 and 2012 all three grades showed some further increase in the proportions of students disapproving of smoking, but no further increase occurred in 2013. Undoubtedly the heavy media coverage of the tobacco issue (the settlement with the states attorneys general, the congressional debate, the congressional testimony of the tobacco executives, the eventual state settlements, etc.) had an important influence on these attitudes and beliefs. However, that coverage diminished considerably in 1998, raising the question of whether those changes in youth attitudes would continue. It may well be, of course, that the removal of certain kinds of cigarette advertising and promotion, combined with national- and state-level antismoking campaigns and subsequent significant increases in cigarette prices, have served to sustain and prolong these changes. In terms of media effects, MTF has shown important changes, including more recent substantial declines, in reported recall by students of antismoking ads resulting from both state and national campaigns.¹³

¹³ Johnston, L. D., Terry-McElrath, Y. M., O'Malley, P. M., & Wakefield, M. (2005). Trends in recall and appraisal of anti-smoking advertising among American youth: National survey results, 1997–2001. *Prevention Science*, 6, 1–19. Also unpublished data.

Age- and Cohort-Related Differences in Cigarette Smoking

- Initiation of smoking occurs most often in grades 6 through 9 (i.e., at modal ages 11–12 to 14–15), although according to the 2013 eighth graders, 8% had already initiated smoking in grade 6 or earlier. The initiation rate trails off considerably by 12th grade, although, as we have shown in our follow-up studies, a number of the light smokers in 12th grade make the transition to heavy smoking in the first two years after high school. Analyses presented in this volume and elsewhere have shown that cigarette smoking evidences a clear cohort effect. That is, if a class (or birth) cohort establishes an unusually high rate of smoking at an early age relative to other cohorts, the rate is likely to remain high throughout the life cycle when compared to that of other birth cohorts at equivalent ages.
- As we reported in “Other Findings from the Study” in the 1986 Volume I in this series, some 53% of 12th graders who were half-pack-a-day (or more) smokers in senior year in 1985 said that they had tried to quit smoking but could not. Of those who had been daily smokers in 12th grade, nearly three quarters were still daily smokers seven to nine years later (based on the 1985 follow-up surveys of the Class of 1985), despite the fact that in high school only 5% thought they would “definitely” be smoking five years hence. A subsequent analysis, based on the 1995 follow-up survey, showed similar results. Nearly two thirds (63%) of those who had been daily smokers in 12th grade were still daily smokers seven to nine years later, although in high school only 3% of them had thought they would “definitely” be smoking five years hence. Clearly, the smoking habit is established at an early age, is difficult to break for those young people who have initiated use, and young people greatly overestimate their own ability to quit. Additional data from 8th- and 10th-grade students show us that younger adolescents are even more likely than older ones to seriously underestimate the dangers of smoking.
- MTF surveys of 8th and 10th graders also show that cigarettes are readily available to teens in 2013, even though perceived availability has been dropping for some years in these two grades; 50% of 8th graders and 71% of 10th graders say that cigarettes would be “fairly easy” or “very easy” for them to get, if they wanted them. There was little change in reported availability between 1992 (when these questions were first asked) and 1997. After that, however, perceived availability of cigarettes decreased significantly for 8th and 10th graders, quite likely reflecting the impact of new regulations and related enforcement efforts aimed at reducing the sale of cigarettes to minors (including the Synar amendment, which required states to pass and enforce laws prohibiting the sale and distribution of tobacco products to persons under 18).¹⁴ (Twelfth graders are not asked this question.)

¹⁴ For a more detailed examination of recent changes in youth access to cigarettes, see Johnston, L. D., O’Malley, P. M., & Terry-McElrath, Y. M. (2004). Methods, locations, and ease of cigarette access for American youth, 1997–2002. *American Journal of Preventive Medicine*, 27, 267–276.

College–Noncollege Differences in Cigarette Smoking

- A striking difference in smoking rates has long existed between college-bound and noncollege-bound 12th graders. For example, in 2013, smoking a half pack or more per day is four times as prevalent among the noncollege-bound 12th graders as among the college bound (8.6% vs. 2.1%). Among respondents of college age (one to four years past high school), those not in college also show dramatically higher rates of half-pack-a-day smoking than those who are in college—11.2% versus 2.4%, respectively. Clearly, these important differences precede college attendance.
- In the first half of the 1990s, smoking rose among college students and their same-age peers, although the increases were not as steep for either group as they were among 12th graders. But in 1998 and 1999, while smoking was declining among secondary school students at all grades, smoking continued to increase among college students and their noncollege age peers, reflecting the cohort effect from earlier, heavier smoking classes of 12th graders moving into the older age groups. Between 1991 and 1999, the 30-day prevalence of cigarette smoking by college students rose from 23% to 31%, or by about one third, and daily smoking rose from 14% to 19%, also by about one third. The year 2000 showed, for the first time in several years, a decline in college student smoking; that continued with a significant decline to 23% in 2003, and another significant decline to 19% in 2006. The rate in 2013 was 14%. (Because of the smaller numbers of cases in the college student samples, the trend lines are not always as smooth as they are for most of the other groups discussed here.) A much more modest decline has also been observed among their noncollege peers, but only since 2001; and the difference between their smoking rates and those of 12th graders in the same year have grown very large. A number of in-depth analyses of MTF panel data have revealed that the differences in smoking rates between those who do and do not attend college are evident by the end of 12th grade and have their roots in earlier educational successes and failures.¹⁵

Male–Female Differences in Cigarette Smoking

- In the 1970s, 12th-grade females caught up to and passed 12th-grade males in rates of *current smoking*. Both genders then showed a decline in use followed by a long, fairly level period, with use by females consistently higher, but with the gender difference diminishing. In the early 1990s, another crossover occurred among the 12th graders when rates rose more among males than

¹⁵ Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates. Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

females; thereafter, males have been consistently slightly higher in rates of current smoking. In the lower grades, the genders have generally had similar smoking rates since their use was first measured in 1991.

- Among college students, females had a slightly higher probability of being daily smokers from 1980 through 1994—although this long-standing gender difference was not seen among their age peers who were not in college. However, a crossover occurred between 1994 and 2001, with college males exceeding college females in daily smoking—an echo of the crossover among 12th graders in 1991. Since about 2001 there has been little consistent gender difference in smoking among college students.

RACIAL/ETHNIC COMPARISONS

The three largest ethnic groups in the population—Whites, African Americans, and Hispanics—are examined here for 8th, 10th, and 12th graders. (Sample size limitations simply do not allow accurate characterization of smaller racial/ethnic groups unless data from a number of years are combined. Separate publications from the study have done just that.) A number of interesting findings emerge from the comparison of these three groups; the reader is referred to chapters 4 and 5 of *Volume I* for a full discussion and to MTF Occasional Paper 81¹⁶ for both tabular and graphic documentation of differences among these three ethnic groups across all drugs.¹⁷

African-American 12th graders have consistently shown lower usage rates than White 12th graders for most drugs, both licit and illicit. At the lower grade levels, where few have yet dropped out of school, African-American students also have generally had

¹⁶ Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Miech, R.A. (2014). *Demographic subgroup trends among adolescents for various classes of licit and illicit drugs, 1975–2013* (Monitoring the Future Occasional Paper No. 81). Ann Arbor, MI: Institute for Social Research, University of Michigan. Available at: <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ81.pdf>.

¹⁷ We periodically publish comparisons that contain a number of the smaller racial/ethnic groups in the population, based on data combined for a number of contiguous years in order to attain adequate sample sizes. The first was Bachman, J. G., Wallace, J. M., Jr., O'Malley, P. M., Johnston, L. D., Kurth, C. L., & Neighbors, H. W. (1991). Racial/ethnic differences in smoking, drinking, and illicit drug use among American high school seniors, 1976–1989. *American Journal of Public Health, 81*, 372–377. More recent articles are: Bachman, J. G., O'Malley, P. M., Johnston, L. D., Schulenberg, J. E., & Wallace, J. M., Jr. (2011). Racial/ethnic differences in the relationship between parental education and substance use among U.S. 8th-, 10th-, and 12th-grade students: Findings from the Monitoring the Future Project. *Journal of Studies on Alcohol and Drugs, 72*(2), 279–285. doi: 10.1037/a0031464; Wallace, J. M., Jr., Bachman, J. G., O'Malley, P. M., Johnston, L. D., Schulenberg, J. E., & Cooper, S. M. (2002). Tobacco, alcohol and illicit drug use: Racial and ethnic differences among U.S. high school seniors, 1976–2000. *Public Health Reports, 117* (Supplement 1), S67–S75; Wallace, J. M., Jr., Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Cooper, S. M., & Johnston, L. D. (2003). Gender and ethnic differences in smoking, drinking, and illicit drug use among American 8th, 10th, and 12th grade students, 1976–2000. *Addictions, 98*, 225–234; and Delva, J., Wallace, J. M., Jr., O'Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2005). The epidemiology of alcohol, marijuana, and cocaine use among Mexican American, Puerto Rican, Cuban American, and other Latin American 8th-grade students in the United States: 1991–2002. *American Journal of Public Health, 95*, 696–702. See also Bachman, J. G., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (2010). *Impacts of parental education on substance use: Differences among White, African-American, and Hispanic students in 8th, 10th, and 12th grades (1999–2008)* (Monitoring the Future Occasional Paper No. 70). Ann Arbor, MI: Institute for Social Research. Available at <http://www.monitoringthefuture.org/pubs/occpapers/occ70.pdf>.

lower usage rates for many drugs, though not all. The differences in the upper grades generally have been quite large for some drugs, including *inhalants, LSD specifically, hallucinogens other than LSD, ecstasy (MDMA), salvia, narcotics other than heroin, OxyContin, Vicodin, amphetamines, Ritalin, Adderall, sedatives (barbiturates), and tranquilizers*. But, in 2013 African American 8th graders have rates of use roughly equivalent to White 8th graders for a number of drugs, and for some drugs African Americans actually have a higher annual prevalence, *marijuana* in particular (14% vs. 9%).

- African-American students currently have a much lower 30-day prevalence rate of *cigarette smoking* than do White students (10% vs. 19% among 12th graders in 2013), partly because the smoking rate among African-American students declined from 1980 to 1992, while the rate for White students remained fairly stable. After 1992, smoking rates rose among both White and African-American 12th graders, but less so among the latter. After 1996 (or 1998 in the case of 12th graders) smoking among White students showed a sharp and continuing decline in all three grades for some years, which considerably narrowed the smoking differences between the races, despite some decline among African Americans as well; nevertheless, there remain substantial differences. Smoking rates among Hispanic students have tended to fall in between the other two groups in the upper grades, and have tracked closely to the White smoking rates at 8th grade.
- In 12th grade, *occasions of heavy drinking* are much less likely to be reported by African-American students (13%) than White (26%) or Hispanic students (22%).
- In 12th grade, of the three racial/ethnic groups, Whites have tended to have the highest rates of use on a number of drugs, including *marijuana, hallucinogens, LSD specifically, hallucinogens other than LSD, salvia, narcotics other than heroin, OxyContin specifically, Vicodin specifically, amphetamines, Ritalin specifically, Adderall specifically, sedatives (barbiturates), tranquilizers, alcohol, getting drunk, cigarettes, and smokeless tobacco*. However, in 2013 Hispanics had the highest levels of *marijuana, salvia, and Vicodin* use at 12th grade.
- Hispanics have tended to have the highest usage rate in terms of annual prevalence in 12th grade for a number of the most dangerous drugs, such as *crack* and *crystal methamphetamine (ice)*. From 2009 to 2011, Whites had the highest for *heroin* use, followed by African Americans in 2012 and 2013. From 2010 to 2011 and again in 2013, African Americans were highest for *heroin use with a needle*. In 2012 they were tied with Whites. The difference between Whites and Hispanics appeared to have been eliminated in 2012 for marijuana, salvia, and Ritalin, as use by Hispanics has risen. In 2013 the difference reappeared for marijuana and salvia. Further, in 8th grade, Hispanics have the highest rate of *illicit drug* use overall and the highest rates for most drugs. For example, in 8th grade, the 2013 annual prevalence of

marijuana use for Hispanics is 17%, versus 9% for Whites and 14% for African Americans; the two-week prevalence of *binge drinking* is 8% for Hispanics, 4% for Whites, and 5% for African Americans. Hispanics have the highest rates of use for many drugs in 8th grade, but not for as many in 12th, which suggests that their considerably higher dropout rate (compared to Whites and African Americans) may change their relative ranking by 12th grade.

- With regard to trends, 12th graders in all three racial/ethnic groups exhibited a decline in *cocaine* use from 1986 through 1992, although the decline was less steep among African-American 12th graders because their earlier increase in use was not as large as the increase among White and Hispanic students.
- For virtually *all of the illicit drugs*, the three groups have tended to trend in parallel at 12th grade. Because White 12th graders had the highest level of use on a number of drugs—including *amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—they also had the largest declines; African Americans have had the lowest rates and, therefore, the smallest declines. As mentioned above, there is a convergence between Whites and Hispanics in 12th grade for *Ritalin*.

For a more detailed consideration of racial/ethnic differences in substance use, see the last section of chapter 5 in *Volume I*.

DRUG USE IN EIGHTH GRADE

It is useful to focus specifically on the youngest age group in the study—the 8th graders, most of whom are 13 or 14 years old—in part because the worrisome levels of both licit and illicit drug use that they report help illustrate the nation’s urgent need to continue to address the substance abuse problems among its youth. Further, it is a well-established fact that the earlier young people start to use drugs, both licit and illicit, the more likely they are to experience adverse outcomes.¹⁸¹⁹²⁰

- Among 8th graders in 2013, more than one in four (28%) reports having tried *alcohol* (more than just a few sips), and about one in eight (12%) indicates having already been *drunk* at least once.

¹⁸ Merline, A.C., O’Malley, P.M., Schulenberg, J.E., Bachman, J.G., & Johnston, L.D. (2004). Substance use among adults 35 years of age: Prevalence, adulthood predictors, and impact of adolescent substance use. *American Journal of Public Health, 94*, 96-102.

¹⁹ Zucker, R. A. (2006). Alcohol use and the alcohol use disorders: A developmental-biopsychosocial systems formulation covering the lifecourse. In D. Cicchetti & D. J. Cohen (Eds.), *Developmental psychopathology: Vol. 3. Risk, disorder, and adaptation* (2nd ed., pp. 620–656). Hoboken, NJ: Wiley.

²⁰ Office of the Surgeon General. (2007). *The Surgeon General’s call to action to prevent and reduce underage drinking*. Rockville, MD: Department of Health and Human Services.

- About one seventh of 8th graders in 2013 (15%) has tried *cigarettes*, and one in twenty-two (4.5%) reports having smoked in the prior month. Shocking to many adults is the fact that only 62% of 8th graders recognize that there is great risk associated with smoking one or more packs of cigarettes per day. While an increasing proportion of youth will recognize the risk by 12th grade, for many this is too late, because they will have developed a smoking habit by then.
- Among 8th grade males in 2013, 10% tried *smokeless tobacco*, 4% used it in the past month, and 0.9% used it daily. Rates are much lower among females.
- One 8th grader in nine (11%) reports ever trying *inhalants*, and one in 43 (2.3%) reports inhalant use in just the month prior to the 2013 survey. This is the only class of drugs for which use is substantially higher in 8th grade than in 10th or 12th grade.
- *Marijuana* has been tried by one in every six 8th graders (17%) and has been used in the prior month by about one in every 14 (7.0%). Some 1.1% use it on a daily or near-daily basis in 8th grade.
- A surprisingly large number of 8th graders (4.2%) say they have tried prescription-type *amphetamines* without medical instruction; 1.4% say they have used them in the prior 30 days.
- For most of the *other illicit drugs*, relatively few 8th graders in 2013 say they have tried them. (This is consistent with the retrospective reports from 12th graders concerning the grades in which they first used the various drugs.) But the proportions having at least some experience with them is not inconsequential. Even a rate as low as 3% represents about one child in every 30-student classroom. The 2013 eighth-grade proportions reporting any lifetime experience with the other illicit drugs are: *tranquilizers* (2.9%), *hallucinogens other than LSD* (1.9%), *ecstasy* (1.8%), *cocaine other than crack, LSD*, and *methamphetamine* (all at 1.4%), *crack* (1.2%), *steroids* (1.1% overall, 1.3% among males), *heroin* (1.0%), and *Rohypnol* (0.7%).
- In total, 26% of all 8th graders in 2013 have tried some *illicit drug* (including inhalants), while 9.3%, or one in eleven, have tried *some illicit drug other than marijuana or inhalants*. Put another way, in an average 30-student classroom of 8th graders, about eight have used some illicit drug other than marijuana, including inhalants; and about three have used some illicit drug other than marijuana or inhalants.
- The very large number of 8th graders who have already begun using the so-called “gateway drugs” (*tobacco, alcohol, inhalants, and marijuana*) suggests that a substantial number are also at risk of proceeding further to such drugs as LSD, cocaine, amphetamines, and heroin.

DRUG USE BY AGE 50

Because we have now followed graduating 12th graders into their 50s, we can characterize the drug-using history of today's 50-year-olds (at least those who are high school graduates). This is important, not only because it shows how use by these respondents has developed over the three decades since they left high school, but also because most of them are now themselves the parents of adolescents and young adults. Their own past experiences with drug use may complicate communications with their children regarding drugs; worse, the continuing active use of substances by some of them may set an unfortunate example. The level of lifetime use they have attained is striking (see chapter 4 of *Volume II* for greater detail and discussion).

- Among 50-year-old high school graduates in 2013, we estimate that about three quarters (76%) have tried *marijuana*, and that about two thirds (62%) have tried an *illicit drug other than marijuana*. (These estimates are adjusted to correct for panel attrition, as described in chapter 4 of *Volume II*.)

Their current behavior is far less extreme than those statistics might suggest, but it is not by any means negligible. One in eight (12%) indicates using *marijuana* in the last 12 months, and one in eleven (9%) indicates using *any other illicit drug* in the same period. Their past-month prevalence rates are lower—7.5% and 4.6%, respectively, for marijuana and any other illicit drug. About one in 43 (2.3%) is a *current daily marijuana* user, though substantially more indicate that they have used marijuana daily at some time in the past.

- High proportions of 50-year-old respondents in 2013 have had some experience during their lifetime of using (outside of medical regimen) several of the specific illicit drugs other than marijuana. These include *cocaine* in any form (40%), *amphetamines* (35%), *tranquilizers* (26%), *narcotics other than heroin* (21%), *sedatives (barbiturates)* (20%), and *hallucinogens* of any type (16%). In sum, today's adults in their 50s tend to be a very drug-experienced segment of the population, as might be expected due to the fact that they graduated from high school near the peak of the drug epidemic. To repeat, 76% have tried marijuana and 62% have tried some illicit drug other than marijuana.
- Illicit drugs other than marijuana that have been used in just the prior 12 months by this age group (outside of medical regimen) include *narcotics other than heroin* (3.1%), *tranquilizers* (4.1%), *cocaine* (1.7%), and *noncrack forms of cocaine* (1.5%). Little active use is reported by these respondents for *amphetamines*, *crack*, or *heroin*. (Of course, we would not expect many heavy users of heroin or crack to have remained in the panel studies for this long.)

- **Alcohol** consumption is relatively high among 50-year-olds, with two thirds (67%) indicating that they consumed at least one alcoholic drink in the prior 30 days, 11% reporting current **daily drinking** (defined as drinking on 20 or more occasions in the prior 30 days), and 20% indicating **recent occasions of heavy drinking** (defined as five or more drinks in a row on at least one occasion in the prior two weeks). The rate of recent occasions of heavy drinking is much lower than was exhibited by members of this cohort when they were of high school and college ages.
- About one in six or seven (15%) 50-year-old high school graduates currently smokes **cigarettes**. Almost all of those are current **daily smokers** (10%).

SUMMARY AND CONCLUSIONS

We can summarize the findings on trends as follows: For more than a decade—from the late 1970s to the early 1990s—the use of a number of **illicit drugs** declined appreciably among 12th-grade students, and declined even more among American college students and young adults. These substantial improvements—which seem largely explainable in terms of changes in attitudes about drug use, beliefs about the risks of drug use, and peer norms against drug use—have some extremely important policy implications. One clear implication is that these various substance-using behaviors among American young people are malleable—they *can* be changed. It has been done before. The second is that demand-side (rather than supply-side) factors appear to have been pivotal in bringing about most of those changes. The levels of **marijuana** availability, as reported by 12th graders, have held fairly steady at high levels throughout the life of the study. (Moreover, among students who abstained from marijuana use, as well as among those who quit, availability and price rank very low on their lists of reasons for their not using.) And, in fact, the perceived availability of **cocaine** was actually rising during the beginning of the sharp decline in cocaine and crack use in the mid- to late- 1980s, which occurred when the perceived risk associated with that drug rose sharply. (See the last section of chapter 9, *Volume I*, for more examples and further discussion of this point.)

However, improvements are surely not inevitable; and when they occur, they should not be taken for granted. Relapse is always possible and, indeed, just such a relapse in the longer term epidemic occurred during the early to mid-1990s, as the country let down its guard on many fronts. (See chapter 8 of *Volume I* for a more detailed discussion.)

Over the years, MTF has demonstrated that changes in perceived risk and disapproval have been important causes of change in the use of a number of drugs. These beliefs and attitudes are almost certainly influenced by the amount and nature of public attention paid to the drug issue in the historical period during which young people are growing up. A substantial decline in attention to this issue in the early 1990s very likely explains why the increases in perceived risk and disapproval among students

ceased and began to backslide. News coverage of the drug issue plummeted between 1989 and 1993 (although it made a considerable comeback as surveys—including MTF—began to document that the nation’s drug problem was worsening again), and the media’s *pro bono* placement of ads from the Partnership for a Drug-Free America also fell considerably. (During that period, MTF 12th graders showed a steady decline in their recalled exposure to such ads, and in the judged impact of such ads on their own drug-taking behavior.)

Also, the deterioration in the drug abuse situation first began among our youngest cohorts—perhaps because as they were growing up they had not had the same opportunities for vicarious learning from the adverse drug experiences of people around them and people portrayed in the media—those we have called the “unfortunate role models.” Clearly, there was a danger that, as the drug epidemic subsided in the 1980s and early 1990s, newer cohorts would have far less opportunity to learn through informal means about the dangers of drugs—that what we have called a generational forgetting of those risks would occur through a process of generational replacement of older, more drug-savvy cohorts with newer, more naive ones. *This suggests that as drug use subsides, as it did by the early 1990s, the nation must redouble its efforts to ensure that such naive cohorts learn these lessons about the dangers of drugs through more formal means—from schools, parents, and focused messages in the media, for example—and that this more formalized prevention effort be institutionalized so that it will endure for the long term.*

Clearly, for the foreseeable future, American young people will be aware of the psychoactive potential of a host of drugs and will continue to have access to them—a situation quite different from the one that preceded the late 1960s. (Awareness and access are two necessary conditions for an epidemic.²¹) That means that each new generation of young people must learn the reasons that they should *not* use drugs. Otherwise, their natural curiosity and desire for new experiences will lead a great many to use.

One lesson evident from the changes of the past decade or so is that the types of drugs most in favor can change substantially over time. The illegal drugs began to decline in use in the late 1990s, while prescription drugs, and even over-the-counter drugs, began to gain favor. Today a good many of the drugs having the highest prevalence rates among teens are of this type, including narcotic drugs other than heroin.

Unfortunately, a second relapse phase in America’s youth epidemic of drug use may now be beginning, as indicated by the upturn in marijuana use in recent years. Perceived risk for marijuana (and for Ecstasy) has been falling, and recalled exposure to anti-drug ads has declined sharply in recent years. To a considerable degree the issue has fallen off the national screen (just as happened in the late 1980s and early 1990s), as other urgent matters (including two wars, the rise of terrorism, and a major

²¹ Johnston, LD (1991). Toward a theory of drug epidemics. In L Donhew, HE Sypher, and WJ Bukiski (Eds.), *Persuasive communication and drug abuse prevention* (pp.93-131). Hillsdale, NJ, Earlbaum. Available at www.monitoringthefuture.org/pubs/chapters/ldj1991theory.pdf

recession) have competed for attention. Indeed, this confluence of events is very reminiscent of the period preceding the first relapse—including a considerable decrease in the levels of drug use, little attention paid to the issue by the media or government, a sharp drop in funding for anti-drug prevention programs and ad campaigns, a war and a recession. While marijuana use, specifically, is now receiving more attention, that attention has been focused on the medical use and full legalization, not so much on the consequences of use.

Another lesson that derives from the MTF epidemiological data is that social influences that tend to reduce the *initiation* of substance use also have the potential to deter *continuation* by those who have already begun to use, particularly if they are not yet habitual users. Chapter 5 of *Volume I* shows how increased quitting rates have contributed importantly to downturns in the use of a number of drugs at different historical periods. The lesson is that primary prevention should not be the only goal of intervention programs; early-stage users may be persuaded to quit when their beliefs and attitudes regarding drugs are changed.

The following facts help to put into perspective the magnitude and variety of substance use problems that presently remain among American young people:

- A quarter (26%) of today's 8th graders have tried an ***illicit drug*** (if inhalants are included as an illicit drug), and half (52%) of 12th graders have done so.
- By their late 20s, two thirds (65%) of today's young adults have tried an ***illicit drug***, and about four in ten (39%) have tried some ***illicit drug other than marijuana***, usually in addition to marijuana. (These figures do not include inhalants.)
- Today, about one in eight young adults (12% in 2013) has tried ***cocaine***, and 4.5% have tried it by their senior year of high school, when they are 17 or 18 years old. One in every 56 twelfth graders (1.8%) has tried ***crack***. Among young adults 29–30 years of age, one in 25 (4.0%) has tried crack.
- One in every 15 twelfth graders (6.5%) in 2013 smokes ***marijuana daily***. Among young adults ages 19 to 28, the percentage is about the same (6.2%). Among those same 12th graders in 2013, nearly one in every six (16%) has been a daily marijuana smoker at some time for at least a month, and among young adults the comparable figure is 18%, about one in six.
- About one in five 12th graders (22%) had ***five or more drinks in a row*** on at least one occasion in the two weeks prior to the survey, and we know that such behavior tends to increase among young adults one to four years past high school—that is, in the peak college years. Indeed, 43% of all male college students report such binge drinking. (The study also has documented evidence of ***extreme binge drinking*** with 8% of 12th graders in 2013 indicating having had 10 or more drinks in a row, and 4.4% indicating 15 or more drinks in a row, in the prior two weeks; see Table 5-5e.)

- Even with considerable improvements in smoking rates among American adolescents since the late 1990s, about one in six (16%) of 12th graders in 2013 currently smoke *cigarettes*, and one in twelve (9%) is already a current *daily smoker*. In addition, we know from studying previous cohorts that many young adults increase their rates of smoking within a year or so after they leave high school.

Despite the substantial improvement in this country's drug situation in the 1980s and early 1990s, and then some further improvement beginning in the late 1990s, American secondary school students and young adults show a level of involvement with *illicit drugs* that is among the highest in the world's industrialized nations.²² Even by longer term historical standards in the U.S. these rates remain extremely high, though in general they are not as high as in the peak years of the epidemic in the late 1970s. *Heavy drinking* also remains widespread and troublesome, though it has been declining gradually over a long period and now is at or near historical lows among teens. Of course, the continuing initiation to *cigarette smoking* of a fair-sized, albeit decreasing proportion of young people remains a matter of great public health concern. The declines in youth smoking have decelerated sharply in all grades in recent years although they are at recent low levels and negative youth attitudes about smoking and smokers leveled off several years ago after rising considerably. The improvements in youth smoking overall may be nearing an end unless there is further change in environmental factors, such as cigarette prices (including taxes), advertising and promotion of cigarettes, places where smoking is permitted, and the availability of quit-smoking services. There was, in fact, an increase in federal taxes on tobacco in 2009, with the final rule becoming effective in August, 2010, which may help to explain why all three grades showed further declines in smoking prevalence since 2011.

E-cigarettes present a new challenge, and MTF will soon be able to provide its first estimates of the prevalence of use for this new product. MTF already provides estimates of use of other newer tobacco products such as snus and hookah smoking.

After a long period of improvement, there was evidence in recent years that the use of *smokeless tobacco* has been on the rise among adolescents. Fortunately, the 2011 and 2012 surveys showed some small though

²² A published report from an international collaborative study, modeled largely after MTF, provides comparative data from national school surveys of 15- to 16-year-olds that was completed in 2011 in 36 European countries. It also includes 2011 MTF data from 10th graders in the United States. See Hibell, B., Guttormsson, U, Ahlström, S., Balakireva, O., Bjarnasson, T., Kokkevi, A., & Kraus, L. (Eds.). (2012). *The 2011 ESPAD report Substance Use among Students in 36 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs, The European Monitoring Centre for Drugs and Drug Addiction, the Council of Europe Co-operation Group to Combat Drug Abuse and Illicit Trafficking in Drugs (the Pompidou Group). See also, Johnston, L. et al., American teens are less likely than European teens to use cigarettes and alcohol, but more likely to use illicit drugs. National press release from the University of Michigan's News and Information Services, June 1, 2012. Available at <http://www.ns.umich.edu/new/releases/20420-american-teens-are-less-likely-than-european-teens-to-use-cigarettes-and-alcohol-but-more-likely-to-use-illicit-drugs>.

nonsignificant declines in all three grades, possibly also as a result of the increase in the federal tobacco tax. The fairly recent rise in smokeless tobacco use may well be a result of the introduction and promotion of new products such as snus and dissolvable tobacco.

- Of particular note, abusable prescription drugs (with the exception of *amphetamines*) showed very limited declines from the mid-1990s into the early 2000s, despite the gradual (and in some cases sharp) declines in the use of many of the illegal drugs during that same period. In 2012 *tranquilizer use* held steady at slightly below recent peak levels, though in 2013 use declined in all three grades. The use of *narcotics other than heroin* among 12th graders (the only grade reported for these drugs) is still near peak levels, though fortunately it has declined a bit over the past three years. *Sedative (barbiturate)* use (also reported for 12th graders only) has shown a gradual decline since 1975, but it did not continue into 2012 or 2013. Perceived risk tends to be relatively low for these prescription-type drugs, which we believe is a major reason why their use is relatively high. Perceived risk is rising for amphetamines, narcotics other than heroin, and sedatives (barbiturates), although only the rise for narcotics other than heroin is large enough to be statistically significant.
- Finally, we note the seemingly unending capacity of pharmacological experts and amateurs to discover new substances with abuse potential that can be used to alter mood and consciousness (e.g., bath salts and synthetic marijuana), and of young people to discover the abuse potential of existing products (such as *Robitussin* and plants like *salvia*) and to rediscover older drugs (such as *LSD* and *heroin*). While as a society we have made significant progress on a number of fronts in the fight against drug abuse, we must remain vigilant against the opening of new fronts, as well as the reemergence of trouble on older ones. In particular, we must guard against generational forgetting in our newest cohorts of adolescents due to a lack of public attention to the issue during the time that they are growing up.

One of the dynamics that keeps the drug epidemic rolling is the emergence of new drugs whose hazards are little known. In 1999 we saw this happen with the drug *ecstasy (MDMA)*. Other drugs like *Rohypnol*, *ketamine*, *GHB*, and *OxyContin* appeared in the 1990s and have been added to the list of drugs under study. Recently, questions on use of *salvia*, *Adderall*, and *Provigil* were added to the questionnaires. In 2011 we added *synthetic marijuana*, which turned out to be the second most used illicit drug after natural marijuana, and in 2012 we added *bath salts*. The spread of such new drugs appears to be facilitated and hastened today by young people's widespread use of web-based social networks. We predict a continuous flow of such new substances onto the scene, and believe that the task of rapidly documenting their emergence, establishing their adverse consequences, and quickly demystifying them will remain an important means by which policymakers, researchers, and educators deal with the continuing threats posed by such drugs. We also

Monitoring the Future

anticipate that there will be rediscoveries of older substances, as has been occurring in recent years with respect to the various psychotherapeutic prescription drugs, including *tranquilizers*, *sedatives (barbiturates)*, and *narcotic drugs*.

The drug problem is not an enemy that can be vanquished. It is more a recurring and relapsing problem that must be contained to the extent possible on an ongoing basis. Therefore, it is a problem that requires an ongoing, dynamic response—one that takes into account the continuing generational replacement of our children, the generational forgetting of the dangers of drugs that can occur with that replacement, and the perpetual stream of new abusable substances that will threaten to lure young people into involvement with drugs.

TABLE 2-1
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Any Illicit Drug ^a																								
8th Grade	18.7	20.6	22.5	25.7	28.5	31.2	29.4	29.0	28.3	26.8	26.8	24.5	22.8	21.5	21.4	20.9	19.0	19.6	19.9	21.4	20.1	18.5	20.3	+1.8
10th Grade	30.6	29.8	32.8	37.4	40.9	45.4	47.3	44.9	46.2	45.6	45.6	44.6	41.4	39.8	38.2	36.1	35.6	34.1	36.0	37.0	37.7	36.8	38.8	+2.0
12th Grade	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	50.4	48.2	46.8	47.4	46.7	48.2	49.9	49.1	50.4	+1.3
College Students	50.4	48.8	45.9	45.5	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	51.4	49.1	49.2	50.5	51.0	+0.4
Young Adults	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	60.4	59.7	59.8	59.3	59.3	58.4	59.1	58.9	60.5	+1.6
Any Illicit Drug other than Marijuana ^{a,b}																								
8th Grade	14.3	15.6	16.8	17.5	18.8	19.2	17.7	16.9	16.3	15.8	17.0	13.7	13.6	12.2	12.1	12.2	11.1	11.2	10.4	10.6	9.8	8.7	9.3	+0.6
10th Grade	19.1	19.2	20.9	21.7	24.3	25.5	25.0	23.6	24.0	23.1	23.6	22.1	19.7	18.8	18.0	17.5	18.2	15.9	16.7	16.8	15.6	14.9	15.7	+0.8
12th Grade	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0	30.7	29.5	27.7	28.7	27.4	26.9	25.5	24.9	24.0	24.7	24.9	24.1	24.7	+0.7
College Students	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	25.6	24.8	24.3	23.8	26.7	+2.9
Young Adults	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3	31.6	32.8	33.9	35.2	34.0	34.8	34.2	34.7	32.8	33.3	33.2	32.8	34.2	+1.4
Any Illicit Drug including Inhalants ^{a,c,d}																								
8th Grade	28.5	29.6	32.3	35.1	38.1	39.4	38.1	37.8	37.2	35.1	34.5	31.6	30.3	30.2	30.0	29.2	27.7	28.3	27.9	28.6	26.4	25.1	25.7	+0.6
10th Grade	36.1	36.2	38.7	42.7	45.9	49.8	50.9	49.3	49.9	49.3	48.8	47.7	44.9	43.1	42.1	40.1	39.8	38.7	40.0	40.6	40.8	40.0	41.3	+1.4
12th Grade	47.6	44.4	46.6	49.1	51.5	53.5	56.3	56.1	56.3	57.0	56.0	54.6	52.8	53.0	53.5	51.2	49.1	49.3	48.4	49.9	51.8	50.3	52.0	+1.7
College Students	52.0	50.3	49.1	47.0	47.0	49.1	50.7	55.4	54.4	54.6	53.1	52.3	54.1	52.9	53.9	53.3	52.5	51.0	51.1	50.0	49.7	52.0	50.5	-1.5
Young Adults	63.4	61.2	61.2	58.5	59.0	58.2	58.4	58.5	58.5	59.5	59.0	59.6	60.6	62.5	61.4	61.2	61.2	60.2	59.3	59.3	59.5	59.5	61.7	+2.3
Marijuana/Hashish																								
8th Grade	10.2	11.2	12.6	16.7	19.9	23.1	22.6	22.2	22.0	20.3	20.4	19.2	17.5	16.3	16.5	15.7	14.2	14.6	15.7	17.3	16.4	15.2	16.5	+1.3
10th Grade	23.4	21.4	24.4	30.4	34.1	39.8	42.3	39.6	40.9	40.3	40.1	38.7	36.4	35.1	34.1	31.8	31.0	29.9	32.3	33.4	34.5	33.8	35.8	+2.1
12th Grade	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8	42.3	41.8	42.6	42.0	43.8	45.5	45.2	45.5	+0.3
College Students	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8	51.2	51.0	49.5	50.7	49.1	49.1	46.9	47.5	46.8	47.5	46.8	46.6	49.1	47.7	-1.4
Young Adults	58.6	56.4	55.9	53.7	53.6	53.4	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	57.0	56.7	56.7	55.9	56.0	55.9	56.3	56.5	57.1	+0.6
Inhalants ^{c,d}																								
8th Grade	17.6	17.4	19.4	19.9	21.6	21.2	21.0	20.5	19.7	17.9	17.1	15.2	15.8	17.3	17.1	16.1	15.6	15.7	14.9	14.5	13.1	11.8	10.8	-1.0
10th Grade	15.7	16.6	17.5	18.0	19.0	19.3	18.3	18.3	17.0	16.6	15.2	13.5	12.7	12.4	13.1	13.3	13.6	12.8	12.3	12.0	10.1	9.9	8.7	-1.2
12th Grade	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	11.4	11.1	10.5	9.9	9.5	9.0	8.1	7.9	6.9	-1.0
College Students	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4	12.9	9.6	7.7	9.7	8.5	7.1	7.4	6.3	4.9	6.9	5.5	3.7	5.7	4.3	-1.4
Young Adults	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	10.3	10.9	9.1	9.5	8.9	7.9	7.2	7.2	6.5	-0.7
Nitrites ^e																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.6	1.5	1.4	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	1.1	1.2	1.2	0.6	1.1	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	1.4	1.2	1.3	1.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens ^{b,f}																								
8th Grade	3.2	3.8	3.9	4.3	5.2	5.9	5.4	4.9	4.8	4.6	5.2	4.1	4.0	3.5	3.8	3.4	3.1	3.3	3.0	3.4	3.3	2.8	2.5	-0.3
10th Grade	6.1	6.4	6.8	8.1	9.3	10.5	10.5	9.8	9.7	8.9	8.9	7.8	6.9	6.4	5.8	6.1	6.4	5.5	6.1	6.1	6.0	5.2	5.4	+0.2
12th Grade	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0	14.7	12.0	10.6	9.7	8.8	8.3	8.4	8.7	7.4	8.6	8.3	7.5	7.6	+0.1
College Students	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8	14.4	14.8	13.6	14.5	12.0	11.0	10.6	9.1	8.5	8.0	7.8	7.4	7.6	7.8	+0.1
Young Adults	15.7	15.7	15.4	15.4	16.1	16.4	16.8	17.4	18.0	18.4	18.3	19.6	19.7	19.3	17.6	17.2	16.0	14.8	14.2	13.9	13.0	12.2	12.4	+0.2

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
LSD																								
8th Grade	2.7	3.2	3.5	3.7	4.4	5.1	4.7	4.1	4.1	3.9	3.4	2.5	2.1	1.8	1.9	1.6	1.6	1.9	1.7	1.8	1.7	1.3	1.4	+0.1
10th Grade	5.6	5.8	6.2	7.2	8.4	9.4	9.5	8.5	8.5	7.6	6.3	5.0	3.5	2.8	2.5	2.7	3.0	2.6	3.0	3.0	2.8	2.6	2.7	+0.1
12th Grade	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	3.5	3.3	3.4	4.0	3.1	4.0	4.0	3.8	3.9	+0.1
College Students	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7	11.8	12.2	8.6	8.7	5.6	3.7	3.5	3.3	4.3	3.3	4.0	3.7	3.1	4.4	+1.4
Young Adults	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	11.2	10.1	9.6	8.1	7.3	7.2	6.1	6.2	6.3	0.0
Hallucinogens other than LSD^b																								
8th Grade	1.4	1.7	1.7	2.2	2.5	3.0	2.6	2.5	2.4	2.3	3.9	3.3	3.2	3.0	3.3	2.8	2.6	2.5	2.4	2.7	2.8	2.3	1.9	-0.4
10th Grade	2.2	2.5	2.8	3.8	3.9	4.7	4.8	5.0	4.7	4.8	6.6	6.3	5.9	5.8	5.2	5.5	5.7	4.8	5.4	5.3	5.2	4.5	4.4	0.0
12th Grade	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9	10.4	9.2	9.0	8.7	8.1	7.8	7.7	7.8	6.8	7.7	7.3	6.6	6.4	-0.2
College Students	6.0	5.7	5.4	4.4	6.5	6.5	7.5	8.7	8.8	8.2	10.7	11.0	12.8	10.1	10.6	10.1	8.5	8.2	7.8	7.1	6.9	7.2	6.8	-0.4
Young Adults	8.4	8.0	7.6	7.4	7.8	7.9	8.5	9.4	9.3	9.9	12.0	15.0	16.4	15.6	15.4	14.9	14.1	13.0	13.0	12.6	12.1	11.1	11.4	+0.3
PCP^g																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	2.4	2.2	2.1	1.8	1.7	1.8	2.3	1.6	1.3	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	2.0	2.4	2.1	2.2	1.6	1.6	1.7	1.1	1.4	+0.3
Ecstasy (MDMA)^h																								
8th Grade	—	—	—	—	—	3.4	3.2	2.7	2.7	4.3	5.2	4.3	3.2	2.8	2.8	2.5	2.3	2.4	2.2	3.3	2.6	2.0	1.8	-0.2
10th Grade	—	—	—	—	—	5.6	5.7	5.1	6.0	7.3	8.0	6.6	5.4	4.3	4.0	4.5	5.2	4.3	5.5	6.4	6.6	5.0	5.7	+0.7
12th Grade	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	5.4	6.5	6.5	6.2	6.5	7.3	8.0	7.2	7.1	-0.1
College Students	2.0	2.9	2.3	2.1	3.1	4.3	4.7	6.8	8.4	13.1	14.7	12.7	12.9	10.2	8.3	6.9	5.4	6.2	6.5	6.2	6.8	8.7	8.1	-0.6
Young Adults	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	14.9	14.4	13.1	13.1	11.5	12.3	11.3	11.4	11.6	+0.3
Cocaine																								
8th Grade	2.3	2.9	2.9	3.6	4.2	4.5	4.4	4.6	4.7	4.5	4.3	3.6	3.6	3.4	3.7	3.4	3.1	3.0	2.6	2.6	2.2	1.9	1.7	-0.2
10th Grade	4.1	3.3	3.6	4.3	5.0	6.5	7.1	7.2	7.7	6.9	5.7	6.1	5.1	5.4	5.2	4.8	5.3	4.5	4.6	3.7	3.3	3.3	3.3	0.0
12th Grade	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	8.0	8.5	7.8	7.2	6.0	5.5	5.2	4.9	4.5	-0.4
College Students	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4	9.1	8.6	8.2	9.2	9.5	8.8	7.7	8.5	7.2	8.1	6.6	5.5	5.2	5.1	-0.1
Young Adults	21.0	19.5	16.9	15.2	13.7	12.9	12.1	12.3	12.8	12.7	13.1	13.5	14.7	15.2	14.3	15.2	14.7	14.8	13.9	13.6	12.5	11.9	12.2	+0.3
Crackⁱ																								
8th Grade	1.3	1.6	1.7	2.4	2.7	2.9	2.7	3.2	3.1	3.1	3.0	2.5	2.5	2.4	2.4	2.3	2.1	2.0	1.7	1.5	1.5	1.0	1.2	+0.1
10th Grade	1.7	1.5	1.8	2.1	2.8	3.3	3.6	3.9	4.0	3.7	3.1	3.6	2.7	2.6	2.5	2.2	2.3	2.0	2.1	1.8	1.6	1.4	1.5	0.0
12th Grade	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	3.5	3.5	3.2	2.8	2.4	2.4	1.9	2.1	1.8	-0.3
College Students	1.5	1.7	1.3	1.0	1.8	1.2	1.4	2.2	2.4	2.5	2.0	1.9	3.1	2.0	1.7	2.3	1.3	1.4	1.0	1.2	0.8	0.7	0.7	0.0
Young Adults	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	4.1	4.4	3.9	4.3	3.3	3.6	2.9	2.7	2.6	-0.1
Other Cocaine^j																								
8th Grade	2.0	2.4	2.4	3.0	3.4	3.8	3.5	3.7	3.8	3.5	3.3	2.8	2.7	2.6	2.9	2.7	2.6	2.4	2.1	2.1	1.8	1.6	1.4	-0.2
10th Grade	3.8	3.0	3.3	3.8	4.4	5.5	6.1	6.4	6.8	6.0	5.0	5.2	4.5	4.8	4.6	4.3	4.8	4.0	4.1	3.4	3.0	3.0	2.9	-0.1
12th Grade	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	7.1	7.9	6.8	6.5	5.3	5.1	4.9	4.4	4.2	-0.2
College Students	9.0	7.6	6.3	4.6	5.2	4.6	5.0	7.4	7.8	8.1	8.3	8.6	8.5	9.3	8.1	6.2	8.0	7.1	7.9	6.7	5.4	5.1	5.2	+0.1
Young Adults	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	13.3	14.4	14.0	13.9	13.5	13.1	12.2	11.8	11.8	0.0

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Heroin^k																								
8th Grade	1.2	1.4	1.4	2.0	2.3	2.4	2.1	2.3	2.3	1.9	1.7	1.6	1.6	1.6	1.5	1.4	1.3	1.4	1.3	1.3	1.2	0.8	1.0	+0.1
10th Grade	1.2	1.2	1.3	1.5	1.7	2.1	2.1	2.3	2.3	2.2	1.7	1.8	1.5	1.5	1.4	1.5	1.2	1.5	1.3	1.2	1.1	1.0	1.0	0.0
12th Grade	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	1.5	1.4	1.5	1.3	1.2	1.6	1.4	1.1	1.0	-0.1
College Students	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9	1.7	1.2	1.0	1.0	0.9	0.5	0.7	0.5	0.7	0.8	0.7	0.6	0.5	0.4	0.0
Young Adults	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	1.7	1.9	1.6	1.9	1.6	1.8	1.7	1.6	1.6	0.0
With a Needle^l																								
8th Grade	—	—	—	—	1.5	1.6	1.3	1.4	1.6	1.1	1.2	1.0	1.0	1.1	1.0	1.0	0.9	0.9	0.9	0.9	0.8	0.6	0.6	0.0
10th Grade	—	—	—	—	1.0	1.1	1.1	1.2	1.3	1.0	0.8	1.0	0.9	0.8	0.8	0.9	0.9	0.7	0.9	0.8	0.8	0.7	0.7	0.0
12th Grade	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.9	0.8	0.7	0.7	0.6	1.1	0.9	0.7	0.7	0.0
College Students	—	—	—	—	0.4	0.1	0.2	0.5	0.8	0.7	0.2	0.3	0.1	0.1	0.3	0.3	0.1	0.0	0.1	0.1	0.3	0.2	0.1	-0.1
Young Adults	—	—	—	—	0.4	0.4	0.3	0.4	0.6	0.4	0.6	0.4	0.5	0.4	0.6	0.6	0.5	0.5	0.5	0.8	0.7	0.5	1.0	+0.4
Without a Needle^l																								
8th Grade	—	—	—	—	1.5	1.6	1.4	1.5	1.4	1.3	1.1	1.0	1.1	1.0	0.9	0.9	0.7	0.9	0.8	0.7	0.7	0.5	0.5	+0.1
10th Grade	—	—	—	—	1.1	1.7	1.7	1.7	1.6	1.7	1.3	1.3	1.0	1.1	1.1	1.0	1.1	0.8	1.0	0.9	0.8	0.8	0.7	-0.1
12th Grade	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	1.3	1.1	1.4	1.1	0.9	1.4	1.3	0.8	0.9	+0.1
College Students	—	—	—	—	0.5	1.0	1.2	2.1	1.0	2.5	1.3	1.2	1.1	1.0	0.3	0.8	0.4	0.7	0.4	0.4	0.4	0.5	0.8	+0.2
Young Adults	—	—	—	—	0.9	1.3	1.5	1.7	1.9	2.1	2.1	1.8	2.2	2.1	1.8	2.4	1.9	2.1	1.9	1.8	1.6	1.7	1.8	+0.1
Narcotics other than Heroin^{m,n}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9†	13.5	13.2	13.5	12.8	13.4	13.1	13.2	13.2	13.0	13.0	12.2	11.1	-1.1
College Students	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7	8.9	11.0‡	12.2	14.2	13.8	14.4	14.6	14.1	12.4	14.0	12.2	12.4	10.3	10.8	+0.5
Young Adults	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5‡	13.9	16.8	17.6	17.8	18.7	18.8	19.5	18.5	19.0	18.2	17.6	17.4	-0.3
Amphetamines^{m,o}																								
8th Grade	10.5	10.8	11.8	12.3	13.1	13.5	12.3	11.3	10.7	9.9	10.2	8.7	8.4	7.5	7.4	7.3	6.5	6.8	6.0	5.7	5.2	4.5	4.2	-0.3
10th Grade	13.2	13.1	14.9	15.1	17.4	17.7	17.0	16.0	15.7	15.7	16.0	14.9	13.1	11.9	11.1	11.2	11.1	9.0	10.3	10.6	9.0	8.9	8.1	-0.8
12th Grade	15.4	13.9	15.1	15.7	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	13.1	12.4	11.4	10.5	9.9	11.1	12.2	12.0	12.4	+0.3	
College Students	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9	12.3	12.4	11.9	12.3	12.7	12.3	10.7	11.2	9.1	11.8	12.1	13.4	14.4	15.3	+1.0
Young Adults	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	14.6	15.6	15.3	14.6	14.9	16.1	16.5	17.4	18.7	+1.3
Methamphetamine^{p,q}																								
8th Grade	—	—	—	—	—	—	—	—	4.5	4.2	4.4	3.5	3.9	2.5	3.1	2.7	1.8	2.3	1.6	1.8	1.3	1.3	1.4	0.0
10th Grade	—	—	—	—	—	—	—	—	7.3	6.9	6.4	6.1	5.2	5.3	4.1	3.2	2.8	2.4	2.8	2.5	2.1	1.8	1.6	-0.2
12th Grade	—	—	—	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	4.5	4.4	3.0	2.8	2.4	2.3	2.1	1.7	1.5	-0.2
College Students	—	—	—	—	—	—	—	—	7.1	5.1	5.3	5.0	5.8	5.2	4.1	2.9	1.9	1.9	1.0	1.1	0.6	0.3	0.9	+0.6
Young Adults	—	—	—	—	—	—	—	—	8.8	9.3	9.0	9.1	8.9	9.0	8.3	7.3	6.7	6.3	4.7	4.3	3.2	3.5	3.1	-0.4
Crystal Methamphetamine (Ice)^q																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	4.0	3.4	3.4	2.8	2.1	1.8	2.1	1.7	2.0	+0.3
College Students	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8	1.3	2.3	2.0	2.9	2.2	2.4	1.7	1.3	1.1	0.7	0.8	0.2	0.6	0.0	-0.6
Young Adults	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	4.4	4.7	3.7	3.6	3.4	2.8	3.1	2.6	2.8	+0.2

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Sedatives																								
(Barbiturates) ^{m,r}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8‡	9.9	10.5	10.2	9.3	8.5	8.2	7.5	7.0	6.9	7.5	+0.6
College Students	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7	6.9	6.0	5.9	5.7	7.2	8.5	6.3	5.9	6.4	6.0	5.3	3.6	3.5‡	5.4	—
Young Adults	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	10.0	9.5	9.8	10.6	9.5	8.6	7.9	7.2‡	9.5	—
Methaqualone ^{m,s}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	1.3	1.2	1.0	0.8	0.7	0.4	0.6	0.8	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{b,m}																								
8th Grade	3.8	4.1	4.4	4.6	4.5	5.3	4.8	4.6	4.4	4.4‡	5.0	4.3	4.4	4.0	4.1	4.3	3.9	3.9	3.9	4.4	3.4	3.0	2.9	-0.1
10th Grade	5.8	5.9	5.7	5.4	6.0	7.1	7.3	7.8	7.9	8.0‡	9.2	8.8	7.8	7.3	7.1	7.2	7.4	6.8	7.0	7.3	6.8	6.3	5.5	-0.8
12th Grade	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9‡	10.3	11.4	10.2	10.6	9.9	10.3	9.5	8.9	9.3	8.5	8.7	8.5	7.7	-0.8
College Students	6.8	6.9	6.3	4.4	5.4	5.3	6.9	7.7	8.2	8.8‡	9.7	10.7	11.0	10.6	11.9	10.0	9.1	8.6	9.2	8.1	7.1	6.4	7.8	+1.4
Young Adults	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5‡	11.9	13.4	13.8	14.9	14.5	15.0	14.5	15.8	13.8	14.3	13.8	13.3	13.2	-0.1
Any Prescription Drug ^{o,t}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	24.0	23.9	22.2	21.5	20.9	21.6	21.7	21.2	21.5	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rohypnol ^u																								
8th Grade	—	—	—	—	—	1.5	1.1	1.4	1.3	1.0	1.1	0.8	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.9	2.0	1.0	0.7	-0.3
10th Grade	—	—	—	—	—	1.5	1.7	2.0	1.8	1.3	1.5	1.3	1.0	1.2	1.0	0.8	1.3	0.9	0.7	1.4	1.2	0.8	1.1	+0.4
12th Grade	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—	—	—	—	—	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^v																								
Any Use																								
8th Grade	70.1	69.3‡	55.7	55.8	54.5	55.3	53.8	52.5	52.1	51.7	50.5	47.0	45.6	43.9	41.0	40.5	38.9	38.9	36.6	35.8	33.1	29.5	27.8	-1.7
10th Grade	83.8	82.3‡	71.6	71.1	70.5	71.8	72.0	69.8	70.6	71.4	70.1	66.9	66.0	64.2	63.2	61.5	61.7	58.3	59.1	58.2	56.0	54.0	52.1	-1.8
12th Grade	88.0	87.5‡	80.0	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	75.1	72.7	72.2	71.9	72.3	71.0	70.0	69.4	68.2	-1.2
College Students	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0	86.6	86.1	86.0	86.2	84.6	86.6	84.7	83.1	85.3	82.6	82.3	80.5	81.0	78.0	-3.0
Young Adults	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	89.1	88.9	87.9	88.4	87.9	87.5	87.4	86.5	86.2	-0.3
Been Drunk ^w																								
8th Grade	26.7	26.8	26.4	25.9	25.3	26.8	25.2	24.8	24.8	25.1	23.4	21.3	20.3	19.9	19.5	19.5	17.9	18.0	17.4	16.3	14.8	12.8	12.2	-0.6
10th Grade	50.0	47.7	47.9	47.2	46.9	48.5	49.4	46.7	48.9	49.3	48.2	44.0	42.4	42.3	42.1	41.4	41.2	37.2	38.6	36.9	35.9	34.6	33.5	-1.1
12th Grade	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	57.5	56.4	55.1	54.7	56.5	54.1	51.0	54.2	52.3	-1.9
College Students	79.6	76.8	76.4	74.4	76.6	76.2	77.0	76.8	75.1	74.7	76.1	75.1	74.9	73.4	72.9	73.1	71.6	72.5	69.1	70.5	67.9	70.0	66.5	-3.4
Young Adults	82.9	81.1	81.4	80.7	82.1	80.7	81.4	79.8	81.6	80.4	81.1	81.2	80.9	80.1	79.9	80.9	80.1	80.1	78.2	79.0	78.9	78.9	77.4	-1.5
Flavored Alcoholic Beverages ^{9,p}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	37.9	35.5	35.5	34.0	32.8	29.4	30.0	27.0	23.5	21.9	-1.6
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	58.6	58.8	58.1	55.7	53.5	51.4	51.3	48.4	46.7	44.9	-1.8
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	71.0	73.6	69.9	68.4	65.5	67.4	62.6	62.4	60.5	58.9	-1.7
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	79.0	84.5	80.9	80.6	78.6	78.1	77.4	76.7	76.6	67.5	-9.1 s
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	83.2	84.6	84.4	84.0	82.6	83.5	81.4	82.2	82.4	80.9	-1.6

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Cigarettes																								
Any Use																								
8th Grade	44.0	45.2	45.3	46.1	46.4	49.2	47.3	45.7	44.1	40.5	36.6	31.4	28.4	27.9	25.9	24.6	22.1	20.5	20.1	20.0	18.4	15.5	14.8	-0.8
10th Grade	55.1	53.5	56.3	56.9	57.6	61.2	60.2	57.7	57.6	55.1	52.8	47.4	43.0	40.7	38.9	36.1	34.6	31.7	32.7	33.0	30.4	27.7	25.7	-2.1 s
12th Grade	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	50.0	47.1	46.2	44.7	43.6	42.2	40.0	39.5	38.1	-1.4
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smokeless Tobacco ^x																								
8th Grade	22.2	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	9.1	9.8	9.6	9.9	9.7	8.1	7.9	-0.2
10th Grade	28.2	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	15.1	12.2	15.2	16.8	15.6	15.4	14.0	-1.5
12th Grade	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	16.3	17.6	16.9	17.4	17.2	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^{y,z}																								
8th Grade	1.9	1.7	1.6	2.0	2.0	1.8	1.8	2.3	2.7	3.0	2.8	2.5	2.5	1.9	1.7	1.6	1.5	1.4	1.3	1.1	1.2	1.2	1.1	-0.1
10th Grade	1.8	1.7	1.7	1.8	2.0	1.8	2.0	2.0	2.7	3.5	3.5	3.5	3.0	2.4	2.0	1.8	1.8	1.4	1.3	1.6	1.4	1.3	1.3	0.0
12th Grade	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	2.6	2.7	2.2	2.2	2.2	2.0	1.8	1.8	2.1	+0.3
College Students	1.4	1.7	1.9	0.5	0.8	0.6	1.6	0.9	1.3	0.6	1.5	1.2	1.2	1.6	1.0	1.9	0.6	1.6	1.3	0.7	1.1	0.4	0.8	+0.4
Young Adults	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	1.8	1.8	1.7	1.8	1.8	1.7	1.3	1.7	1.2	-0.5

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 2-4

TABLE 2-2
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Any Illicit Drug^a																								
8th Grade	11.3	12.9	15.1	18.5	21.4	23.6	22.1	21.0	20.5	19.5	19.5	17.7	16.1	15.2	15.5	14.8	13.2	14.1	14.5	16.0	14.7	13.4	14.9	+1.5 s
10th Grade	21.4	20.4	24.7	30.0	33.3	37.5	38.5	35.0	35.9	36.4	37.2	34.8	32.0	31.1	29.8	28.7	28.1	26.9	29.4	30.2	31.1	30.1	31.8	+1.6
12th Grade	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	36.5	35.9	36.6	36.5	38.3	40.0	39.7	40.3	+0.6
College Students	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	36.6	33.9	35.0	35.2	36.0	35.0	36.3	37.3	38.9	+1.7
Young Adults	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	33.3	33.2	34.7	34.0	36.3	+2.3 s
Any Illicit Drug other than Marijuana^{a,b}																								
8th Grade	8.4	9.3	10.4	11.3	12.6	13.1	11.8	11.0	10.5	10.2†	10.8	8.8	8.8	7.9	8.1	7.7	7.0	7.4	7.0	7.1	6.4	5.5	5.8	+0.3
10th Grade	12.2	12.3	13.9	15.2	17.5	18.4	18.2	16.6	16.7	16.7†	17.9	15.7	13.8	13.5	12.9	12.7	13.1	11.3	12.2	12.1	11.2	10.8	10.9	0.0
12th Grade	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4†	21.6	20.9	19.8	20.5	19.7	19.2	18.5	18.3	17.0	17.3	17.6	17.0	17.3	+0.3
College Students	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6†	16.4	16.6	17.9	18.6	18.5	18.1	17.3	15.3	16.9	17.1	16.8	17.1	19.0	+1.9
Young Adults	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9†	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	17.4	18.5	17.6	17.2	18.1	+0.9
Any Illicit Drug including Inhalants^{a,c,d}																								
8th Grade	16.7	18.2	21.1	24.2	27.1	28.7	27.2	26.2	25.3	24.0	23.9	21.4	20.4	20.2	20.4	19.7	18.0	19.0	18.8	20.3	18.2	17.0	17.5	+0.5
10th Grade	23.9	23.5	27.4	32.5	35.6	39.6	40.3	37.1	37.7	38.0	38.7	36.1	33.5	32.9	31.7	30.7	30.2	28.8	31.2	31.8	32.5	31.5	33.0	+1.5
12th Grade	31.2	28.8	32.5	37.6	40.2	41.9	43.3	42.4	42.8	42.5	42.6	42.1	40.5	39.1	40.3	38.0	37.0	37.3	37.6	39.2	41.5	40.2	41.8	+1.7
College Students	29.8	31.1	31.7	31.9	33.7	35.1	35.5	39.1	37.4	37.0	38.2	37.7	36.0	35.9	37.9	35.5	36.8	35.7	35.0	34.5	36.5	36.9	37.2	+0.3
Young Adults	27.8	29.2	28.9	29.2	30.4	30.2	30.1	30.6	30.6	31.2	33.2	32.4	32.7	34.9	32.8	32.6	33.2	33.5	33.1	33.3	34.2	34.2	36.3	+2.0
Marijuana/Hashish																								
8th Grade	6.2	7.2	9.2	13.0	15.8	18.3	17.7	16.9	16.5	15.6	15.4	14.6	12.8	11.8	12.2	11.7	10.3	10.9	11.8	13.7	12.5	11.4	12.7	+1.2
10th Grade	16.5	15.2	19.2	25.2	28.7	33.6	34.8	31.1	32.1	32.2	32.7	30.3	28.2	27.5	26.6	25.2	24.6	23.9	26.7	27.5	28.8	28.0	29.8	+1.8
12th Grade	23.9	21.9	26.0	30.7	34.7	35.8	37.5	37.8	36.5	37.0	36.2	34.9	34.3	33.6	31.5	31.7	32.4	32.8	34.8	36.4	36.4	36.4	36.4	0.0
College Students	26.5	27.7	27.9	29.3	31.2	33.1	31.6	35.9	35.2	34.0	35.6	34.7	33.7	33.3	33.3	30.2	31.8	32.3	32.8	32.7	33.2	34.9	35.5	+0.6
Young Adults	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	28.2	27.7	28.5	28.6	29.3	28.7	31.0	30.2	32.2	+2.0 s
Synthetic Marijuana^{p,q}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Inhalants^{c,d}																								
8th Grade	9.0	9.5	11.0	11.7	12.8	12.2	11.8	11.1	10.3	9.4	9.1	7.7	8.7	9.6	9.5	9.1	8.3	8.9	8.1	8.1	7.0	6.2	5.2	-1.1 s
10th Grade	7.1	7.5	8.4	9.1	9.6	9.5	8.7	8.0	7.2	7.3	6.6	5.8	5.4	5.9	6.0	6.5	6.6	5.9	6.1	5.7	4.5	4.1	3.5	-0.6
12th Grade	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	5.0	4.5	3.7	3.8	3.4	3.6	3.2	2.9	2.5	-0.4
College Students	3.5	3.1	3.8	3.0	3.9	3.6	4.1	3.0	3.2	2.9	2.8	2.0	1.8	2.7	1.8	1.5	1.5	1.1	1.2	1.7	0.9	1.5	0.5	-1.0
Young Adults	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	1.3	1.3	0.8	1.4	0.9	1.2	0.8	1.1	0.5	-0.6 s
Nitrites^e																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.9	0.5	0.9	1.1	1.1	1.6	1.2	1.4	0.9	0.6	0.6	1.1	0.9	0.8	0.6	0.5	0.8	0.6	0.9	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.2	0.1	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}																								
8th Grade	1.9	2.5	2.6	2.7	3.6	4.1	3.7	3.4	2.9	2.8†	3.4	2.6	2.6	2.2	2.4	2.1	1.9	2.1	1.9	2.2	2.2	1.6	1.6	0.0
10th Grade	4.0	4.3	4.7	5.8	7.2	7.8	7.6	6.9	6.9	6.1†	6.2	4.7	4.1	4.1	4.0	4.1	4.4	3.9	4.1	4.2	4.1	3.5	3.4	-0.1
12th Grade	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1†	9.1	6.6	5.9	6.2	5.5	4.9	5.4	5.9	4.7	5.5	5.2	4.8	4.5	-0.3
College Students	6.3	6.8	6.0	6.2	8.2	6.9	7.7	7.2	7.8	6.7†	7.5	6.3	7.4	5.9	5.0	5.6	4.9	5.1	4.7	4.9	4.1	4.5	4.5	0.0
Young Adults	4.5	5.0	4.5	4.8	5.6	5.6	5.9	5.2	5.4	5.4†	5.4	4.7	5.2	4.7	4.5	4.1	3.8	3.8	3.9	4.2	3.7	3.6	3.9	+0.3

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change	
LSD																									
8th Grade	1.7	2.1	2.3	2.4	3.2	3.5	3.2	2.8	2.4	2.4	2.2	1.5	1.3	1.1	1.2	0.9	1.1	1.3	1.1	1.2	1.1	0.8	1.0	+0.2	
10th Grade	3.7	4.0	4.2	5.2	6.5	6.9	6.7	5.9	6.0	5.1	4.1	2.6	1.7	1.6	1.5	1.7	1.9	1.8	1.9	1.9	1.8	1.7	1.7	-0.1	
12th Grade	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	1.8	1.7	2.1	2.7	1.9	2.6	2.7	2.4	2.2	-0.2	
College Students	5.1	5.7	5.1	5.2	6.9	5.2	5.0	4.4	5.4	4.3	4.0	2.1	1.4	1.2	0.7	1.4	1.3	2.6	2.0	2.1	2.0	1.9	2.6	+0.7	
Young Adults	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	0.8	1.2	1.1	1.4	1.7	1.5	1.7	1.6	2.0	+0.5	
Hallucinogens other than LSD^b																									
8th Grade	0.7	1.1	1.0	1.3	1.7	2.0	1.8	1.6	1.5	1.4†	2.4	2.1	2.1	1.9	2.0	1.8	1.6	1.6	1.5	1.8	1.8	1.3	1.2	-0.1	
10th Grade	1.3	1.4	1.9	2.4	2.8	3.3	3.3	3.4	3.2	3.1‡	4.3	4.0	3.6	3.7	3.5	3.7	3.8	3.3	3.5	3.5	3.5	3.0	2.7	-0.3	
12th Grade	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4‡	5.9	5.4	5.4	5.6	5.0	4.6	4.8	5.0	4.2	4.8	4.3	4.0	3.7	-0.4	
College Students	3.1	2.6	2.7	2.8	4.0	4.1	4.9	4.4	4.5	4.4‡	5.5	5.8	7.1	5.6	5.0	5.4	4.7	4.4	4.1	4.4	3.4	3.9	3.7	-0.2	
Young Adults	1.7	1.9	1.9	2.0	2.5	2.8	3.1	3.0	3.0	3.4‡	3.5	4.0	4.9	4.5	4.2	3.8	3.6	3.4	3.3	3.7	3.2	2.9	3.2	+0.3	
PCP^g																									
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
12th Grade	1.4	1.4	1.4	1.6	1.8	2.6	2.3	2.1	1.8	2.3	1.8	1.1	1.3	0.7	1.3	0.7	0.9	1.1	1.0	1.0	1.3	0.9	0.7	-0.2	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Young Adults	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	0.6	0.2	0.3	0.4	0.1	0.2	0.3	*	0.2	+0.2	
Ecstasy (MDMA)^h																									
8th Grade	—	—	—	—	—	2.3	2.3	1.8	1.7	3.1	3.5	2.9	2.1	1.7	1.7	1.4	1.5	1.7	1.3	2.4	1.7	1.1	1.1	0.0	
10th Grade	—	—	—	—	—	4.6	3.9	3.3	4.4	5.4	6.2	4.9	3.0	2.4	2.6	2.8	3.5	2.9	3.7	4.7	4.5	3.0	3.6	+0.6	
12th Grade	—	—	—	—	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	3.0	4.1	4.5	4.3	4.3	4.5	5.3	3.8	4.0	+0.2	
College Students	0.9	2.0	0.8	0.5	2.4	2.8	2.4	3.9	5.5	9.1	9.2	6.8	4.4	2.2	2.9	2.6	2.2	3.7	3.1	4.3	4.2	5.8	5.3	-0.5	
Young Adults	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	3.0	3.0	2.5	3.3	3.1	3.5	3.6	4.1	4.2	+0.2	
Salvia^{p,q}																									
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.7	1.6	1.4	1.2	-0.2
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.7	3.9	2.5	2.3	-0.2
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.7	5.5	5.9	4.4	3.4	-1.0 s
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.8	3.5	3.1	1.5	1.0	-0.5
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.5	3.6	2.2	1.4	0.9	-0.5
Cocaine																									
8th Grade	1.1	1.5	1.7	2.1	2.6	3.0	2.8	3.1	2.7	2.6	2.5	2.3	2.2	2.0	2.2	2.0	2.0	1.8	1.6	1.6	1.4	1.2	1.0	-0.2	
10th Grade	2.2	1.9	2.1	2.8	3.5	4.2	4.7	4.7	4.9	4.4	3.6	4.0	3.3	3.7	3.5	3.2	3.4	3.0	2.7	2.2	1.9	2.0	1.9	-0.1	
12th Grade	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	5.1	5.7	5.2	4.4	3.4	2.9	2.9	2.7	2.6	-0.1	
College Students	3.6	3.0	2.7	2.0	3.6	2.9	3.4	4.6	4.6	4.8	4.7	4.8	5.4	6.6	5.7	5.1	5.4	4.4	4.2	3.5	3.3	3.1	2.7	-0.4	
Young Adults	6.2	5.7	4.7	4.3	4.4	4.1	4.7	4.9	5.4	5.4	5.8	5.8	6.6	7.1	6.9	6.6	6.2	6.0	5.2	4.7	4.7	4.1	3.9	-0.2	
Crackⁱ																									
8th Grade	0.7	0.9	1.0	1.3	1.6	1.8	1.7	2.1	1.8	1.8	1.7	1.6	1.6	1.3	1.4	1.3	1.3	1.1	1.1	1.0	0.9	0.6	0.6	0.0	
10th Grade	0.9	0.9	1.1	1.4	1.8	2.1	2.2	2.5	2.4	2.2	1.8	2.3	1.6	1.7	1.7	1.3	1.3	1.3	1.2	1.0	0.9	0.8	0.8	+0.1	
12th Grade	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	1.9	2.1	1.9	1.6	1.3	1.4	1.0	1.2	1.1	-0.2	
College Students	0.5	0.4	0.6	0.5	1.1	0.6	0.4	1.0	0.9	0.9	0.9	0.4	1.3	1.3	0.8	1.0	0.6	0.5	0.3	0.4	0.3	0.3	0.3	0.0	
Young Adults	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	1.2	1.1	1.0	0.9	0.7	0.5	0.6	0.5	0.3	-0.1	
Other Cocaine^j																									
8th Grade	1.0	1.2	1.3	1.7	2.1	2.5	2.2	2.4	2.3	1.9	1.9	1.8	1.6	1.6	1.7	1.6	1.5	1.4	1.3	1.3	1.1	1.0	0.8	-0.2	
10th Grade	2.1	1.7	1.8	2.4	3.0	3.5	4.1	4.0	4.4	3.8	3.0	3.4	2.8	3.3	3.0	2.9	3.1	2.6	2.3	1.9	1.7	1.8	1.6	-0.2	
12th Grade	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	4.5	5.2	4.5	4.0	3.0	2.6	2.6	2.4	2.4	0.0	
College Students	3.2	2.4	2.5	1.8	3.3	2.3	3.0	4.2	4.2	4.1	4.1	5.0	5.1	6.3	5.0	3.8	5.3	4.2	4.2	4.0	3.0	3.0	2.8	-0.1	
Young Adults	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	6.3	5.9	5.6	5.5	5.0	4.8	4.3	4.0	3.7	-0.3	

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Heroin^k																								
8th Grade	0.7	0.7	0.7	1.2	1.4	1.6	1.3	1.3	1.4	1.1	1.0	0.9	0.9	1.0	0.8	0.8	0.8	0.9	0.7	0.8	0.7	0.5	0.5	0.0
10th Grade	0.5	0.6	0.7	0.9	1.1	1.2	1.4	1.4	1.4	1.4	0.9	1.1	0.7	0.9	0.9	0.9	0.8	0.8	0.9	0.8	0.8	0.6	0.6	0.0
12th Grade	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	0.8	0.8	0.9	0.7	0.7	0.9	0.8	0.6	0.6	-0.1
College Students	0.1	0.1	0.1	0.1	0.3	0.4	0.3	0.6	0.2	0.5	0.4	0.1	0.2	0.4	0.3	0.3	0.2	0.3	0.4	0.2	0.1	0.1	0.3	+0.2
Young Adults	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.5	0.5	0.5	0.6	+0.2
With a Needle^l																								
8th Grade	—	—	—	—	0.9	1.0	0.8	0.8	0.9	0.6	0.7	0.6	0.6	0.7	0.6	0.5	0.6	0.5	0.5	0.6	0.5	0.4	0.3	0.0
10th Grade	—	—	—	—	0.6	0.7	0.7	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.5	0.4	0.5	+0.1
12th Grade	—	—	—	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.5	0.4	0.4	0.3	0.7	0.6	0.4	0.4	0.0
College Students	—	—	—	—	0.1	*	0.1	0.2	0.1	0.1	0.1	*	0.1	0.1	0.3	0.3	*	0.0	0.1	0.0	0.2	0.2	0.1	-0.1
Young Adults	—	—	—	—	0.1	0.1	0.1	0.1	0.1	*	0.3	*	*	0.1	0.2	0.3	0.1	0.1	0.1	0.2	0.4	0.3	0.3	0.0
Without a Needle^l																								
8th Grade	—	—	—	—	0.8	1.0	0.8	0.8	0.9	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.6	0.4	0.5	0.4	0.3	0.3	+0.1
10th Grade	—	—	—	—	0.8	0.9	1.1	1.0	1.1	1.1	0.7	0.8	0.5	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.5	0.4	0.4	-0.1
12th Grade	—	—	—	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	0.8	0.6	1.0	0.5	0.6	0.8	0.7	0.4	0.4	+0.1
College Students	—	—	—	—	0.0	0.8	0.4	0.9	0.3	0.8	0.6	0.2	0.1	0.6	0.2	0.3	0.2	0.3	0.1	0.3	0.2	0.1	0.5	+0.4
Young Adults	—	—	—	—	0.3	0.4	0.4	0.7	0.6	0.5	0.9	0.2	0.4	0.3	0.4	0.5	0.3	0.4	0.6	0.4	0.2	0.4	0.7	+0.4
Narcotics other than Heroin^{m,n}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7‡	9.4	9.3	9.5	9.0	9.0	9.2	9.1	9.2	8.7	8.7	7.9	7.1	-0.8
College Students	2.7	2.7	2.5	2.4	3.8	3.1	4.2	4.2	4.3	4.5	5.7‡	7.4	8.7	8.2	8.4	8.8	7.7	6.5	7.6	7.2	6.2	5.4	5.4	0.0
Young Adults	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0‡	7.1	8.5	9.0	8.7	9.1	8.7	9.1	8.4	9.0	7.9	7.3	7.0	-0.3
OxyContin^{m,p,aa,bb}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	1.3	1.7	1.7	1.8	2.6	1.8	2.1	2.0	2.1	1.8	1.6	2.0	+0.4
10th Grade	—	—	—	—	—	—	—	—	—	—	—	3.0	3.6	3.5	3.2	3.8	3.9	3.6	5.1	4.6	3.9	3.0	3.4	+0.4
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.0	4.5	5.0	5.5	4.3	5.2	4.7	4.9	5.1	4.9	4.3	3.6	-0.7
College Students	—	—	—	—	—	—	—	—	—	—	—	1.5	2.2	2.5	2.1	3.0	2.8	3.6	5.0	2.3	2.4	1.2	2.3	+1.2
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.9	2.6	3.1	3.1	3.1	2.9	3.9	5.2	3.2	2.8	2.3	2.8	+0.5
Vicodin^{m,p,aa,bb}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	2.5	2.8	2.5	2.6	3.0	2.7	2.9	2.5	2.7	2.1	1.3	1.4	0.0
10th Grade	—	—	—	—	—	—	—	—	—	—	—	6.9	7.2	6.2	5.9	7.0	7.2	6.7	8.1	7.7	5.9	4.4	4.6	+0.2
12th Grade	—	—	—	—	—	—	—	—	—	—	—	9.6	10.5	9.3	9.5	9.7	9.6	9.7	9.7	8.0	8.1	7.5	5.3	-2.2 SS
College Students	—	—	—	—	—	—	—	—	—	—	—	6.9	7.5	7.4	9.6	7.6	6.7	6.7	8.4	4.9	5.8	3.8	4.4	+0.6
Young Adults	—	—	—	—	—	—	—	—	—	—	—	8.2	8.6	8.9	9.3	9.1	8.9	9.1	8.9	7.8	7.1	6.3	6.2	-0.1
Amphetamines^{m,o}																								
8th Grade	6.2	6.5	7.2	7.9	8.7	9.1	8.1	7.2	6.9	6.5	6.7	5.5	5.5	4.9	4.9	4.7	4.2	4.5	4.1	3.9	3.5	2.9	2.6	-0.3
10th Grade	8.2	8.2	9.6	10.2	11.9	12.4	12.1	10.7	10.4	11.1	11.7	10.7	9.0	8.5	7.8	7.9	8.0	6.4	7.1	7.6	6.6	6.5	5.9	-0.6
12th Grade	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	8.6	8.1	7.5	6.8	6.6	7.4	8.2	7.9	8.7	+0.8
College Students	3.9	3.6	4.2	4.2	5.4	4.2	5.7	5.1	5.8	6.6	7.2	7.0	7.1	7.0	6.7	6.0	6.9	5.7	7.5	9.0	9.3	11.1	10.6	-0.6
Young Adults	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	5.1	5.6	5.6	5.3	6.0	7.1	7.2	7.8	7.8	0.0
Ritalin^{m,p,q,bb}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	2.9	2.8	2.6	2.5	2.4	2.6	2.1	1.6	1.8	1.5	1.3	0.7	1.1	+0.4
10th Grade	—	—	—	—	—	—	—	—	—	—	4.8	4.8	4.1	3.4	3.4	3.6	2.8	2.9	3.6	2.7	2.6	1.9	1.8	-0.1
12th Grade	—	—	—	—	—	—	—	—	—	—	5.1	4.0	4.0	5.1	4.4	4.4	3.8	3.4	2.1	2.7	2.6	2.6	2.3	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	5.7	4.7	4.7	4.2	3.9	3.7	3.2	1.7	1.9	2.3	1.8	3.6	+1.8
Young Adults	—	—	—	—	—	—	—	—	—	—	—	2.9	2.9	2.7	2.5	2.6	2.4	2.4	1.7	1.7	1.5	1.6	2.0	+0.4

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Adderall^{m,p,q,bb}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.0	2.3	1.7	1.7	1.8	+0.1
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.7	5.3	4.6	4.5	4.4	-0.1
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.4	6.5	6.5	7.6	7.4	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	10.2	9.0	9.8	9.0	10.7	+1.6
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.8	7.0	6.6	7.4	7.0	-0.4
Provigil^{m,q}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.8	1.3	1.5	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.2	0.0	0.2	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.5	0.3	—	—	—
Methamphetamine^{p,q}																								
8th Grade	—	—	—	—	—	—	—	—	3.2	2.5	2.8	2.2	2.5	1.5	1.8	1.8	1.1	1.2	1.0	1.2	0.8	1.0	1.0	0.0
10th Grade	—	—	—	—	—	—	—	—	4.6	4.0	3.7	3.9	3.3	3.0	2.9	1.8	1.6	1.5	1.6	1.6	1.4	1.0	1.0	0.0
12th Grade	—	—	—	—	—	—	—	—	4.7	4.3	3.9	3.6	3.2	3.4	2.5	2.5	1.7	1.2	1.2	1.0	1.4	1.1	0.9	-0.2
College Students	—	—	—	—	—	—	—	—	3.3	1.6	2.4	1.2	2.6	2.9	1.7	1.2	0.4	0.5	0.3	0.4	0.2	0.0	0.4	+0.4
Young Adults	—	—	—	—	—	—	—	—	2.8	2.5	2.8	2.5	2.7	2.8	2.4	1.9	1.5	1.0	0.9	0.7	0.5	1.0	0.6	-0.3
Crystal Methamphetamine (Ice)^q																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.0	2.0	2.1	2.3	1.9	1.6	1.1	0.9	0.9	1.2	0.8	1.1	+0.3
College Students	0.1	0.2	0.7	0.8	1.1	0.3	0.8	1.0	0.5	0.5	0.6	0.8	0.9	1.1	1.4	0.6	0.7	0.1	0.1	0.5	0.1	0.6	0.0	-0.6
Young Adults	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	1.6	1.1	1.1	0.8	0.8	0.5	0.5	0.6	0.8	+0.2
Sedatives																								
(Barbiturates)^{m,r}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0†	6.5	7.2	6.6	6.2	5.8	5.2	4.8	4.3	4.5	4.8	+0.3
College Students	1.2	1.4	1.5	1.2	2.0	2.3	3.0	2.5	3.2	3.7	3.8	3.7	4.1	4.2	3.9	3.4	3.6	3.7	3.1	2.5	1.7	2.2†	2.7	—
Young Adults	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	4.2	3.9	4.2	4.7	3.8	3.3	3.2	2.7†	3.4	—
Methaqualone^{m,s}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	0.2	0.8	0.7	1.1	1.0	1.1	1.1	0.3	0.8	0.9	0.6	0.8	0.9	0.8	0.5	0.5	0.6	0.3	0.3	0.4	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}																								
8th Grade	1.8	2.0	2.1	2.4	2.7	3.3	2.9	2.6	2.5	2.6†	2.8	2.6	2.7	2.5	2.8	2.6	2.4	2.4	2.6	2.8	2.0	1.8	1.8	0.0
10th Grade	3.2	3.5	3.3	3.3	4.0	4.6	4.9	5.1	5.4	5.6†	7.3	6.3	5.3	5.1	4.8	5.2	5.3	4.6	5.0	5.1	4.5	4.3	3.7	-0.6
12th Grade	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7†	6.9	7.7	6.7	7.3	6.8	6.6	6.2	6.2	6.3	5.6	5.6	5.3	4.6	-0.7
College Students	2.4	2.9	2.4	1.8	2.9	2.8	3.8	3.9	3.8	4.2†	5.1	6.7	6.9	6.7	6.4	5.8	5.5	5.0	5.4	4.9	4.2	3.4	4.4	+1.0
Young Adults	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6†	5.5	7.0	6.8	7.4	6.7	6.5	7.1	6.8	6.4	6.3	5.9	5.3	5.4	+0.1
Any Prescription Drug^{o,t}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17.1	16.8	15.8	15.4	14.4	15.0	15.2	14.8	15.0	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Over-the-counter Cough/Cold Medicines^{p,q}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.2	4.0	3.6	3.8	3.2	2.7	3.0	2.9	-0.1
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.3	5.4	5.3	6.0	5.1	5.5	4.7	4.3	-0.5
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	5.8	5.5	5.9	6.6	5.3	5.6	5.0	-0.5
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change	
Rohypnol^u																									
8th Grade	—	—	—	—	—	1.0	0.8	0.8	0.5	0.5	0.7	0.3	0.5	0.6	0.7	0.5	0.7	0.5	0.4	0.5	0.8	0.4	0.4	0.0	
10th Grade	—	—	—	—	—	1.1	1.3	1.2	1.0	0.8	1.0	0.7	0.6	0.7	0.5	0.5	0.7	0.4	0.4	0.6	0.6	0.5	0.6	+0.1	
12th Grade	—	—	—	—	—	1.1	1.2	1.4	1.0	0.8	0.9‡	1.6	1.3	1.6	1.2	1.1	1.0	1.3	1.0	1.5	1.3	1.5	0.9	-0.6	
College Students	—	—	—	—	—	—	—	—	—	—	—	0.7	0.4	0.3	0.1	0.2	0.1	0.3	0.0	—	—	—	—	—	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.3	0.5	0.1	0.1	0.2	0.3	0.2	0.1	—	—	—	—	—	
GHB^{p,cc}																									
8th Grade	—	—	—	—	—	—	—	—	—	1.2	1.1	0.8	0.9	0.7	0.5	0.8	0.7	1.1	0.7	0.6	0.6	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	1.1	1.0	1.4	1.4	0.8	0.8	0.7	0.6	0.5	1.0	0.6	0.5	—	—	—	
12th Grade	—	—	—	—	—	—	—	—	—	1.9	1.6	1.5	1.4	2.0	1.1	1.1	0.9	1.2	1.1	1.4	1.4	1.4	1.0	-0.4	
College Students	—	—	—	—	—	—	—	—	—	—	—	0.6	0.3	0.7	0.4	*	0.1	0.2	0.0	0.1	0.1	0.0	0.1	+0.1	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.8	0.6	0.5	0.3	0.2	0.4	0.3	0.2	0.3	0.3	0.4	0.3	0.0	
Ketamine^{p,dd}																									
8th Grade	—	—	—	—	—	—	—	—	—	1.6	1.3	1.3	1.1	0.9	0.6	0.9	1.0	1.2	1.0	1.0	0.8	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	2.1	2.1	2.2	1.9	1.3	1.0	1.0	0.8	1.0	1.3	1.1	1.2	—	—	—	
12th Grade	—	—	—	—	—	—	—	—	—	2.5	2.5	2.6	2.1	1.9	1.6	1.4	1.3	1.5	1.7	1.6	1.7	1.5	1.4	-0.1	
College Students	—	—	—	—	—	—	—	—	—	—	—	1.3	1.0	1.5	0.5	0.9	0.2	0.4	0.1	0.7	0.6	0.4	0.9	+0.5	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.2	0.9	0.6	0.5	0.5	0.3	0.4	0.5	0.8	0.5	0.8	0.5	-0.2	
Alcohol^v																									
Any Use																									
8th Grade	54.0	53.7‡	45.4	46.8	45.3	46.5	45.5	43.7	43.5	43.1	41.9	38.7	37.2	36.7	33.9	33.6	31.8	32.1	30.3	29.3	26.9	23.6	22.1	-1.5	
10th Grade	72.3	70.2‡	63.4	63.9	63.5	65.0	65.2	62.7	63.7	65.3	63.5	60.0	59.3	58.2	56.7	55.8	56.3	52.5	52.8	52.1	49.8	48.5	47.1	-1.4	
12th Grade	77.7	76.8‡	72.7	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	68.6	66.5	66.4	65.5	66.2	65.2	63.5	63.5	62.0	-1.5	
College Students	88.3	86.9	85.1	82.7	83.2	82.9	82.4	84.6	83.6	83.2	83.0	82.9	81.7	81.2	83.0	82.1	80.9	82.1	79.4	78.6	77.4	79.2	75.6	-3.6 s	
Young Adults	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	83.8	84.4	84.0	83.6	83.8	82.7	83.5	82.5	82.5	0.0	
Been Drunk^w																									
8th Grade	17.5	18.3	18.2	18.2	18.4	19.8	18.4	17.9	18.5	18.5	16.6	15.0	14.5	14.5	14.1	13.9	12.6	12.7	12.2	11.5	10.5	8.6	8.4	-0.1	
10th Grade	40.1	37.0	37.8	38.0	38.5	40.1	40.7	38.3	40.9	41.6	39.9	35.4	34.7	35.1	34.2	34.5	34.4	30.0	31.2	29.9	28.8	28.2	27.1	-1.2	
12th Grade	52.7	50.3	49.6	51.7	52.5	51.9	53.2	52.0	53.2	51.8	53.2	50.4	48.0	51.8	47.7	47.9	46.1	45.6	47.0	44.0	42.2	45.0	43.5	-1.6	
College Students	69.1	67.3	65.6	63.1	62.1	64.2	66.8	67.0	65.4	64.7	68.8	66.0	64.7	67.1	64.2	66.2	64.8	66.8	61.5	63.8	60.1	61.5	57.9	-3.5	
Young Adults	62.0	60.9	61.1	58.8	61.6	59.9	63.2	59.6	63.2	60.6	63.1	61.8	62.9	63.8	63.5	65.7	65.8	66.0	65.5	64.8	64.0	64.6	63.1	-1.6	
Flavored Alcoholic Beverages^{9,p,ee}																									
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	30.4	27.9	26.8	26.0	25.0	22.2	21.9	19.2	17.0	15.7	-1.3	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	49.7	48.5	48.8	45.9	43.4	41.5	41.0	38.3	37.8	35.6	-2.2	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	55.2	55.8	58.4	54.7	53.6	51.8	53.4	47.9	47.0	44.4	44.2	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	63.2	67.0	63.5	62.6	65.0	66.1	60.3	63.0	58.1	57.6	-0.6	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	62.7	58.4	58.5	58.9	58.3	57.0	52.0	56.3	54.8	54.1	-0.7	
Alcoholic Beverages containing Caffeine^{p,w}																									
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11.8‡	10.9	10.2	-0.7	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	22.5‡	19.7	16.9	-2.8 s	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	26.4‡	26.4	23.5	-2.8 s	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.6‡	33.8	39.1	+5.4	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.1‡	36.7	36.9	+0.2	
Cigarettes																									
Any Use																									
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
College Students	35.6	37.3	38.8	37.6	39.3	41.4	43.6	44.3	44.5	41.3	39.0	38.3	35.2	36.7	36.0	30.9	30.7	30.0	29.9	28.1	25.8	23.4	23.2	-0.2	
Young Adults	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.1	39.1	38.6	39.0	39.1	36.9	36.2	35.0	33.9	33.0	31.5	29.8	29.8	0.0	

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change		
Bidis^{p,ff}																										
8th Grade	—	—	—	—	—	—	—	—	—	3.9	2.7	2.7	2.0	1.7	1.6	—	—	—	—	—	—	—	—	—		
10th Grade	—	—	—	—	—	—	—	—	—	6.4	4.9	3.1	2.8	2.1	1.6	—	—	—	—	—	—	—	—	—		
12th Grade	—	—	—	—	—	—	—	—	—	9.2	7.0	5.9	4.0	3.6	3.3	2.3	1.7	1.9	1.5	1.4	—	—	—	—		
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Kreteks^{p,ff}																										
8th Grade	—	—	—	—	—	—	—	—	—	—	2.6	2.6	2.0	1.9	1.4	—	—	—	—	—	—	—	—	—		
10th Grade	—	—	—	—	—	—	—	—	—	6.0	4.9	3.8	3.7	2.8	—	—	—	—	—	—	—	—	—	—		
12th Grade	—	—	—	—	—	—	—	—	—	10.1	8.4	6.7	6.5	7.1	6.2	6.8	6.8	5.5	4.6	2.9	3.0	1.6	-1.4 s			
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Tobacco using a Hookah^s																										
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17.1	18.5	18.3	21.4	+3.1 s		
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	27.9	25.7	26.1	+0.3		
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	20.1	19.1	20.4	+1.3		
Small Cigars^s																										
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	23.1	19.5	19.9	20.4	+0.5	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	23.6	20.3	19.0	-1.3	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	19.2	18.0	18.4	+0.5	
Dissolvable Tobacco^{p,s}																										
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.0	1.1	+0.1	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.6	1.2	-0.4	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.5	1.6	1.9	+0.2	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.0	0.3	0.2	0.0	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.6	0.3	-0.3	
Snus^{p,s}																										
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.4	2.0	-0.4	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	5.2	-1.7 s	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.9	7.9	-0.2	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.5	4.7	4.8	+0.1
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.1	5.7	4.8	-0.9
Steroids^{y,z}																										
8th Grade	1.0	1.1	0.9	1.2	1.0	0.9	1.0	1.2	1.7	1.7	1.6	1.5	1.4	1.1	1.1	0.9	0.8	0.9	0.8	0.5	0.7	0.6	0.6	0.0		
10th Grade	1.1	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.7	2.2	2.1	2.2	1.7	1.5	1.3	1.2	1.1	0.9	0.8	1.0	0.9	0.8	0.8	0.0		
12th Grade	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	1.5	1.8	1.4	1.5	1.5	1.5	1.2	1.3	1.5	+0.2		
College Students	0.6	0.2	0.9	0.2	0.4	0.2	0.7	0.2	0.9	0.1	0.6	0.5	0.3	0.6	0.5	0.8	0.6	0.1	0.7	0.3	0.2	0.3	0.8	+0.5		
Young Adults	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.7	0.4	0.7	0.8	0.2	0.4	0.5	+0.1		
Bath Salts (Synthetic stimulants)^{p,q}																										
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	1.0	+0.1	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.9	+0.3	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.3	0.9	-0.4	
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.1	-0.2	
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.4	-0.1	

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 2-4.

TABLE 2-3
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Any Illicit Drug^a																								
8th Grade	5.7	6.8	8.4	10.9	12.4	14.6	12.9	12.1	12.2	11.9	11.7	10.4	9.7	8.4	8.5	8.1	7.4	7.6	8.1	9.5	8.5	7.7	8.5	+0.9
10th Grade	11.6	11.0	14.0	18.5	20.2	23.2	23.0	21.5	22.1	22.5	22.7	20.8	19.5	18.3	17.3	16.8	16.9	15.8	17.8	18.5	19.2	18.6	19.4	+0.8
12th Grade	16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	23.1	21.5	21.9	22.3	23.3	23.8	25.2	25.2	25.5	+0.3
College Students	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	20.7	19.2	21.4	22.3	22.5	+0.2
Young Adults	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	19.8	18.9	20.6	19.9	21.8	+1.9 s
Any Illicit Drug other than Marijuana^{a,b}																								
8th Grade	3.8	4.7	5.3	5.6	6.5	6.9	6.0	5.5	5.5	5.6†	5.5	4.7	4.7	4.1	4.1	3.8	3.6	3.8	3.5	3.5	3.4	2.6	3.3	+0.6 s
10th Grade	5.5	5.7	6.5	7.1	8.9	8.9	8.8	8.6	8.6	8.5†	8.7	8.1	6.9	6.9	6.4	6.3	6.9	5.3	5.7	5.8	5.4	5.0	5.1	+0.1
12th Grade	7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4†	11.0	11.3	10.4	10.8	10.3	9.8	9.5	9.3	8.6	8.6	8.9	8.4	8.4	-0.1
College Students	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9†	7.5	7.8	8.2	9.1	8.2	8.2	8.1	7.3	8.4	8.1	8.2	7.8	8.2	+0.4
Young Adults	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4†	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	8.5	8.6	8.4	7.8	8.3	+0.5
Any Illicit Drug including Inhalants^{a,c,d}																								
8th Grade	8.8	10.0	12.0	14.3	16.1	17.5	16.0	14.9	15.1	14.4	14.0	12.6	12.1	11.2	11.2	10.9	10.1	10.4	10.6	11.7	10.5	9.5	9.9	+0.3
10th Grade	13.1	12.6	15.5	20.0	21.6	24.5	24.1	22.5	23.1	23.6	23.6	21.7	20.5	19.3	18.4	17.7	18.1	16.8	18.8	19.4	20.1	19.3	20.1	+0.8
12th Grade	17.8	15.5	19.3	23.0	24.8	25.5	26.9	26.6	26.4	26.4	26.5	25.9	24.6	23.3	24.2	22.1	22.8	22.8	24.1	24.5	26.2	25.2	26.6	+1.4
College Students	15.1	16.5	15.7	16.4	19.6	18.0	19.6	21.0	21.8	22.6	21.9	21.9	21.6	21.7	19.0	19.7	18.1	18.9	21.3	20.5	20.6	20.0	22.5	+2.5
Young Adults	15.4	15.3	15.1	16.1	16.1	16.4	16.9	16.7	17.4	18.8	19.2	19.5	20.1	19.6	18.0	18.4	19.1	19.3	20.3	19.6	20.3	19.1	22.4	+3.2 ss
Marijuana/Hashish																								
8th Grade	3.2	3.7	5.1	7.8	9.1	11.3	10.2	9.7	9.7	9.1	9.2	8.3	7.5	6.4	6.6	6.5	5.7	5.8	6.5	8.0	7.2	6.5	7.0	+0.5
10th Grade	8.7	8.1	10.9	15.8	17.2	20.4	20.5	18.7	19.4	19.7	19.8	17.8	17.0	15.9	15.2	14.2	14.2	13.8	15.9	16.7	17.6	17.0	18.0	+0.9
12th Grade	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	19.8	18.3	18.8	19.4	20.6	21.4	22.6	22.9	22.7	-0.2
College Students	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	20.0	20.2	19.7	19.3	18.9	17.1	16.7	16.8	17.0	18.5	17.5	19.4	20.5	20.6	+0.2
Young Adults	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	15.8	15.7	16.0	16.0	17.0	16.1	18.3	17.7	19.0	+1.3
Inhalants^{c,d}																								
8th Grade	4.4	4.7	5.4	5.6	6.1	5.8	5.6	4.8	5.0	4.5	4.0	3.8	4.1	4.5	4.2	4.1	3.9	4.1	3.8	3.6	3.2	2.7	2.3	-0.4
10th Grade	2.7	2.7	3.3	3.6	3.5	3.3	3.0	2.9	2.6	2.6	2.4	2.4	2.2	2.4	2.2	2.3	2.5	2.1	2.2	2.0	1.7	1.4	1.3	-0.1
12th Grade	2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	2.0	1.5	1.2	1.4	1.2	1.4	1.0	0.9	1.0	+0.1
College Students	0.9	1.1	1.3	0.6	1.6	0.8	0.8	0.6	1.5	0.9	0.4	0.7	0.4	0.4	0.3	0.4	0.1	0.4	0.1	0.5	0.3	0.2	0.1	-0.2
Young Adults	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	0.2	0.3	0.2	0.4	0.2	0.1	0.1	0.3	0.1	-0.1
Nitrites^e																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.6	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	*	0.1	0.2	0.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}																								
8th Grade	0.8	1.1	1.2	1.3	1.7	1.9	1.8	1.4	1.3	1.2†	1.6	1.2	1.2	1.0	1.1	0.9	1.0	0.9	0.9	1.0	1.0	0.6	0.8	+0.1
10th Grade	1.6	1.8	1.9	2.4	3.3	2.8	3.3	3.2	2.9	2.3†	2.1	1.6	1.5	1.6	1.5	1.5	1.7	1.3	1.4	1.6	1.4	1.2	1.1	-0.1
12th Grade	2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6†	3.3	2.3	1.8	1.9	1.9	1.5	1.7	2.2	1.6	1.9	1.6	1.6	1.4	-0.2
College Students	1.2	2.3	2.5	2.1	3.3	1.9	2.1	2.1	2.0	1.4†	1.8	1.2	1.8	1.3	1.2	0.9	1.3	1.7	1.0	1.4	1.2	1.1	1.0	-0.1
Young Adults	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2†	1.2	0.9	1.2	0.9	0.8	0.7	0.9	0.9	0.8	1.0	0.9	0.6	1.0	+0.4 s
LSD																								
8th Grade	0.6	0.9	1.0	1.1	1.4	1.5	1.5	1.1	1.1	1.0	1.0	0.7	0.6	0.5	0.5	0.4	0.5	0.5	0.5	0.6	0.5	0.3	0.5	+0.1
10th Grade	1.5	1.6	1.6	2.0	3.0	2.4	2.8	2.7	2.3	1.6	1.5	0.7	0.6	0.6	0.6	0.7	0.7	0.7	0.5	0.7	0.7	0.5	0.6	0.0
12th Grade	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	0.7	0.6	0.6	1.1	0.5	0.8	0.8	0.8	0.8	0.0
College Students	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	0.9	1.0	0.2	0.2	0.2	0.1	0.3	0.3	0.8	0.3	0.7	0.5	0.4	0.4	-0.1
Young Adults	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.1	0.2	0.2	0.4	0.2	0.4	0.3	0.3	0.4	+0.2

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Hallucinogens other than LSD^b																								
8th Grade	0.3	0.4	0.5	0.7	0.8	0.9	0.7	0.7	0.6	0.6‡	1.1	1.0	1.0	0.8	0.9	0.7	0.7	0.7	0.7	0.8	0.7	0.5	0.5	0.0
10th Grade	0.4	0.5	0.7	1.0	1.0	1.0	1.2	1.4	1.2	1.2‡	1.4	1.4	1.2	1.4	1.3	1.3	1.4	1.0	1.1	1.2	1.1	0.9	0.8	-0.1
12th Grade	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7‡	1.9	2.0	1.5	1.7	1.6	1.3	1.4	1.6	1.4	1.5	1.2	1.3	1.0	-0.3
College Students	0.6	0.7	1.1	0.8	1.6	1.2	1.2	0.7	1.2	0.8‡	0.8	1.1	1.7	1.2	1.1	0.7	1.1	1.3	0.8	1.2	0.8	0.7	0.8	+0.1
Young Adults	0.3	0.5	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.7‡	0.6	0.8	1.2	0.9	0.8	0.6	0.8	0.7	0.7	0.8	0.6	0.4	0.7	+0.2
PCP^g																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	0.7	0.4	0.5	0.6	0.5	0.8	0.8	0.5	0.4	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.1	0.2	0.2	0.1	*	0.1	0.1	0.2	0.2	*	*	0.1	0.1	0.1	*	*	*	0.1	*	0.0	0.1	0.0	0.2	+0.2
Ecstasy (MDMA)^h																								
8th Grade	—	—	—	—	—	1.0	1.0	0.9	0.8	1.4	1.8	1.4	0.7	0.8	0.6	0.7	0.6	0.8	0.6	1.1	0.6	0.5	0.5	0.0
10th Grade	—	—	—	—	—	1.8	1.3	1.3	1.8	2.6	2.6	1.8	1.1	0.8	1.0	1.2	1.2	1.1	1.3	1.9	1.6	1.0	1.2	+0.2
12th Grade	—	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	1.0	1.3	1.6	1.8	1.8	1.4	2.3	0.9	1.5	+0.5
College Students	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	2.5	1.5	0.7	1.0	0.7	0.8	0.6	0.4	0.6	0.5	1.0	0.7	1.4	0.8	-0.6
Young Adults	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	0.6	0.7	0.5	0.6	0.6	0.8	0.7	1.0	1.1	+0.1
Cocaine																								
8th Grade	0.5	0.7	0.7	1.0	1.2	1.3	1.1	1.4	1.3	1.2	1.2	1.1	0.9	0.9	1.0	1.0	0.9	0.8	0.8	0.6	0.8	0.5	0.5	0.0
10th Grade	0.7	0.7	0.9	1.2	1.7	1.7	2.0	2.1	1.8	1.8	1.3	1.6	1.3	1.7	1.5	1.5	1.3	1.2	0.9	0.9	0.7	0.8	0.8	0.0
12th Grade	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	2.3	2.5	2.0	1.9	1.3	1.3	1.1	1.1	1.1	0.0
College Students	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	1.4	1.9	1.6	1.9	2.4	1.8	1.8	1.7	1.2	1.3	1.0	1.2	1.1	0.9	-0.3
Young Adults	2.0	1.8	1.4	1.3	1.5	1.2	1.6	1.7	1.9	1.7	2.2	2.2	2.4	2.2	2.2	2.3	2.1	1.9	1.8	1.4	1.5	1.3	1.5	+0.2
Crackⁱ																								
8th Grade	0.3	0.5	0.4	0.7	0.7	0.8	0.7	0.9	0.8	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.4	0.5	0.3	0.3	+0.1
10th Grade	0.3	0.4	0.5	0.6	0.9	0.8	0.9	1.1	0.8	0.9	0.7	1.0	0.7	0.8	0.7	0.7	0.5	0.5	0.4	0.5	0.4	0.4	0.4	0.0
12th Grade	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	1.0	0.9	0.9	0.8	0.6	0.7	0.5	0.6	0.6	0.0
College Students	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.1	0.3	0.4	0.4	0.1	*	0.1	0.1	0.1	0.1	0.1	0.0	0.3	+0.3
Young Adults	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2	0.1	0.2	0.1	0.1	+0.1
Other Cocaine^j																								
8th Grade	0.5	0.5	0.6	0.9	1.0	1.0	0.8	1.0	1.1	0.9	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.7	0.5	0.6	0.3	0.3	0.0
10th Grade	0.6	0.6	0.7	1.0	1.4	1.3	1.6	1.8	1.6	1.6	1.2	1.3	1.1	1.5	1.3	1.3	1.1	1.0	0.8	0.7	0.6	0.7	0.7	0.0
12th Grade	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	2.0	2.4	1.7	1.7	1.1	1.1	1.0	1.0	0.9	-0.1
College Students	1.0	0.9	0.6	0.3	0.8	0.6	1.3	1.5	1.0	0.9	1.5	1.4	1.9	2.2	1.8	1.3	1.6	1.1	1.2	1.0	1.2	1.3	0.9	-0.3
Young Adults	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	1.9	1.9	2.0	1.7	1.6	1.5	1.4	1.3	1.3	0.0
Heroin^k																								
8th Grade	0.3	0.4	0.4	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.5	0.3	0.4	0.4	0.4	0.4	0.4	0.2	0.3	0.0
10th Grade	0.2	0.2	0.3	0.4	0.6	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	-0.1
12th Grade	0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.0
College Students	0.1	*	*	*	0.1	*	0.2	0.1	0.1	0.2	0.1	*	*	0.1	0.1	0.2	0.1	*	0.1	0.0	0.0	0.1	0.2	+0.1
Young Adults	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	+0.2
With a Needle^l																								
8th Grade	—	—	—	—	0.4	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.0
10th Grade	—	—	—	—	0.3	0.3	0.3	0.4	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.0
12th Grade	—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.1	0.4	0.4	0.3	0.2	0.0
College Students	—	—	—	—	*	*	0.1	*	0.1	0.1	*	*	0.1	0.1	0.1	0.1	*	0.0	0.1	0.0	0.0	0.2	0.1	-0.1
Young Adults	—	—	—	—	*	*	0.1	*	0.1	*	0.2	*	*	0.1	0.1	0.1	*	*	0.1	0.1	0.2	0.2	0.3	+0.1

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Without a Needle^l																								
8th Grade	—	—	—	—	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	+0.1
10th Grade	—	—	—	—	0.3	0.3	0.4	0.5	0.5	0.4	0.2	0.4	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.3	0.2	0.2	0.2	-0.1
12th Grade	—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	0.5	0.3	0.4	0.2	0.3	0.4	0.4	0.2	0.2	+0.1
College Students	—	—	—	—	*	0.1	0.2	0.2	0.3	0.4	0.3	*	*	0.3	*	0.2	0.1	0.1	0.1	0.0	0.0	0.1	0.3	+0.2
Young Adults	—	—	—	—	0.1	*	0.1	0.2	0.2	0.2	0.4	*	0.1	0.1	0.1	0.3	0.2	*	0.3	0.1	0.1	0.1	0.4	+0.2
Narcotics other than Heroin^{m,n}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0†	4.0	4.1	4.3	3.9	3.8	3.8	3.8	4.1	3.6	3.6	3.0	2.8	-0.3
College Students	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	1.7	1.7†	3.2	2.3	3.0	3.1	3.1	2.2	2.3	2.7	2.3	2.1	2.2	1.5	-0.7
Young Adults	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7†	2.9	2.9	3.0	3.5	3.2	3.4	3.6	3.2	3.4	2.9	2.7	2.6	-0.1
Amphetamines^{m,o}																								
8th Grade	2.6	3.3	3.6	3.6	4.2	4.6	3.8	3.3	3.4	3.4	3.2	2.8	2.7	2.3	2.3	2.1	2.0	2.2	1.9	1.8	1.8	1.3	1.4	+0.1
10th Grade	3.3	3.6	4.3	4.5	5.3	5.5	5.1	5.1	5.0	5.4	5.6	5.2	4.3	4.0	3.7	3.5	4.0	2.8	3.3	3.3	3.1	2.8	2.8	0.0
12th Grade	3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	3.9	3.7	3.7	2.9	3.0	3.3	3.7	3.3	4.1	+0.8
College Students	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	2.9	3.3	3.0	3.1	3.2	2.9	2.5	3.1	2.8	3.4	4.1	4.5	4.6	5.3	+0.6
Young Adults	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	2.1	2.2	2.3	2.2	2.5	2.9	3.0	3.2	3.2	0.0
Methamphetamine^{p,q}																								
8th Grade	—	—	—	—	—	—	—	—	1.1	0.8	1.3	1.1	1.2	0.6	0.7	0.6	0.6	0.7	0.5	0.7	0.4	0.5	0.4	-0.1
10th Grade	—	—	—	—	—	—	—	—	1.8	2.0	1.5	1.8	1.4	1.3	1.1	0.7	0.4	0.7	0.6	0.7	0.5	0.6	0.4	-0.2
12th Grade	—	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	0.9	0.9	0.6	0.6	0.5	0.5	0.6	0.5	0.4	-0.1
College Students	—	—	—	—	—	—	—	—	1.2	0.2	0.5	0.2	0.6	0.2	0.1	0.2	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0
Young Adults	—	—	—	—	—	—	—	—	0.8	0.7	1.0	1.0	0.7	0.6	0.7	0.5	0.6	0.3	0.3	0.2	0.3	0.4	0.2	-0.2
Crystal Methamphetamine (Ice)^q																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	0.9	0.7	0.6	0.6	0.5	0.6	0.6	0.4	0.8	+0.4
College Students	*	*	0.3	0.5	0.3	0.1	0.2	0.3	*	*	0.1	*	0.3	0.1	0.2	*	0.1	0.0	0.0	0.2	0.0	0.3	0.0	-0.3
Young Adults	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.4	0.4	0.6	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.4	+0.1
Sedatives (Barbiturates)^{m,r}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9†	2.9	3.3	3.0	2.7	2.8	2.5	2.2	1.8	2.0	2.2	+0.2
College Students	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	1.5	1.7	1.7	1.5	1.3	1.3	1.4	1.4	1.2	0.6	0.8	0.8†	0.9	—	
Young Adults	0.5	0.5	0.6	0.6	0.8	0.8	0.9	1.1	1.3	1.7	1.5	1.5	1.8	1.7	1.5	1.6	1.9	1.2	1.1	1.1	1.1†	1.2	—	
Methaqualone^{m,s}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	0.5	0.4	0.4	0.2	0.3	0.2	0.2	0.3	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}																								
8th Grade	0.8	0.8	0.9	1.1	1.2	1.5	1.2	1.2	1.1	1.4†	1.2	1.2	1.4	1.2	1.3	1.3	1.1	1.2	1.2	1.2	1.0	0.8	0.9	+0.2
10th Grade	1.2	1.5	1.1	1.5	1.7	1.7	2.2	2.2	2.2	2.5†	2.9	2.9	2.4	2.3	2.3	2.4	2.6	1.9	2.0	2.2	1.9	1.7	1.6	-0.1
12th Grade	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6†	2.9	3.3	2.8	3.1	2.9	2.7	2.6	2.6	2.7	2.5	2.3	2.1	2.0	-0.1
College Students	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	2.0†	1.5	3.0	2.8	2.7	2.2	2.1	1.8	1.6	2.2	1.3	1.6	1.1	1.2	+0.1
Young Adults	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8†	2.1	2.8	2.4	2.7	2.6	2.3	2.8	2.7	2.8	2.2	2.3	1.9	1.9	0.0

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Any Prescription Drug ^{o,t}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.6	8.1	7.8	7.2	7.3	6.9	7.2	7.0	7.0	+0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rohypnol ^u																								
8th Grade	—	—	—	—	—	0.5	0.3	0.4	0.3	0.3	0.4	0.2	0.1	0.2	0.2	0.4	0.3	0.1	0.2	0.2	0.6	0.1	0.1	0.0
10th Grade	—	—	—	—	—	0.5	0.5	0.4	0.5	0.4	0.2	0.4	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.2	0.1	-0.1
12th Grade	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^y																								
Any Use																								
8th Grade	25.1	26.1	24.3	25.5	24.6	26.2	24.5	23.0	24.0	22.4	21.5	19.6	19.7	18.6	17.1	17.2	15.9	15.9	14.9	13.8	12.7	11.0	10.2	-0.8
10th Grade	42.8	39.9	38.2	39.2	38.8	40.4	40.1	38.8	40.0	41.0	39.0	35.4	35.4	35.2	33.2	33.8	33.4	28.8	30.4	28.9	27.2	27.6	25.7	-1.9
12th Grade	54.0	51.3	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	47.0	45.3	44.4	43.1	43.5	41.2	40.0	41.5	39.2	-2.3 s
College Students	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	67.4	67.0	68.9	66.2	67.7	67.9	65.4	66.6	69.0	65.8	65.0	63.5	67.7	63.1	-4.6 s
Young Adults	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.0	68.3	67.0	68.4	68.6	68.7	69.5	68.9	69.4	68.4	68.8	69.5	68.7	-0.8
Been Drunk ^w																								
8th Grade	7.6	7.5	7.8	8.7	8.3	9.6	8.2	8.4	9.4	8.3	7.7	6.7	6.7	6.2	6.0	6.2	5.5	5.4	5.4	5.0	4.4	3.6	3.5	-0.1
10th Grade	20.5	18.1	19.8	20.3	20.8	21.3	22.4	21.1	22.5	23.5	21.9	18.3	18.2	18.5	17.6	18.8	18.1	14.4	15.5	14.7	13.7	14.5	12.8	-1.6 s
12th Grade	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	30.2	30.0	28.7	27.6	27.4	26.8	25.0	28.1	26.0	-2.1
College Students	45.0	45.0	43.8	42.8	37.9	40.3	46.4	44.3	44.6	43.9	44.7	44.4	40.4	47.4	43.1	47.6	46.8	45.3	42.4	43.6	39.9	40.1	40.2	+0.2
Young Adults	35.4	35.6	34.2	34.3	33.0	33.2	35.6	34.2	37.7	35.7	36.8	37.1	37.8	39.0	39.0	42.1	41.4	40.7	40.5	39.4	39.5	39.1	37.7	-1.4
Flavored Alcoholic Beverages ^{9,p}																								
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	14.6	12.9	13.1	12.2	10.2	9.5	9.4	8.6	7.6	6.3	-1.3 s
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	25.1	23.1	24.7	21.8	20.2	19.0	19.4	15.8	16.3	15.5	-0.7
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	31.1	30.5	29.3	29.1	27.4	27.4	24.1	23.1	21.8	21.0	-0.7
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	34.1	30.9	26.2	27.5	35.8	32.3	31.5	29.5	31.3	29.1	-2.2
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	27.6	24.9	25.9	26.7	24.4	24.5	23.8	26.1	25.4	-0.7
Cigarettes																								
Any Use																								
8th Grade	14.3	15.5	16.7	18.6	19.1	21.0	19.4	19.1	17.5	14.6	12.2	10.7	10.2	9.2	9.3	8.7	7.1	6.8	6.5	7.1	6.1	4.9	4.5	-0.5
10th Grade	20.8	21.5	24.7	25.4	27.9	30.4	29.8	27.6	25.7	23.9	21.3	17.7	16.7	16.0	14.9	14.5	14.0	12.3	13.1	13.6	11.8	10.8	9.1	-1.7 s
12th Grade	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	23.2	21.6	20.4	20.1	19.2	18.7	17.1	16.3	-0.9	
College Students	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	28.2	25.7	26.7	22.5	24.3	23.8	19.2	19.9	17.9	17.9	16.4	15.2	12.5	14.0	+1.5
Young Adults	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	28.6	27.0	26.2	24.6	23.3	22.4	21.3	19.7	20.0	+0.4
Smokeless Tobacco ^x																								
8th Grade	6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	3.2	3.5	3.7	4.1	3.5	2.8	2.8	+0.1
10th Grade	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	6.5	7.5	6.6	6.4	6.4	0.0
12th Grade	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	8.4	8.5	8.3	7.9	8.1	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^{y,z}																								
8th Grade	0.4	0.5	0.5	0.5	0.6	0.4	0.5	0.5	0.7	0.8	0.7	0.8	0.7	0.5	0.5	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.0
10th Grade	0.6	0.6	0.5	0.6	0.6	0.5	0.7	0.6	0.9	1.0	0.9	1.0	0.8	0.8	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.0
12th Grade	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	0.9	1.1	1.0	1.0	1.0	1.1	0.7	0.9	1.0	+0.1
College Students	0.3	0.2	0.2	0.2	0.1	*	0.2	0.2	0.4	*	0.3	*	0.1	*	*	*	0.1	*	0.2	0.0	0.2	0.0	0.0	0.0
Young Adults	0.2	0.1	*	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.2	0.3	0.5	0.2	0.1	0.1	0.0

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 2-4.

TABLE 2-4
Trends in 30-Day Prevalence of Daily Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
Marijuana/Hashish																								
Daily ⁹⁹																								
8th Grade	0.2	0.2	0.4	0.7	0.8	1.5	1.1	1.1	1.4	1.3	1.3	1.2	1.0	0.8	1.0	1.0	0.8	0.9	1.0	1.2	1.3	1.1	1.1	0.0
10th Grade	0.8	0.8	1.0	2.2	2.8	3.5	3.7	3.6	3.8	3.8	4.5	3.9	3.6	3.2	3.1	2.8	2.8	2.7	2.8	3.3	3.6	3.5	4.0	+0.5
12th Grade	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	5.0	5.0	5.1	5.4	5.2	6.1	6.6	6.5	6.5	0.0
College Students	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	4.6	4.5	4.1	4.7	4.5	4.0	4.3	3.5	3.9	4.9	4.4	4.7	4.8	5.1	+0.4
Young Adults	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	4.9	5.0	5.0	5.1	5.4	5.3	6.1	5.6	6.2	+0.6
Alcohol ^{v,99}																								
Any Daily Use																								
8th Grade	0.5	0.6‡	1.0	1.0	0.7	1.0	0.8	0.9	1.0	0.8	0.9	0.7	0.8	0.6	0.5	0.5	0.6	0.7	0.5	0.5	0.4	0.3	0.3	-0.1
10th Grade	1.3	1.2‡	1.8	1.7	1.7	1.6	1.7	1.9	1.9	1.8	1.9	1.8	1.5	1.3	1.3	1.4	1.4	1.0	1.1	1.1	0.8	1.0	0.9	-0.1
12th Grade	3.6	3.4‡	3.4	2.9	3.5	3.7	3.9	3.9	3.4	2.9	3.6	3.5	3.2	2.8	3.1	3.0	3.1	2.8	2.5	2.7	2.1	2.5	2.2	-0.4
College Students	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	3.6	4.7	5.0	4.3	3.7	4.6	4.8	4.3	4.0	4.3	3.6	3.8	3.9	3.6	-0.3
Young Adults	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	5.2	5.4	5.6	5.3	5.3	4.6	5.2	5.5	5.1	-0.4
Been Drunk																								
Daily ^{w,99}																								
8th Grade	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.0
10th Grade	0.2	0.3	0.4	0.4	0.6	0.4	0.6	0.6	0.7	0.5	0.6	0.5	0.5	0.4	0.4	0.5	0.5	0.3	0.4	0.3	0.2	0.4	0.3	-0.1
12th Grade	0.9	0.8	0.9	1.2	1.3	1.6	2.0	1.5	1.9	1.7	1.4	1.2	1.6	1.8	1.5	1.6	1.3	1.4	1.1	1.6	1.3	1.5	1.3	-0.1
College Students	0.5	0.2	0.3	0.8	0.5	0.1	1.3	0.8	1.0	0.7	0.5	0.8	1.1	0.8	0.5	0.6	0.7	0.5	0.7	0.3	1.3	0.4	0.5	+0.1
Young Adults	0.5	0.4	0.4	0.5	0.3	0.4	0.9	0.5	0.9	0.5	0.4	0.6	0.8	0.7	0.5	0.6	0.6	0.5	1.0	0.7	0.7	0.4	0.5	+0.1
5+ Drinks in a Row																								
in Last 2 Weeks																								
8th Grade	10.9	11.3	11.3	12.1	12.3	13.3	12.3	11.5	13.1	11.7	11.0	10.3	9.8	9.4	8.4	8.7	8.3	8.1	7.8	7.2	6.4	5.1	5.1	0.0
10th Grade	21.0	19.1	21.0	21.9	22.0	22.8	23.1	22.4	23.5	24.1	22.8	20.3	20.0	19.9	19.0	19.9	19.6	16.0	17.5	16.3	14.7	15.6	13.7	-1.9 s
12th Grade	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	27.1	25.4	25.9	24.6	25.2	23.2	21.6	23.7	22.1	-1.6
College Students	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	39.3	40.9	40.1	38.5	41.7	40.1	40.2	41.1	40.0	36.9	37.0	36.1	37.4	35.2	-2.1
Young Adults	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	37.0	37.6	37.8	37.9	36.7	35.9	36.5	35.5	35.1	-0.4
Cigarettes																								
Any Daily Use																								
8th Grade	7.2	7.0	8.3	8.8	9.3	10.4	9.0	8.8	8.1	7.4	5.5	5.1	4.5	4.4	4.0	4.0	3.0	3.1	2.7	2.9	2.4	1.9	1.8	-0.1
10th Grade	12.6	12.3	14.2	14.6	16.3	18.3	18.0	15.8	15.9	14.0	12.2	10.1	8.9	8.3	7.5	7.6	7.2	5.9	6.3	6.6	5.5	5.0	4.4	-0.5
12th Grade	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	13.6	12.2	12.3	11.4	11.2	10.7	10.3	9.3	8.5	-0.8
College Students	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	17.8	15.0	15.9	13.8	13.8	12.4	9.2	9.3	9.2	8.0	7.6	7.3	5.2	5.6	+0.4
Young Adults	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	19.6	18.6	17.3	16.7	15.0	14.8	13.8	12.8	12.1	-0.7
1/2 Pack+/Day																								
8th Grade	3.1	2.9	3.5	3.6	3.4	4.3	3.5	3.6	3.3	2.8	2.3	2.1	1.8	1.7	1.7	1.5	1.1	1.2	1.0	0.9	0.7	0.6	0.7	+0.1
10th Grade	6.5	6.0	7.0	7.6	8.3	9.4	8.6	7.9	7.6	6.2	5.5	4.4	4.1	3.3	3.1	3.3	2.7	2.0	2.4	2.4	1.9	1.5	1.5	0.0
12th Grade	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0	6.9	5.9	5.7	5.4	5.0	4.7	4.3	4.0	3.4	-0.6
College Students	8.0	8.9	8.9	8.0	10.2	8.4	9.1	11.3	11.0	10.1	7.8	7.9	7.6	6.8	6.7	4.9	4.3	4.3	3.8	3.9	2.5	2.4	2.4	0.0
Young Adults	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	12.5	11.9	11.1	10.2	9.3	9.3	7.5	7.6	7.0	-0.6
Smokeless Tobacco																								
Daily *																								
8th Grade	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	0.8	0.8	0.8	0.9	0.8	0.5	0.5	0.0
10th Grade	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	1.6	1.4	1.9	2.5	1.7	2.0	1.9	-0.2
12th Grade	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	2.8	2.7	2.9	3.1	3.1	3.2	3.0	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

Source: The Monitoring the Future study, the University of Michigan.

See footnotes on the next page.

Footnotes for Tables 2-1 through 2-4

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. ' — ' indicates data not available.

' * ' indicates less than 0.05% but greater than 0%. ' ‡ ' indicates some change in the question. See relevant footnote for that drug.

See relevant figure to assess the impact of the wording changes. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

Approximate Weighted <i>N</i> s	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002
8th Graders	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100
10th Graders	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300
12th Graders	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900
College Students	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260
Young Adults	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300

Approximate Weighted <i>N</i> s	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
8th Graders	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	14,600	14,600
10th Graders	15,800	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	12,900	12,900
12th Graders	14,600	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	12,600	12,600
College Students	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090
Young Adults	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,630	4,580	4,360

^aFor 12th graders, college students, and young adults only: Use of any illicit drug includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders. For 8th and 10th graders only: The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because these younger respondents appear to overreport use (perhaps because they include the use of nonprescription drugs in their answers).

^bIn 2001 the question text was changed on half of the questionnaire forms for each age group. Other psychedelics was changed to other hallucinogens and shrooms was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. For 8th, 10th, and 12th graders only: The 2001 data presented here are based on the changed forms only; *N* is one half of *N* indicated. In 2002 the remaining forms were changed to the new wording. The data are based on all forms beginning in 2002. Data for any illicit drug other than marijuana and data for hallucinogens are also affected by these changes and have been handled in a parallel manner.

^cFor 12th graders, college students, and young adults only: Data based on five of six forms in 1991–1998; *N* is five sixths of *N* indicated. Data based on three of six forms beginning in 1999; *N* is three sixths of *N* indicated.

^dInhalants are unadjusted for underreporting of amyl and butyl nitrites.

^eFor 12th graders and young adults only: Data based on one of six forms; *N* is one sixth of *N* indicated. Questions about nitrite use were dropped from the young adult questionnaires in 1995 and from the 12th-grade questionnaires in 2010.

^fHallucinogens are unadjusted for underreporting of PCP.

^gFor 12th graders, college students, and young adults only: Data based on one of six forms; *N* is one sixth of *N* indicated. For 12th graders only: In 2011 the flavored alcoholic beverage question text was changed. Skyy Blue and Zima were removed from the list of examples. An examination of the data did not show any effect from the wording change.

^hFor 8th and 10th graders only: Data based on one of two forms in 1996; *N* is one half of *N* indicated. Data based on one third of *N* indicated in 1997–2001 due to changes in the questionnaire forms. Data based on two of four forms beginning in 2002; *N* is one half of *N* indicated. For 12th graders only: Data based on one of six forms in 1996–2001; *N* is one sixth of *N* indicated. Data based on two of six forms beginning in 2002; *N* is two sixths of *N* indicated. For college students and young adults only: Data based on two of six forms in 1991–2001; *N* is two sixths of *N* indicated. Data based on three of six forms beginning in 2002; *N* is three sixths of *N* indicated.

ⁱFor college students and young adults only: Data based on five of six forms beginning in 2002; *N* is five sixths of *N* indicated.

^jFor 12th graders only: Data based on four of six forms; *N* is four sixths of *N* indicated. For college students and young adults only: Data based on four of six forms; *N* is four sixths of *N* indicated.

^kIn 1995, the heroin question was changed in one of two forms for 8th and 10th graders, in three of six forms for 12th graders, and in two of six forms for college students and young adults. Separate questions were asked for use with and without injection. In 1996, the heroin question was changed in all remaining 8th- and 10th-grade forms. Data presented here represent the combined data from all forms.

^lFor 8th and 10th graders only: Data based on one of two forms in 1995; *N* is one half of *N* indicated. Data based on all forms beginning in 1996. For 12th graders only: Data based on three of six forms; *N* is three sixths of *N* indicated. For college students and young adults only: Data based on two of six forms; *N* is two sixths of *N* indicated.

^mOnly drug use not under a doctor's orders is included here.

ⁿFor 12th graders, college students, and young adults only: In 2002 the question text was changed in half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced with Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; *N* is one half of *N* indicated. In 2003, the remaining forms were changed to the new wording. The data are based on all forms beginning in 2003. In 2013 the list of examples was changed on one form: MS Contin, Roxycodone, Hydrocodone (Lortab, Lorcet, Norco), Suboxone, Tylox, and Tramadol were added to the list. An examination of the data did not show any affect from the wording change.


^oIn 2009, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. In 2010 the remaining forms were changed in a like manner. For 12th graders only: In 2011 the introduction to the question was changed slightly in one of six forms. Bennies, Benzedrine, and Methedrine were deleted from the list of examples. An examination of the data did not show any effect from the wording change. In 2013 the question wording was changed slightly in two of the 8th and 10th grade questionnaires and in four of the 12th grade questionnaires. Vyvanse was also added to the list of examples in one of the 12th grade forms. 2013 data are based on the unchanged forms only; for 8th and 10th graders *N* is one half of *N* indicated, for 12th graders *N* is two sixths of *N* indicated. Data for prescription drug use are affected by these changes and have been handled in a parallel manner.

(Footnotes continued on next page.)

Footnotes for Tables 2-1 through 2-4 (cont.)

- ^pFor 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated. In 2011 the flavored alcoholic beverage question text was changed. Skyy Blue and Zima were removed from the list of examples. An examination of the data did not show any effect from the wording change.
- ^qFor 12th graders only: Data based on two of six forms; N is two sixths of N indicated. Provigil was dropped from the study in 2012. For college students and young adults only: Beginning in 2009 Salvia data based on one of six forms; N is one sixth of N indicated. Data based on two of six forms in 2010 and 2011; N is two sixths of N indicated. Data based on three of six forms beginning in 2012; N is three sixths of N indicated. For Synthetic Marijuana data based on two of six forms in 2011; N is two sixths of N indicated. Data based on three of six forms beginning in 2012; N is three sixths of N indicated. For Bath Salts data based on three of six forms; N is three sixths of N indicated.
- ^rFor 12th graders only: In 2004 the question text was changed in half of the questionnaire forms. Barbiturates was changed to sedatives, including barbiturates. Goofballs, yellows, reds, blues, and rainbows were deleted from the list of examples; Phenobarbital, Tuinal, Nembutal, and Seconal were added. An examination of the data did not show any effect from the wording change. In 2005 the remaining forms were changed in a like manner. In 2013 the question text was changed in all forms: Tuinal, Nembutal, and Seconal were replaced with Ambien, Lunesta, and Sonata. In one form the list of examples was also changed: Tuinal was dropped from the list and Dalmane, Restoril, Halcion, Intermezzo, and Zolpimist were added. An examination of the data did not show any effect from the wording change. In 2013 the college student and young adult questionnaires were changed in a like manner. An examination of the data showed an affect from the wording change. For this reason 2012 and 2013 data are not comparable.
- ^sFor 12th graders only: Data based on one of six forms; N is one sixth of N indicated. Methaqualone was dropped from the study in 2013. For college students and young adults only: Data based on three of six forms. N is three sixths of N indicated.
- ^tThe use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers...without a doctor telling you to use them.
- ^uFor 8th and 10th graders only: Data based on one of two forms in 1996; N is one half of N indicated. Data based on three of four forms in 1997–1998; N is two thirds of N indicated. Data based on two of four forms in 1999–2001; N is one third of N indicated. Data based on one of four forms beginning in 2002; N is one sixth of N indicated. For 12th graders only: Data based on one of six forms in 1996–2001; N is one sixth of N indicated. Data based on two of six forms in 2002–2009; N is two sixths of N indicated. Data for 2001 and 2002 are not comparable due to changes in the questionnaire forms. Data based on one of six forms beginning in 2010; N is one sixth of N indicated. For college students and young adults only: Data based on two of six forms; N is two sixths of N indicated.
- ^vFor 8th, 10th, and 12th graders only: In 1993, the question text was changed slightly in half of the forms to indicate that a drink meant more than just a few sips. The 1993 data are based on the changed forms only; N is one half of N indicated for these groups. In 1994 the remaining forms were changed to the new wording. The data are based on all forms beginning in 1994. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005. For college students and young adults: The revision of the question text resulted in rather little change in the reported prevalence of use. The data for all forms are used to provide the most reliable estimate of change.
- ^wFor all grades: In 2012 the alcoholic beverage containing caffeine (like Four Loko or Joose) question text was changed to alcoholic beverage mixed with an energy drink (like Red Bull). The data in 2011 and 2012 are not comparable due to this question change. For 12th graders only: Data based on two of six forms; N is two sixths of N indicated. For college students and young adults only: been drunk data based on three of six forms; N is three sixths of N indicated. Alcoholic beverages containing caffeine data based on two of six forms; N is two sixths of N indicated.
- ^xFor 8th and 10th graders only: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated. For 12th graders only: Data based on one of six forms; N is one sixth of N indicated. For 8th, 10th, and 12th graders only: Snus and dissolvable tobacco were added to the list of examples in 2011. An examination of the data did not show any effect from the wording change. For college students and young adults only: Questions about smokeless tobacco use were dropped from the analyses in 1989.
- ^yFor 8th and 10th graders only: In 2006, the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining forms were changed in a like manner. For 12th graders only: Data based on two of six forms in 1991–2005; N is two sixths of N indicated. In 2006 a slightly altered version of the question was added to a third form. An examination of the data did not show any effect from the wording change. Data based on three of six forms beginning in 2006; N is three sixths of N indicated. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in two of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining form was changed in a like manner.
- ^zFor college students and young adults only: Data based on two of six forms in 1990–2009; N is two sixths of N indicated. In 2008, the question text was changed slightly.
- ^{aa}For 12th graders only: Data based on two of six forms in 2002–2005; N is two sixths of N indicated. Data based on three of six forms beginning in 2006; N is three sixths of N indicated.
- ^{ab}For college students and young adults only: Data based on two of six forms through 2009; N is two sixths of N indicated. Data based on three of six forms beginning in 2010; N is three sixths of N indicated.
- ^{ac}For 12th graders only: Data based on two of six forms in 2000; N is two sixths of N indicated. Data based on three of six forms in 2001; N is three sixths of N indicated. Data based on one of six forms beginning in 2002; N is one sixth of N indicated. For college students and young adults only: Data based on two of six forms; N is two sixths of N indicated. Data based on three of six forms beginning in 2010; N is three sixths of N indicated. Data based on two of six forms beginning in 2012; N is two sixths of N indicated.
- ^{ad}For 12th graders only: Data based on two of six forms in 2000; N is two sixths of N indicated. Data based on three of six forms in 2001–2009; N is three sixths of N indicated. Data based on two of six forms beginning in 2010; N is two sixths of N indicated. For college students and young adults only: Data based on two of six forms; N is two sixths of N indicated. Data based on three of six forms beginning in 2010; N is three sixths of N indicated.
- ^{ae}For 12th graders only: The 2003 flavored alcoholic beverage data were created by adjusting the 2004 data to reflect the observed 2003 to 2004 change in a slightly different version of the flavored alcoholic beverage question. In 2004 the original question was revised to include wine coolers among the examples—a change that had very little effect on the observed prevalence-of-use rate.
- ^{af}For 12th graders only: Data based on two of six forms in 2000–2008; N is two sixths of N indicated. Beginning in 2009 data based on one of six forms; N is one sixth of N indicated.
- ^{ag}Daily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes and smokeless tobacco, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

FIGURE 2-1
Trends in Annual Prevalence of an Illicit Drug Use Index
across 5 Populations



Source. The Monitoring the Future study, the University of Michigan.

Notes. Illicit drug use index includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin which is not under a doctor's orders, stimulants, sedatives (barbiturates), methaqualone (excluded since 1990), or tranquilizers. Beginning in 1982, the question about stimulant use (i.e., amphetamines) was revised to get respondents to exclude the inappropriate reporting of nonprescription stimulants. The prevalence rate dropped slightly as a result of this methodological change.

Chapter 3

STUDY DESIGN AND PROCEDURES

Monitoring the Future (MTF) incorporates several types of surveys into one study, yielding analytic power beyond the sum of those component parts. The components include cross-sectional studies, repeated cross-sectional studies, and panel studies of individual cohorts or sets of cohorts. The annual cross-sectional surveys provide point estimates of various behaviors and conditions in any given year for a number of subpopulations (e.g., 8th graders, 10th graders, 12th graders, college students, all young adult high school graduates ages 19–30, 35-year-olds, 40-year-olds, etc.), as well as point estimates for various subgroups within these different populations. Repeating these annual cross-sectional surveys over time allows an assessment of change across history in consistent age segments of the population, as well as among subgroups. The panel study feature permits the examination of developmental change in the same individuals as they assume adult responsibilities, enter and leave various adult roles and environments, and continue further into adulthood. It also permits an assessment of a number of outcomes later in life that may be linked to substance use in adolescence and beyond.

Finally, with a *series* of panel studies of sequential graduating class cohorts, in what is known as a cohort-sequential design, we are able to offer distinctions among and explanations for three fundamentally different types of change: period, age, and cohort. It is this feature that creates the synergistic effect in terms of analytic and explanatory power.²³²⁴

RESEARCH DESIGN AND PROCEDURES FOR THE TWELFTH GRADE SURVEYS

Twelfth graders have been surveyed in the spring of each year since 1975. Each year's data collection has taken place in between 120 and 146 public and private high schools selected to provide an accurate representative cross-section of 12th graders throughout the coterminous United States (see Figure 3-1).

²³ For a more detailed description of the study design, see Bachman, J. G., Johnston, L. D., O'Malley, P. M., & Schulenberg, J. E. (2011). *The Monitoring the Future project after thirty-seven years: Design and procedures* (Monitoring the Future Occasional Paper No.76). Ann Arbor, MI: Institute for Social Research. Available online at <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ76.pdf>.

²⁴ For a more detailed description of the full range of research objectives of Monitoring the Future, see Johnston, L. D., O'Malley, P. M., Schulenberg, J. E., & Bachman, J. G. (2006). *The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2006* (Monitoring the Future Occasional Paper No. 65). Ann Arbor, MI: Institute for Social Research. Available online at <http://www.monitoringthefuture.org/pubs/occpapers/occ65.pdf>.

The Population under Study

Senior year of high school is an optimal point at which to monitor drug use and related attitudes of youth. First, completion of high school represents the end of an important developmental period in this society, demarcating both the end of universal education and, for many, the end of living full time in the parental home. Therefore, it is a logical point at which to take stock of cumulated influences. Further, completion of high school represents a jumping-off point—a point from which young people diverge into widely differing social environments and experiences. Thus senior year is a good time to take a “before” measure, allowing the subsequent calculation of changes that may be attributable to the environmental and role transitions occurring in young adulthood, including college attendance, military service, and so on. Finally, there are some important practical advantages built into the original system of data collections around samples of 12th graders. The need for systematically repeated, large-scale samples from which to make reliable estimates of change requires that considerable emphasis be put on cost efficiency as well as feasibility. The last year of high school constitutes the final point at which a reasonably good national sample of an age-specific cohort can be drawn and studied economically.

The Omission of Dropouts

One limitation in the MTF study design is the exclusion of individuals who drop out of high school before graduation—approximately 9–15% of each age cohort nationally, according to U.S. Census statistics. (The dropout rate has been declining in recent years; 9% is the most recent estimate.) Clearly, the omission of high school dropouts introduces biases in the estimation of certain characteristics of the entire age group; however, for most purposes, the small proportion of students who drop out sets outer limits on the bias. Further, since the bias from missing dropouts should remain relatively constant from year to year, their omission should introduce little or no bias in *change* estimates. Indeed, we believe the changes observed over time for those who are surveyed in the 12th grade are likely to parallel the changes for dropouts in most instances. Appendix A in Volume I addresses in detail the likely effects of the exclusion of dropouts (as well as absentees from school) on estimates of drug use prevalence and trends among the entire age cohort.

Sampling Procedures and Sample Weights

A multistage random sampling procedure is used to secure the nationwide sample of 12th graders each year. Stage 1 is the selection of particular geographic areas, Stage 2 is the selection (with probability proportionate to size) of one or more high schools in each area, and Stage 3 is the selection of 12th graders within each high school. Up to about 350 twelfth graders in each school may be included. In schools with fewer 12th graders, the usual procedure is to include all of them in the data collection, though a smaller sample is sometimes taken (either by randomly sampling entire classrooms or by some other unbiased, random method) to accommodate the needs of the school. Weights are assigned to compensate for differential probabilities of selection at each stage of sampling. Final weights are normalized to average 1.0 (so that the weighted

number of cases equals the unweighted number of cases overall). In order to be able to check observed trends in any given one-year interval, schools participate in the study for two consecutive years on a staggered schedule, with one half of them being replaced with a new random half-sample of schools each year. Therefore in any given year about half of the schools in the sample are participating for the first time and the other half are participating for their second and final year. This three-stage sampling procedure, with annual replacement of half of the sample of schools each year, has yielded the numbers of participating schools and students shown in Table 3-1.

Questionnaire Administration

About three weeks prior to the questionnaire administration date, parents of the target respondents are sent a letter by first-class mail, usually from the principal, announcing and describing the MTF study and providing parents an opportunity to decline participation of their son or daughter if they wish. A flyer outlining the study in more detail is enclosed with the letter. Copies of the flyers are also given to the students by teachers in the target classrooms in advance of the date of administration. The flyers make clear that participation is entirely voluntary. Local Institute for Social Research representatives and their assistants conduct the actual questionnaire administrations following standardized procedures detailed in an instruction manual. The questionnaires are administered in classrooms during a normal class period whenever possible; however, circumstances in some schools require the use of larger group administrations. Teachers are asked to remain present in the classroom to help maintain order, but to remain at their desks so that they cannot see students' answers.

Questionnaire Format

Because many questions are needed to cover all of the topic areas in the MTF study, much of the questionnaire content for 12th graders is divided into six different questionnaire forms distributed to participants in an ordered sequence that ensures six virtually identical random subsamples. (Five questionnaire forms were used between 1975 and 1988.) About one third of each form consists of key, or "core," variables common to all forms. All demographic variables, and nearly all of the drug *use* variables included in this report, are contained in this core set of measures. Many questions on attitudes, beliefs, and perceptions of relevant features of the social environment are in fewer forms, and data are thus based on fewer cases—a single form would have one fifth of the total number of cases in 1975–1988 (approximately 3,300 per year) and one sixth of the total beginning in 1989 (approximately 2,500 per year). All tables in this report list the sample sizes upon which the statistics are based, stated in terms of the weighted number of cases (which, as explained above, is roughly equivalent to the actual number of cases).

RESEARCH DESIGN AND PROCEDURES FOR THE EIGHTH- AND TENTH-GRADE SURVEYS

In 1991, MTF was expanded to include nationally representative samples of 8th- and 10th-grade students surveyed on an annual basis. Separate samples of schools and students are drawn at each grade level. In general, the procedures used for the annual in-school surveys of 8th- and 10th-grade students closely parallel those used for 12th graders, including the selection of schools and students, questionnaire administration, and questionnaire format. A major exception is that only two different questionnaire forms were used from 1991 to 1996, expanding to four forms beginning in 1997. The same four questionnaire forms are used for both 8th and 10th graders; most of the content is drawn from the 12th-grade surveys, including the core section. Thus, key demographic variables and measures of drug use and related attitudes and beliefs are generally identical for all three grades. Many fewer questions about other values and attitudes are included in the 8th- and 10th-grade forms, in part because we think that many of them are likely to be more fully formed by 12th grade and, therefore, are best monitored there.

About 16,000 eighth-grade students in approximately 150 schools (mostly middle schools) and about 13,000 to 15,000 tenth-grade students in approximately 125 schools are surveyed each year (see Table 3-1).

Mode of Administration

From 1991 to 1993, follow-ups for 8th and 10th graders were administered similarly to those for 12th graders.²⁵ When follow-up surveys of new 8th- and 10th-grade cohorts were discontinued, the collection of personal identification information was no longer necessary. (For confidentiality reasons, this personal information had been gathered on a tear-off sheet at the back of each questionnaire.) We believed that there were potential advantages in moving toward a fully anonymous procedure for these grade levels, including the following: (a) school cooperation might be easier to obtain; (b) any suppression effect on self-reported substance use that the confidential mode of administration might have could be both eliminated and quantified; and (c) if there *were* any mode of administration effect, it would be removed from the national data, which are widely compared with results of state and local surveys (nearly all of which use anonymous questionnaires), thus making those comparisons more valid. Therefore, in 1998, the half sample of schools beginning their two-year participation in MTF received fully anonymous questionnaires, while the half sample participating for their second and final year continued to get confidential questionnaires. In 1999 and thereafter, all questionnaires administered to 8th and 10th graders have been fully anonymous.

²⁵ A book reporting results from analyses of these younger panels was published in 2008. See Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education-drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

A careful examination of the 1998 results, based on the two equivalent half samples at grades 8 and 10, revealed that there was no effect of this methodological change among 10th graders and only a very modest effect, if any, in self-reported substance use rates among 8th graders (with prevalence rates slightly higher in the anonymous condition).²⁶ All tables and figures in *Volume I* combine data from both half samples of 8th graders surveyed in a given year. This is also true for 10th graders, for whom we found no methodological effect, and 12th graders, for whom we assumed no such effect since none was found for 10th graders. (See this chapter's later section entitled "Representativeness and Sample Accuracy" for a further discussion of half samples among all three grades.)

Questionnaire Forms and Sample Proportions

A benefit of not interlocking the 8th- and 10th-grade samples was that we could consider having more forms of the questionnaire. Beginning in 1997, the number of forms was expanded to four, although they are not distributed in equal numbers. Forms 1, 2, 3, and 4 are assigned to one third, one third, one sixth, and one sixth of the students, respectively. Thus, if a question appears on only one form, it is administered to either one third or one sixth of the sample. A question in two forms may be assigned to one third of the sample (one sixth plus one sixth), one half of the sample (one third plus one sixth), or two thirds of the sample (one third plus one third). No questions appear on exactly three forms. Footnotes to the tables indicate what proportion of all respondents in each grade complete the question, if that proportion is other than the entire sample. All of the samples, whether based on one or more forms, are random samples.

The two additional forms were introduced to allow for more questions. The new Forms 1 and 2 substantially follow the content of the previous Forms 1 and 2, but each is now assigned to a third of the sample instead of half. Form 3 builds on Form 1, with some questions omitted to make room for more content; and Form 4 builds on the content of Form 2 in a similar manner. Much of the new content was placed in both of the two new forms (Forms 3 and 4), each of which is administered to one sixth of the sample, in order to assign one third of the total sample to those new measures.

²⁶ We have examined in detail the effects of administration mode using multivariate controls to assess the effects of the change on 8th-grade self-report data. Our findings generally show even less effect than is to be found without such controls. See O'Malley, P. M., Johnston, L. D., Bachman, J. G., & Schulenberg, J. E. (2000). A comparison of confidential versus anonymous survey procedures: Effects on reporting of drug use and related attitudes and beliefs in a national study of students. *Journal of Drug Issues*, 30, 35–54.

RESEARCH DESIGN AND PROCEDURES FOR THE TWELFTH-GRADE FOLLOW-UP SURVEYS

Beginning with the graduating class of 1976, some members of each 12th-grade class have been selected to be surveyed by mail after high school. From the 13,000–19,000 twelfth graders originally surveyed in a given senior class, a representative sample of 2,400 is randomly chosen for follow-up. In order to ensure that drug-using populations are adequately represented in the follow-up surveys, 12th graders reporting 20 or more occasions of marijuana use in the previous 30 days (i.e., daily users), or *any* use of the other illicit drugs in the previous 30 days, are selected with higher probability (by a factor of 3.0) than the remaining 12th graders. Differential weighting is then used in all follow-up analyses to compensate for these differential sampling probabilities. Because those in the drug-using stratum receive a weight of only 0.33 in the calculation of all statistics to correct for their overrepresentation at the selection stage, there are actually more follow-up respondents than are reported in the weighted *Ns* given in the tables; and in recent years actual numbers average about 23% higher than the weighted numbers. The 2,400 participants selected from each 12th-grade class are randomly split into two groups of 1,200 each—one group to be surveyed on even-numbered calendar years in a series of biannual follow-up surveys, and the other group to be surveyed on odd-numbered years also in a series of biannual follow-up surveys. This two-year cycle is intended to reduce respondent burden, thus yielding better retention rates. By alternating the two half samples, MTF collects data from every graduating class each year (through age 30), even though any given respondent participates only every other year.

Until 2002, each respondent was surveyed biennially up to seven times; at the seventh follow-up, which would occur either 13 or 14 years after graduation, the respondents had reached modal age 31 or 32. In 2002, as a cost-saving measure, the seventh biennial follow-up was discontinued, and since then each respondent is surveyed every other year until modal age 29 or 30. Additional follow-ups then occur at modal ages 35, 40, 45, 50, and beginning in 2013, age 55. Data like these, gathered on representative national samples over such a large portion of the life span, are extremely rare and can provide needed insight into the etiology and life-course history of substance use and relevant behaviors, including those related to HIV transmission.

Follow-Up Procedures

Using information provided by 12th-grade respondents on a tear-off card (requesting the respondent's name, address, phone number(s), and recent email address), mail contact is maintained with the subset of people selected for inclusion in the follow up panels. Newsletters are sent to them each year, providing a short summary of results on a variety of survey topics. Name and address corrections are requested from both the U.S. Postal Service and the individual. Questionnaires are sent in the spring to each individual biennially through age 30, then at 5-year intervals. A check, made

payable to the respondent, is attached to the front of each questionnaire.²⁷ Reminder letters and postcards are sent at fixed intervals thereafter; telephone callers attempt to gather up-to-date location information for those respondents with whom we are trying to make contact; and, finally, those whom we can contact but who have not responded receive a prompting phone call from the Survey Research Center's phone interviewing facility in Ann Arbor, Michigan. If requested, a second copy of the questionnaire is sent. No questionnaire content is administered by phone. If a respondent asks not to be contacted further, that wish is honored.

Follow-Up Questionnaire Format

The questionnaires used in the follow-up surveys of 19- to 30-year-olds parallel those used in 12th grade. Many of the questions are the same (including the core section dealing with drug use), and respondents are consistently mailed the same version (or form) of the questionnaire that they first received in 12th grade, so that *changes over time* in their behaviors, attitudes, experiences, and so forth can be measured. Questions specific to high school status and experiences are dropped in the follow-ups, and questions relevant to post-high school status and experiences are added (mostly in the core section). The post-high school questions deal with issues such as college attendance, military service, civilian employment, marriage, and parenthood. In the study's early follow-ups (1975-1988), the sample size for a question appearing on a single form was one fifth of the total sample. A sixth form was introduced in 12th grade beginning with the class of 1989 and extended a year later to the follow-up surveys. Therefore, since 1990, a question appearing on a single form has been administered to one sixth of the total sample in the 19-30 age band. Single-form data from a single cohort are too small to make reliable estimates; therefore, in most cases where they are reported, single-form data from several adjacent cohorts are combined.

For the five-year surveys beginning at age 35, both half-samples from a class cohort are surveyed simultaneously and only one questionnaire form is used. Much of the questionnaire content is maintained but streamlined with a focus on the major family and work issues relevant to respondents ages 35 to 55; we have also added measures of substance use disorders and health outcomes.

²⁷ Until 1991, the follow-up checks were for \$5. After an experiment indicated that an increase was warranted, the check amount was raised to \$10 beginning with the class of 1992. The check amount was raised to \$20 in 2006, and to \$25 beginning in 2008.

REPRESENTATIVENESS AND SAMPLE ACCURACY

School Participation

Schools are invited to participate in the MTF study for a two-year period. For each school that declines to participate, a similar school (in terms of size, geographic area, urbanicity, etc.) is recruited as a replacement. In 2013, either an original school or a replacement school was obtained in 95% of the sample units. With very few exceptions, each school participating in the first year has agreed to participate in the second year as well. Figure 3-2 provides the year-specific school participation rates and the percentage of units filled since 1977. As shown in the figure, replacements for declining schools are obtained in the vast majority of cases.

Two questions are sometimes raised with respect to school participation rates: (a) Are participation rates sufficient to ensure the representativeness of the sample? (b) Does variation in participation rates over time contribute to changes in estimates of drug use?

With respect to participation rates ensuring that the sample is representative, the selection of a comparable replacement school, demographically close to the original school, occurs in practically all instances in which an original school refuses. This almost entirely removes problems of bias in region, urbanicity, and the like that might result from certain schools refusing to participate. Other potential biases could be more subtle, however. If, for example, it turned out that most schools with “drug problems” refused to participate, the sample would be seriously biased. And if any other single factor were dominant in most refusals, that reason for refusal might also suggest a source of serious bias. However, the reasons given for a school refusing to participate tend to be varied and are often a function of happenstance events specific to that particular year; only a very few schools, if any, object specifically to the drug-related survey content.

If it were the case that schools differed substantially in drug use, then which particular schools participated could have a greater effect on estimates of drug use. However, the great majority of variance in drug use lies *within* schools, not between schools.²⁸ For example, between 1991 and 2002, the between-schools variance for annual marijuana use was 4.0–5.3% of the total variance for each of the three grades; for inhalant use, 1.6–2.7%; for cocaine use, 1.2–2.2%; for alcohol use, 3.5–6.1%; and for cigarette use, 2.1–5.2%. To the extent that schools tend to be fairly similar in drug use, which particular schools participate (within a selection framework that seeks national representation) has a small effect on estimates of drug use.²⁹ Further, some,

²⁸ O'Malley, P. M., Johnston, L. D., Bachman, J. G., Schulenberg, J. E., & Kumar, R. (2006). How substance use differs among American secondary schools. *Prevention Science*, 7, 409–420.

²⁹ Among participating schools, there is very little difference in substance use rates between the schools that were original selections, taken as a set, and the schools that were replacements. Averaged over the years 1991 through 2000, for grades 8, 10, and 12 combined, the difference between original schools and replacement schools averaged 0.03 percentage points in the observed prevalence rates averaged across a number of drug use measures: two indexes of annual illicit drug use, the annual

if not most, of the between-schools variance is due to differences related to factors such as region and urbanicity, which remain well controlled in the present sampling design.

With respect to participation rates and changes in estimates of drug use, it is extremely unlikely that results have been significantly affected by changes in school participation rates. If changes in participation rates seriously affected prevalence estimates, there would be noticeable bumps up or down in concert with the changing rates. But this series of surveys produces results that are very smooth and generally change in an orderly fashion from one year to the next. Moreover, different substances trend in distinctly different ways. We have observed, for example, marijuana use decreasing while cocaine use was stable (in the early 1980s), alcohol use declining while cigarette use held steady (in the mid- to late 1980s), ecstasy use rising sharply while cocaine use showed some decline (late 1990s, early 2000s); and marijuana use continuing to rise while alcohol use hit historic lows (2011). All of these patterns are explainable in terms of psychological, social, and cultural factors and cannot be explained by the common factor of changes in school participation rates.

Of course, there could be some sort of constant bias across the years; but even in the unlikely event that there is, it seems highly improbable that it would be of much consequence for policy purposes, given that it would not affect trends and likely would have a very modest effect on prevalence rates. Thus we have a high degree of confidence that school refusal rates have not seriously biased the survey results.

Nevertheless, securing the cooperation of schools has become more difficult in recent years. This is a problem common to the field, not specific to MTF. Therefore, beginning with the 2003 survey, we have provided payment to schools as a means of increasing their incentive to participate. (By that time, several other ongoing school-based survey studies already were using payments to schools.)

At each grade level, half of each year's sample comprises schools that started their participation the previous year, and half comprises schools that began participating in the current year. (Both samples are national replicates, meaning that each is drawn to be nationally representative by itself.) This staggered half sample design is used to check on possible errors in the year to year trend estimates due to school turnover. For example, separate sets of one year trend estimates are computed based on students in the half sample of schools that participated in *both 2011 and 2012*, then based on the students in the half sample that participated in *both 2012 and 2013*, and so on. Thus, each one-year *matched half sample* trend estimate derived in this way is based on a constant set of schools (about 65 in 12th grade, for example, over a given one-year interval). When the trend data derived from the matched half sample (examined separately for each class of drugs) are compared with trends based on the

prevalence of each of the major illicit drug classes, and several measures of alcohol and cigarette use. For the individual drugs and drug indexes, the differences between the original and replacement schools, averaged across grades and years, fell within +0.9%.

total sample of schools, the results are usually highly similar, indicating that the trend estimates are affected little by school turnover or shifting participation rates. As would be expected, the *absolute* prevalences for a given year are not as accurately estimated using just the half sample because the sample size is only half as large.

Student Participation

In 2013, completed questionnaires were obtained from 90% of all sampled students in 8th grade, 88% in 10th grade, and 82% in 12th grade (see Table 3-1 for response rates in earlier years). In the large majority of cases, students are missed due to absence from class at the time of data collection; for reasons of cost efficiency, we typically do not schedule special follow up data collections for absent students. Because students with fairly high rates of absenteeism also report above-average rates of drug use, some degree of bias is introduced into the prevalence estimates by missing the absentees. Much of that bias could be corrected through the use of special weighting based on the reported absentee rates provided by the students who did respond; however, we decided not to use such a weighting procedure because the bias in overall drug use estimates was determined to be quite small *and* the necessary weighting procedures would have introduced greater sampling variance in the estimates.³⁰ Appendix A in this report illustrates the changes in trend and prevalence estimates that would result if corrections for absentees had been included. Of course, some students simply refuse, when asked, to complete a questionnaire. However, the proportion of explicit refusals amounts to less than 1.7% of the target sample for each grade.

Sampling Accuracy of the Estimates

Confidence intervals (95%) are provided in Tables 4-1a through 4-1d for lifetime, annual, 30-day, and daily prevalence of use for 8th-, 10th-, and 12th-grade students. As can be seen in Table 4-1a, confidence intervals for lifetime prevalence for 12th graders average less than $\pm 1.4\%$ across a variety of drug classes. That is, if we took a large number of samples of this size from the universe of all schools containing 12th graders in the coterminous United States, 95 times out of 100 the sample would yield a result that would be less than 1.4 percentage points divergent from the result we would get from a comparable massive survey of *all* 12th graders in *all* schools. This is a high level of sampling accuracy, permitting detection of fairly small changes from one year to the next. Confidence intervals for the other prevalence periods (last 12 months, last 30 days, and current daily use) are generally smaller than those for lifetime use. In general, confidence intervals for 8th and 10th graders are very similar to those observed for 12th graders. Some drugs (smokeless tobacco, PCP, and others, as indicated in the footnotes for Tables 2-1 to 2-4) are measured on only one or two questionnaire forms; these drugs will have somewhat larger confidence intervals because they are based on smaller sample sizes. Appendix C provides information on

³⁰ See Appendix A in the following publication for a discussion of this point: Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975–1983* (DHHS (ADM) 85 1374). Washington, DC: U.S. Government Printing Office.

how to calculate confidence intervals around other point estimates, as well as information needed to compare trends across time or to test the significance of differences between subgroups in any given year.

PANEL RETENTION

We discuss here the nature of the panel attrition problem generally, the response rates for MTF panel surveys in recent years, and evidence relevant to assessing the impact of attrition on the study's research results.

The Problem of Panel Attrition

Virtually all longitudinal studies of drug use experience attrition, which is often differential with respect to substance use.³¹ In addition, survey response rates in general have been declining over the past few decades,³² highlighting an important challenge in the conduct of population-based research.

A vital feature of the MTF panel studies is their very low cost per respondent. There are many advantages to collecting panel data through low-cost mail surveys, as we have done since the outset of the study. Indeed, given the number of questionnaires sent each year (roughly 18,500) across the entire coterminous United States and elsewhere in the world, using low-cost mail surveys has been our best (and really the only) cost-effective option. Now that internet use is widespread among young adults, we are conducting experiments within the ongoing surveys to test web-based survey options as a means to engage more respondents in the panel studies and reduce data collection costs.)

One disadvantage of data collection by mail is that attrition rates for mailed surveys tend to be higher than those for surveys obtained with much more expensive methods, such as intensive personal tracking and interviewing. Certainly there exist a few large epidemiological/etiological surveys that have better retention rates, but their procedures are extremely expensive and not realistic for an ongoing effort of the scale of MTF. Nevertheless, our retention rates compare favorably with those of most longitudinal studies (including interview studies) reported in the field.

Response Rates

The MTF survey data on American college students—an important subgroup in the panel surveys—now encompasses 34 years. We know about our respondents' actual college attendance only from those who are invited to and do complete follow-up

³¹ McGuigan, K. A., Ellickson, P. L., Hays, R. D., & Bell, R. M. (1997). Adjusting for attrition in school-based samples: Bias, precision, and cost trade-off of three methods. *Evaluation Review*, 21, 554–567.

³² Kim, J., Gershenson, C., Glaser, P., & Smith, T.W. (2011). The polls—trends: Trends in surveys on surveys. *Public Opinion Quarterly*, 75(1), 165-191; Groves, R.M. (2006). Nonresponse rates and nonresponse bias in household surveys. *Public Opinion Quarterly*, 70(5), 646-675.

questionnaires; however, we can use 12th-grade questionnaire answers (i.e., college intentions/expectations and program of study) to predict college attendance with a high degree of accuracy. MTF's retention of 12th graders who identified themselves as "college-bound" remains reasonably good. Among those participants in high school who were targeted for follow-up, *and* who reported planning to attend college and being enrolled in a college-prep curriculum, the follow-up retention rates for the three most recent classes surveyed at each follow-up point were: 51% in the first follow-up, one to two years past high school (based on the classes of 2011–2012); 50% in the second follow-up, three to four years past high school (based on the classes of 2009–2010); and 54% in the third follow-up, five to six years past high school (based on the classes of 2007–2008). These rates compare well with another national survey of substance use among college students, the Harvard College Alcohol Study, which had cross-sectional response rates of 59% in 1997 and 1999, but of 52% by 2001.³³ To date in *Volume II*, we have reported only on college students who are one to four years past high school graduation. As the average age of attendance rises, having the extended age coverage will be of growing importance.

Retention rates in the biennial follow-ups of *all* panel members modal ages 19–30 (corresponding to the first six follow-ups) decline with the length of the follow-up interval, of course. For the five surveys from 2009 to 2013, the response rate in the first follow-up (corresponding to one to two years past high school) averaged 50%; and for the second through sixth follow-ups (corresponding to 3–12 years past high school) response rates averaged 46%. Among long-term respondents—the 35-, 40-, 45-, and 50-year-olds—the retention rates are quite good, apparently because some of the decline with age in retention rates reflects cohort differences. Among the 35-year-old respondents surveyed from 2009 to 2013, corresponding to 17 years past high school, the average response rate was 44%. Among 40-year-old respondents surveyed from 2009 to 2013, corresponding to a 22-year follow-up interval, the average retention rate was 43%. Among 45-year-olds surveyed in 2009 to 2013, the average retention rate was 48%; among 50-year-olds surveyed in 2009 to 2013, the response rate averaged 55%; and for 55-year-olds surveyed for the first time in 2013, the response rate was 52%. In sum, the response rates attained under the current design range from respectable to good, especially when the low-cost nature of the procedures, the very long time intervals involved, and the substantial length of the questionnaires are taken into account. More importantly, the evidence leaves us confident that the data resulting from these follow-up panels are reasonably accurate, which brings us to our adjustments for panel attrition and the comparison of our results with those from other sources.

³³ Wechsler, H., Lee, J. E., Kuo, M., Seibring, M., Nelson, T. F., & Lee, H. (2002). Trends in college binge drinking during a period of increased prevention efforts: Findings from 4 Harvard School of Public Health College Alcohol Study surveys: 1993–2001. *Journal of American College Health, 50*, 203–217.

The Impact of Panel Attrition on Research Results

An important purpose of the MTF follow-ups is to allow estimation of drug prevalence among American high school graduates at various age levels. Thus, we have always been concerned about making the appropriate adjustments to account for panel attrition. In essence, our standard adjustment process is a poststratification procedure in which we reweight the data obtained from the participating follow-up samples so that their *12th-grade* distribution of answers on a given drug reproduces the original distribution of use observed for that drug, which was based on *all* participating 12th graders. This procedure is carried out separately for cigarettes, smokeless tobacco, alcohol, and marijuana, as well as other illicit drugs (combined). As expected, it produces prevalence estimates that are somewhat higher than those uncorrected for attrition, indicating that there is indeed some positive association between drug use and panel attrition. However, the adjustments are relatively modest, as documented next.

One reason the adjustments are modest is that attrition rates do not differ greatly by levels of 12th-grade substance use; they differ some, but less than one might expect. For example, among all respondents who had never used marijuana, an average of 79% of the classes of 1976–1998 participated in the first follow-up. The proportion responding is somewhat lower among those who had used marijuana just once or twice in the last 12 months: 75%. This proportion decreases gradually with increasing levels of marijuana use; but even among those who used marijuana on 20 or more occasions in the last 30 days in 12th grade, 67% participated in the first follow-up. The corresponding participation rates for the same drug use strata at the fourth follow-up (i.e., at modal ages 25–26) were 66%, 63%, and 56%, respectively. Thus, even among those who were quite heavy users of marijuana in high school, response rates at the fourth follow-up were only 10 percentage points lower than among those who had never used marijuana by 12th grade. That is not to say that we assume all types of drug users remain in the panels at comparably high rates. We believe that people who become dependent on or addicted to heroin or cocaine are unlikely to be retained in reasonable proportions. That is why we are careful not to quantify or characterize these special segments of the population. But we note that they constitute very low proportions of the drug-using portion of the population, and even lower proportions of the entire adult population.

The National Survey on Drug Use and Health (NSDUH) provides the best available data against which to validate the estimates generated for adult age groups in MTF, because it is also based on national samples but uses cross-sectional surveys that do not carry the burden of panel attrition. Their results, of course, may be affected by their own nonresponse rates; but that will be true of any comparison survey. The overall response rate for NSDUH in 2011 was 74%.

In some earlier analyses, we compared the prevalence rates on a set of drugs—cigarettes, alcohol, marijuana, and cocaine—for which there was reasonable similarity in question wording across the two studies. The comparisons that follow are for the age group 19–28 in the MTF panel data, and for 19–29 in the NSDUH

cross-sectional data (the closest age break reported by NSDUH). We used the most recent readily available comparable data at the time (2009), but similar results were found in a number of prior years. NSDUH would be expected to have higher rates than MTF because it includes school dropouts. In fact, however, the MTF estimates for 30-day marijuana and 12-month cocaine use, when the post-stratification weights are applied, are actually higher than the NSDUH estimates: 17.0% versus 15.8% for marijuana, and 5.2% versus 5.1% for cocaine. Even when the post-stratification weights are not applied, the MTF estimates are only slightly lower than the NSDUH estimates: 15.3% versus 15.8% for marijuana, and 4.8% versus 5.1% for cocaine. The fact that the MTF estimates for both marijuana and cocaine are similar to those observed in NSDUH suggests that attrition does not produce substantially lower estimates of drug use than would be obtained if response rates were higher—particularly after our poststratification adjustments are applied.

Comparisons for alcohol and cigarettes show larger differences, with alcohol use consistently higher in MTF and cigarette use consistently higher in NSDUH. We believe it likely that both are due to definitional differences in the exact question wording. In 2009, MTF estimate of 30-day alcohol prevalence was 69.1% (69.4% with poststratification) versus 65.9% in NSDUH. For cigarettes, the 30-day MTF prevalence estimate was 21.0% (23.3% with poststratification), versus 36.7% in NSDUH. (Because cigarette smoking rates are particularly high among dropouts, some of this difference should be explainable by differences in the populations covered by the two studies.) It is worth noting that the nature and magnitude of the differences between MTF and NSDUH estimates tend to be quite consistent for each of the four drugs at least as far back as 1992.

Even with attrition, substantial proportions of recent drug users remain in the MTF follow-up samples. In recent years, about 15–18% of the 19- to 28-year-old respondents reported marijuana use in just the prior 30 days, and about 4–7% reported cocaine use in the past 12 months. These proportions and the underlying numbers of actual cases are quite adequate for many analytic purposes.

A point worth emphasizing here is that, in the MTF panel, attrition is not as great a problem as is non-response in a cross-sectional study, because much is already known about each of the follow-up nonrespondents, including their substance use, based on extensive questionnaire responses in 12th grade (and, for many, in subsequent years as well). Thus, adjustments can be made utilizing data that are highly informative about the missing individuals.

Effects on Relational Analyses

While differential attrition (uncorrected) may contribute to some bias in point estimates and other univariate statistics, such attrition tends to have less influence on bivariate and multivariate statistics. This was found to be true in a secondary analyses of data from seven panel studies that followed adolescents over time,³⁴ and we have

³⁴ Cordray, S., & Polk, K. (1983). The implication of respondent loss in panel studies of deviant behavior. *Journal of Research in Crime and Delinquency*, 20, 214–242.

found this to be true in MTF panel analyses³⁵ and in analyses with other panel data sets.³⁶ Thus, differential attrition may be of less concern in multivariate panel analyses focused on understanding the course, causes, and consequences of substance use. Still, as we summarized above, correcting for attrition can be important, and we continue to do so using these and other correction procedures in our scientific publications (e.g., data imputation, FIML).

VALIDITY OF MEASURES OF SELF-REPORTED DRUG USE

Are sensitive behaviors such as drug use honestly reported? Like most studies dealing with sensitive behaviors, we have no direct, totally objective validation of the present measures; however, the considerable amount of existing inferential evidence strongly suggests that the MTF self report questions produce largely valid data. Here we briefly summarize this evidence.³⁷

First, using a three-wave panel design, we established that the various measures of self reported drug use have a high degree of reliability—a necessary condition for validity.³⁸ In essence, respondents were highly consistent in their self-reported behaviors over a three- to four-year time interval. Second, we found a high degree of consistency among logically related measures of use within the same questionnaire administration. Third, the proportion of 12th graders reporting some illicit drug use has reached two thirds of all respondents in peak years and over 80% in some follow up years, constituting *prima facie* evidence that the degree of underreporting must be very limited. Fourth, 12th graders' reports of use by their unnamed friends—about whom they would presumably have considerably less reason to conceal information about use—have been highly consistent with self reported use in the aggregate, in

³⁵ Bryant, A. L., Schulenberg, J. E., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (2000). Understanding the links among school misbehavior, academic achievement, and cigarette use: A national panel study of adolescents. *Prevention Science, 1*(2), 71–87; Schulenberg, J. E., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (1994). High school educational success and subsequent substance use: A panel analysis following adolescents into young adulthood. *Journal of Health and Social Behavior, 35*, 45–62.

³⁶ Bachman, J. G., O'Malley, P. M., & Johnston, J. (1978). *Youth in Transition: Vol. 6. Adolescence to adulthood: A study of change and stability in the lives of young men*. Ann Arbor, MI: Institute for Social Research; Schulenberg, J. E., Bryant, A. L., & O'Malley, P. M. (2004). Taking hold of some kind of life: How developmental tasks relate to trajectories of well-being during the transition to adulthood. *Development and Psychopathology, 16*, 1119–1140.

³⁷ A more complete discussion may be found in: Johnston, L. D., & O'Malley, P. M. (1985). Issues of validity and population coverage in student surveys of drug use. In B. A. Rouse, N. J. Kozel, & L. G. Richards (Eds.), *Self report methods of estimating drug use: Meeting current challenges to validity* (NIDA Research Monograph No. 57 (ADM) 85 1402). Washington, DC: U.S. Government Printing Office; Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975–1983* (DHHS (ADM) 85 1374). Washington, DC: U.S. Government Printing Office; Wallace, J. M., Jr., & Bachman, J. G. (1993). Validity of self-reports in student-based studies on minority populations: Issues and concerns. In M. de LaRosa (Ed.), *Drug abuse among minority youth: Advances in research and methodology* (NIDA Research Monograph No. 130). Rockville, MD: National Institute on Drug Abuse.

³⁸ O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self reports of drug use. *International Journal of the Addictions, 18*, 805–824.

terms of both prevalence *and* trends in prevalence, as discussed in chapter 9. Fifth, we have found self reported drug use to relate in consistent and expected ways based on theory to a number of other attitudes, behaviors, beliefs, and social situations—strong evidence of construct validity. Sixth, the missing data rates for the self reported use questions are only very slightly higher than for the preceding nonsensitive questions, in spite of explicit instructions to respondents immediately preceding the drug section to leave blank those questions they feel they cannot answer honestly. Seventh, an examination of consistency in reporting of lifetime use conducted on the long-term panels of graduating seniors found quite low levels of recanting of earlier reported use of the illegal drugs.³⁹ There was a higher level of recanting for the psychotherapeutic drugs, suggesting that adolescents may actually overestimate their use of some drugs because of misinformation about definitions, and this misinformation is corrected as they get older. Finally, the great majority of respondents, when asked, say they *would* answer such questions honestly if they were users.⁴⁰

As an additional step to assure the validity of the data, we check for logical inconsistencies in the answers to the triplet of questions about use of each drug (i.e., lifetime, annual, and 30-day use), and if a respondent exceeds a minimum number of inconsistencies across the set of drug use questions, his or her record is deleted from the data set. Similarly, we check for improbably high rates of use of multiple drugs and delete such cases, assuming that the respondents are not taking the task seriously. Fortunately, very few cases have to be eliminated for these reasons.

This is not to argue that self-reported measures of drug use are necessarily valid in all studies. In MTF we have gone to great lengths to create a situation and set of procedures in which respondents recognize that their confidentiality will be protected. We have also tried to present a convincing case as to why such research is needed. The evidence suggests that a high level of validity has been obtained. Nevertheless, insofar as any remaining reporting bias exists, we believe it to be in the direction of underreporting. Thus, with the possible exception of the psychotherapeutic drugs, we believe our estimates to be lower than their true values, even for the obtained samples, but not substantially so.

³⁹ Johnston, L. D., & O'Malley, P. M. (1997). The recanting of earlier reported drug use by young adults. In L. Harrison (Ed.), *The validity of self-reported drug use: Improving the accuracy of survey estimates* (NIDA Research Monograph No. 167, pp. 59–80). Rockville, MD: National Institute on Drug Abuse.

⁴⁰ For a discussion of reliability and validity of student self-report measures of drug use like those used in MTF across varied cultural settings, see Johnston, L. D., Driessen, F. M. H. M., & Kokkevi, A. (1994). *Surveying student drug misuse: A six-country pilot study*. Strasbourg, France: Council of Europe. Available at http://monitoringthefuture.org/pubs/monographs/surveying_student_drug_misuse_1994.pdf

Consistency and Measurement of Trends

MTF is designed to be sensitive to changes from one time period to another. A great strength of this study is that the measures and procedures have been standardized and applied consistently across many years. To the extent that any biases remain because of limits in school and/or student participation, and to the extent that there are distortions (lack of validity) in the responses of some students, it seems very likely that such problems will exist in much the same proportions from one year to the next. In other words, biases in the survey estimates will tend to be consistent from one year to another, meaning that our measurement of *trends* should be affected very little. The smooth and consistent nature of most trend curves reported for the various drugs provides rather compelling empirical support for this

TABLE 3-1
Sample Sizes and Response Rates

Grade:	Number of Public Schools			Number of Private Schools			Total Number of Schools				Total Number of Students				Student Response Rate (%)		
	8th	10th	12th	8th	10th	12th	8th	10th	12th	Total	8th	10th	12th	Total	8th	10th	12th
1975	—	—	111	—	—	14	—	—	125	—	—	—	15,791	—	—	—	78
1976	—	—	108	—	—	15	—	—	123	—	—	—	16,678	—	—	—	77
1977	—	—	108	—	—	16	—	—	124	—	—	—	18,436	—	—	—	79
1978	—	—	111	—	—	20	—	—	131	—	—	—	18,924	—	—	—	83
1979	—	—	111	—	—	20	—	—	131	—	—	—	16,662	—	—	—	82
1980	—	—	107	—	—	20	—	—	127	—	—	—	16,524	—	—	—	82
1981	—	—	109	—	—	19	—	—	128	—	—	—	18,267	—	—	—	81
1982	—	—	116	—	—	21	—	—	137	—	—	—	18,348	—	—	—	83
1983	—	—	112	—	—	22	—	—	134	—	—	—	16,947	—	—	—	84
1984	—	—	117	—	—	17	—	—	134	—	—	—	16,499	—	—	—	83
1985	—	—	115	—	—	17	—	—	132	—	—	—	16,502	—	—	—	84
1986	—	—	113	—	—	16	—	—	129	—	—	—	15,713	—	—	—	83
1987	—	—	117	—	—	18	—	—	135	—	—	—	16,843	—	—	—	84
1988	—	—	113	—	—	19	—	—	132	—	—	—	16,795	—	—	—	83
1989	—	—	111	—	—	22	—	—	133	—	—	—	17,142	—	—	—	86
1990	—	—	114	—	—	23	—	—	137	—	—	—	15,676	—	—	—	86
1991	131	107	117	31	14	19	162	121	136	419	17,844	14,996	15,483	48,323	90	87	83
1992	133	106	120	26	19	18	159	125	138	422	19,015	14,997	16,251	50,263	90	88	84
1993	126	111	121	30	17	18	156	128	139	423	18,820	15,516	16,763	51,099	90	86	84
1994	116	116	119	34	14	20	150	130	139	419	17,708	16,080	15,929	49,717	89	88	84
1995	118	117	120	34	22	24	152	139	144	435	17,929	17,285	15,876	51,090	89	87	84
1996	122	113	118	30	20	21	152	133	139	424	18,368	15,873	14,824	49,065	91	87	83
1997	125	113	125	27	18	21	152	131	146	429	19,066	15,778	15,963	50,807	89	86	83
1998	122	110	124	27	19	20	149	129	144	422	18,667	15,419	15,780	49,866	88	87	82
1999	120	117	124	30	23	19	150	140	143	433	17,287	13,885	14,056	45,228	87	85	83
2000	125	121	116	31	24	18	156	145	134	435	17,311	14,576	13,286	45,173	89	86	83
2001	125	117	117	28	20	17	153	137	134	424	16,756	14,286	13,304	44,346	90	88	82
2002	115	113	102	26	20	18	141	133	120	394	15,489	14,683	13,544	43,716	91	85	83
2003	117	109	103	24	20	19	141	129	122	392	17,023	16,244	15,200	48,467	89	88	83
2004	120	111	109	27	20	19	147	131	128	406	17,413	16,839	15,222	49,474	89	88	82
2005	119	107	108	27	20	21	146	127	129	402	17,258	16,711	15,378	49,347	90	88	82
2006	122	105	116	29	18	20	151	123	136	410	17,026	16,620	14,814	48,460	91	88	83
2007	119	103	111	32	17	21	151	120	132	403	16,495	16,398	15,132	48,025	91	88	81
2008	116	103	103	28	19	17	144	122	120	386	16,253	15,518	14,577	46,348	90	88	79
2009	119	102	106	26	17	19	145	119	125	389	15,509	16,320	14,268	46,097	88	89	82
2010	120	105	104	27	18	22	147	123	126	396	15,769	15,586	15,127	46,482	88	87	85
2011	117	105	110	28	21	19	145	126	129	400	16,496	15,382	14,855	46,733	91	86	83
2012	115	107	107	27	19	20	142	126	127	395	15,678	15,428	14,343	45,449	91	87	83
2013	116	103	106	27	17	20	143	120	126	389	15,233	13,262	13,180	41,675	90	88	82

Source. The Monitoring the Future study, the University of Michigan.


FIGURE 3-1
Schools included in 1 Year's Data Collection
8th, 10th, and 12th Grades



Source. The Monitoring the Future study, the University of Michigan.

Note. One dot equals one school.

FIGURE 3-2
School Participation Rates



Percent of slots

filled by...	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05	'06	'07	'08	'09	'10	'11	'12	'13
Original	59	63	62	63	71	71	66	72	67	66	72	71	68	70	59	55	60	53	52	53	51	51	57	62	56	49	53	62	63	59	58	53	54	58	56	53	54
Replacements	39	36	35	32	25	26	32	26	29	33	26	26	30	29	39	43	39	44	44	43	47	48	42	35	42	48	37	34	40	39	43	44	39	40	43	41	
Total	98	99	97	95	96	97	99	98	96	99	98	99	99	98	98	98	99	97	96	96	98	99	99	97	98	97	98	99	97	96	98	97	96	96	95		

Source: The Monitoring the Future study, the University of Michigan.

Chapter 4

PREVALENCE OF DRUG USE IN EARLY AND MIDDLE ADULTHOOD

Longitudinal panel studies, tracking the same individuals across years, are typically used to examine developmental changes with age, as is evident in many of our publications. At the same time, the multiple cohort feature of the MTF design provides a useful snapshot of each age group in a given year, showing the prevalence of use of various substances for each age group in that year, thus enabling us to compare these prevalence rates with those of the same age in earlier years. This chapter highlights such prevalence data for the age groups captured by MTF, starting right after high school and moving through middle adulthood. Each age group is defined by the modal age for its graduating high school class cohort.⁴¹ We will see that recent use tends to be higher in the early post-high school age groups, corresponding to the new freedoms associated with leaving high school and often moving away from the parental home.⁴² But sometimes there are also strong cohort effects that underlie differences among age groups at a given point in time; in this chapter we will see evidence of both age and cohort effects.

Estimates of drug use in the adult population are most often generated through household survey interviews of cross-sections of the general population. In the present study, our estimates come from self-completed mail questionnaires from respondents in the follow-up surveys. These are representative samples of previous classes of high school students who started their participation in MTF in their senior year. As described in more detail in chapter 3, MTF has conducted ongoing panel studies on representative samples from each graduating high school senior class beginning with the class of 1976. From each class, two matched subpanels of roughly 1,200 students each are randomly selected to comprise long-term follow-up panels—one of these two panels is surveyed every even-numbered year after graduation, and the other is surveyed every odd-numbered year, up through age 30, after which data collection occurs at five-year intervals. So, while each *cohort* participates every year up through age 30, each individual *respondent* participates only every other year. This alternating panel design was chosen to reduce the repetitiveness (and burden) of participating in the panel study while still allowing for full age coverage between 19 and 30. Thus, in a given year, the study includes respondents age 19-30 from one of

⁴¹High school seniors have a modal age (the most common age) of 18; therefore in a follow-up conducted 12 years later they would have a modal age of 30.

⁴²Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates; see also Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education-drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

the two panels from each of the last 12 senior classes previously participating in MTF.⁴³

In 2013, representative samples of the classes of 2001 through 2012—modal ages 19 to 30—were surveyed, using the standard young adult survey instruments. For brevity, we refer to this 19- through 30-year-old age group as “young adults” in this chapter.

To build on these important national panels of young adults, we extend the surveys into middle adulthood. The middle adulthood surveys are conducted at modal age 35 (that is, 17 years after high school graduation) and at five-year intervals thereafter. In each of these later follow-ups, the two half panels from the relevant graduating class are *both* surveyed, using a single questionnaire form instead of six forms. The content of the questionnaires is revised to some degree across age to be more relevant to the different developmental periods. The results of the 2013 follow-up surveys characterize the population of high school graduates of modal ages 19–30, 35, 40, 45, 50, and 55. The high school dropout segment, perhaps 11%–15% missing from the senior year surveys, is missing from all of the follow-up surveys as well. Thus, the results presented here are not necessarily generalizable to that segment of the population.

Figures 4-1 through 4-21 contain 2013 *prevalence* data by age, corresponding to respondents ages 19–30, as well as 35-, 40-, 45-, 50-, and 55-year-olds. For comparison purposes, data are also included for the 2013 high school senior class, listed as 18-year-olds. Figures provided in chapter 5 contain the *trend* data for each of these age groups derived from the repeated cross-sectional surveys, including 12th graders and high school graduates through age 50. In the figures in chapters 4 and 5, age groups spanning the young adult years have been paired into two-year intervals in order to increase the number of cases, and thus the precision, for each point estimate. The data for ages 35, 40, 45, 50, and 55 are, of necessity, based on a single age in each case. As indicated above, *both* half samples from a given class cohort are included in each year’s samples of 35-, 40-, 45-, 50-, and 55-year-olds. In 2013 the paired half samples come from the high school graduating classes of 1996, 1991, 1986, 1981, and 1976, respectively. The respective weighted numbers of cases are 844, 833, 838, 828, and 870. (Actual unweighted numbers are somewhat higher, because those from the oversampled drug-using stratum in high school are counted as only one-third of a case in the weighted data.)

⁴³Through 2001, the follow-ups also included modal ages 31 and 32. This seventh follow-up was dropped in 2002 because we believed that the costs were no longer justified by the marginal benefits of having these follow-up data, given that an age-35 survey was being conducted. Throughout the time between surveys, we send annual newsletters to respondents in order to help maintain contact.

REPLICABILITY OF FINDINGS

It is worth noting that the pattern of age-related differences showing up in any one year can be checked in an adjacent year (i.e., the previous year's volume or the succeeding year's) for replicability, because two nonoverlapping half samples of follow-up respondents in the 19-to-30 age band are surveyed on alternating years. In the case of the 35-, 40-, 45-, 50-, and 55-year-olds, two entirely different graduating classes make up the samples for any two adjacent, chronological years of the survey results.

A NOTE ON ADJUSTED LIFETIME PREVALENCE ESTIMATES

In Figures 4-1 through 4-21, two different estimates of *lifetime* prevalence are provided. One estimate is based on the respondent's *most recent* (i.e., 2013) statement of whether he or she ever used the drug in question (the light gray bar). The other estimate takes into account the respondent's answers regarding lifetime use gathered in *all* of the previous data collections in which he or she participated (the white bar). *To be categorized as one who has used the drug based on all past answers regarding that drug, the respondent must have reported either lifetime use in the most recent data collection and/or reported some use in his or her lifetime on at least two earlier data collections.* Because respondents of ages 18 through 20 cannot have their responses adjusted on the basis of two earlier data collections, adjusted prevalence rates are reported only for ages 21 and up. Most other epidemiological studies can present only an unadjusted estimate because they have data from a single cross-sectional survey. An adjusted estimate of the type used here is possible only when panel data have been gathered so that a respondent can be classified as having used a drug at some time in his or her life, based on earlier answers, even though he or she no longer indicates lifetime use in the most recent survey.

The divergence of these two estimates as a function of age shows that there is more inconsistency as time passes. Obviously, there is more opportunity for inconsistency as the number of data collections increases. Our judgment is that the truth lies somewhere between the two estimates: the lower estimate may be depressed by tendencies to forget, forgive, or conceal earlier use, and the upper estimate may include earlier response errors or incorrect definitions of drugs that respondents appropriately corrected in later surveys as they became more knowledgeable. It should be noted that a fair proportion of those giving inconsistent answers across time had earlier reported having used the given drug only once or twice in their lifetime.

As we have reported elsewhere, the cross-time stability of self-reported usage measures, taking into account both prevalence and frequency of self-reported use, is still very high.⁴⁴ Note that the divergence between the two lifetime prevalence

⁴⁴O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self-reports of drug use. *International Journal of the Addictions, 18*, 805-824.

estimates is greatest for the psychotherapeutic drugs and for the derivative index of “use of an illicit drug other than marijuana,” which is heavily affected by the estimates of psychotherapeutic drug use. We believe this is due to respondents having greater difficulty accurately categorizing psychotherapeutic drugs (usually taken in pill form) with a high degree of certainty—especially if such a drug was used only once or twice. We expect higher inconsistency across time when the event—and in many of these cases, a single event—is reported with a relatively low degree of certainty at quite different points in time. Those who have gone beyond simple experimentation with one of these drugs would undoubtedly be able to categorize them with a higher degree of certainty. Also, those who have experimented more recently, in the past month or year, should have a higher probability of recall, as well as fresher information for accurately categorizing the drug.

We provide both estimates to make clear that a full use of respondent information provides a possible range for lifetime prevalence estimates, not a single point. However, by far the most important use of the prevalence data is to track *trends* in *current* (as opposed to lifetime) use. Thus, we are much less concerned about the nature of the variability in the lifetime estimates than we might otherwise be. The lifetime prevalence estimates are of importance primarily in showing the degree to which a drug class has penetrated the general population overall as well as particular cohorts; we believe that the evidence from the lifetime estimates suggests that cross-sectional surveys of adults are subject to underreporting, and that to a degree such underreporting increases with age, because adolescence and early adulthood are the periods in the life course during which most drug use occurs.⁴⁵

The reweighting procedures used to adjust the panel data for the effects of panel attrition are described in chapter 3.

PREVALENCE OF DRUG USE ACROSS AGE GROUPS⁴⁶

Figures 4-1 through 4-21 provide 2013 prevalence rates for each class of drugs, covering respondents ages 18 to 55. For virtually all drugs, available age comparisons show much higher *lifetime* prevalence for the older age groups, as would be expected.

⁴⁵For a more detailed analysis and discussion, see Johnston, L. D., & O’Malley, P. M. (1997). The recanting of earlier-reported drug use by young adults. In L. Harrison & A. Hughes (Eds.), *The validity of self-reported drug use: Improving the accuracy of survey estimates* (NIDA Research Monograph No. 97-4147). Washington, DC: National Institute on Drug Abuse. Available at http://archives.drugabuse.gov/pdf/monographs/monograph167/059-080_Johnston.pdf

⁴⁶This section discusses differences in the current year as a function of age, but it should be noted that these age differences are confounded with cohort differences. Thus, although the discussion is accurate with respect to age differences, it is not necessarily the case that the age differences would be similar in other time periods. In fact, our recent evidence, including many findings provided in Chapter 5, suggests similarities and differences by age across cohorts. See Jager, J., Schulenberg, J. E., O’Malley, P. M., & Bachman, J. G. (2013). Historical variation in drug use trajectories across the transition to adulthood: The trend toward lower intercepts and steeper, ascending slopes. *Development and Psychopathology*, 25(2), 527-543. doi:10.1017/S0954579412001228.

In fact, the figures reach surprisingly high levels among adults in their early 30s through their 50s.

- The *adjusted lifetime prevalence* figures are most striking for today's 50-year-olds (the high school class of 1981), who were passing through adolescence near the peak of the drug epidemic. Some 88% reported trying *an illicit drug* (lifetime prevalence, adjusted), leaving only 12%, or about one in every eight, who reported never having done so (Figure 4-1). Four out of five 50-year-olds (81%) said they had tried *marijuana*, and nearly three quarters (72%) said they had tried some *other illicit drug*, including 46% who had tried *cocaine* specifically. The adjusted and unadjusted lifetime prevalence figures for the 55-year-olds are similar to those for the 50-year-olds. Clearly, the parents of today's teenagers and young adults are themselves a very drug-experienced generation which, among other things, may help to explain the acceptance of medical marijuana in a number of states and full legalization of adult marijuana use in two states so far.
- In 2013, the adjusted lifetime prevalence figures among 29- to 30-year-olds reached 71% for *any illicit drug*, 66% for *marijuana*, 47% for *any illicit drug other than marijuana*, and 18% for *cocaine*. Put another way, even among young Americans who graduated from high school in 2001 and 2002—after the peak of the larger drug epidemic, but near the peak of the relapse phase in the epidemic—fewer than one third (29%) *never* tried an illegal drug.
- The 2013 survey responses, *unadjusted* for previous answers, show somewhat lower lifetime prevalence for 29- to 30-year-olds: 65% for *any illicit drug*, 62% for *marijuana*, 39% for *any illicit drug other than marijuana*, and 17% for *cocaine*.
- Despite the higher lifetime prevalence rates among older age groups, these groups generally show *annual* or *30-day* prevalence rates that are no higher than those of today's 12th graders. Indeed, for a number of drugs, the levels reported by older respondents are lower—sometimes considerably lower—suggesting that the incidence of quitting more than offsets the incidence of initiating use of these drugs during the years after high school.

In analyses published elsewhere, we looked closely at patterns of change in drug use with age and identified post high school experiences that contribute to declining levels of annual or current use of drugs as respondents grow older. For example, the likelihood of marriage increases with age, and we have found that marriage is consistently associated with declines in *alcohol* use, *heavy drinking*, *marijuana* use, and *cocaine* use, and most likely just about all of the other illicit drugs as well.⁴⁷

⁴⁷Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates; and Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002).

- For use of **any illicit drug** (Figure 4-1), 2013 *lifetime* prevalence (unadjusted) is about 66-67% (64% to 70%) among 25- to 40-year-olds versus 50% among 12th graders. *Annual* prevalence is highest among 21- to 22-year-olds in 2013(41%), and lowest among the older age groups, reaching 27% among 29- to 30-year-olds and 17% among 50- and 55-year-olds. Current (30-day) prevalence shows the rate generally declining with each age band, from 25% among 12th graders to 16% among 29- to 30-year-olds. For the age groups above age 30, current use is not much below that of 29- to 30-year-olds, ranging from 10% to 14%.
- *Lifetime* prevalence rates for **marijuana** generally are higher with each increase in age level, except for a dip among 35- and 40-year-olds, from the classes of 1996 and 1991, respectively, who graduated from high school at relatively low points in the recent history of teenage marijuana use. (This pattern is true for several other illicit drugs.) However, *annual* and *30-day* prevalence rates for marijuana generally decline from ages 22 to 45, and then remain fairly level through age 55. Thus, it is clear that greater proportions of the older cohorts have discontinued use (Figures 4-1, 4-2, and 4-3). The annual and 30-day prevalence rates were fairly equivalent for 18-, 19- to 20-, and 21- to 22-year-olds in 2013.
- Current **daily marijuana** use shows considerable variation across age (as shown in Figure 5-3c in chapter 5). In 2013 prevalence ranges from 5.8% – 7.8% for those respondents 18 to 26 years old to about 2–3% for the age groups from 40 to 55 years old. (Clearly, most respondents who were daily users at some point in their teenage years are no longer daily users.)
- **Synthetic marijuana** refers to a set of substances that contain synthetic cannabinoids that are meant to mimic the effects of cannabinoids found in natural marijuana; synthetic cannabinoids are created artificially and typically sprayed on herbal and plant material, which is then smoked. These substances have been sold over-the counter in head shops, gas stations, and other venues under various brand names like “spice” and “K-2.” In 2011 the Drug Enforcement Administration scheduled five of the chemicals most commonly used, but chemists often are able to adjust the chemical composition just enough to evade the prohibitions. The American Association of Poison Control Centers has reported a substantial increase since 2010 in calls about

The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs. Mahwah, NJ: Lawrence Erlbaum Associates. See also Schulenberg, J. E., O’Malley, P. M., Bachman, J. G., & Johnston, L. D. (2000). “Spread your wings and fly”: The course of well-being and substance use during the transition to young adulthood. In L. J. Crockett & R. K. Silbereisen (Eds.), *Negotiating adolescence in times of social change* (pp. 224–255). New York: Cambridge University Press. And see O’Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2004). Studying the transition from youth to adulthood: Impacts on substance use and abuse. In J. S. House, F. T. Juster, R. L. Kahn, H. Schuman, & E. Singer (Eds.), *A telescope on society: Survey research and social science at the University of Michigan and beyond* (pp. 305–329). Ann Arbor, MI: The University of Michigan Press.

adverse reactions to synthetic marijuana.⁴⁸ A question about the use of these products was added to both the MTF in-school and follow-up surveys in 2011. Some 7.9% of the 2013 seniors report using synthetic marijuana in the prior 12 months (down significantly from 11.3% in 2012). Among all young adults ages 19 to 30 years, the annual prevalence is 3.0% (also down significantly from 4.8% in 2012), with little difference and no identifiable pattern related to age (Table 4-3).

- Another important class of synthetic drugs called *bath salts* was added to the MTF questionnaires in 2012. Fortunately, the rates of use of these dangerous over-the-counter stimulants, which are intended to mimic the effects of amphetamines, are quite low at this point. In 2013, the high school seniors had an annual prevalence rate of 0.9%, and all of the other age groups had rates of 0.5% or less with the single exception of 23- to 24-year-olds (1.1%). Data from the national poison control centers suggest that use of these drugs peaked prior to 2012, quite possibly as the result of heavy media coverage of their adverse effects.
- Adjusted rates for lifetime use of *any illicit drug other than marijuana* (Figure 4-2) show an appreciable rise with age, reaching 47% among the 29- to 30-year-old age group and 74% among 55-year-olds. In other words, about three quarters of all 55-year-olds have tried some illicit drug other than marijuana, and close to half of today's 30-year-olds have done so. *Current (past 30-day)* use shows a slight rise and then a decline across age bands; rates range from 8% among 12th graders to a high of 11% among 21- to 22-year-olds, and then down to 4-5% among the age groups above 35. After ages 21-22, *annual* use is generally lower with increased age of the respondent. A number of the individual drugs that comprise this general category show lower rates of use at higher ages for annual prevalence, usually with the highest rate observed at ages 18-22. This is particularly true for *amphetamines, hallucinogens, LSD* specifically, *hallucinogens other than LSD*, and *inhalants*. The falloff with age is not as great nor as consistent for *cocaine, crack, other cocaine, crystal methamphetamine (ice), heroin, narcotics other than heroin, sedatives (barbiturates),* and *tranquilizers*, though in general usage rates are somewhat lower among those in their 30s than among those in their early 20s. Several of these classes of drugs are discussed individually next.
- *Inhalants* show some very interesting differences across age strata (Figure 4-13). There is a modest difference across age in contemporaneously reported lifetime prevalence, but a considerable difference in the lifetime prevalence figure *adjusted* for previous reporting of use. The adjusted pattern—an increase with age—is the one we have come to expect, and we believe is the

⁴⁸American Association of Poison Control Centers, Synthetic marijuana data updated April 16, 2012. <http://www.aapcc.org/alerts/synthetic-marijuana/>.

more accurate one. *Annual* prevalence rates drop off with age, while *30-day* rates already are quite low by 12th grade and can drop only a little. Clearly, current use of inhalants is almost absent beyond about age 20, and we know from data presented in *Volume I* that much of the decline in use with age has already occurred by the time young people have reached 10th grade. Questions on inhalant use are not included in the surveys administered to respondents over the age of 30, given the negligible rates of current use reported for those over age 24.

- For *amphetamines* used without a doctor's orders, lifetime prevalence is much higher among the older age groups—reflecting in part the addition of new users who initiate use in their 20s, but also reflecting some cohort differences (Figure 4-4). As is true for most psychotherapeutic drugs, corrected lifetime prevalence and contemporaneously reported lifetime prevalence diverge considerably. However, more recent use, as reflected in the annual prevalence figure, is considerably lower among the older age groups. This has not always been true; the present pattern reflects a sharper historic decline in use among older respondents than has occurred among 12th graders, as well as cohort differences in having ever used these drugs. These trends are discussed in the next chapter.
- *Ritalin*, an amphetamine widely prescribed for the treatment of attention deficit hyperactivity disorder or ADHD, shows a considerable drop-off in annual prevalence of non-medical use from 3.4% at ages 19–20 to 0.3% at ages 29–30.
- *Adderall*, a similar and newer drug, shows a higher annual prevalence of non-medical use in 2013, compared to Ritalin, and also drops off in use with age, from 9–10% among 19- to 22-year-olds to 3.0% among 29- to 30-year-olds. The higher rates of use among those in their early 20s are consistent with the interpretation that initially Ritalin and perhaps now Adderall are used by college students attempting to enhance their academic performance (Table 4-3). Respondents over age 30 are not asked about Ritalin and Adderall use.
- Questions on the use of *methamphetamine* are contained in only two of the six questionnaire forms for young adults, so estimates are less reliable than those based on all six forms. Unadjusted lifetime use increases with age, from 1.6% for 19- to 20-year-olds to 6.0% for 27- to 28-year-olds, and adjusted figures are only slightly higher. This suggests that much initiation of methamphetamine use occurs after high school, though more recent graduating cohorts have been reporting considerably lower levels of use. *Annual* prevalence does not vary with age, however, remaining at 0–1% from ages 18–30 in this population of high school graduates (Table 4-3 and Figure 4-5.) Respondents over age 30 are not asked about methamphetamine use.
- *Crystal methamphetamine (ice)* is also included on only two questionnaire forms through age 30 and is not asked of older respondents. Among the 19- to 30-year-old respondents *combined*, only 0.9% now report any use in the prior

year—slightly lower than the 1.1% reported by 12th graders (Table 4-3 and Figure 4-6).

- **Sedatives (barbiturates)** show lifetime prevalence rates for non-medically prescribed use that rise with age stratum, but are roughly similar across the age band 25 through 40; they then continue to rise with each age band thereafter. This is consistent with the sharp falloff in sedative use among 12th graders in the early years of the study, giving rise to a cohort effect. Annual use is fairly similar across ages 19 to 40 at about 3-4%, then decreases slightly in the older age bands to 2% (Figure 4-14). At present, past 30-day usage rates are quite low at all ages; it is highest for age 18 at 2% and declines to 1% in nearly all older age bands. Because of the substantial long-term decline in sedative (barbiturate) use over the life of MTF, the 55-year-olds have, by far, the highest adjusted lifetime prevalence rate (34%); but they are not any more likely to be currently using than the younger age groups.⁴⁹
- Annual prevalence of non-medically prescribed use of **narcotics other than heroin** is highest in the 18–35 age range (at 6–7%), compared to 2–3% among 45-, 50-, and 55-year-olds. *Thirty-day* prevalence shows a somewhat similar profile, with rates at 2–3% up through age 35 and then 12% in all age strata thereafter. Figure 4-15 shows the cohort effects of the upturn in the use of narcotic drugs in recent years, with those ages 27–30 having higher lifetime prevalence rates than 35- and 40-year-olds, at least up through 2012.
- Similarly, non-medically prescribed use of **tranquilizers** shows a general increase with age in lifetime prevalence although it dips some among those ages 35 and 40, again reflecting the increased use among adolescents in the 1990s. The highest lifetime rates are found among those over 40, with the oldest cohorts having much higher levels. Despite that fact, there is some modest decrease with age in annual prevalence after age 30. *Thirty-day* prevalence is 1–2% across nearly all ages (Figure 4-16).
- **Cocaine** had generally presented a unique case among the illicit drugs, in that lifetime, annual, and current prevalence rates have *all* tended to rise with age into the 20s. By 1994, however, 30-day cocaine use had reached such low levels that it varied rather little by age. Following the resurgence of cocaine use in the 1990s, some differences by age in annual prevalence emerged, though there are still rather few differences for current prevalence (Figure 4-7). Annual prevalence is now highest among respondents ages 18 through 35 (at 3–5%) and falls off to 1–2% in the age groups beyond 35. The cohort differences in lifetime cocaine use are particularly vivid, with the 50- and 55-

⁴⁹Barbiturates were the dominant form of sedatives in use when these questions were first introduced. In the intervening years, a number of nonbarbiturate sedatives have entered the market and largely displaced barbiturates. We believe that a number of users of nonbarbiturate sedatives are reporting them in answer to this question, which also defines them in terms of the conditions for which they are prescribed. In recognition of this fact, we now label them as “sedatives (barbiturates).” The rewording of the question was made in half of the questionnaire forms in 2004 and in the other half in 2005.

year-olds both showing 46% adjusted lifetime prevalence rates in 2013, respectively, compared to 10% among 21- to 22-year-olds. Very few (1.1% or less) of the 35- to 55-year-olds today are current users of cocaine, despite the fact that so many of them have used it at least once in their lifetime. Among 55-year olds, 46% used cocaine at some time in their life, but only 0.5% reported using it in the past month.

- In 2013, lifetime prevalence of **crack** use (Figure 4-8) rises fairly steadily between ages 19 to 20 (1%) and ages 25 to 26 (4%, unadjusted); it is considerably higher among 50- and 55-year-olds (11%, unadjusted), reflecting something of a cohort effect due to the rather transient popularity of crack in the early to mid-1980s, and a brief resurgence in the mid-1990s. Current (30-day) prevalence is 0.6% or below in all of these age groups. Annual prevalence is highest among 18-year-olds at 1.1% and between 0.0% and 0.7% for all other age groups. We believe that the omission of high school dropouts is likely to have a greater than average impact on the prevalence estimates for crack. It also seems likely that any panel respondents who become dependent on crack (or other illicit drugs like heroin) would be less likely than average to respond to the questionnaires; therefore, such extreme users are no doubt underrepresented among the panel respondents.
- **Ecstasy (MDMA)** was added to two of the six follow-up questionnaire forms in 1989 to assess how widespread its use had become among young adults. It was added to a third form in 2002 (Figure 4-17). Questions about its use were not asked of high school students until 1996, primarily because of concern that its alluring name might stimulate interest in the drug. That concern diminished considerably after the drug had become more widely known. Ecstasy use is not asked of respondents 35 years of age and older.

In 2013, among all 19- to 30-year-olds combined, 12% said they have tried ecstasy, compared to 7.1% of the 12th graders. The age differences in *lifetime* ecstasy use are quite dramatic, with adjusted prevalence increasing sharply with age stratum and showing the highest rate (16%) at ages 27–30 (Figure 4-17). This very likely reflects the rapid rise in ecstasy use among high school seniors between 1997 and 2001, and the subsequent sharp decline. Because *annual* prevalence is highest among 21- to 22-year-olds (at 6%) and lowest among 29- to 30-year-olds (at 2%), there clearly has been a high degree of non-continuation among those in their late 20s. *Past-month* ecstasy use is now at 1% or less for all age groups between 19 and 30, except for the 23- to 24-year-olds, who are at 2%.

- A question about the use of **salvia** was introduced into one questionnaire form in 2009 as a single tripwire question asking only the frequency of use in the past year (Table 4-3). Salvia is not currently regulated by the federal government; however, as of 2012, about 25 states had made the drug illegal. Salvia has some mild hallucinogenic properties, and there has been considerable media attention to its potential for harm. Annual prevalence for ages 19 through 30 stands at 0.8% in 2013. It appears that salvia is slightly

more popular among the younger of the young adults, with annual prevalence rates in 2013 between 1% and 2% among 19- to 22-year-olds versus less than 1.0% (0.3–0.7%) among those ages 25–30.

- *Note: The question about Provigil was dropped in 2012.* Another drug introduced for the first time in the 2009 survey was **Provigil**, a prescription stay-awake drug. It appears that this drug had not made serious inroads in the young adult population as of 2011, because annual prevalence, as determined by a single tripwire question, was only 0.0–0.4% among those ages 19–30, and showed no clear relationship to age. As a result, the question about Provigil use was dropped in 2012.
- All **alcohol** prevalence rates are higher among young adults than among 12th graders, and they generally increase for the first three to six years after high school, through age 25 or 26 (Figures 4-20a and 4-20b). Prevalence rates vary only modestly among the older age groups. *Lifetime* prevalence changes very little after ages 25 to 26, due in large part to a “ceiling effect.” *Current* 30-day use rises from 39% among 18-year olds to a peak of 76% among 25-26 year olds, then declines slowly through age 50, down to 52% among 55-year-olds. *Current daily drinking* (Figure 4-20b) increases from 2% among 18-year-olds to 7% among 25- to 26-year-olds, holds at about 7% across the age span 25 to 35, is slightly higher among those in their forties, and is highest among 50- and 55-year olds (11%). Among the various measures of alcohol consumption, **occasions of heavy drinking** (i.e., having five or more drinks in a row on at least one occasion in the two weeks prior to the survey, sometimes called “binge drinking”) show some considerable differences by age (Figure 4-20b). There is a large difference between 18-year-olds (22%) and 21- to 24-year-olds, who have the highest prevalence of such heavy drinking (38–40%). There is a falloff at each subsequent age level above age 22, dropping to 31% among 29- to 30-year-olds and down to 17% by age 55. We have interpreted this curvilinear relationship as reflecting an age effect—not a cohort effect—because it seems to replicate across different graduating class cohorts, and also because it has been linked directly to age-related events such as leaving the parental home (which increases heavy drinking) and marriage (which decreases it), both of which are, in turn, related to attending college.⁵⁰ Clearly, binge drinking is most popular among people in their twenties and falls off after that. Still, substantial fractions of older age groups engage in such drinking (24% at ages 35 and 40 and 17% at age 55).

Extreme binge drinking is a concept that was introduced into the MTF measurement package in 2005. Two measures are used; drinking *10 or more drinks* on one or more occasions in the prior two weeks and drinking *15 or*

⁵⁰O’Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321. See also Bachman, J. G., Wadsworth, K. N., O’Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

more drinks on one or more occasions in the prior two weeks. Because these two measures have been included in only one of the six questionnaire forms used with young adults, the numbers of cases are very limited, ranging from 123 to 186 weighted cases per year for each two-year age band from 19 to 30. Therefore, we have combined young adult data from years 2005 through 2013 in order to generate more reliable estimates of these behaviors.⁵¹ The prevalence rate among all young adults of having 10 or more drinks on at least one occasion in the two weeks prior to the survey was 11.4% across the years 2005 to 2013 *combined* or roughly one in every eight to nine respondents. The comparable prevalence for having 15 or more drinks was 4.3% or roughly one in twenty-three respondents. As with binge drinking at the 5+ drink level, there are differences across age groups among young adults in their extreme binge drinking rates. Both measures of extreme binge drinking are highest among those 21- to 22-years of age (14.4% for 10+ drinks and 6.1% for 15+ drinks): they decline with each higher age band (reaching 9.0% and 2.5%, respectively by ages 29 to 30.)

- **Cigarette smoking** shows an unusual pattern of age-related differences, influenced to some extent by cohort differences (Figure 4-21). **Current (30-day) smoking** rates used to be about the same for 12th graders and those in their early 20s, partly because most initiation of cigarette use happens in high school. Beginning in 2005, however, current smoking was somewhat lower among 12th graders, almost surely due to the sharp drop in smoking that has been occurring among secondary school students—a cohort effect evident as early as 8th grade. Smoking at heavier levels such as a half pack daily is (and has been) higher among those in their 20s than among 12th graders, as many light or moderate smokers in high school transition into a pattern of heavier use after high school.⁵² At later ages, a rising proportion of current smokers—that is, those reporting any smoking in the past 30 days—also report smoking at the daily and half-pack-a-day or more rates. Through age 26 a good proportion (about 60%) of those indicating any smoking in the prior year were not *daily* smokers; the proportion then declines with age, so that by age 55 only about a quarter of those who smoked in the prior year were not daily smokers.

The prevalence of smoking a *half pack or more* of cigarettes per day rises from 3% among 18-year olds to 8% by ages 27–28 and then increases slightly to 9–11% among those 35 and older. The proportion of current smokers who smoke a half pack or more per day also rises with age: about one fifth among

⁵¹The weighted combined sample of 18-30 year-olds for 2005 through 2013 has 8,418 observations, with some individuals contributing more than one observation because of their continued participation in the panel study.

⁵²Because age is confounded with class cohort, and because we have established that cigarette smoking shows strong cohort effects (enduring differences among cohorts), one must be careful in interpreting age-related differences in a cross-sectional sample as if they were due only to age effects—that is, changes with age consistently observable across cohorts. However, multivariate analyses conducted on MTF panel data from multiple cohorts do show a consistent age effect of the type mentioned here (see O'Malley, Bachman, & Johnston, 1988, in previous footnote).

18 year olds (3% half pack or more, 16% 30-day smokers), about two fifths among 29-30 year olds (7% half pack or more, 17% 30-day smokers), and about three fourths among 55-year olds (11% half pack or more, 14% 30-day smokers).

- Past 30-day prevalence of *smokeless tobacco* use stands at 7.7% among all young adults (most of it by males, as will be discussed below). Daily prevalence is 2.8% (Tables 4-4 and 4-5).
- Questions have been added recently on the consumption of tobacco in various specific forms other than cigarettes. Tripwire questions were added for these forms of tobacco use in 2011, providing only annual prevalence data (Table 4-3). Past year prevalence of use in 2013 among 19- to 30-year-olds is 19% for using a *hookah to smoke tobacco*, 18% for smoking *small cigars*, 5% for using *snus*, and only 0.3% for using *dissolvable tobacco*. Among young adults, hookah smoking is highest among 19- to 20-year-olds and declines steadily to 11% at ages 27 to 30. Annual prevalence of smoking small cigars is also highest among 19- to 20-year-olds and is lower at increasing ages, dropping to 14% in the 27- to 30-year-old age bands. Annual prevalence of use of snus is at 5.1% among 19–30 year olds and does not vary much by age. Annual prevalence of dissolvable tobacco use is only 0.3% among 19- to 30-year-olds, with all age groups at 1% or less.
- Questions on *steroid* use (Figure 4-18) were added to one questionnaire form in 1989 and to an additional form in 1990, making it difficult to determine age-related differences with much accuracy due to limited sample sizes. Overall, 1.3% of all 19- to 30-year-olds in 2013 report having used steroids in their lifetime and 0.4% in the prior 12 months. Questions about steroid use are not asked of respondents over age 30.

In sum, lifetime prevalence rates in some of the older age groups studied here, who passed through adolescence in the heyday of the drug epidemic, show remarkably high lifetime rates of illicit drug use—particularly when lifetime prevalence is corrected for the recanting of previously reported use. However, *current* use of most illicit drugs is substantially lower among those over age 30 than among those in their late teens to early 20s. For the two licit drugs, alcohol and cigarettes, the picture is different, with less falloff in active use with age and higher levels of daily alcohol use in the older ages.

PREVALENCE COMPARISONS FOR SUBGROUPS OF YOUNG ADULTS

Subgroup differences for 19- to 30-year-olds are presented in Tables 4-1 through 4-5. While Table 4-1 provides only gender differences, the remaining tables have prevalence rates by gender, age, region of the country, and population density. Each of these subgroup dimensions is discussed separately below.

Gender Differences

In general, most of the gender differences in drug use that were observed in high school students may be found in the young adult (19- to 30-year-old) sample as well. See Tables 4-3 for the full set of gender comparisons.

- Among the full young adult sample ages 19 to 30, more males than females report past-year use of *any illicit drug* (38% vs. 33%), *marijuana* (35% vs. 28%), and *any illicit drug other than marijuana* (20% vs. 16%). Males have higher annual prevalence rates for nearly all illicit drugs—with ratios of two times greater or more for *synthetic marijuana, inhalants, hallucinogens, LSD, hallucinogens other than LSD, crack, powder cocaine, heroin with a needle, heroin without a needle, GHB, ketamine, bath salts, and steroids* (Table 4-3).
- All three measures of *cocaine* use showed higher rates of use by male than female 19- to 30-year-olds. Annual *cocaine* use is reported by 5.3% of males and 2.8% of females, *powder cocaine* use by 5.2% of males and 2.6% of females, and *crack* use by 0.5% of males and 0.2% of females.
- Other large gender differences among 19- to 30-year-olds are found in *daily marijuana* use (8.5% for males vs. 3.8% for females), *daily alcohol* use (8.3% vs. 3.2%), and occasions of drinking *five or more drinks in a row* in the prior two weeks (44% vs. 28%). The gender difference in occasions of drinking five or more drinks in a row is larger among young adults than among 12th graders, where it is 26% for males versus 18% for females. There is a particularly large gender difference in the measures of *extreme binge drinking*: the rate for having 10 or more drinks on at least one occasion in the prior two weeks was 19.5 % for males vs 6.5% for females. The rate for having 15 or more drinks in that interval was 5.6% for males and 1.7% for females.
- *Synthetic marijuana* use is higher among young adult males than females (4.5% vs. 2.0%), as is use of *bath salts* (0.7% vs. 0.3%).
- *Ecstasy (MDMA)* use is slightly higher among males than among females with annual prevalence of 4.8% and 3.2%, respectively.
- Annual prevalence of use of *narcotics other than heroin* outside of medical supervision is also slightly higher among males than females (8.1% versus 6.1%). Use of *Vicodin*, one of the most widely used drugs in the class, is 7.1% and 5.2%, respectively. The gender difference for *OxyContin* is 3.6% for males vs. 1.9% for females.
- The use of *amphetamines* is slightly higher among males than among females with annual prevalence of 8.3% and 6.5%, respectively.

- In the 1980s, there were few differences between males and females in their rates of *cigarette* use. By the early 1990s, however, males had slightly higher rates of use. In 2013, 19- to 30-year-old males are more likely than females to have smoked in the past month (23% vs. 18%), to have smoked daily in the past month (13% vs. 11%), and to have smoked half a pack or more per day in the past month (7.7% vs. 6.5%). These gender differences reflect a cohort effect in which the differences between the genders in high school during the 1990s are carried up the age spectrum.
- Among young adults there is a large gender difference in 2013 in the use of *smokeless tobacco*, with males much more likely than females to have used in the prior month (14.6% vs. 2.9%). Almost all past year use of *snus* occurs among males (11.1% vs. 0.8% among females) as is true for *dissolvable tobacco* (0.5% vs. 0.1%, respectively). Males are nearly three times as likely to have smoked *small cigars* in the past year as females (29% vs. 10%). There is less gender difference in the use of *hookah* pipes (23% vs. 16%, respectively).
- *Steroid* use among young adults is much more prevalent among males than females, as is true for 12th graders. Among 12th graders in 2013, 2.2% of males reported steroid use in the past year versus 0.7% of females. These statistics are much lower among 19- to 30-year-olds, but use by males remains higher (0.9% for males vs. 0.1% for females).

Regional Differences

Follow-up respondents are asked in what state they currently reside. States are then grouped into the same regions used in the analysis of high school data.⁵³ Tables 4-2 through 4-5 present regional differences in lifetime, annual, 30-day, and current daily prevalence for 19- to 30-year-olds combined.

- There exist some regional differences in the annual prevalence of *marijuana* use, with rates being higher in the Northeast (35%) and West (35%) than the Midwest (30%) and the South (26%). Likewise, regarding annual prevalence of any illicit drug use, rates are higher in the Northeast (39%) and West (38%) than in the Midwest (33%) and South (31%). The same is true for the use of *any illicit drug other than marijuana* (Table 4-3).
- The annual prevalence rate for *synthetic marijuana* does not differ significantly by region (Table 4-3).

⁵³States are grouped into regions as follows: *Northeast*—Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania; *Midwest*—Ohio, Indiana, Illinois, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas; *South*—Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas; *West*—Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, and California.

- The West and Northeast continue to have higher rates than the other two regions for **hallucinogen** use, though the regional differences are not large (Table 4-3). **LSD** use is highest in the Northeast while the use of **hallucinogens other than LSD** also is highest in the West (3.6%) and Northeast (3.5%).
- For **ecstasy (MDMA)**, annual prevalence is higher in the West (5.5%) and Northeast (5.1%) than in the other two regions (2.7% to 3.1%).
- For the remaining **illicit drugs**, current regional differences are not substantial (Tables 4-3 and 4-4).
- Prevalence rates for **alcohol use** are typically somewhat higher in the Northeast and Midwest regions than in the South and West; this pattern has generally been true among 12th graders as well. For **binge drinking** among 19- to 30-year-olds, the Northeast and Midwest are at 38% and 40% respectively, with the West at 30% and the South at 31%. Self-reported **drunkenness** shows a similar pattern, as would be expected.
- **Cigarette smoking** among young adults is now fairly evenly distributed across the four regions, with 30-day prevalence ranging from 17% in the West to 21% in the Midwest (Table 4-4). Smoking a half pack or more per day is distinctly lower in the West at 3.6% versus a range of 7.0% to 8.4% in the other three regions (Table 4-5).
- Use of **small cigars** in the past year is highest in the Midwest (at 21%) and stands at 16% to 18% in the other three regions (Table 4-3).
- The 30-day prevalence of **smokeless tobacco** use among young adults is highest in the Midwest (Table 4-4). The use of **snus** in the prior year is also highest in the Midwest, while use of a **hookah** to smoke tobacco is highest in the West and lowest in the South (Table 4-3).

Population Density Differences

Population density is measured by asking respondents to select the response category that best describes the size and nature of the community where they lived during March of the year in which they completed the follow-up questionnaire. The various categories are listed in Tables 4-2 through 4-5; the population sizes given to the respondent to help define each level are provided in a footnote to each table. An examination of the 1987 and 1988 drug use data for the two most urban strata revealed that the modest differences in prevalence rates between the suburbs and their corresponding cities were not worth the complexity of reporting them separately; accordingly, since then these categories have been merged to increase sample sizes. See Tables 4-2 through 4-5 for the tabular results on 19- to 30-year-olds combined.

- Differences in illicit drug use by population density tend to be very modest, perhaps more modest than is commonly supposed. Among the general population, use of most illicit drugs is broadly distributed among all areas from rural to urban. To the extent that there are variations, almost all of the associations are positive, with rural/country areas having the lowest levels of use, and small towns having the next lowest. Medium-sized cities, large cities, and very large cities tend to be higher, with only small variations among them. Positive associations with population density exist for annual prevalence of *any illicit drug, any illicit drug other than marijuana, marijuana, hallucinogens other than LSD, ecstasy (MDMA), cocaine, powder cocaine, amphetamines, Adderall, tobacco using a hookah, and small cigars*. The association is strongest for *cocaine*, where the annual prevalence rate in the very large cities (5.9%) is nearly two times that in farm/country (3.3%).
- The use of *bath salts, salvia, synthetic marijuana, tranquilizers, and steroids* does not vary significantly by population density.
- In 2013 annual prevalence rates of *methamphetamine* and *crystal methamphetamine* use are highest in the farm/country stratum with practically no difference among the other strata (Table 4-3).
- Among young adults, the lifetime and annual *alcohol* use measures all show a slight positive association with population density, while 30-day use has a somewhat stronger positive association, with 58% of the farm/country stratum reporting use in the prior 30 days versus 81% of those in very large cities.

Occasions of heavy drinking are positively associated with population density as well (Table 4-5), with 29% of those in the farm/country stratum indicating having had five or more drinks in a row at least once in the prior two weeks compared to 41% of those in the very large cities. *Daily alcohol use* in the prior month is also positively associated with population density in 2013 with 4.1% of young adults in the farm/country stratum indicating daily use versus 7.8% in the very large cities.

- Contrary to what we find for almost all other substances, there exists a negative association between population density and *daily cigarette smoking*, which is highest in the farm/country stratum and lowest in the very large cities (daily prevalence rates of 16% and 10%, respectively). Smoking at the half-pack-a-day level is about two and one-half times as high in the farm/country stratum as in very large cities (12% vs. 5%, respectively; Table 4-5).
- The use of *small cigars* in the prior year is very evenly distributed across all population density strata (ranging from 16.4% to 18.4%).
- *Hookah* smoking is higher in the cities of all sizes (from 21% to 23% using in the prior year) than in the small town (17%) or farm/country strata (7%).

Monitoring the Future

- On the other hand, *smokeless tobacco* use is far higher in the farm/country stratum (with a 24% 30-day prevalence rate) versus all other strata (from 4% to 7%). See Table 4-4.
- The use of *snus* by young adults also is more concentrated in the farm/country stratum (10.4% annual prevalence) versus 2.8% in the large cities and 4.4–4.7% in the other strata (Table 4-3).

TABLE 4-1
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N =</i>	<u>Males</u>	<u>Females</u>	<u>Total</u>
		2,100	3,100	5,200
Any Illicit Drug ^a				
Annual		37.7	32.8	34.7
30-Day		25.2	18.0	20.9
Any Illicit Drug other than Marijuana ^a				
Annual		20.2	15.9	17.7
30-Day		9.1	7.2	7.9
Marijuana				
Annual		34.5	28.0	30.6
30-Day		23.2	14.8	18.2
Daily ^b		8.5	3.8	5.7
Inhalants ^c				
Annual		0.7	0.3	0.5
30-Day		0.1	0.1	0.1
Hallucinogens				
Annual		6.0	2.2	3.7
30-Day		1.3	0.6	0.9
LSD				
Annual		3.0	1.0	1.8
30-Day		0.6	0.2	0.4
Hallucinogens other than LSD				
Annual		5.0	1.8	3.1
30-Day		0.9	0.5	0.6
PCP ^d				
Annual		0.5	0.3	0.4
30-Day		0.3	*	0.1
Ecstasy (MDMA) ^c				
Annual		4.8	3.2	3.8
30-Day		1.1	0.9	1.0
Cocaine				
Annual		5.3	2.8	3.8
30-Day		1.7	1.1	1.4
Crack ^e				
Annual		0.5	0.2	0.3
30-Day		0.1	0.1	0.1
Other Cocaine ^f				
Annual		5.2	2.6	3.7
30-Day		1.5	1.1	1.2
Heroin				
Annual		0.8	0.5	0.6
30-Day		0.5	0.2	0.3
With a Needle ^g				
Annual		0.4	0.1	0.3
30-Day		0.4	0.1	0.2
Without a Needle ^g				
Annual		1.0	0.4	0.6
30-Day		0.4	0.3	0.3

(Table continued on next page.)

TABLE 4-1 (cont.)
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<u>Males</u>	<u>Females</u>	<u>Total</u>
<i>Approximate Weighted N =</i>	2,100	3,100	5,200
Narcotics other than Heroin^h			
Annual	8.1	6.1	6.9
30-Day	3.2	2.2	2.6
Amphetamines, Adjusted^{h,j}			
Annual	8.3	6.5	7.2
30-Day	3.4	2.6	2.9
Methamphetamine^g			
Annual	0.8	0.6	0.7
30-Day	0.4	0.2	0.3
Crystal Methamphetamine (Ice)^g			
Annual	1.1	0.7	0.9
30-Day	0.9	0.1	0.4
Sedatives (Barbiturates)^h			
Annual	3.3	3.3	3.3
30-Day	1.2	1.3	1.2
Tranquilizers^h			
Annual	5.7	5.2	5.4
30-Day	1.9	1.9	1.9
Alcohol			
Annual	84.0	82.5	83.1
30-Day	72.5	66.8	69.1
Daily ^b	8.3	3.2	5.2
5+ Drinks in a Row in Last 2 Weeks	44.2	27.8	34.4
10+ Drinks in a Row in Last 2 Weeks ^{d,i}	17.1	5.5	10.1
15+ Drinks in a Row in Last 2 Weeks ^{d,k}	7.5	1.5	3.8
Been Drunk^c			
Annual	67.4	59.7	62.9
30-Day	41.8	33.2	36.8
Daily ^b	0.8	0.2	0.5
Flavored Alcoholic Beverages^d			
Annual	47.9	54.1	51.6
30-Day	22.0	25.0	23.8
Cigarettes			
Annual	32.3	26.8	29.0
30-Day	22.5	17.5	19.5
Daily	13.0	11.3	12.0
1/2 Pack+/Day	7.7	6.5	7.0
Smokeless Tobacco^d			
Lifetime	39.6	15.3	25.3
30-Day	14.6	2.9	7.7
Daily	5.4	1.0	2.8
Steroids^g			
Annual	0.9	0.1	0.4
30-Day	0.1	*	0.1

(Table continued on next page.)

TABLE 4-1 (cont.)
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2013

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.

^cThis drug was asked about in three of the six questionnaire forms. Total *N* is approximately 2,600.

^dThis drug was asked about in one of the six questionnaire forms. Total *N* is approximately 900.

^eThis drug was asked about in five of the six questionnaire forms. Total *N* is approximately 4,300.

^fThis drug was asked about in four of the six questionnaire forms. Total *N* is approximately 3,500.

^gThis drug was asked about in two of the six questionnaire forms. Total *N* is approximately 1,700.

^hOnly drug use that was not under a doctor's orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^jErratum corrected on Nov. 10th, 2015. These are the correct values, rather than the 19.5%, 6.5%, and 11.7% previously contained here.

^kErratum corrected on Nov. 10th, 2015. These are the correct values, rather than the 8.6%, 1.7%, and 4.5% previously contained here.

TABLE 4-2
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013
(Entries are percentages.)

	Approximate Weighted N	Any Illicit Drug ^a	Any Illicit Drug other than				Hallucinogens		Ecstasy			
			Marijuana ^a	Marijuana	Inhalants ^b	Hallucinogens	LSD	other than LSD	PCP ^c	(MDMA) ^b	Cocaine	Crack ^d
Total	5,200	61.3	34.9	57.9	7.3	13.1	6.8	12.0	1.9	12.1	13.0	2.8
Gender												
Male	2,100	63.1	37.4	60.5	9.7	17.9	9.4	16.8	1.9	13.9	15.2	3.1
Female	3,100	60.1	33.1	56.1	5.8	10.0	5.0	8.8	1.9	10.9	11.5	2.6
Modal Age												
19–20	900	48.7	24.3	47.0	5.0	8.2	4.7	7.0	0.8	7.7	4.7	1.0
21–22	900	58.3	33.3	54.5	6.9	10.3	5.5	9.3	0.3	11.3	9.6	1.8
23–24	900	62.2	34.4	58.4	6.3	12.9	7.0	12.0	1.3	11.3	11.7	1.9
25–26	800	64.7	39.3	60.8	7.1	14.6	7.0	13.7	2.3	13.6	17.4	4.3
27–28	800	69.7	40.6	65.7	7.5	16.6	7.2	15.8	2.3	14.5	18.9	4.3
29–30	800	65.5	38.6	62.1	11.6	16.8	9.6	14.9	3.9	14.3	16.9	4.0
Region												
Northeast	1,000	64.5	35.8	62.3	7.4	15.3	8.5	13.7	1.4	13.9	14.8	2.5
Midwest	1,400	60.0	35.1	56.9	6.2	13.6	8.1	12.2	3.2	10.6	11.6	3.1
South	1,600	59.4	33.1	54.8	6.7	10.2	5.0	9.5	1.3	10.2	10.9	3.1
West	1,100	62.1	35.9	59.2	10.3	14.7	6.4	13.7	1.2	15.2	16.1	2.6
Population Density^e												
Farm/Country	500	54.6	32.4	50.0	4.1	10.6	4.8	9.7	*	7.9	11.3	3.2
Small Town	1,300	57.7	33.2	54.5	7.6	12.6	6.8	11.7	2.7	11.8	12.0	2.4
Medium City	1,300	59.7	33.8	56.9	6.9	12.5	6.8	11.5	1.8	11.0	11.5	2.9
Large City	1,200	65.6	35.3	62.3	8.5	13.4	7.0	12.2	1.7	13.1	13.6	2.5
Very Large City	700	68.1	40.6	63.8	8.2	16.5	7.9	15.2	2.4	16.8	17.5	3.6

(Table continued on next page.)

TABLE 4-2 (cont.)
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013
 (Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Other Cocaine^f</i>	<i>Heroin</i>	<i>Heroin with a Needle^g</i>	<i>Heroin without a Needle^g</i>	<i>Narcotics other than Heroin^h</i>	<i>Amphetamines^{h,i}</i>	<i>Methamphetamine^g</i>	<i>Crystal Methamphetamine (Ice)^g</i>
Total	5,200	12.6	1.7	0.9	1.8	18.2	18.5	3.4	3.1
Gender									
Male	2,100	15.2	2.2	1.3	2.5	20.4	20.8	3.8	3.7
Female	3,100	10.8	1.3	0.6	1.4	16.8	17.0	3.1	2.8
Modal Age									
19–20	900	4.3	0.8	0.4	1.2	11.0	13.8	1.6	0.7
21–22	900	9.4	1.2	0.4	1.0	16.4	18.4	1.8	0.5
23–24	900	11.7	1.9	1.0	2.6	16.1	18.7	3.2	2.6
25–26	800	16.5	2.4	1.9	2.4	21.4	23.0	3.5	3.2
27–28	800	18.2	1.7	1.1	1.9	22.9	20.3	6.0	7.5
29–30	800	16.4	2.1	0.3	1.9	22.8	17.5	4.8	4.6
Region									
Northeast	1,000	15.3	2.3	0.7	1.9	18.7	18.9	2.6	0.2
Midwest	1,400	10.5	1.8	0.8	1.9	20.1	19.1	3.3	3.1
South	1,600	10.3	1.0	0.4	1.3	15.8	18.3	2.8	3.6
West	1,100	15.8	1.9	1.5	2.3	19.0	17.7	5.3	5.2
Population Density^e									
Farm/Country	500	11.7	1.8	1.1	2.0	17.9	14.4	4.5	4.1
Small Town	1,300	11.4	1.5	0.4	1.7	17.9	17.7	3.3	2.1
Medium City	1,300	10.8	1.7	0.8	1.9	17.9	17.6	3.4	3.2
Large City	1,200	13.4	1.5	0.9	1.5	19.1	19.5	2.9	3.0
Very Large City	700	16.9	1.8	1.1	2.1	18.9	22.7	3.7	4.3

(Table continued on next page.)

TABLE 4-2 (cont.)
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Sedatives (Barbiturates)^h</i>	<i>Tranquilizers^h</i>	<i>Alcohol</i>	<i>Been Drunk^b</i>	<i>Flavored Alcoholic Beverages^c</i>	<i>Cigarettes</i>	<i>Smokeless Tobacco^c</i>	<i>Steroids^g</i>
Total	5,200	9.6	14.2	87.2	79.1	81.6	—	25.3	1.3
Gender									
Male	2,100	10.2	14.0	87.2	80.4	79.3	—	39.6	2.7
Female	3,100	9.2	14.4	87.1	78.2	83.2	—	15.3	0.4
Modal Age									
19–20	900	5.8	7.9	72.1	60.4	66.0	—	21.9	0.7
21–22	900	8.4	10.1	85.2	76.1	71.0	—	9.0	0.4
23–24	900	9.2	12.8	88.8	82.1	84.3	—	26.2	2.3
25–26	800	12.8	17.3	91.1	81.4	91.4	—	28.4	1.0
27–28	800	11.6	18.6	94.7	88.1	92.7	—	43.3	1.6
29–30	800	10.6	19.7	92.3	87.2	86.0	—	21.8	1.6
Region									
Northeast	1,000	8.9	15.0	90.3	82.8	82.9	—	20.6	0.9
Midwest	1,400	10.4	13.9	90.1	84.2	84.1	—	34.1	0.8
South	1,600	9.8	14.3	85.5	75.8	80.1	—	24.4	1.2
West	1,100	9.2	13.7	83.0	73.9	78.5	—	17.4	2.0
Population Density^e									
Farm/Country	500	9.4	10.4	81.9	74.8	78.4	—	39.6	1.2
Small Town	1,300	9.6	14.6	85.6	78.3	75.1	—	23.4	0.7
Medium City	1,300	9.5	13.1	85.9	75.9	82.1	—	27.3	1.6
Large City	1,200	10.6	15.3	89.8	80.9	85.3	—	22.3	1.4
Very Large City	700	9.0	16.5	91.9	85.8	89.3	—	16.6	1.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N is approximately 2,600.

^cThis drug was asked about in one of the six questionnaire forms. Total N is approximately 900.

^dThis drug was asked about in five of the six questionnaire forms. Total N is approximately 4,300.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000.

Within each level of population density, suburban and urban respondents are combined.

^fThis drug was asked about in four of the six questionnaire forms. Total N is approximately 3,500.

^gThis drug was asked about in two of the six questionnaire forms. Total N is approximately 1,700.

^hOnly drug use that was not under a doctor's orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

TABLE 4-3
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	Any Illicit Drug ^a	Any Illicit Drug other than Marijuana ^a	Marijuana	Synthetic Marijuana ^c	Inhalants ^c	Hallucinogens	LSD	Hallucinogens other than LSD	PCP ^d	Ecstasy (MDMA) ^c	Salvia ^c	Cocaine	Crack ^e
Total	5,200	34.7	17.7	30.6	3.0	0.5	3.7	1.8	3.1	0.4	3.8	0.8	3.8	0.3
Gender														
Male	2,100	37.7	20.2	34.5	4.5	0.7	6.0	3.0	5.0	0.5	4.8	1.1	5.3	0.5
Female	3,100	32.8	15.9	28.0	2.0	0.3	2.2	1.0	1.8	0.3	3.2	0.6	2.8	0.2
Modal Age														
19–20	900	37.5	17.3	35.5	3.4	0.4	5.0	2.9	4.1	*	5.0	1.4	2.6	0.3
21–22	900	40.9	21.6	36.7	3.4	0.9	4.7	2.9	3.7	*	5.9	1.3	4.8	0.7
23–24	900	37.1	18.3	34.3	4.5	0.8	4.1	2.0	3.4	0.2	4.9	0.7	4.2	0.2
25–26	800	34.6	17.4	28.4	3.1	0.2	2.7	1.5	2.2	*	3.3	0.6	4.4	0.5
27–28	800	30.8	15.8	25.2	1.4	0.2	2.7	0.7	2.4	0.8	2.0	0.3	3.5	*
29–30	800	26.6	15.2	22.4	2.0	0.3	2.6	0.7	2.3	1.0	1.6	0.3	3.4	0.3
Region														
Northeast	1,000	38.7	19.3	34.7	3.0	0.8	4.2	2.4	3.5	0.2	5.1	1.2	6.1	0.8
Midwest	1,400	33.3	18.3	29.5	4.0	0.2	3.7	1.7	3.2	0.9	3.1	0.7	2.6	0.3
South	1,600	31.2	15.6	26.1	2.5	0.6	2.9	1.5	2.3	0.2	2.7	0.3	2.9	0.1
West	1,100	37.6	17.9	34.5	2.1	0.4	4.2	1.7	3.6	*	5.5	1.2	4.7	0.3
Population Density ^j														
Farm/Country	500	25.4	13.5	20.6	2.1	0.9	2.3	1.2	1.9	*	1.3	0.6	2.2	0.3
Small Town	1,300	32.8	17.1	28.7	3.3	0.4	3.2	1.6	2.7	0.6	4.1	0.9	3.5	0.2
Medium City	1,300	34.5	17.7	30.5	3.2	0.5	3.8	2.1	3.0	0.2	3.4	0.7	3.2	0.2
Large City	1,200	37.7	18.3	33.5	3.0	0.3	3.5	1.6	3.1	0.5	3.6	0.7	4.0	0.3
Very Large City	700	39.6	20.1	35.9	2.6	0.6	5.2	2.0	4.5	0.3	6.7	1.1	6.0	0.7

(Table continued on next page.)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Other Cocaine^f</i>	<i>Heroin</i>	<i>Heroin with a Needle^b</i>	<i>Heroin without a Needle^b</i>	<i>Narcotics other than Heroin^g</i>	<i>OxyContin^{c,g}</i>	<i>Vicodin^{c,g}</i>	<i>Amphetamines^{g,h}</i>	<i>Ritalin^{g,h}</i>	<i>Adderall^{c,g}</i>	<i>Methamphetamine^b</i>	<i>Crystal Methamphetamine (Ice)^b</i>
Total	5,200	3.7	0.6	0.3	0.6	6.9	2.6	6.0	7.2	1.8	6.4	0.7	0.9
Gender													
Male	2,100	5.2	0.8	0.4	1.0	8.1	3.6	7.1	8.3	2.3	7.9	0.8	1.1
Female	3,100	2.6	0.5	0.1	0.4	6.1	1.9	5.2	6.5	1.4	5.3	0.6	0.7
Modal Age													
19–20	900	2.5	0.5	0.1	0.6	7.1	3.3	5.3	8.7	3.4	8.8	0.4	0.3
21–22	900	4.6	0.8	*	1.0	6.9	2.8	7.4	11.1	2.7	9.5	0.4	0.2
23–24	900	3.9	0.6	0.6	0.9	7.2	2.8	5.8	7.4	1.4	5.9	1.3	1.2
25–26	800	4.4	0.8	0.8	0.7	6.9	2.7	7.9	6.3	1.2	5.4	0.5	1.2
27–28	800	3.2	0.6	*	0.5	6.8	2.3	4.5	5.1	1.3	5.1	0.5	1.0
29–30	800	3.4	0.4	*	*	6.5	1.5	5.2	4.5	0.3	3.0	0.9	1.2
Region													
Northeast	1,000	6.0	1.5	0.4	1.3	7.5	3.2	6.6	6.9	2.3	5.8	0.6	0.1
Midwest	1,400	2.3	0.6	0.1	0.6	7.4	2.7	7.3	8.5	2.3	7.9	0.2	1.2
South	1,600	2.7	0.2	*	0.1	5.7	2.3	3.6	6.6	1.3	5.7	0.9	1.2
West	1,100	4.8	0.4	0.4	0.7	7.6	2.4	7.4	6.8	1.3	5.9	1.0	0.7
Population Density^j													
Farm/Country	500	2.1	0.9	0.3	0.6	6.0	1.6	4.7	3.9	1.0	3.5	1.8	1.9
Small Town	1,300	3.8	0.6	*	0.3	6.8	2.8	4.0	7.0	0.9	5.8	0.7	0.8
Medium City	1,300	2.9	0.6	0.3	0.8	7.4	3.2	7.5	7.7	2.7	7.3	0.8	1.1
Large City	1,200	4.0	0.5	0.2	0.8	6.8	2.2	6.5	7.5	1.1	6.5	0.3	0.5
Very Large City	700	5.2	0.5	0.2	0.4	7.0	2.7	7.5	8.1	3.0	7.6	0.2	0.6

(Table continued on next page.)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Bath Salts (synthetic stimulants)^c</i>	<i>Sedatives (Barbiturates)^g</i>	<i>Tranquilizers^g</i>	<i>GHB^b</i>	<i>Ketamine^b</i>	<i>Alcohol</i>	<i>Been Drunk^c</i>	<i>Flavored Alcoholic Beverages^d</i>	<i>Alcoholic Beverages containing Caffeine^b</i>	<i>Cigarettes</i>	<i>Tobacco using a Hookah^c</i>	<i>Small Cigars^c</i>	<i>Dissolvable Tobacco^b</i>	<i>Snus^b</i>	<i>Steroids^g</i>
Total	5,200	0.4	3.3	5.4	0.3	0.6	83.1	62.9	51.6	35.3	29.0	18.8	17.7	0.3	5.1	0.4
Gender																
Male	2,100	0.7	3.3	5.7	0.5	0.8	84.0	67.4	47.9	43.4	32.3	23.1	29.0	0.5	11.1	0.9
Female	3,100	0.3	3.3	5.2	0.1	0.4	82.5	59.7	54.1	29.4	26.8	15.9	9.8	0.1	0.8	0.1
Modal Age																
19–20	900	0.3	3.1	4.8	0.4	0.7	68.4	48.6	51.5	31.6	28.0	28.3	22.7	1.0	8.6	0.7
21–22	900	0.2	3.6	4.8	0.3	1.0	82.8	67.6	62.9	42.5	31.2	24.7	19.3	*	4.8	0.3
23–24	900	1.1	3.6	6.8	0.7	0.1	84.7	67.1	55.2	41.1	32.0	21.2	17.3	0.5	4.6	0.6
25–26	800	0.5	4.2	4.6	*	0.8	87.9	63.1	53.6	36.8	29.6	15.9	18.5	*	4.1	0.4
27–28	800	0.1	2.7	6.1	0.1	0.2	89.6	69.6	48.6	32.5	27.9	10.5	13.7	*	1.3	0.5
29–30	800	0.4	2.6	5.5	*	0.6	86.6	62.2	37.8	27.0	24.9	10.5	14.0	0.1	6.7	*
Region																
Northeast	1,000	*	3.5	6.3	0.3	0.9	87.4	69.6	50.1	38.3	29.2	17.6	16.1	0.1	2.9	*
Midwest	1,400	0.4	3.5	5.7	0.2	0.4	86.8	68.5	55.3	38.6	30.8	19.6	20.6	0.4	8.8	0.3
South	1,600	0.7	3.3	4.7	0.4	0.6	80.9	58.3	50.0	31.0	27.8	14.8	15.7	0.2	3.8	0.6
West	1,100	0.6	2.9	5.1	0.2	0.2	77.8	55.8	50.1	35.6	27.7	24.5	18.1	0.4	3.5	0.8
Population Density^l																
Farm/Country	500	0.3	4.0	3.9	0.2	0.5	75.6	54.6	51.6	28.9	32.2	6.7	18.4	0.4	10.4	*
Small Town	1,300	0.4	3.0	5.6	0.2	0.1	80.6	61.4	47.8	34.5	30.4	17.0	18.0	*	4.7	0.2
Medium City	1,300	0.5	3.9	5.4	0.1	1.1	82.2	61.5	53.6	37.4	28.4	22.8	18.5	*	4.4	0.7
Large City	1,200	0.6	3.1	5.5	0.5	0.5	86.9	66.5	53.3	35.9	28.3	20.7	17.1	0.9	4.6	0.6
Very Large City	700	0.3	2.8	5.9	0.4	0.7	89.0	67.6	50.8	36.4	26.0	22.1	16.4	*	2.8	0.4

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in two of the six questionnaire forms. Total N is approximately 1,700.

^cThis drug was asked about in three of the six questionnaire forms. Total N is approximately 2,600.

^dThis drug was asked about in one of the six questionnaire forms. Total N is approximately 900.

^eThis drug was asked about in five of the six questionnaire forms. Total N is approximately 4,300.

^fThis drug was asked about in four of the six questionnaire forms. Total N is approximately 3,500.

^gOnly drug use that was not under a doctor's orders is included here.

^hBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^lA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000.

Within each level of population density, suburban and urban respondents are combined.

TABLE 4-4
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	Approximate Weighted N	Any Illicit Drug ^a					Hallucinogens other than			Ecstasy (MDMA) ^b	Cocaine	Crack ^d
		Any Illicit Drug ^a	other than Marijuana	Marijuana	Inhalants ^b	Hallucinogens	LSD	LSD	PCP ^c			
Total	5,200	20.9	7.9	18.2	0.1	0.9	0.4	0.6	0.1	1.0	1.4	0.1
Gender												
Male	2,100	25.2	9.1	23.2	0.1	1.3	0.6	0.9	0.3	1.1	1.7	0.1
Female	3,100	18.0	7.2	14.8	0.1	0.6	0.2	0.5	*	0.9	1.1	0.1
Modal Age												
19–20	900	22.5	7.8	21.8	*	1.6	0.6	1.4	*	0.8	1.0	*
21–22	900	25.9	10.5	23.0	0.3	1.0	0.5	0.5	*	1.2	2.2	0.5
23–24	900	22.9	8.6	20.0	0.3	1.2	0.5	0.7	*	1.7	1.5	0.1
25–26	800	19.8	7.8	15.8	*	0.7	0.4	0.4	*	0.9	1.2	*
27–28	800	17.4	6.7	13.9	0.1	0.5	0.1	0.4	0.8	0.6	1.7	*
29–30	800	16.0	6.1	13.7	*	0.4	0.1	0.4	*	0.7	0.6	0.1
Region												
Northeast	1,000	22.5	8.9	20.0	0.1	1.4	0.6	0.9	*	1.9	2.4	0.2
Midwest	1,400	19.7	7.5	17.2	0.1	0.5	0.2	0.4	0.4	0.8	0.8	0.1
South	1,600	18.8	7.6	15.7	0.1	0.8	0.3	0.7	*	0.6	0.9	0.2
West	1,100	23.4	7.8	20.9	0.3	1.0	0.5	0.6	*	1.2	1.8	*
Population Density ^e												
Farm/Country	500	15.0	6.9	11.4	*	0.9	0.7	0.6	*	0.2	0.5	*
Small Town	1,300	19.6	7.6	17.4	*	1.0	0.4	0.7	*	0.8	1.2	0.1
Medium City	1,300	20.4	8.2	17.8	0.2	0.8	0.4	0.5	*	0.4	1.3	*
Large City	1,200	23.0	7.7	19.9	0.2	1.0	0.4	0.7	0.5	1.1	1.4	0.2
Very Large City	700	23.9	8.3	21.5	0.1	0.9	0.2	0.8	*	2.9	2.2	0.2

(Table continued on next page.)

TABLE 4-4 (cont.)
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Other Cocaine^f</i>	<i>Heroin</i>	<i>Heroin With Needle^g</i>	<i>Heroin Without Needle^g</i>	<i>Narcotics other than Heroin^h</i>	<i>Amphetamines^{h,j}</i>	<i>Methamphetamine^g</i>	<i>Crystal Methamphetamine (Ice)^g</i>
Total	5,200	1.2	0.3	0.2	0.3	2.6	2.9	0.3	0.4
Gender									
Male	2,100	1.5	0.5	0.4	0.4	3.2	3.4	0.4	0.9
Female	3,100	1.1	0.2	0.1	0.3	2.2	2.6	0.2	0.1
Modal Age									
19–20	900	0.9	0.2	0.1	0.4	2.2	4.3	*	*
21–22	900	1.8	0.3	*	*	2.8	5.3	*	0.2
23–24	900	1.4	0.5	0.6	0.6	3.1	2.8	0.3	0.3
25–26	800	1.1	0.5	0.7	0.5	2.5	1.8	0.3	0.5
27–28	800	1.3	0.3	*	0.4	2.5	1.8	0.5	1.0
29–30	800	0.9	0.0	*	*	2.3	1.4	0.4	0.6
Region									
Northeast	1,000	2.5	0.7	0.4	0.3	2.7	3.2	*	*
Midwest	1,400	0.5	0.3	0.1	0.6	2.5	3.1	0.1	0.3
South	1,600	0.7	0.1	*	*	2.4	2.8	0.3	0.6
West	1,100	1.9	0.2	0.3	0.3	2.7	2.9	0.6	0.7
Population Density^e									
Farm/Country	500	0.5	0.3	0.3	*	2.7	2.4	0.7	0.9
Small Town	1,300	1.5	0.2	*	0.1	2.5	3.0	0.1	0.2
Medium City	1,300	0.9	0.4	0.3	0.6	2.6	3.0	0.6	0.7
Large City	1,200	1.2	0.2	0.1	0.2	2.5	2.7	0.1	0.4
Very Large City	700	1.9	0.3	0.2	0.2	2.3	3.4	*	*

(Table continued on next page.)

TABLE 4-4 (cont.)
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Sedatives (Barbiturates)^h</i>	<i>Tranquilizers^h</i>	<i>Alcohol</i>	<i>Been Drunk^b</i>	<i>Flavored Alcoholic Beverages^c</i>	<i>Cigarettes</i>	<i>Smokeless Tobacco^c</i>	<i>Steroids^g</i>
Total	5,200	1.2	1.9	69.1	36.8	23.8	19.5	7.7	0.1
Gender									
Male	2,100	1.2	1.9	72.5	41.8	22.0	22.5	14.6	0.1
Female	3,100	1.3	1.9	66.8	33.2	25.0	17.5	2.9	*
Modal Age									
19–20	900	1.1	1.3	51.5	32.1	26.3	18.4	11.0	*
21–22	900	1.3	1.6	70.5	43.1	31.3	20.8	0.7	*
23–24	900	1.4	2.1	72.7	41.7	27.3	21.4	6.5	*
25–26	800	1.7	1.6	75.9	34.4	23.7	19.5	9.6	0.4
27–28	800	0.6	2.9	73.9	37.0	18.9	20.0	9.7	0.1
29–30	800	1.1	1.9	71.1	32.5	14.9	16.5	8.0	*
Region									
Northeast	1,000	1.4	1.8	74.8	43.7	23.3	18.3	5.0	*
Midwest	1,400	1.2	1.8	73.1	39.9	26.2	21.4	14.7	0.1
South	1,600	1.2	2.0	65.0	30.9	22.6	19.6	6.3	0.2
West	1,100	1.1	1.7	64.6	34.5	23.2	17.2	2.8	*
Population Density^e									
Farm/Country	500	2.0	1.7	57.5	28.5	17.3	22.5	23.8	*
Small Town	1,300	1.0	1.5	63.3	34.6	21.4	21.2	5.7	*
Medium City	1,300	1.4	1.8	68.6	35.5	30.3	19.2	6.0	0.2
Large City	1,200	0.9	2.3	74.7	40.7	20.9	18.2	6.5	0.1
Very Large City	700	1.1	1.9	80.5	41.8	23.9	16.3	3.7	*

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total *N* is approximately 2,600.

^cThis drug was asked about in one of the six questionnaire forms. Total *N* is approximately 900.

^dThis drug was asked about in five of the six questionnaire forms. Total *N* is approximately 4,300.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000.

Within each level of population density, suburban and urban respondents are combined.

^fThis drug was asked about in four of the six questionnaire forms. Total *N* is approximately 3,500.

^gThis drug was asked about in two of the six questionnaire forms. Total *N* is approximately 1,700.

^hOnly drug use that was not under a doctor's orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

TABLE 4-5
Thirty-Day Prevalence of Daily Use ^a of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2013

(Entries are percentages.)

	<i>Approximate Weighted N</i>	<i>Marijuana Daily</i>	<i>Alcohol Daily</i>	<i>Alcohol: 5+ Drinks in a Row in Last 2 Weeks</i>	<i>Cigarettes Daily</i>	<i>Cigarettes: 1/2 Pack+ per Day</i>	<i>Smokeless Tobacco^c</i>
Total	5,200	5.7	5.2	34.4	12.0	7.0	2.8
Gender							
Male	2,100	8.5	8.3	44.2	13.0	7.7	5.4
Female	3,100	3.8	3.2	27.8	11.3	6.5	1.0
Modal Age:							
19–20	900	6.2	2.7	27.2	10.8	5.4	3.2
21–22	900	7.8	4.9	40.2	12.0	6.5	0.3
23–24	900	6.2	4.9	37.7	13.1	8.1	4.0
25–26	800	5.8	6.9	37.0	10.9	6.4	4.1
27–28	800	5.1	6.5	33.6	13.8	8.5	3.2
29–30	800	2.9	5.8	30.9	11.3	7.1	1.8
Region							
Northeast	1,000	6.1	6.0	37.5	12.1	7.0	2.7
Midwest	1,400	5.6	4.4	39.6	13.7	8.0	5.1
South	1,600	4.7	5.2	31.0	13.1	8.4	1.8
West	1,100	6.8	5.7	29.6	7.9	3.6	1.4
Population Density ^b							
Farm/Country	500	3.7	4.1	29.0	16.4	11.5	9.1
Small Town	1,300	5.9	4.6	30.7	14.9	9.4	4.2
Medium City	1,300	5.4	4.1	33.8	11.0	6.2	0.7
Large City	1,200	6.3	6.4	37.9	9.5	4.8	1.5
Very Large City	700	6.0	7.8	40.9	9.5	4.6	1.7


Source. The Monitoring the Future study, the University of Michigan.

^aDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.

^bA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000. Within each level of population density, suburban and urban respondents are combined.

^cThis drug was asked about in one of the six questionnaire forms. Total N is approximately 900.

FIGURE 4-1
ANY ILLICIT DRUG^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding, some bars with the same number may have uneven height.

^aThe questions on hallucinogen use are not included in the age 55 questionnaire. Therefore, the data presented here include hallucinogens for ages 18 to 50, but not for age 55.

FIGURE 4-2
ANY ILLICIT DRUG OTHER THAN MARIJUANA^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aThe questions on hallucinogen use are not included in the age 55 questionnaire. Therefore, the data presented here include hallucinogens for ages 18 to 50, but not for age 55.


FIGURE 4-3
MARIJUANA
Lifetime, Annual, 30-Day, and Daily Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding some bars with the same number may have uneven height.

FIGURE 4-4
AMPHETAMINES^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aUsed without a doctor's orders.

FIGURE 4-5
METHAMPHETAMINE
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aQuestions about the use of methamphetamines were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-6
CRYSTAL METHAMPHETAMINE (ICE)
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aQuestions about the use of crystal methamphetamine were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.


FIGURE 4-7
COCAINE
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding some bars with the same number may have uneven height.


FIGURE 4-8
CRACK COCAINE
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding, some bars with the same number may have uneven height.


FIGURE 4-9
OTHER COCAINE
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding, some bars with the same number may have uneven height.

FIGURE 4-10
HALLUCINOGENS^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50^b
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.


Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aUnadjusted for the possible underreporting of PCP.

^bQuestions about the use of hallucinogens were not included in the questionnaires for 55-year-olds.

FIGURE 4-11
LSD
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aQuestions about the use of LSD were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-12
HALLUCINOGENS OTHER THAN LSD ^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30 ^b
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.
 Due to rounding, some bars with the same number may have uneven height.

^aUnadjusted for the possible underreporting of PCP.

^bQuestions about the use of hallucinogens other than LSD were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-13
INHALANTS^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^b
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.


Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding some, bars with the same number may have uneven height.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites.

^bQuestions about the use of inhalants were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-14
SEDATIVES (BARBITURATES) ^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aUsed without a doctor's orders.

FIGURE 4-15
NARCOTICS OTHER THAN HEROIN^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aUsed without a doctor's orders.

FIGURE 4-16
TRANQUILIZERS^a
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aUsed without a doctor's orders.

FIGURE 4-17
ECSTASY (MDMA)
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

Due to rounding, some bars with the same number may have uneven height.

^aQuestions about the use of ecstasy were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-18
STEROIDS
Lifetime, Annual, and 30-Day Prevalence^a
among Respondents of Modal Ages 18 through 30^b
by Age Group, 2013




Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.
 Due to rounding, some bars with the same number may have uneven height.

^aUsed without a doctor's orders.

^bQuestions about the use of steroids were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.


FIGURE 4-19
HEROIN
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.

Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding some bars with the same number may have uneven height.


FIGURE 4-20a
ALCOHOL
Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.


Notes. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion. Due to rounding some bars with the same number may have uneven height.

FIGURE 4-20b
ALCOHOL
2-Week Prevalence of 5 or More Drinks in a Row and
30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.
Notes. Due to rounding some bars with the same number may have uneven height.

FIGURE 4-21
CIGARETTES
Annual, 30-Day, Daily, and Half-Pack-a-Day Prevalence
among Respondents of Modal Ages 18 through 55
by Age Group, 2013



Source. The Monitoring the Future study, the University of Michigan.
 Notes. Due to rounding some bars with the same number may have uneven height.

Chapter 5

TRENDS IN DRUG USE IN EARLY AND MIDDLE ADULTHOOD

In this chapter, we examine historical trends through 2013 in substance use for various age bands covering early and middle adulthood, ages 19 through 55. We use MTF panel data from graduating high school seniors over more than three and a half decades. Although panel data are typically used to study stability and change in the same individuals over time, we use the data here cross-sectionally to consider how substance use has varied across the years by age group, much as we use the repeated cross-sectional surveys of secondary school students to track changes in behaviors over time for particular grade levels (8, 10, and 12) in Volume 1. In the early 1990s, we began to document large and important increases among secondary school students in the use of several substances, particularly marijuana and cigarettes. The increases continued among 12th graders through 1997, as discussed in Volume I. One of the important issues addressed in this chapter is whether such increases occurred only among adolescents or whether those higher-using graduating classes have carried their higher levels of drug use with them as they moved into young adulthood. In other words, are they exhibiting lasting differences across class cohorts or “cohort effects.” These would be indicated by the inflection points in the cross-time trends (turning either up or down) coming sequentially across the age strata with a time lag between adjacent strata.

Figures 5-1 through 5-19c present separate trend lines for two-year age strata, that is, respondents who are one to two years beyond high school, three to four years beyond high school, and so on. These two-year age strata are used to reduce the random fluctuations that would be seen with one-year strata.⁵⁴ Each data point through age 32 in these figures is based on approximately 1,200 weighted cases drawn from two adjacent high school classes; actual (unweighted) numbers of cases are somewhat higher than those shown in the tables.⁵⁵ Figures 5-1 through 5-19c also present trend data from respondents at modal ages 35, 40, 45, 50, and 55 based on follow-up data collected at those ages. Modal age 55 was first added to the survey in 2013, so as yet it has no trend results. The older age strata are constituted in a slightly different way, in that the two half samples from a *single* graduating class (which through age 30 had been surveyed in alternating years) are now both surveyed in the same year.⁵⁶ The

⁵⁴ Strictly speaking, these two-year strata are not age strata, because they are based on all respondents in the given year from two adjacent high school classes, and they do not take into account the minor differences in individual respondents’ ages within each graduating class; however, they are close approximations to age strata, and we characterize them by the modal age of the respondents as ages 19 to 20, 21 to 22, and so on.

⁵⁵ For the 2013 data, the 19- to 20-year-old stratum is composed of participating respondents from the high school graduating classes of 2012 and 2011, respectively; the 21- to 22-year-old stratum contains data from the classes of 2010 and 2009, respectively; and so on.

⁵⁶ In 2013, the 35-year-olds are graduates from the high school class of 1995 (weighted $N = 844$), the 40-year-olds from the high school class of 1991 (weighted $N = 833$), the 45-year-olds from the high school class of 1986 (weighted $N = 838$), the 50-year-

figures also include trend data for 18-year-olds for comparison purposes. The page following the figure for each drug contains a table of values for each point in the trend lines separately for the various age strata.

Tables 5-1 through 5-5 are derived from the same data but presented in tabular form for 19- to 28-year-olds *combined*. Data are given for each year in which they are available for that full age band (i.e., from 1986 onward). Respondents ages 29 and over are omitted from the tables because their inclusion would shorten the time period over which trends can be examined. However, the full data for them are contained in Figures 5-1 through 5-19c.

RECENT TRENDS IN DRUG USE AMONG YOUNG ADULTS AGE 19-28

In 2013 the primary changes from the previous year in substance use among young adults ages 19 to 28 combined (shown in Tables 5-2 through 5-4) are as follows:

- **Marijuana** use showed a statistically significant rise in annual prevalence to 32.2%—an increase of 2.0 percentage points from 2012. This is up from 27.7% in 2006—the recent low point. Largely due to this increase in marijuana use, the percent of young adults indicating use of **any illicit drug** in the prior year also showed a significant increase—up 2.3 percentage points to 36.3%. This is up from 32.1% in 2006.
- **Daily marijuana** use also rose in 2013 (0.6 percentage points, a non-significant increase over 2012) to reach 6.2%. Daily use is up from 5.0% in 2006 and from 2.3% in 1992, the low point since estimates for this age group first became available in 1986. The 2013 rate is the highest level of daily use observed in this population since 1986..
- **Synthetic marijuana** showed a significant decline in 2013—down 2.1 percentage points to 3.2%. This is also down appreciably from the 7.4% prevalence observed in 2011, when use of this drug was first measured; and it parallels a sharp decline in use among secondary school students.
- **All other illicit drugs** showed minor changes in 2013.
- There were no significant changes in the various measures of **alcohol** use, although most of these measures showed some modest decline in 2013.
- **Cigarette smoking** did not show any significant changes in 2013. *Thirty-day* prevalence rose slightly (up 0.4 percentage points) to 20.0%, while *daily* and

olds are graduates from the high school class of 1981 (weighted $N = 828$), and the 55-year-olds are graduates from the high school class of 1976 (weighted $N=870$). The unweighted actual N s are somewhat higher.

half-pack-a-day smoking both declined some to reach 12.1% and 7.0%, respectively, in 2013.

LONGER-TERM TRENDS IN EARLY AND MIDDLE ADULTHOOD

The trend results are as follows:

- Longer term declines among young adults in the annual prevalence of several drugs appeared to end in 1992 or 1993 (Table 5-2). Among the 19- to 28-year-old young adult sample, this was true for the use of *any illicit drug, marijuana, any illicit drug other than marijuana, hallucinogens, narcotics other than heroin, crack, amphetamines, sedatives (barbiturates), and tranquilizers*. In 1994, annual prevalence for most drugs remained steady. *Cocaine other than crack* reached its low point in 1994 after a period of substantial decline that began in the late 1980s. In 1995 there were modest increases (a percentage point or less) in the annual prevalence of almost all of the drug classes in Table 5-2, some of which were statistically significant.

Thus, it is clear that by 1992 or 1993 the downward secular trend (i.e., period effect) observable in all of these age strata (as well as among adolescents) had ended.⁵⁷ What has happened since then, however, is more of a cohort effect, reflecting an interaction between age and period such that only adolescents showed an increase in illicit drug use initially, and then they carried those new (higher) levels of drug use with them as they entered older age bands. Figure 5-1 shows the effects of generational replacement as the teens of the early 1990s reached their 20s. While all age groups generally moved in parallel through about 1992, the youngest age bands first showed signs of increase in their overall level of illicit drug use. The 18-year-olds shifted up first, followed by the 19- to 20-year-olds in 1994, the 21- to 22-year-olds in 1996, the 23- to 26-year-olds in 1999, the 29- to 30-year-olds in 2004, and the 35-year-olds in 2008. So far, the 40-, 45-, and 50-year-olds have not shown much increase. (In fact, the 8th graders, who are not included in these graphs, but are described in *Volume I*, actually began an increase in use a year earlier than the 12th graders, showing that the cohort effect was already underway before use turned upward among 12th graders.)

To summarize, in the earlier decline phase of the drug epidemic, annual prevalence of use of *any illicit drug* moved in parallel for all age strata, as illustrated in Figure 5-1; this pattern reflects a secular trend, because a similar change is observed simultaneously across different age levels. After 1992—in what we have called the “relapse phase” of the popular drug epidemic that began in the 1960s—a quite different pattern emerged: 8th graders increased

⁵⁷ Actually, the downturn ended at least a year earlier among 8th graders, who showed the beginning of an increase in 1992 (Table 2-2).

their drug use first, followed by 10th and 12th graders; then the next oldest age group increased use, but with a little delay; the next oldest then increased use, but with a longer delay; and the oldest three groups (40-, 45- and 50-year-olds) are not yet showing an increase. This pattern reflects a classic cohort effect, in which different age groups are not all moving in parallel; rather, different age groups show increases when the cohorts (i.e., high school classes) having heavier use at an earlier stage in development reach the relevant age level. In addition, note that the slopes of the age bands are successively less steep in the older age groups, suggesting that some of the cohort effect may be dissipating with maturation. But we think it unlikely that only cohort effects are occurring (in addition to the long-established age effects); period effects also likely play a role.

- Use of *marijuana* shows an almost identical pattern to the illicit drug use index—not surprising given the fact that marijuana, by far the most prevalent of the illicit drugs, tends to drive the index (Figure 5-3a). After a long and steady decline from the late 1970s to the early 1990s, annual marijuana use leveled for a while among young adults before beginning a gradual increase. Virtually all of this increase was attributable to the two youngest age bands (18 and 19 to 20) until 1996, when the 21- to 22-year-olds began to show a rise. The older age bands then tended to show increases fairly sequentially, with 29- to 30- and 35-year-olds showing significant increases in 2008. The 18-year-olds' use of marijuana in the prior 12 months declined after 1997 and, later, several of the succeeding age bands through age 26 began to show declines in a pattern that again suggests lasting cohort differences. Since about 2006, however, use has been rising not only among the 18-year-olds, but also among all age bands through about 35 years old, thus indicating a secular trend. This strongly suggests an impact on use by culture-wide events to which all of the age bands are exposed.
- A similar pattern emerged for current *daily marijuana* use (Figure 5-3c). In the mid- to late 1990s, daily marijuana use among 35- and 40-year-olds was as high as or higher than use among some younger age groups, suggesting a lasting cohort effect on this behavior, because the cohorts comprising those older age strata grew up in a period of particularly high adolescent marijuana use. However, in more recent years, the 35-, 40-, 45-, 50-, and 50-year-olds were similar to respondents ages 27 to 30, who had among the lowest levels of daily use. An important finding shown in Figure 5-3c is that, although the various age groups had been moving in parallel for many years at fairly similar levels of prevalence, the trends diverged considerably in the 1990s in a staggered fashion, such that the 18- to 30-year-olds came to have distinctly higher levels of daily marijuana use (around 3% to 7% in 2013) than the older age groups (around 2% to 3%), again reflecting stable cohort differences. In 2010 the upturn in daily marijuana use that had been occurring at younger ages (best seen in the table accompanying Figure 5-3c) reached the age-35 stratum, with a significant increase from their 2009 prevalence rate. In 2012 and 2013 there were no significant changes in daily marijuana use for any adult age group.

- The index of using *any illicit drug other than marijuana* has shown a similar transition in the pattern of change. Period effects seemed to predominate until about 1992, but a cohort-related pattern of change emerged thereafter (Figure 5-2). And, while annual use leveled by 1997 among 18-year-olds, it began rising in 1999 among 19- to 20-year-olds, in 2000 among 21- to 22-year-olds, in 2002 among 23- to 24-year-olds, in 2005 among 29- to 30-year-olds, and so on. The primary difference from the picture for marijuana is that the increases were not as sharp in the 1990s for most of the age groups. (Compare Figure 5-2 with Figure 5-1 to see the difference.) Since about 2008 the levels of use in all age groups have remained fairly steady.
- A question about *synthetic marijuana* was added in 2011 (Table 5-2). Annual prevalence was 7.4% among the 19- to 28-year-olds but dropped significantly to 5.0% in 2012 and dropped significantly again to just 3.2% in 2013, reflecting a decline of more than half over just two years.
- In the 1980s and 1990s, *LSD* use also increased among those in their teens and early 20s much more than among the older strata, as Figure 5-6 illustrates. Over the interval 1985 to 1996, there was a gradual but considerable increase in annual LSD use among respondents ages 18 to 24, which was sharpest among 12th graders and 19- to 20-year-olds. The increase did not seem to radiate up the age spectrum beyond age 26. A turnaround began among 12th graders after 1995 and then among the older age groups in a somewhat staggered fashion, again indicative of a cohort effect. Declines in the years since then have been greatest among 18- to 24-year-olds, who had attained the highest rates of LSD use. Use declined considerably from 2001 to 2003 in all age bands (including 8th and 10th graders), and then leveled through 2007 at historically low rates, suggesting that an important secular trend may have set in, quite possibly related to decreased availability of the drug. Since 2007 there has been some evidence of a very gradual increase in use, particularly among the younger age bands.
- Several drug classes exhibited a faster decline in use among the older age groups than among 12th graders during the earlier period of decline in the 1980s (see Figures 5-1 through 5-19c). These included *any illicit drug*, *any illicit drug other than marijuana*, *amphetamines*, *hallucinogens* (until 1987), *LSD* (through 1989), and *methaqualone*, but not *marijuana*.
- In fact, a crossover is evident for some drugs when 12th graders are compared to young adults. In earlier years 12th graders had lower usage levels, but for some years after 1993 they tended to have higher levels than post-high school respondents for use of *any illicit drug*, *marijuana*, *hallucinogens*, *LSD* specifically, *crack cocaine*, *tranquilizers*, and *crystal methamphetamine (ice)*.
- *Cocaine* (Figure 5-9) gives quite a dramatic picture of change. Unlike most other drugs, active use of cocaine has generally tended to rise with age after high school, usually peaking three to four years past graduation. This is a

classic example of an age effect. Despite the large age differences in absolute prevalence, all age strata moved in a fairly parallel way through 1991, indicating that a secular trend was taking place in addition to the age effect. All age strata began a sharp and sustained decline in use after 1986—again reflecting a period effect. The two youngest strata (12th graders and 19- to 20-year-olds) leveled by 1992, whereas use continued a decelerating decline for a few years beyond that in the older age groups, signaling the beginning of a cohort effect. From 1994 to 1999, annual prevalence of cocaine use rose some in the five youngest strata (i.e., those younger than 27) on a somewhat staggered basis, with the three older groups still decreasing a bit more over that same period. This, to some degree, reversed the age differences that were so prominent in the 1970s and 1980s. Cohort-related change appears to have predominated in the 1990s, quite possibly as the result of “generational forgetting” of the cocaine-related casualties so evident in the early to mid-1980s. In other words, those in the older cohorts retained that learning experience, but those in the newer cohorts never had it. The fact that from 1994 to 1996 the 35-year-olds had higher lifetime prevalence levels of cocaine use than some of the younger age groups also suggests some lasting cohort-related differences established during the peak years of the cocaine epidemic. During the past few years there has been some decline in cocaine use among those ages 18 to 28 but less among those 29 years and older, who were at quite low levels already.

- **Crack** use was added to the 12th graders’ questionnaires in 1986 and to the follow-up questionnaires in 1987. The decline in annual crack use, which began right after the introduction of these questions, ended in 1991 among 12th graders, and by 1994 it had ended among young adults (Figure 5-10 and Table 5-2). Among 19- to 28-year-olds, the annual prevalence rate held at about 1%, which was down from the peak levels of just over 3% in 1986 through 1988. As was true for a number of other drugs, crack use began to rise after 1993 among 12th graders, at the beginning of the relapse phase in the epidemic, but not in the older age strata until years later, when increases were observed in a somewhat staggered pattern going up the age scale. Again, a cohort effect due to generational replacement seems to have been occurring. Since 1994, 18-year-olds have had the highest reported rates of use, though they have shown considerable decline since 1999. After another significant decline of 0.4 percentage points in 2011, the 1.0% annual prevalence in 2011 was the lowest since crack use was first measured in 1986, when it was 4.1%. In 2013 use among young adults was at 1.1%. Importantly, all of the other age groups now have annual prevalence rates at or below 0.7%.
- With regard to *inhalants*, the large separation of trend lines for the younger age groups in Figure 5-4 shows that, across many cohorts, use has dropped consistently and sharply with age, particularly in the first few years after high school. In fact, of all the populations covered by MTF, the 8th graders (not shown in Figure 5-4) have had the highest rate of use, indicating that the decline in use with age starts at least as early as 8th or 9th grade. Like cocaine,

inhalants have shown a strong age effect, but unlike cocaine, use of inhalants declines rather than increases with age.

Figure 5-4 also shows that, until the mid-1990s, there was a long-term gradual increase in annual inhalant use (unadjusted for underreporting of nitrite inhalants), one which was greatest among 12th graders, next greatest among 19- to 20-year-olds, and next greatest among 21- to 22-year-olds. Respondents more than six years past high school, who historically have had a negligible rate of use, did not exhibit the increases in use seen among the younger respondents, which began at least as early as 1977 among 12th graders and in 1983 among 19- to 20-year-olds. There was some subsequent increase among 21- to 22-year-olds and later still an increase among 23- to 24-year-olds. After 1995, this long-term trend, reflecting a cohort effect, began to reverse in the two youngest age strata (coincident with an anti-inhalant media campaign by the Partnership for a Drug-Free America) as well as among several other age strata, suggesting a period effect. Subsequently, further declines among several age strata are suggestive of a cohort effect. The older age strata have generally shown very low rates of inhalant use throughout the course of the study.

- The annual prevalence for *ecstasy* (*MDMA*) use among the entire young adult sample (ages 19 to 28) was at about 1.5% in 1989 and 1990 (Table 5-2 and Figure 5-8). After 1991 it dropped to around 0.8% for several years before starting to rise significantly in 1995. Ecstasy use then began to rise in all of the young adult age strata, most notably in the younger age bands (19 through 26) through 2001. Use among 12th graders, which was not measured until 1996, was by then the highest of any of the age groups at 4.6% annual prevalence. Twelfth graders' use declined by a full percentage point through 1998 before jumping significantly—by two full percentage points—in 1999. (Use by 10th graders also jumped significantly in 1999.) Thus it appears that young people from their mid-teens to mid-20s discovered ecstasy after some years of low and relatively level use. In 2000 the sharp increase in use continued among ages 15 to 16 (10th graders) through age 26, with highs of over 10% among 19- to 22-year-olds, and also showed up among 8th graders for the first time. By 2001 the increase had slowed and even begun to reverse among ages 18 to 26, even as the 31- to 32-year-olds showed their first appreciable increase in ecstasy use. We attributed the deceleration in 2001 to a fairly sharp increase in perceived risk of ecstasy use in that year, and predicted a turnaround in use in 2002. In 2002, and again in 2003, perceived risk increased sharply and, as Figure 5-8 illustrates, all age bands showed a reversal, with a sharp decrease in use. Clearly, the decrease has been sharpest in the younger age bands, perhaps because a cohort effect is at work in the upper ages, helping to offset a downward secular trend. In 2013 annual ecstasy (*MDMA*) prevalence remains at 5 to 6% for 19- to 24-year-olds and between 3.3 and 1.6% for those 25 to 30 (Table 4-3).
- A tripwire question asking about use of *salvia* in the past 12 months has been included in the study since 2009. Annual use of *salvia* has declined steadily

among the 19- to 28-year-olds, from 3.5% in 2009 to 0.9% in 2013 (Table 5-2).

- In the late 1970s, *amphetamine* use rose some with age beyond high school; but after a long period of secular decline in use from 1981 to the early 1990s, this relationship had reversed (see Figure 5-13). The declines were greatest in the older strata and least among 12th graders, even though use decreased substantially in all groups. As was true for many illicit drugs, amphetamine use began to rise among 12th graders after 1992, and eventually among the 19- to 24-year-olds; but there has only recently been a small increase among 25- to 30-year-old respondents. In other words, another cohort-related pattern of change seems to have emerged in the 1990s for amphetamines, though in this case it may be dissipating quickly after respondents reach their early 20s. While amphetamine use declined a fair amount among 12th graders between 2002 and 2009 (from 11.1% to 6.6%), there was less proportional decline in the older age groups, and little or no decline among those ages 29 to 35. After 2009 there was some resurgence in use, particularly among the younger age groups. At present the age differences in amphetamine use through age 55 are of considerable magnitude and mostly ordinal (with 18- to 24-year-olds showing the highest and mostly undifferentiated rates of use and those over 30 much lower rates).
- Since 1990, when it was first measured, use of *crystal methamphetamine (ice)* has remained at fairly low rates in this young adult population (Figure 5-14). However, among 19- to 28-year-olds combined, annual prevalence rose from 0.4% in 1992 to 1.6% by 2005, before falling back to 0.8% in 2013 (Table 5-2). Use had been rising among 12th graders and 19- to 20-year-olds specifically between 2000 and 2002, but since then their use has declined to low levels. General *methamphetamine* use was first measured in 1999, and its use until 2005 was stable among 19- to 28-year-olds, with annual prevalence fluctuating between 2.4% and 2.8%. Use has declined since to 0.6% by 2013 (Table 5-2).
- Use of *heroin* increased appreciably in 1995 among 12th graders and young adults ages 19 to 24, but not among the older age bands (Figure 5-11). It remained at this higher plateau in these younger age bands through 2000 or 2001, before falling off some, particularly among 12th graders. Among young adults, generally, annual use had previously been quite stable from at least as far back as 1986 through 1994 (Table 5-2), and it stabilized again at a higher level after 1994—a level roughly twice as high as the previous one. Use among 12th graders has declined since 2000, among 19- to 20-year-olds since 2001, and among the 21 to 22-year-olds since 2006, but remains fairly stable (at a very low rate of use) among the older age groups.
- Among 19- to 28-year-olds, use of *narcotics other than heroin* leveled after 1991, following a long period of slow, fairly steady decline (Figure 5-12 and Table 5-2). After 1992 twelfth graders showed an appreciable increase in use, which continued into 2004, while 19- to 20-year-olds showed some increase

after 1994, 21- to 22-year-olds after 1996, 23- to 24-year-olds after 1997, and the older age groups after 2000. Thus, cohort-related change appears to have been occurring during the 1990s and beyond for this class of drugs as well, following a long period of secular trends. In 2002, the question text was changed on three of the six questionnaire forms to update the list of examples of narcotic drugs other than heroin. Talwin, laudanum, and paregoric, each of which had negligible rates of use by 2001, were replaced by Vicodin, OxyContin, and Percocet. As a consequence of this revision, reported use rates increased in 2002. Data presented here for 2002 are from three of the six questionnaire forms with the new wording (which showed higher prevalence rates than the older question did). All six questionnaire forms contained the new wording beginning in 2003, so the data presented for 2003 onward are based on all forms. Although the older version of the question showed no significant changes occurring in 2002, there was a significant increase in narcotics use observed in 2003 (based on the new question in both 2002 and 2003). Among 19- to 28-year-olds, annual prevalence reached a peak level of 9.1% in 2006; it has since fallen some, to 7.0% in 2013. Some turnaround was observed among 19- to 22-year-olds after 2004 in the use of this important class of drugs, but use continued to rise in some of the older age bands through 2007 to 2009, likely reflecting a cohort effect. Use of these drugs outside of medical supervision remains relatively high in all age groups studied here, with rates of around 7% among those ages 18 to 30, also about 7% among those age 35, and 2.5% to 4% among those ages 40 to 55. The most important finding is that use of this important class of drugs has either leveled or declined in recent years in all of the age groups covered in the study.

- The annual prevalence rates for *Vicodin* and *OxyContin*, which were first measured in 2002 (separately from the general question about narcotics other than heroin), were appreciable (8.2% and 1.9%, respectively) for 19- to 28-year-olds. Increases were observed for these two drugs in subsequent years. Among 19- to 28-year-olds (Table 5-2), the annual prevalence of *OxyContin* use rose from 1.9% in 2002 to 3.1% in 2004 through 2006—changes that were fairly parallel to those observed among 12th graders over the same interval (when their annual prevalence rose from 4.0% in 2002 to 5.5% in 2005). The increases in *OxyContin* use between 2002 and 2005 were significant for both 12th graders and 19- to 28-year-olds. Annual prevalence was stable from 2004 to 2007 at about 3% for young adults, increased to 5.2% in 2009, but is down to 2.8% in 2013. *Vicodin* use rose by less, but started from a higher base, with annual prevalence increasing slightly among 19- to 28-year-olds, from 8.2% in 2002 to 8.9% in 2004; it remained at about 9% through 2009, followed by a decline to 6.2% by 2013. Thus, since 2009 the annual prevalence of *OxyContin* use among young adults has declined by about half, while the annual prevalence of *Vicodin* use has declined by about one third. Given the widespread concern about these narcotic drugs, which are among those most cited in overdose deaths, this downturn is very good news.

- **Sedative (barbiturate)** use (Figure 5-15) showed a long-term parallel decline in all age groups covered through the late 1970s and 1980s, leveling by about 1988. While use remained low and quite level for most of the age bands for about five years, it began to rise by 1993 among 18-year-olds, by 1995 among 19- to 20-year-olds, by 1997 among 21- to 22-year-olds, by 1998 among 23- to 24-year-olds, by 2001 among 25- to 28-year-olds, and by 2005 among 29- to 30-year-olds. The same cohort-related pattern of change seen during the 1990s for many other drugs also exists for sedatives (barbiturates); like most other drugs, this pattern was preceded by a long period of secular change during which all age-groups moved in parallel. While use leveled off among most age groups by 2005, the 35-, 40-, and 45-year-olds all showed increases in sedative (barbiturate) use between 2006 and 2008. However, their usage rates leveled after 2008. In 2013 the annual usage rates for the 35- to 55-year-olds were about 2% to 3%. Sedative use among 18-year-olds declined steadily since 2005, among 19- to 20-year-olds since 2008, and among 21- to 22-year-olds since 2009, suggesting another cohort effect. In 2012 the usage rates leveled off for the 18- to 22-year-olds.
- **Tranquilizers** (Figure 5-16) follow a similar pattern to that just described for sedatives (barbiturates). One difference is that the 12th graders' annual prevalence rate has not always been the highest among the various age groups, as was the case for sedatives (barbiturates), although it was highest between 1994 and 2000 as a result of a greater increase in tranquilizer use among the 12th graders than in the young adult strata. Since about 2004, however, as use rose and then leveled among those in their early 20s, the 12th graders no longer stand out as having the highest rate of tranquilizer use. In fact, the 21- to 22-year-olds or 23- to 24-year-olds had the highest rate in 2005 through 2009; in 2011, the 25- to 26-year-olds had the highest rate; and in 2012 the 27- to 28-year-olds had the highest rate. This was another clear example of a cohort-related pattern of change. Now that tranquilizer use has leveled and begun to decline among 12th graders and 19- to 22-year-olds, it is likely that we will see this pattern echoed in the older age groups as a new cohort effect plays out.
- Use of **anabolic steroids** has been substantially lower after high school than during 12th grade (Figure 5-17), ever since measures were first introduced in 1991 (in two follow-up questionnaire forms). The age-related differences are not consistent; prevalence rates among the young adult strata are all quite low and do not appear to trend in any systematic way. In general, it seems that the rise in steroid use from 1999 to 2003 among 8th and 10th graders and from 2001 to 2004 among 12th graders was largely specific to those age groups. Annual prevalence rates are now very low for respondents in all young adult strata of ages 19 and older (ranging from <.05% to 0.7%).
- **Alcohol** trends for the older age groups (see Figures 5-18a-d) have been somewhat different than for the younger age groups in some interesting ways. For **30-day** prevalence and **occasions of heavy drinking**, the declines for the two youngest age strata (12th graders and those one to two years past high

school) during the 1980s were greater than for the older age groups. These differential trends were due in part to the effects of changes in minimum drinking age laws in many states, changes that would be expected to affect primarily the age groups under age 21. However, because similar (though weaker) trends were evident among 12th graders in states that maintained a constant minimum drinking age of 21, the changed laws cannot account for all the downward trends, suggesting that there was also a more general downward trend in alcohol consumption during the 1980s.⁵⁸ By 1994, these declines in 30-day prevalence had slowed or discontinued for virtually all age groups until 1997, when they began to turn downward again for 12th graders, and 1999, when they began to decline among 19- to 20-year-olds. The long term declines in the 30-day prevalence of alcohol use have been very substantial—from 72% in 1980 to 39% in 2013 among 18-year-olds, and from 77% in 1981 to 52% in 2013 among 19- to 20-year-olds. Since about 1997, as the declines continued in the under-21 groups (that is, those under the minimum legal drinking age), no such declines occurred among the 21 and older groups; in fact, there have been some modest increases in all these groups. These trends have resulted in substantial differences in 30-day drinking rates between those 18 to 20 years of age (40% to 52%) versus those 21 and over (62% to 76%).

- Those respondents three to four years past high school show the smallest downward trend in *occasions of heavy drinking* or *binge drinking* (i.e., having five or more drinks in a row at least once in the past two weeks) since the early 1980s (see Figure 5-18d). One important segment of that age stratum is composed of college students, who have shown less decline in alcohol use over the past quarter century (see chapter 9). In 2013 the binge drinking rate for 19- to 20-year-olds (who are, of course, under the legal age for alcohol possession and, in many states, consumption) is considerably lower (27%) than for those just one to two years older (40%).

The older age groups generally showed no recent decline in binge drinking rates. Note that the binge drinking trend lines for different age groups (Figure 5-18d) are spread out on the vertical dimension, reflecting large and persisting age differentials (age effects) in this behavior. The relationship with age is curvilinear, however. In recent years the 21- to 24-year-olds have consistently shown the highest rates of binge drinking, and the 25- to 26-year-olds have just about joined them after some years of increasing use going back to the mid-1990s. Binge drinking also had been gradually increasing since the early 2000s among 25- to 30-year-olds, perhaps reflecting a cohort effect that emerged during the period of increasing adolescent binge drinking in the early 1990s.

From the early 1980s through the mid-1990s, rates of *daily drinking* (Figure 5-18c) fell by considerable proportions in all age strata for which we have

⁵⁸ O'Malley, P. M., & Wagenaar, A. C. (1991). Effects of minimum drinking age laws on alcohol use, related behaviors, and traffic crash involvement among American youth: 1976–1987. *Journal of Studies on Alcohol*, 52, 478–491.

data, reflecting an important change in drinking patterns in the culture. Among 19- to 28-year-olds combined, daily drinking declined from 1987 (6.6%) to 2000 (4.1%), and has increased a little to 5.1% by 2013 (Table 5-4). Daily drinking rates have generally been highest for the 40 and older groups in recent years; and they have declined substantially among 18-year-olds and 19- to 20-year-olds over the life of the study. Despite these declines for young adults up through age 20, daily drinking among 21- to 22-year-olds (which also declined from the early 1980s through the mid-1990s) has increased some since 1995. In 2013 there is a considerable difference among the age strata in rates of daily drinking, ranging from 2% among 18-year-olds to 11% among 50- and 55-year-olds.

It is worth noting that the 35-, 40-, 45-, 50-, and 55-year-olds have had among the lowest rates of binge drinking but among the highest rates of daily drinking in recent years for which we have data available. These patterns—particularly the high rate of daily drinking—likely reflect age effects as well as perhaps some enduring cohort differences (because these cohorts had considerably higher rates of daily drinking when they were in high school). They may also reflect medical findings that suggest that one or two drinks per day for males and one per day for females have some benefits for heart health.^{59,60} That may be a more salient message for those who are in their forties or above than for younger people.

Questions about *extreme binge drinking*—having 10 or more or 15 or more drinks on at least one occasion in the two weeks prior to the survey—were added to the MTF questionnaires in 2005. Because these questions are included in only a single questionnaire form, the annual samples are relatively small for making accurate change estimates. However, it appears from the trend data that are available for the 19- to 28-year-olds that the 10+ drinks measure may have peaked around 2008 to 2009 (at about 12.8%) and has been gradually declining since then (to about 10.2% in 2012 and 2013 combined). The 15+ drinks measure averaged about 5.0% from 2005 to 2008—a period of fairly level use—and then dropped to an average of 4.1% in 2012 and 2013 combined.

- The prevalence rates for *cigarette smoking* show more complex trends than most other substances, due to the long-term presence of both cohort and age effects, plus slightly different patterns of such effects on the several different measures of smoking during the past 30 days (one or more cigarettes per month, one or more cigarettes per day, and a half pack or more of cigarettes per day).

⁵⁹ Manttari, M., Tenkanen, L., Alikoski, T., & Manninen, V. (1997). Alcohol and coronary heart disease: The roles of HDL-cholesterol and smoking. *Journal of Internal Medicine*, 241, 157–63.

⁶⁰ Savolainen, M. J., & Kesaniemi, Y. A. (1995). Effects of alcohol on lipoproteins in relation to coronary heart disease. *Current Opinions in Lipidology*, 6, 243–50.

In the earlier years of MTF, the curves across time were of the same general shape for each age band (Figures 5-19a–c), but each of those curves tended to be displaced to the right of the immediately preceding age group, which was two years younger. The pattern is clearest in Figure 5-19c (half pack plus per day). This pattern is very similar to the one described in *Volume I* for lifetime smoking rates for various grade levels *below* senior year; it is the classic pattern exhibited by a cohort effect,⁶¹ and we believe that the persisting cohort differences are due to the dependence-producing characteristics of cigarette smoking.

The declining levels of cigarette smoking across cohorts at age 18, which were observed when the classes of 1978 through 1981 became 12th graders, were later observable in the early-30s age band, as those same high school graduating classes grew older (see Figures 5-19b and c). This was true at least through about 1991. By then there had been a considerable convergence of rates across age groups, largely because there were few cohort differences among the senior classes who graduated from the early to mid-1980s through the early 1990s—a period of fairly level cigarette use in high school.

In addition to these cohort differences, there are somewhat different age trends in which, as respondents grow older, the proportion smoking at all in the past 30 days declines some, while the proportion smoking a half pack per day actually increases. Put another way, many of the light smokers in high school either transition to heavier smoking or quit smoking.⁶²⁻⁶⁴

The picture was further complicated in the 1990s when it appears that a new cohort effect emerged, with smoking among adolescents rising sharply (beginning after 1991 for 8th and 10th graders and after 1992 for 12th graders). The 19- to 20-year-olds soon showed a rise at the beginning of the 1990s—perhaps responding to some of the same social forces as the adolescents (including the Joe Camel advertising campaign); but 21- to 24-year-olds did not show an increase until about 1995, and 25- to 26-year-olds until about 1996. Those young adults over age 26 showed a modest increase from 1997 through 2004, but a decline in use since then; it is quite possible

⁶¹ O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321.

⁶² To illustrate, in the graduating class cohort of 1976, 39% were 30-day smokers in senior year, 39% by ages 19 to 20, but only 28% by ages 31 to 32—a net drop of 11 percentage points over the entire interval. By way of contrast, 19% of that class was half-pack-a-day smokers in senior year, 24% by ages 19 to 20, and 21% at ages 31 to 32—a net gain of five percentage points and two percentage points over the respective intervals.

⁶³ Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁶⁴ Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates

that an upward cohort effect was at least partially offset by a downward secular trend during this period.

After about 1999, smoking rates among nearly all age groups leveled or declined, suggesting that societal forces may be affecting all age groups in a similar way, giving rise to a secular trend. Large increases in the price of cigarettes and a great deal of adverse publicity for the tobacco industry are highly plausible candidates, as are the introduction of the national anti-smoking campaign of the American Legacy Foundation, an increase in state and national antismoking advertising, the demise of the Joe Camel campaign and all billboard advertising, and the imposition of no-smoking regulations in many public and workplace settings by states and municipalities. Since 2003, thirty-day, daily, and half-pack smoking have all declined among 35-, 40-, and 45-year-olds (Figures 5-19a through 5-19c).

- Apart from cigarettes, none of the other drugs included in the study showed a clear long-term pattern of enduring cohort differences in the earlier years of MTF (the 1970s and 1980s), despite wide variations in their use by different cohorts at a given age. There was one exception; a modest cohort effect was observable for *daily marijuana* use (Figure 5-3c) during the late 1970s and early 1980s.⁶⁵ (But as subsequent classes leveled at lower rates of use, evidence for the cohort effect faded.) The emergence in the 1990s of a new epidemic of marijuana use among teens once again yielded a strong pattern of cohort effects. As can be seen in Figure 5-3c, daily use rose sharply among 12th graders and 19- to 20-year-olds after 1992, among 21- to 22-year-olds after 1993 with a sharp rise occurring in 1997, among 23- to 24-year-olds after 1998, among 25- to 26-year-olds after 2000, among 27- to 28-year-olds in 2003, among 29- to 30-year-olds in 2005, among 35- and 40-year-olds in 2006, and among 45-year-olds in 2007. This is not unlike the pattern of change for cigarette smoking that occurred in the 1990s (Figure 5-19a). The cohort effect for daily marijuana use may be attributable, in part, to the very strong association between that behavior and regular cigarette smoking. The net effect of all of this is that a considerable age difference has emerged in current daily marijuana use since the early 1990s, when there was practically no difference. The cohort effect resulting from the rise in use among 18-year-olds in the latter half of the 1990s has been working its way up the age spectrum, and in 2010 was observable in the form of a significant increase among 35-year-olds. In 2013, among the 40-, 45-, 50-, and 55-year-olds in the study, 2.1–2.7% reported that they currently smoked marijuana on a daily basis (though many more of them did so in the past) while among those ages 19 to 24, between 6.2% and 7.8% did so.

⁶⁵ O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321.

- In sum, except for *cigarettes* and *alcohol*, substance use prior to 1992 among 12th graders and young adults had shown longer term trends that were highly parallel across age groups, indicating that general secular trends predominated in that period. Since 1992, however, there has been considerable divergence in the trends for different age bands on a number of drugs as use among adolescents rose sharply, followed by subsequent rises among 19- to 20-year-olds, 21- to 22-year-olds, and so on. This divergence indicates a new cohort effect, quite possibly reflecting a generational forgetting of the dangers of drugs by the cohorts who reached senior year in the early to mid-1990s. Data discussed in chapter 6, “Attitudes and Beliefs about Drugs among Young Adults,” provide additional evidence for this interpretation.

The resurgence of illicit substance use in the past few years among secondary school students is showing up among the age bands 19–28, but generally not those above age 28.

TRENDS FOR IMPORTANT SUBGROUPS OF YOUNG ADULTS

Four-year age bands have been used here to examine subgroup trends in order to yield sufficiently large numbers of cases to permit reliable estimates for the various subgroups being examined. Subgroup data for respondents of each gender and for respondents from communities of different sizes are available for 19- to 22-year-olds since 1980, 23- to 26-year-olds since 1984, and 27- to 30-year-olds since 1988. Beginning in 1987, a question about state of residence was added to all follow-up questionnaires, permitting trend data to be calculated for the four regions of the country since then. These various subgroup data are not presented in tables or figures here because of the substantial amount of space they would require. However, for the reader interested in more detail, these are available in a separate *MTF Occasional Paper 80* at <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ80.pdf> . A verbal synopsis of this information is presented below.

Gender Differences in Trends

- Over the long term, gender differences narrowed for some drugs among young adults in each of these three age bands, primarily when a steeper decline in use among males (who generally had higher rates of use) occurred in the 1980s. The overall picture, though, is one of parallel trends, with use among males remaining consistently and modestly higher for most drugs, including the indexes of *any illicit drug* use and use of *any illicit drug other than marijuana* in the prior year (see Table 5-5, for example).

The downward trend in *marijuana* use among 19- to 22-year-olds between 1980 and 1989 was also a bit sharper among males than females, narrowing the gap between the two groups. Annual prevalence fell by 22 percentage points (to 34%) among males, compared to a drop of 14 percentage points (to 31%) among females, leaving a difference of three percentage points. Since 1995, the gender gap has averaged about 5 to 9 percentage points in all three

age groups—that is, for 19- to 22-year-olds, 23- to 26-year-olds, and 27- to 30-year-olds.

Similarly, between 1980 and 1993, *daily marijuana* use for the 19–22 age group fell from 12.9% to 2.9% among males, and 6.1% to 1.7% among females, narrowing the rather large gap that existed in the early 1980s. As use rose after 1993, the gap widened again. Among 23- to 26-year-olds, as daily use first began to increase in 1998 and 1999, the gap between the genders began to widen. In the oldest age group (ages 27–30), the difference had been fairly constant, with daily marijuana use among males generally being about two percentage points higher than among females.

- In all three age bands, use of *synthetic marijuana* by males tends to be higher than use by females. In 2011, when use was first measured, it was highest among the 19- to 22-year-olds with males higher than females; but it has fallen sharply over the past two years for both genders and closing the gap between them considerably. Annual prevalence in 2013 for the 19-22 age group was 4.5% for males and 2.5% for females. The two older age bands started out with considerably lower rates in 2011, but also have shown some decline since, narrowing the gender difference among 27- to 30-year-olds considerably (to 2.3% versus 1.3%).
- For *LSD*, males have consistently had higher rates of use than females. Among 19- to 22-year-olds, the male–female differences tended to diminish as use declined (from 1980 to 1985 and again from 1999 to 2004) and expand as use increased (1986–1995). In the two older age bands there was less change in use, and differences had been relatively consistent (with males higher) since data have been available, beginning in 1984 for 23- to 26-year-olds and in 1988 for 27- to 30-year-olds. After 1999 and 2001 for the two groups, respectively, LSD use dropped, substantially narrowing the gender differences. Males began to show these declines first, and both genders have moved to almost no use of LSD since about 2003.
- Use of *hallucinogens other than LSD* taken as a group has consistently been higher among males in all three age strata with the difference growing larger when use increased some in the late 1990s and early 2000s.
- *Ecstasy (MDMA)* exhibited little or no gender difference in any of the three age bands before use began to grow in the late 1990s. Between then and 2009, there was little gender difference in ecstasy use among 19- to 22-year-olds. In recent years, males have had slightly higher rates of use. In the older age groups a gender difference opened up after 1997, with males fairly consistently having higher rates of use among both 23- to 26-year-olds and 27- to 30-year-olds.
- The annual prevalence of *salvia* use was much higher among males in the 19- to 22-year-olds when first measured in 2009, and somewhat higher in the two older age groups. However, use by males has dropped dramatically in the

years since then; there is no gender difference as of 2013, even though use by females also dropped some during that interval in the two younger age groups.

- Males have had higher rates of *cocaine* use than females since MTF began. During the period of sharp decline from the peak levels in annual cocaine prevalence (1986–1993), use dropped more among males than females, narrowing the gender differences that existed. In the 19- to 22-year-old age band, annual prevalence for males declined by 16 percentage points (to 4.5%) versus 13 percentage points among females (to 2.8%) by 1993. In the 23- to 26-year-old age band, there was also a narrowing of the gender difference between 1986 and 1993, with annual prevalence down 19 percentage points among males (to 6.9%) and 13 percentage points among females (to 4.2%). Use in the 27- to 30-year-old group also dropped faster among males between 1988 (when data were first available) and 1997—down 13 percentage points versus 7 among females. In sum, during the period of sharp decline in overall cocaine use, the gender differences—which had been fairly large—narrowed considerably in all three of these age bands. During the resurgence in cocaine use of the 1990s and into the early 2000s, which has occurred on a somewhat staggered basis over the years, the gap between genders expanded only slightly. In 2013 there were only modest differences in cocaine use between the genders.
- *Crack* followed a similar pattern during the earlier period of decline, though the proportional difference between the two genders has consistently been higher than for cocaine overall. With crack, though, there was some gender convergence (between 1992 and 1998) among 19- to 22-year-olds, as use among males declined slightly and use among females rose gradually. Since 1999, there has been no consistent change in differences between males and females. In all three age bands, males consistently had slightly higher crack usage rates, at least until a greater decline among males in recent years has nearly eliminated the differences.
- During the period from 1982 through about 1994, there was little gender difference in annual prevalence for *heroin* use. (By 2008, the small gender difference that had existed in the oldest age band had disappeared.) After 1994, use increased some among the younger groups, with males generally attaining slightly higher rates than females. A similar small increase, slightly higher for males, happened among the 27- to 30-year-olds between 2000 and 2007 but the gender differences have diminished in recent years.
- Among 19- to 22-year-olds, both genders showed some decline in their use of *narcotics other than heroin* between 1980 and 1991, with a near elimination of previous gender differences (males had been higher). Beginning in 1994, use by males began to rise in this age group, while use by females began to rise a year later. Some gender differences developed as use increased, with use by males being higher; after 2006, as use declined, there was less of a difference, with annual prevalence in 2012 at 7.4% for males

and 6.3% for females. In 2013 there was a bump up among males, appearing to reinstate the gender difference. The picture for 23- to 26-year-olds is very similar except that the increase in use occurred a few years later and there was no divergence between the genders in 2013. The gender difference (males higher) had been eliminated by 1988, but re-emerged after 1995 as use increased more among males. Among 27- to 30-year-olds, there has been a smaller gender difference and the least increase in use in the 2000s. Still, use increased for both genders after 1999, with males emerging with modestly higher rates of use.

- Since 2002, the first year in which *OxyContin* data were gathered, the use of OxyContin has generally been higher among males than females for all three age bands. Both genders showed some increase in use between 2002 and 2009 or 2010, followed by some fall-off since then, though there was an increase observed among males in both of the younger age bands in 2013, the 2.6 percentage point increase among 19- to 22-year-olds being statistically significant.
- *Vicodin* use, first measured in 2002, also has been higher among males in most years. There was more of an increase in use among males in all age bands, initially; but the males began to trend down earlier than the females, reducing the disparities in use such that by 2012 the gender difference was virtually eliminated among adults ages 23 to 30. However, as with OxyContin, there was a significant increase in 2013 of 3.9 percentage points among males in the 19- to 22-year-old age band, reopening a gender difference.
- In general, there have been no appreciable gender differences in *amphetamine* use for some years in any of these three young adult age bands. Between 1981 and 1991, rates of amphetamine use were similar for males and females and showed substantial and parallel downward trends for both genders. Among 19- to 22-year-olds, annual prevalence of use dropped 22 percentage points for males (to 5.2% in 1991) and 21 percentage points for females (to 4.7% in 1991). There were small increases in annual prevalence for both genders in the 19- to 22-year-old age group after 1991, in the 23- to 26-year-old age group after 1995, and in the 27- to 30-year-old age band after 2000, but the genders diverged only slightly (with males higher); amphetamine use has been drifting up slowly in all three age bands since about 2008, with males emerging slightly higher than females, particularly among the 19- to 22-year-olds.
- Nonmedical use of *Ritalin*, a prescription stimulant used in the treatment of ADHD, was added to MTF questionnaires in 2002. Its use generally has been slightly higher among males than females, but the differences nearly disappeared by 2011. The gap reappeared after 2011 among the 19- to 22-year-olds, as use by males increased.

- Like Ritalin, nonmedical use of *Adderall* (another prescription stimulant) has generally been slightly higher among males than females since the question was added in 2009 with the largest difference being among 19- to 22-year-olds, which includes most of those in college. The gender differences generally have been small among those over age 22, though a significant decline among 26-year-old females in 2013 enlarged the gap.
- A question on *methamphetamine* use was introduced in 1999. The youngest age stratum had the highest rate of use for both males and females but also showed the greatest decline in the years since then. By 2011 their rates of use were at or below 1.0% among males and females in all three age bands, and have been 1.8% or less since then.
- *Crystal methamphetamine* (“ice”) was added to the study’s coverage in 1990. In the early 1990s, use was low and very similar for both genders in all three young adult age bands. In the mid-1990s the younger two age bands showed a greater increase in use among males—opening a gender gap. The gap then narrowed, though males on average were slightly more likely to report use of crystal methamphetamine until 2005. Since 2009 the gender differences have been small. It should be noted that the estimates are a bit unstable for this drug due to limited sample sizes.
- Questions about the use of “*bath salts*”—stimulant designer drugs meant to mimic the effects of amphetamines—were first introduced in 2012, so there are as yet very little data on trends in their use. Among 19- to 22-year-olds in 2012 there was a large gender difference in use (annual prevalence of 3.0% among males vs. 0.5% among females); however, there was virtually no gender difference in the two older age bands (0.7% vs. 0.6% respectively among 23- to 26-year-olds and less than 0.5% for both genders among 27- to 30-year-olds). In 2013 the large gap between the genders among the 19- to 22-year-olds disappeared as males that age showed a significant 2.8 percentage point decline in use. This decline coincided with a dramatic 18 percentage point increase in the perceived risk of trying bath salts. A similar change in perceived risk occurred among both older groups, as well, no doubt serving to hold their usage rates very low.
- As *sedative (barbiturate)* use declined through the 1980s, the modest gender differences (males were higher) were virtually eliminated in all three age bands. Beginning in the early 1990s, a staggered increase in use by both genders emerged across all three age groups, with males increasing more than females, thereby again opening a small difference in the late 1990s and 2000s. After declines in the past few years, the differences have essentially been eliminated.
- For *tranquilizers*, both genders showed a long, gradual decline (and very similar rates of use) from 1980 through about 1993 in all three age bands. Beginning in 1995, use increased for both genders in the 19- to 22-year-old group, followed by an increase beginning after 1997 among 23- to 26-year-

olds and after 1999 among 27- to 30-year-olds, again reflecting cohort effects driven by generational replacement. Some gender difference emerged during these periods of increase and during the subsequent decrease after 2002 and 2003 for the two younger age bands. Males generally reported somewhat higher usage rates, though the differences have tended to narrow in recent years. Among the 27- to 30- year-olds, males generally had somewhat higher rates from 2004 to 2009.

- **Inhalant** use has generally been quite a bit higher among males than females, particularly in the younger age groups. The 19- to 22-year-old group showed a gradual upward shift from 1980 to 1988, followed by a leveling for some years for both genders. In 1997, female inhalant use began to decline among 19- to 22-year-olds, followed by males in 2001; however, the gender gap did not diminish much with this decline until 2005, when there was a convergence. Among 23- to 26-year-olds the gender gap widened as use by males increased between 1992 and 1999, though a subsequent decline in use among males narrowed the gap, almost eliminating it by 2005. It then re-emerged by 2008. In the oldest age stratum, use among males has consistently been slightly higher, though the prevalence of inhalant use is very low by this age.
- Use of three “club drugs”—*GHB*, *ketamine*, and *Rohypnol*—has tended to be a bit more concentrated among males in all three age strata. But here again, the estimates are not very stable because of the limited numbers of cases upon which they are based. By 2009, prevalence rates were very low for all three drugs, and gender differences were small; this continued to be the case in the years since then. Rohypnol was dropped from the study in 2010 because of the low numbers of users, at which point no gender difference remained in any of the three age groups.
- For **alcohol, 30-day prevalence** rates exhibited a gradual, parallel decline from 1981 through 1992 for both genders in the 19- to 22-year-old age group. Thirty-day prevalence fell from 83% to 72% among males and from 75% to 62% among females by 1992. There has been a convergence since then, because use by males has declined slightly while use by females increased slightly through 2008, before dropping a bit. The increasing proportion of women attending college may help to explain this convergence at least in part. The gender difference was virtually eliminated in this age group by 2004 and use has remained quite level since for both genders. In the two older age bands, there was a more modest, parallel decline for both genders, from 1985 through 1992 in the case of 23- to 26-year-olds, and at least from 1988 (when data were first available) to 1991 or 1992 in the case of 27- to 30-year-olds. After 1992, use among males in the older two age bands showed fairly level rates of use; but use among females rose gradually, almost eliminating the difference among 23- to 26-year-olds and narrowing the difference somewhat in the oldest young adult group.

There was also a general long-term decline in *daily drinking* from about 1981 or 1982 through about 1992, with daily use falling more among males, considerably reducing, but far from eliminating, what had been a large gender difference among 19- to 22-year-olds. To illustrate, in 1981, 11.8% of males reported daily use versus 4.0% of females; the comparable 1992 statistics were 5.3% and 2.7%. After 1995, daily drinking began to increase among 19- to 22-year-olds for both genders, but leveled a few years later. From 2002 to 2005 daily use was rising among males and falling among females, increasing their differences, but since 2005 there has been a considerable convergence with daily use among males falling and use among females increasing modestly. In 2013 there still is a gender difference for daily drinking among the 19- to 22-year-old age group—5.6% for males versus 2.4% for females—but not nearly as large as it was in 1981 (11.8% vs. 4.0%). The gender differences have been larger for the older age groups with little sign of convergence. In fact, among the 27- to 30-year-olds the gender difference has increased since 2000, with use rising among both genders, but to a greater extent among males; in 2013 their daily drinking rates were 10.4% for males vs. 3.6% for females.

There are also long-established and large gender differences in all age groups in the prevalence of *occasions of heavy drinking*. Males in the 19- to 22-year-old band have shown some longer term decline in this statistic, from 54% in 1986 to 45% in 1995 to 39% in 2013. Use by females declined less, from 33% in 1981 to 28% in 1995 before rising some to 34% in 2006, and then back to 29% in 2013. Thus, the gender gap has narrowed (from 24 percentage points in 1986 to 17 percentage points in 1995 to 10 percentage points by 2013). In the two older age bands (23- to 26-year-olds and 27- to 30-year-olds), the sizable gender differences remained mostly stable as the binge drinking rates drifted steadily upward in both genders since the early 1990s. However, these rates have leveled or even declined a bit in both genders among 19- to 22-year-olds for about the past six years, among 23- to 26-year-olds over the past four years, and among the 27- to 30-year-olds in the past three years. The sustained increase for 23- to 26-year-olds up through 2009 is striking, and may be due at least in part to the increased rate of college attendance and decreased rate of marriage among those in this age group.^{66,67}

- Most striking for *cigarette smoking* by young adults are the similarities between the genders in both absolute levels and trends. All three age groups showed a long-term decline in *daily smoking* rates for both males and females after data were first available for each: 19- to 22-year-olds from

⁶⁶ Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁶⁷ Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates.

1980 to 1990, 23- to 26-year-olds from 1984 to 1992, and 27- to 30-year-olds from 1988 to 1999—again reflecting a cohort effect moving up the age scale. Male and female **30-day smoking** rates have also been very close over most of the time for which data are available, particularly in the two older age groups. Among 19- to 22-year-olds there was a crossover after 1993—before that point, females had slightly higher 30-day prevalence rates, whereas males did from 1994 onward, primarily because use was rising faster among males through 1999. Both genders in this age group have shown considerable declines since 1999. Among 23- to 26-year-olds, the genders had very similar smoking rates until males started reporting higher 30-day smoking rates from 1998 on. Males declined less after 1998, opening up a modest gap; however, this gap has narrowed over the past five years as smoking declined more among males. In the oldest age band, the two genders were quite close until males opened the gap in 2002, and their rate has generally remained somewhat higher since then; in fact, the gap has widened some as use among females has declined more than among males. Indeed, in recent years smoking among females in all three age bands has declined more than has smoking among males. In 2013 the rates for the oldest age band are 22% among males and 16% among females.

Regional Differences in Trends

The respondent's current state of residence was first asked in the 1987 follow-up survey; thus trend data by region exist only for the interval since then. In this case, changes have been examined for all 19- to 28-year-olds combined to increase estimate reliability. Because gender, for example, crosscuts all regions, it has less sampling error than when the sample is divided into four separate regions. (Each region is represented by between 800 and 2,200 cases in all years.) By combining across all ages, we lose the ability to see the cohort effects that have occurred with many drugs, but we are able to see whether overall trends are similar across regions. In general, the changes that have occurred since 1987 have been fairly consistent across regions, particularly in terms of the direction of change. The four regions of the country—Northeast, Midwest, South, and West—have generally moved in parallel. Rather than include the large number of tables or figures necessary to show regional trends, we provide a verbal synopsis instead. The detailed information on subgroup trends is available in graphic and tabular forms in *MTF Occasional Paper 80* which may be accessed at <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ80.pdf>.

- There were substantial drops among young adults in all four regions between 1987 (the initial measurement point) and 1991 for **any illicit drug, marijuana, any illicit drug other than marijuana, cocaine, crack, and amphetamines**. After 1991, most or all regions showed some increase and then a leveling in the use of these drugs (except cocaine, which continued to decline through the mid-1990s, inched up thereafter, remained fairly level through 2006, and has declined since).

The proportions of 19- to 28-year-olds using **any illicit drug** have been consistently lowest in the South and highest in the West and Northeast. For

marijuana use, the South has usually been lowest, and for the most part the Midwest has been second lowest. Generally, the other two regions have been fairly close to one another. However, the differences have generally not been great. The 2013 prevalence rates range from 59.3% to 62.8%.

For the use of *any illicit drug other than marijuana*, until 2000 the West stood out as consistently highest, with the other three regions being very similar; since 2001, use in the Northeast generally has been about as high as in the West, though the regional differences are not large and the regions have moved in parallel.

- Data on use of *synthetic marijuana* have been gathered only since 2011. The data from respondents one to ten years past high school have been combined to minimize random fluctuations resulting from small samples. These data show some decline between 2011 and 2012 in all four regions, with the 4.3 percentage-point decline in the Midwest being statistically significant. In 2013 there was further decline in all four regions, with the 2.9 percentage-point decline in the South being statistically significant. There remains little difference among the regions in annual prevalence, which ranges from 2.5% to 4.1% across the four regions in 2013.
- From 1987, the first data point, through 2001 the West had the highest level of *lifetime* prevalence for *LSD*. From 1991 through 1995, the West had slightly higher *annual* prevalence rates of LSD use than the other three regions among young adults (use dropped in 1995 in the West). Otherwise the usage rates have been quite similar in all four regions; all showed sharp declines in LSD use after 2001, though use had been declining some in all regions for several years prior to that. In the past five years the Northeast has emerged as having a slightly higher annual prevalence than the other three regions.
- *Salvia*, which was first measured with a tripwire question in 2009, and for which we combine data for all respondents one to ten years past high school, showed a continuous decline from 2009 through 2013 in the West (which started out highest) and the South. Use began to decline in the Midwest after 2010 and in the Northeast after 2011. Use is quite low in all regions by 2013 at less than 1.6% annual prevalence, compared to 2.5% to 5.4% in the four regions in 2009.
- Questions about *ecstasy (MDMA)* were added to the follow-up surveys of young adults in 1989. Through 1993, rates were highest in the West and South and lower in the Northeast and Midwest regions. Subsequently, use in the Northeast began to increase (as was true among 12th graders), exceeding levels of use found in the South and West from 1999 to 2001. The Midwest has quite consistently had a much lower level of ecstasy use than the other three regions, although it was joined by the South in recent years. In 2000 all four regions showed a sharp and fairly parallel increase in ecstasy use; the rise decelerated in 2001 and began to decline thereafter in all regions. As

discussed elsewhere, we believe that this decrease may be caused by growing concern about the hazards of ecstasy use. By 2003, little regional difference remained in annual prevalence, largely because the declines in use were most pronounced in the Northeast and West. By 2007, use was down a little more in all regions; but since 2007 ecstasy use generally has been increasing in the West, and since 2010 in the Northeast as well, thereby opening a regional difference. In 2013 annual prevalence rates among young adults were 6.1% in the Northeast, 5.7% in the West, 3.5% in the Midwest, and 3.0% in the South.

- The considerable declines in *cocaine* use, observed in all regions between 1987 and 1991, were greatest in the two regions that had attained the highest levels of use by the mid-1980s—the West and Northeast. Thus, regional differences had diminished considerably by 1992. Similar to the finding for 12th graders, in 1992 the decline in annual prevalence stalled in all regions except the Northeast. A gradual further decline then occurred in all regions through 1996 (1997 for the West) before a slight rise began to occur, likely reflecting the effects of generational replacement. Regional variability in annual cocaine prevalence was minimal for some years after the mid-1990s, but since 2005, use in the Midwest and South has declined more than in the West and Northeast, enlarging the regional differences. Annual prevalence for the young adult age band in 2013 was 6.0% in the Northeast, 5.0% in the West, 3.0% in the South, and 2.6% in the Midwest.
- All four regions also exhibited an appreciable drop in *crack* use between 1987 and 1991, again with the greatest declines in the West and Northeast, where prevalence had been the highest. Use then generally leveled in all regions except the South, where it continued a gradual decline through 1997. As was true for cocaine generally, annual prevalence rates among the regions have converged, ranging from 0.2% to 0.8%. (Lifetime prevalence of crack use generally has been highest in the West since crack use was first measured in 1987, as is true for cocaine in general.)
- From 1987 (when data were first available) through 1994, rates of *inhalant* use remained relatively stable, quite low, and about equal in all four regions among 19- to 28-year-olds. Annual use then rose in the Northeast in 1995 and 1996 and remained higher than in the other regions through 2000, before dropping back to rates comparable to the other three regions. Except for that divergence, the regions have moved very much in parallel for this class of drugs. Annual prevalence in 2013 is at low levels in this age group, ranging between 0.1% in the Midwest and 0.8% in the Northeast.
- The regions have trended fairly similarly in their prevalence of *amphetamine* use by young adults. The only modest exception was that use declined more in the Northeast (which started out lowest) in the period 1987 to 1992, giving it a substantially lower rate than the other three regions; it remained lowest until 1998. The West has fairly consistently had the highest rate, but not by much. By the late 1990s, the Northeast had caught up to the Midwest and

South, making the regional differences very small; there have been essentially no regional differences since 2000. In 2013 the annual prevalence rate ranges between 7.0% in the West and 8.9% in the Midwest.

- **Methamphetamine** use has been measured only since 1999 (though crystal methamphetamine, discussed next, has been in the study for a longer time). It shows some differences in rates among the regions and some differential trending, with a gradual decline in annual prevalence in the Northeast (where use has generally been lowest) and a gradual increase in the West (where use has usually been highest) from 2000–2004, after which use declined in the West. Use in the other two regions remained fairly flat until 2006, when both showed some decline. Use in the West has fallen since 2006, leaving very little variability among regions by 2013. (Lifetime prevalence reached particularly high levels in the West, starting at 16% in 1999, and declining fairly steadily to 5% in 2013.) Annual prevalence now ranges from 0.2% in the Midwest to 1.1% in the West.
- The West has also consistently had the highest rates for **crystal methamphetamine (ice)** use, and the regional differences have been very substantial, particularly in terms of lifetime use. The Northeast has generally had the lowest rates. When data were first available on crystal methamphetamine in 1990, the West had a lifetime prevalence of 5.1% versus a range of 1.7% to 2.3% in the other three regions. By 2006, the lifetime prevalence rate in the West had increased to 8.8%, and lifetime prevalence in the Midwest and South grew quite steadily over that interval. This strongly suggests that crystal methamphetamine use among young adults diffused from the West primarily to the South and Midwest regions, but diffused much less to the Northeast, which has had the lowest prevalence since 1998. The annual prevalence figures tell a similar story, but also show that there was a spike in past-year use in the West from 1991 to 1995 before use there declined and then stabilized at around 2% from 1997 through 2001. Rates then rose again in the West between 2001 and 2003 and stabilized at a higher level around 2.7%. Since 2006, use in the West declined, narrowing the differences among regions. In 2013, annual use of crystal methamphetamine stands between 0.6% and 1.1% across all regions.
- **Bath salts** (synthetic stimulants sold over the counter) were first included in the study in 2012 and showed some regional variation, though all regions had an annual prevalence of use below 1.7%. Use by young adults was highest in the Northeast at 1.6% in 2012, but in 2013 it fell to less than 0.5%—a highly significant drop. There was little change in use in the other regions.
- The annual prevalence for **sedatives (barbiturates)** remained flat, and at about equivalent levels, in all four regions of the country from 1987 through 1994. Rates then rose gradually and in parallel in all regions for a number of years, followed by some leveling; regional differences have been consistently small. In 2013 annual prevalence ranges from 2.6% to 3.9%.

- The picture for *tr tranquilizers* is similar to that for sedatives (barbiturates). Annual prevalence generally held fairly steady in all regions from 1987 through 1993. Since then there has been some increase in all regions, with the South experiencing the most increase through 2004, after which all regions showed a leveling in use, followed by a gradual decline in use since about 2007. The regional differences have been small, though they grew a bit larger during the period of increasing use in the late 1990s. Annual prevalence rates in 2013 range from 4.8% in the West to 6.0% in the Northeast.
- Levels and trends in *heroin* use have been quite comparable across the four regions since 1987, at least until recently. All regions had low and stable rates through the early 1990s. A gradual increase was observed from about 1993 through 2000, and annual prevalence was fairly stable in all regions through roughly 2004. After that there was a steady increase in heroin use in the Northeast from 0.4% in 2004 to 1.1% in 2009, and also an increase in the West, from 0.3% in 2004 to 0.8% in 2009. After 2009 young adults in these two regions continued to have the highest prevalence of heroin use through 2012. In 2013, use continued to rise in the Northeast bringing its annual prevalence up to 1.8%, compared to 0.2% to 0.5% in the other three regions. This rise in the Northeast is consistent with statements by governors in the Northeast that they are facing a rising level of heroin use. The rate in the West fell back to 0.5% in 2013.
- Trends in prevalence of the use of *narcotics other than heroin* have also been quite parallel for the four regions. After a period of slight decline between 1987 and 1993 in all regions, a gradual, long-term, and substantial increase occurred from the mid-1990s through 2003 or 2004, depending on the region, with little systematic change through 2010, at which point use began to decline gradually in all regions. The South has tended to have the lowest prevalence of use since 2003, with the other three regions being tightly grouped.
- The annual prevalence of *OxyContin* use was highest in the Northeast and lowest in the West in 2002, when it was first measured. Use rose some in all regions through about 2009, and has shown a substantial decline in all regions since then. The Midwest had the lowest prevalence rate from 2010 through 2012, but showed a significant increase of 2.0 percentage points in 2013, bringing its annual prevalence up to 3.1%. The other three regions remained unchanged in 2013, with annual prevalence ranging from 2.4% in the South to 3.4% in the Northeast. In general, regional differences have not appeared very consistent due to the limited sample sizes.
- Annual prevalence of use *for Vicodin* has shown some variation among the regions. The West and Midwest generally have had the highest rates, with the South the lowest and the Northeast in between. However, the West and Midwest have shown declines in use since 2007 and 2006, respectively, narrowing the differences; the South continues to have the lowest level of

use. Annual prevalence rates now range from 3% in the South to 6.5% in the Northeast to 8% in the Midwest and West. (It should be noted that the sample sizes are more limited than usual for Vicodin and OxyContin, because questions about their use are contained on only two of the six questionnaire forms. As a result, the trends are more difficult to discern with confidence.)

- When two club drugs, *ketamine* and *GHB*, were first measured in 2002, the Northeast stood out as having a higher rate of annual use (especially so for ketamine); but use in the Northeast dropped over the next two years, bringing that region's usage rates down to the same very low levels as the other three regions. There appears to have been a little resurgence of ketamine use between 2008 and 2010 in all regions except the Midwest, followed by a decline in all regions in 2011. In 2012 and 2013 ketamine use stood higher in the Northeast than in the other regions. GHB use also appeared to rise in the Northeast in 2012, but use then fell back in 2013. No region has an annual prevalence above 1.1% in 2013 for either drug.
- *Note: Questions about the use of Rohypnol were discontinued in 2010. Rohypnol* use remained very low in all four regions from 2002, when it was first measured, through 2009, not reaching 1% in any region. For this reason, questions about its use were dropped from the surveys in 2010 to make room for other drugs.
- With respect to *alcohol* use, there were modest declines in 30-day prevalence in all four regions between 1987 (when the first measurement was available for 19- to 28-year-olds) and 1992. The rates for 30-day prevalence then leveled in all regions. The West and South have consistently had lower rates of 30-day use than the Northeast and Midwest (as has generally been true among 12th graders).

Current *daily use* of alcohol also showed a decline from the first (1987) data collection through about 1994 or 1995 in all regions. The proportional declines were substantial—on the order of 40–50%. (This decline corresponds to a period of appreciable decline in daily drinking among 12th graders, though we can tell from their longer-term data that their decline started in 1980; thus the decline may well have begun earlier among 19- to 28-year-olds as well.) After the mid-1990s there was some upward trending in daily prevalence in all regions through about 2007 or 2008, followed by a leveling. In 2013 the four regions have rates of daily alcohol use between 4.4% and 6.2%.

Occasions of heavy drinking (or binge drinking) was fairly level in all regions between 1987 and the late 1990s or early 2000s. There were then some modest increases through about 2006, followed by a leveling and even a slight decline, particularly in the West. The rates have consistently been appreciably higher in the Northeast (39% in 2013) and Midwest (41%) than in the South (32%) and the West (29%).

- There have been highly consistent regional differences among young adults in **cigarette smoking** since data were first available in 1987—they exist for monthly, daily, and half-pack-daily prevalence rates. The West has consistently had the lowest rates of *daily prevalence* (e.g., 8% in 2013, compared to 13–14% in the other three regions), with the Northeast and Midwest generally a little higher than in the South. The South, however, showed less decline in recent years than the Midwest and the Northeast, eliminating the difference with the Northeast by 2008, and by 2013 it had the highest level of daily smoking among the four regions. After some slight decline in *30-day prevalence* in all regions between 1987 and 1989, rates leveled off for about five years (roughly through 1994). There then followed a very gradual increase of a few percentage points through 1998, followed by a gradual decrease—likely reflecting cohort effects resulting from the changes in smoking during the high school years. Daily use showed a very similar pattern. For *half-pack-a-day smoking*, the decline phase was longer (from 1987 through about 1992 or 1993), likely reflecting the lag between smoking initiation and regular heavy smoking. Since 1998, half-pack smoking rates have gradually declined in all four regions. The gradual declines in daily smoking have cumulated and have been important. Between 2004 and 2013, daily prevalence fell among young adults from 23% to 13% in the Northeast, 24% to 13% in the Midwest, 20% to 14% in the South, and 16% to 8% in the West. In general, all of these measures have shown parallel movements across regions, suggesting that the forces accounting for whatever changes there have been are nationwide in scope. It should be remembered that, as illustrated earlier in this chapter, there are strong cohort effects in smoking that are obscured to a considerable degree when we combine age groups across a 10-year age span, as we have done in the present analyses.

Population Density Differences in Trends

The analyses presented here for population density return to the use of four-year age groupings, which allows a longer time interval to be examined for the younger strata and for cross-age comparisons of the trends. Among young adults, five levels of population density are distinguished based on the respondent's answer to the question, "During March of this year did you live mostly . . .?"; answer alternatives were in a very large city (over 500,000 people), in a large city (100,000 to 500,000), in a medium-sized city (50,000 to 100,000), in a small city or town (under 50,000), or on a farm/in the country? Data on the suburbs of cities of each size were combined with the corresponding city. These various subgroup data are not presented in tables or figures here because of the substantial amount of space they would require. Rather, a verbal synopsis of what they contain is presented. More detailed information on subgroup trends is available in graphic and tabular form in *MTF Occasional Paper 80*, which may be accessed at <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ80.pdf>.

- The proportions of young adults using **any illicit drug** have moved in parallel among the various community-size strata. In general, the farm/country

stratum has tended to have the lowest use. The other four community-size strata have tended to differ little from one another, though the very large cities have generally ranked at the top. In 2013, the proportions of 19- to 22-year-olds reporting use of an illicit drug in the past year are 22% for the farm/country stratum, 38% for small towns, 41% for medium-sized cities, 42% for large-sized cities, and 46% for very large cities.

- Trends in the use of *any illicit drug other than marijuana* tell a similar story. There was a long period of fairly parallel declines along with some convergence of usage rates among the community-size strata at all three age levels (among 19- to 22-year-olds it was between 1981 and 1992) followed by an increase in use and more recently a leveling. In general, medium, large, and very large cities have all tended to have about the same rates, and the farm/country stratum has tended to have the lowest rates, particularly prior to 1990; the differences by population density have been quite small since about 2000, ranging from 16% to 20% across the different community-size strata among 19- to 22-year-olds in 2012, for example. However, in 2013 there was a significant decline of 8.5 percentage points in the farm/country stratum and simultaneously a significant 8.4 percentage point increase in the very large cities, resulting in a much larger range of differences. (It remains to be seen if this sharp divergence is replicated in the 2014 results.) The two older age strata do not show a similar divergence.
- *Marijuana* use has moved pretty much in parallel among the various community-size strata over the time intervals for which data exist. Among 19- to 22-year-olds, the annual prevalence rates have been quite close among all communities, except for the farm/country stratum. This most rural stratum has fairly consistently had the lowest rate of marijuana use, and use fell less in the decline period and rose more slowly in the subsequent increase than in the other community-size strata, first narrowing and then increasing the gap. Among the two older age strata annual prevalence of marijuana use is ordinarily related to population density. For example, among 23- to 26-year-olds it is 26% among the farm/country stratum and rises to 39% in the very large cities.
- *Daily marijuana use* has also moved largely in parallel among the five population-density strata within each age band. The population-density strata all showed a decline in daily use from 1980 through about 1992, an increase from 1992 through 1995, a leveling for a couple of years, then a decline from about 2003 through 2007. Since 2007 (2005 in the case of the rural areas), however, there has been some upward trending in marijuana use among the 19- to 22-year-olds but not in the older strata.
- *Synthetic marijuana*, such as “K-2” and “Spice,” was added to the study in 2011, so only three years of data exist so far. The farm-country stratum had the highest annual prevalence initially in the two youngest age bands, but their use fell sharply and significantly in the two years since. In 2013 the annual prevalence rate ranges from 4.0% in the medium size cities to 1.8% in

the farm/country stratum. In fact, use in all community size strata has declined among the 19- to 22-year-olds. In the older age groups, use started from a lower level and generally has fallen in all community size strata, as well, with the one exception of the large cities among 23- to 26-year-olds.

- In general, there have not been large differences in *LSD* use among young adults as a function of community size since 1983. Among 19- to 22-year-olds (the young adult age group with by far the highest rates of *LSD* use prior to 2003), use in communities of all sizes declined appreciably in the early to mid-1980s, particularly in the urban strata, eliminating modest prior differences by 1984. From around 1989 through 1996, there was some increase in *LSD* use in all population-density strata among 19- to 22-year-olds, with the most rural region generally continuing to have the lowest prevalence of use. After 1997, there was some decline in *LSD* use in all community-size strata among 19- to 22-year-olds, followed by a sharp decline occurring from 2001 to 2003, with all strata moving in concert. The 23- to 26-year-old respondents had some modest increases after 1989 in all community-size strata, though the increases had virtually ended by 1995; since about 1999, there have been declines in all strata, with the largest decline in 2001–2003 in most strata. In *Volume I* in this series, we discussed how a sharp decline in supply may be responsible for the sizable decline in use among all ages after 2001. In the oldest age group, *LSD* use has remained very low and for the most part quite stable, but also with some decline after 2001. There has been very little difference among strata.
- The use of *hallucinogens other than LSD*, taken as a class, has also shown considerably higher rates in the youngest age band compared to the two older ones, suggesting a consistent sharp falloff in use with age—an age effect. (The drug most often reported in this general class has been psilocybin or “shrooms,” as is true among 12th graders as well.) Use of this general class of drugs fell in communities of all sizes among young adults between 1980 and about 1988. Then there was a leveling of use for a few years, followed by an extended increase in use among all community-size strata in the 19- to 22-year-old age band. By 2003 the rates attained by each stratum exceeded those originally observed in 1980; there have been declines since then, and very large cities (which had attained the highest usage rates) were the last to show the decline. The 23- to 26-year-old group showed slightly rising rates of use between 1998 and 2004, followed by a leveling. The sharpest increase occurred in the very large cities in 1999 and 2000 and again in 2010. The 27- to 30-year-olds have generally had low rates of use, and the trend lines have been very flat with only minor community-size stratum differences, until 2001 when all strata, especially the very large cities, began to increase before leveling after 2005. All three age groups have shown higher rates of use than previously observed—since 1994 for 19- to 21-year-olds, since about 2000 for 22- to 26-year-olds, and since about 2002 for 27- to 30-year-olds—in part reflecting a cohort effect in the use of these drugs, but also reflecting the change in question wording to include “shrooms” as an example. The

farm/country stratum has generally had the lowest prevalence among the two younger strata and the very large cities the highest.

- *Salvia* (or *salvia divinorum*) use was first measured in 2009 and has shown somewhat irregular trend lines since then. The overall picture appears to be one of decreasing use since 2009 in the youngest age group and since 2010 in the middle age group. Annual prevalence rates are now very low for this drug—from less than 0.5% to 1.6% among 19- to 22-year-olds, and from less than 0.6% to 1.6% among the 23- to 26-year-olds. The case counts are too low to make reliable estimates of change by population density strata.
- *Ecstasy (MDMA) use was first* measured in 1989, and since then has shown the largest increase of any drug among young adults. Among 19- to 22-year-olds use in 1989 was highest in the very large cities (5% annual prevalence); but prevalence declined in all population-density strata between 1989 and 1994 (to 1.6% or less). By 1998, use had begun to increase in all community-size strata within this age band, except in the farm/country stratum. The farm/country stratum moved up sharply in 1999, but then the three most urban strata jumped sharply in 2000, opening a fair gap in use, with large and very large cities having rates nearly twice as high as any of the other strata in 2002. All community-size strata showed large declines in ecstasy use since 2000 or 2001, which lasted through 2004, narrowing the differences among them. In 2011 ecstasy use in the very large cities rose to nearly 10% annual prevalence and has stayed there since then—nearly twice as high as any of the other population density strata. Among 23- to 26-year-olds, use began to increase a little later, and again the most urban stratum showed the most increase, particularly in 2000; but use began to decline after 2001 (in the urban areas) or 2002 (in the rural and small town strata), which had the effect of narrowing the differences among strata. Among the 23- to 26-year-olds, all population-density strata continued to decline, or at least remain level, through 2003, and then stayed level through 2008. As with the younger age group, ecstasy use rose sharply, this time in 2012, to roughly 8%, where it remained in 2013. Considerably less increase in ecstasy use occurred among 27- to 30-year-olds, though there was some increase in the largest cities starting after 1996 and in the large and medium-sized cities after 1999. From 1997 through 2005 the very large cities stood out as having the highest rates of ecstasy use, but the differences were modest through 2012. In 2013 the very large cities were the only ones to show any increase in use in this age band. *It thus appears that over the past two to three years ecstasy use has made a comeback among young adults in the country's very large cities.*

Ecstasy use trends tell an interesting story. In the very large cities use peaked in all three age bands in 2000 and then began to decline. The medium-sized cities were beginning to level or decline by 2001 in the two younger age bands. The small town and farm/country strata peaked in 2001 in all age groups. These data support our belief, based on school-level analyses of secondary schools, that the presence of this drug was still diffusing geographically—in this case, from more urban to more rural areas—and,

were it not for this continued diffusion, ecstasy use would actually have begun to decline nationally a year earlier. The data from 12th graders on perceived risk provide the clue as to the most likely cause of this turnaround; they showed a large jump in the level of perceived risk associated with ecstasy use from 2000 through 2003. Unlike most other drugs discussed here, the pattern of change since the mid-1990s appears to reflect secular trends, with all age groups moving pretty much in parallel, more than cohort effects.

- In the early 1980s, *cocaine* use was positively correlated with population density, with the highest use in the very large cities. The important drop in cocaine use that began after 1986 slowed considerably after 1992 or 1993 in all three age strata and in communities of all sizes, by which time the positive association with population density had been virtually eliminated. Among 19- to 22-year-olds there was a sustained increase in cocaine use among all community-size strata after about 1993 or 1994, and among 23- to 26-year-olds after about 1998. There was some decline in the mid-2000s in all strata except large cities, which showed a decline in more subsequent years. As just stated, usage rates among the population-density strata tended to converge considerably during the period of decline; this convergence remains, except for the very large cities, which since 2007 have shown rates of cocaine use somewhat higher than the less densely populated areas in all three age bands. In the 27- to 30-year-old age group, a gradual increase in use emerged in nearly all population-density strata after 2000, no doubt reflecting a cohort effect working its way up the age spectrum. By 2004, all of the strata in the oldest age band leveled or declined from their peak rates; the single exception was very large cities, where use remained relatively high until 2010, when it declined a bit. In 2013 the very large cities had the highest prevalence rates across all three age groups.
- *Crack* use among all age groups peaked in 1987 or 1988 (strongly suggesting a secular trend at work at that time) and then, after declining appreciably, bottomed out in all population-density strata for several years. Use reported in these young adult samples at all three age levels has borne practically no systematic association with community size, and for the most part the strata have all tended to move in parallel. In 2013, annual prevalence is at or below 1.3% across all strata in the older two age bands; but among 19- to 22-year-olds use is at or below 1.0% in all strata. In the youngest age band of young adults crack use has tended to be highest in the farm-country stratum.
- *Amphetamine* use shows virtually no differences associated with urbanicity in any of the three age groups, and this has generally been true since 1983. The trend curves were highly similar for all levels of population density within each age stratum, with the single exception that among the 23- to 26-year-olds the three urban strata exhibited the greatest increase in amphetamine use of the five strata after 2008; they were joined by the small town stratum in 2012, leaving the farm/country stratum with the lowest rate of use.

- Differences in use of *crystal methamphetamine (ice)* as a function of population density have been quite erratic across time in all four age groups, particularly in the earlier years of collecting such data, due to limited sample sizes. Since 2007, reported rates of use have been relatively low in all strata and age bands, though use in the farm/country stratum did rise to 2.7% in 2013 in both of the two older age groups.
- The use of *methamphetamine* in any form has been measured only since 1999. In general, the farm/country stratum has had higher than average rates of use in the two youngest age groups; otherwise there has been little systematic difference. Among 19- to 22-year-olds, all community-size strata have shown substantial declines in use since 2003 or 2004, reaching very low levels by 2007 and thereafter. Use has declined some over the same interval among 23- to 26-year-olds. Among 27- to 30-year-olds use generally declined from 2002 to 2006 in all population-density strata; after 2009, this group showed a slight rebound in use, particularly in the farm-country stratum. (Use has also risen some in the farm/country stratum since 2011.)
- *Bath salts* were added to the study in 2012, so trends are available for only one year. They showed a high rate of use (6.5% annual prevalence) in 2012 in the farm/country stratum among 19- to 22-year-olds, but a significant decline in 2013 such that there are now practically no differences among the different levels of population density. Usage rates are very low among the two older age bands with little variability by population density. These findings suggest that this type of drug use tended to be concentrated among younger people and in more rural areas, but also that bath salts are no longer much of a threat to public health.
- *Note: Methaqualone was dropped from the study in 1990. Methaqualone* use, which in 1981 was strongly positively associated with population density, dropped to annual prevalence rates of 0.8% or below in all community-size strata for all three age bands by 1989. For that reason, its use is no longer measured in MTF.
- Unlike methaqualone, *sedatives (barbiturates)* have never shown much variation by population density, at least as far back as 1980. This remains true in all three age bands, and the trends have been similar within each age band.
- *Tranquilizer* use among young adults has also had little or no association with population density over this time interval. Like sedatives (barbiturates), there was an earlier period of decline, staggered inflection points, a long period of gradual increase, and then a leveling staggered up the age band from about 2003–2005.
- From 1980 to 1995, annual *heroin* prevalence was less than 1.0%—usually much less—in all population-density strata for all three age bands. After 1994, use among 19- to 22-year-olds in all community-size strata rose and

reached 1.0% in the three most urban strata by 1998. In fact, in the very large cities, it reached 2.1% in 2000 (vs. 0.3–0.6% in the other strata). Use levels have been lower among 23- to 26-year-olds and lower still among 27- to 30-year-olds, making it difficult to discern systematic differences among the population-density strata in those age bands. In 2013 the annual prevalence of heroin is 1.1% or lower in all community-size strata for all three young adult age bands, and it is much lower in most. Over the past four or five years for nearly all community-size strata, there was some increase in use among the 23- to 26-year-olds (particularly in 2010). In 2013 there was some increase in use in the farm/country stratum in the oldest age band and in the middle age band as well, though neither was statistically significant.

- The annual use of *narcotics other than heroin* had some positive association with population density among 19- to 22-year-olds through the early 1990s; however, it has shown rather little association since then. Use of narcotics other than heroin increased substantially in all community-size strata after 1993 in the case of 19- to 22-year-olds, after about 1996 in the case of 23- to 26-year-olds, and after about 1998 in the case of 27- to 30-year-olds; however, no systematic differentiation by community size was evident during those periods of increasing use. Clearly a cohort effect was at work, and the increasing use of these drugs was quite widespread. Use tended to level off since about 2004 in the youngest age band, 2006 in the middle age band, and 2007 in the oldest age band.
- Unfortunately, sample sizes for two of the narcotic drugs of particular interest, *OxyContin* and *Vicodin*, are not sufficient to estimate population-density differences or trends with a reasonable degree of accuracy.
- The absolute levels of *inhalant* use have remained low in these age groups, particularly above age 22. However, during the mid- to late 1980s, there was a gradual increase in use among 19- to 22-year-olds in all community-size strata. No strong or consistent association with population density has appeared, though the very large cities have not infrequently had higher rates than the other areas among 19- to 22-year-olds, particularly in the period 1998 through 2000. Among both the 19- to 22-year-olds and the 23- to 26-year-olds, there has been some falloff in inhalant use since the late 1990s in all population-density strata.
- There have been few differences as a function of population density in the 30-day prevalence of drinking *alcohol* among 19- to 22-year-olds since data were first available in 1980, except that the farm/country stratum has tended to have lower-than-average use. In the two older age bands, however, there has been a fairly consistent positive correlation between population density and use of alcohol in the past 30 days—though not a very strong one. So, for example, in 2013, 59% of 27- to 30-year-olds in the farm/country stratum have had alcohol in the prior 30 days, compared to 83% of those in very large cities. Trends have been fairly parallel for all strata in all age bands. There

have also been no consistent trend differences in current *daily drinking* associated with urbanicity in any of the three age bands.

- For *occasions of heavy drinking*, all community-size strata have been fairly close across time at all three age levels, with two exceptions: The farm/country stratum has fairly consistently shown a slightly lower rate of binge drinking in the youngest two age bands, and such drinking has tended to be highest in the very large cities in the two upper age bands (and more often than not in the oldest age band). However, in the upper two age bands, the differences among the communities of different size have gradually expanded since about 2001, when there were practically no differences. So, for example, in 2013 among 23- to 26-year-old respondents, 29% in the farm/country stratum indicate they engaged in occasions of heavy drinking, versus 42% in the very large cities.
- *Cigarette smoking* has generally been negatively associated with population density in all three age strata, without much evidence of differential trends related to population density. There is one exception: Among 19- to 22-year-olds, all smoking prevalence measures rose from 1997 through 1999 in the farm/country and small town strata, while in most other strata they remained level. The differences in 1999 were most striking for half-pack-a-day smoking among the 19- to 22-year-olds—24% for farm/country, 19% for small town, 15% for both medium-sized and large cities, and 10% for very large cities—compared with 1985, when there was virtually no difference in half-pack-a-day smoking rates among these strata (all were at 18% or 19%). Thus, smoking among those in their early 20s became more concentrated in the nonurban populations. In fact, among 19- to 22-year-olds, the farm/country stratum has usually had the highest rate of daily smoking since 1986, and the small town stratum has generally ranked second since then. As smoking has declined in all strata in the youngest group, this difference has diminished; but not in the older two age bands. Among the two older age groups, the farm/country stratum has been highest more often than not. Among 19- to 22-year-olds, there has been a decline in 30-day prevalence in most population density strata since about 2000 or 2001, and among 23- to 26-year-olds since 2005. Continuing declines in smoking among 12th graders would lead us to predict still further declines among young adults as well, given the strength of cohort effects on smoking rates.

TABLE 5-1
Trends in Lifetime Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28
(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,600	4,600	4,400		
Any Illicit Drug ^a	70.5	69.9	67.9	66.4	64.5	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	60.4	59.7	59.8	59.3	59.3	58.4	59.1	58.9	60.5	+1.6
Any Illicit Drug other than Marijuana ^a	48.4	47.0	44.6	42.7	40.8	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3	31.6	32.8	33.9	35.2	34.0	34.8	34.2	34.7	32.8	33.3	33.2	32.8	34.2	+1.4
Marijuana	66.5	66.0	63.8	62.8	60.2	58.6	56.4	55.9	53.7	53.6	53.5	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	57.0	56.7	56.7	55.9	56.0	55.9	56.3	56.5	57.1	+0.6
Inhalants ^b	12.3	12.7	12.6	13.2	12.5	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	10.3	10.9	9.1	9.5	8.9	7.9	7.2	7.2	6.5	-0.7
Nitrites ^c	2.6	6.9	6.2	—	1.9	1.4	1.2	1.3	1.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens ^d	18.5	17.1	17.0	15.9	16.1	15.7	15.7	15.4	15.4	16.1	16.4	16.7	17.4	18.0	18.4	18.3	19.6	19.7	19.3	17.6	17.2	16.0	14.8	14.2	13.9	13.0	12.2	12.4	+0.2
LSD	14.6	13.7	13.8	12.7	13.5	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	11.2	10.1	9.6	8.1	7.3	7.2	6.1	6.2	6.3	0.0
Hallucinogens other than LSD ^d	12.6	11.4	10.6	9.4	9.1	8.4	8.0	7.6	7.4	7.8	7.9	8.4	9.4	9.3	9.9	12.0	15.0	16.4	15.6	15.4	14.9	14.1	13.0	13.0	12.6	12.1	11.1	11.4	+0.3
PCP ^e	8.4	4.8	5.0	—	2.5	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	2.0	2.4	2.1	2.2	1.6	1.6	1.7	1.1	1.4	+0.3
Ecstasy (MDMA) ^f	—	—	—	3.3	3.7	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	14.9	14.4	13.1	13.1	11.5	12.3	11.3	11.4	11.6	+0.3
Cocaine	32.0	29.3	28.2	25.8	23.7	21.0	19.5	16.9	15.2	13.7	12.9	12.0	12.3	12.8	12.7	13.1	13.5	14.7	15.2	14.3	15.2	14.7	14.8	13.9	13.6	12.5	11.9	12.2	+0.3
Crack ^g	—	6.3	6.9	6.1	5.1	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	4.1	4.4	3.9	4.3	3.3	3.6	2.9	2.7	2.6	-0.1
Other Cocaine ^h	—	28.2	25.2	25.4	22.1	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	13.3	14.4	14.0	13.9	13.5	13.1	12.2	11.8	11.8	0.0
Heroin	1.3	1.3	1.1	1.0	0.9	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	1.7	1.9	1.6	1.9	1.6	1.8	1.7	1.6	1.6	0.0
With a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.4	0.4	0.3	0.4	0.6	0.4	0.6	0.4	0.5	0.4	0.6	0.6	0.5	0.5	0.5	0.8	0.7	0.5	1.0	+0.4
Without a Needle ⁱ	—	—	—	—	—	—	—	—	—	1.0	1.4	1.5	1.7	1.9	2.1	2.1	1.8	2.2	2.1	1.8	2.4	1.9	2.1	1.9	1.8	1.6	1.7	1.8	+0.1
Narcotics other than Heroin ^{j,k}	10.7	10.6	9.8	9.6	9.4	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5	13.9	16.8	17.6	17.8	18.7	18.8	19.5	18.5	19.0	18.2	17.6	17.4	-0.3
Amphetamines, Adjusted ^{l,j}	32.3	30.8	28.8	25.3	24.4	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	14.6	15.6	15.3	14.6	14.9	16.1	16.5	17.4	18.7	+1.3
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	8.8	9.3	9.0	9.1	8.9	9.0	8.3	7.3	6.7	6.3	4.7	4.3	3.2	3.5	3.1	-0.4
Crystal Methamphetamine (Ice) ⁱ	—	—	—	—	2.5	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	4.4	4.7	3.7	3.6	3.4	2.8	3.1	2.6	2.8	+0.2

(Table continued on next page.)

TABLE 5-1 (cont.)
Trends in Lifetime Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

Crystal Methamphetamine (Ice)¹

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change		
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,600	4,600	4,400				
Sedatives (Barbiturates) ^{1,u}	11.1	9.7	8.9	7.9	8.7	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	10.0	9.5	9.8	10.6	9.5	8.6	7.9	7.2	9.5	—		
Sedatives, Adjusted ^{1,m}	16.7	15.0	13.2	12.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methaqualone ¹	13.1	11.6	9.7	8.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{d,j}	17.6	16.5	15.1	13.5	12.9	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5	11.9	13.4	13.8	14.9	14.5	15.0	14.5	15.8	13.8	14.3	13.8	13.3	13.2	13.2	-0.1	
Alcohol ⁿ	94.8	94.9	94.8	94.5	94.3	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	89.1	88.9	87.9	88.4	87.9	87.5	87.4	86.5	86.2	86.2	-0.3	
Been Drunk ^o	—	—	—	—	—	82.9	81.1	81.4	80.7	82.1	80.7	81.4	79.8	81.6	80.4	81.1	81.2	80.9	80.1	79.9	80.9	80.1	80.1	78.2	79.0	78.9	78.9	77.4	77.4	-1.5	
Flavored Alcoholic Beverages ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	83.2	84.6	84.4	84.0	82.6	83.5	81.4	82.2	82.4	80.9	80.9	-1.6	
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Steroids ^q	—	—	—	1.1	1.2	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	1.8	1.8	1.7	1.8	1.8	1.7	1.3	1.7	1.2	1.2	-0.5	

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 5-4.

TABLE 5-2
Trends in Annual Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change	
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400		
Any Illicit Drug ^a	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	33.3	33.2	34.7	34.0	36.3	+2.3 s	
Any Illicit Drug other than Marijuana ^a	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	17.4	18.5	17.6	17.2	18.1	+0.9	
Marijuana	36.5	34.8	31.8	29.0	26.1	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	28.2	27.7	28.5	28.6	29.3	28.7	31.0	30.2	32.2	+2.0 s	
Synthetic Marijuana ^v	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.4	5.3	3.2	-2.1 sss	
Inhalants ^b	1.9	2.1	1.8	1.9	1.9	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	1.3	1.3	0.8	1.4	0.9	1.2	0.8	1.1	0.5	-0.6 s	
Nitrites ^c	2.0	1.3	1.0	—	0.4	0.2	0.1	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Hallucinogens ^d	4.5	4.0	3.9	3.6	4.1	4.5	5.0	4.5	4.8	5.6	5.6	5.8	5.2	5.4	5.4	5.4	4.7	5.2	4.7	4.5	4.1	3.8	3.8	3.9	4.2	3.7	3.6	3.9	+0.3	
LSD	3.0	2.9	2.9	2.7	3.3	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	0.8	1.2	1.1	1.4	1.7	1.5	1.7	1.6	2.0	+0.5	
Hallucinogens other than LSD ^d	2.5	2.1	1.9	1.8	1.7	1.7	1.9	1.9	2.0	2.5	2.8	3.1	3.0	3.0	3.4	3.5	4.0	4.9	4.5	4.2	3.8	3.6	3.4	3.3	3.7	3.2	2.9	3.2	+0.3	
PCP ^e	0.8	0.4	0.4	—	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	0.6	0.2	0.3	0.4	0.1	0.2	0.3	0.0	0.2	+0.2	
Ecstasy (MDMA) ^f	—	—	—	1.4	1.5	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	3.0	3.0	2.5	3.3	3.1	3.5	3.6	4.1	4.2	+0.2	
Salvia ^w	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.5	3.6	2.2	1.4	0.9	-0.5
Cocaine	19.7	15.7	13.8	10.8	8.6	6.2	5.7	4.7	4.3	4.4	4.1	4.6	4.9	5.4	5.4	5.8	5.8	6.6	7.1	6.9	6.6	6.2	6.0	5.2	4.7	4.7	4.1	3.9	-0.2	
Crack ^g	3.2	3.1	3.1	2.5	1.6	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	1.2	1.1	1.0	0.9	0.7	0.5	0.6	0.5	0.3	-0.1	
Other Cocaine ^h	—	13.6	11.9	10.3	8.1	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	6.3	5.9	5.6	5.5	5.0	4.8	4.3	4.0	3.7	-0.3	
Heroin	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.5	0.5	0.5	0.6	+0.2	
With a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.1	0.1	0.1	0.1	0.1	*	0.3	*	*	0.1	0.2	0.3	0.1	0.1	0.1	0.2	0.4	0.3	0.3	0.0	
Without a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.3	0.4	0.4	0.6	0.6	0.5	0.9	0.2	0.4	0.3	0.4	0.5	0.3	0.4	0.6	0.4	0.2	0.4	0.7	+0.4	
Narcotics other than Heroin ^{jk}	3.1	3.1	2.7	2.8	2.7	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0	7.1	8.5	9.0	8.7	9.1	8.7	9.1	8.4	9.0	7.9	7.3	7.0	-0.3	
OxyContin ^{lr}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.9	2.6	3.1	3.1	3.1	2.9	3.9	5.2	3.2	2.8	2.3	2.8	+0.5	
Vicodin ^{lr}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.2	8.6	8.9	9.3	9.1	8.9	9.1	8.9	7.8	7.1	6.3	6.2	-0.1	
Amphetamines, Adjusted ^{ll}	10.6	8.7	7.3	5.8	5.2	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	5.1	5.6	5.6	5.3	6.0	7.1	7.2	7.8	7.8	0.0	
Ritalin ^{lr}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.9	2.9	2.7	2.5	2.6	2.4	2.4	1.7	1.7	1.5	1.6	2.0	+0.4	
Adderall ^{lr}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.8	7.0	6.6	7.4	7.0	-0.4
Provigil ^{lr}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.5	0.3	—	—	—
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.8	2.5	2.8	2.7	2.8	2.4	1.9	1.5	1.0	0.9	0.7	0.5	1.0	0.6	-0.3	
Crystal Methamphetamine (Ice) ⁱ	—	—	—	—	0.4	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	1.7	1.5	1.1	1.1	0.8	0.8	0.5	0.5	0.6	0.8	+0.2

(List of drugs continued.)

(Table continued on next page.)

TABLE 5-2 (cont.)
Trends in Annual Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28
(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change	
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400		
Bath Salts (synthetic stimulants) ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.4	-0.1
Sedatives (Barbiturates) ^{j,u}	2.3	2.1	1.8	1.7	1.9	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	4.2	3.9	4.2	4.7	3.8	3.3	3.2	2.7	3.4	—	
Sedatives, Adjusted ^{l,m}	3.0	2.5	2.1	1.8	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methaqualone ^l	1.3	0.9	0.5	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{d,j}	5.4	5.1	4.2	3.7	3.7	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6	5.5	7.0	6.8	7.4	6.7	6.5	7.1	6.8	6.4	6.3	5.9	5.3	5.4	+0.1	
Rohypnol ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.5	0.1	0.1	0.2	0.3	0.2	0.1	—	—	—	—	—	
GHB ^x	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.6	0.5	0.3	0.2	0.4	0.3	0.2	0.3	0.3	0.4	0.3	0.0	
Ketamine ^x	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.2	0.9	0.6	0.5	0.5	0.3	0.4	0.5	0.8	0.5	0.8	0.5	-0.2	
Alcohol ⁿ	88.6	89.4	88.6	88.1	87.4	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	83.8	84.4	84.0	83.6	83.8	82.7	83.5	82.5	82.5	0.0	
Been Drunk ^o	—	—	—	—	—	62.0	60.9	61.1	58.8	61.6	59.9	63.2	59.6	63.2	60.6	63.1	61.8	62.9	63.8	63.5	65.7	65.8	66.0	65.5	64.8	64.0	64.6	63.1	-1.6	
Flavored Alcoholic Beverages ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	62.7	58.4	58.5	58.9	58.3	57.0	52.0	56.3	54.8	54.1	-0.7	
Alcoholic Beverages containing Caffeine ^{l,t}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.1	36.7	36.9	+0.2	
Cigarettes	40.1	40.3	37.7	38.0	37.1	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.1	39.1	38.6	39.0	39.1	36.9	36.2	35.0	33.9	33.0	31.5	29.8	29.8	0.0	
Small Cigars ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	19.2	18.0	18.4	+0.5	
Tobacco using a Hookah ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	20.1	19.1	20.4	+1.3	
Dissolvable Tobacco ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.6	0.3	-0.3	
Snus ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.1	5.7	4.8	-0.9
Steroids ^q	—	—	—	0.5	0.3	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.7	0.4	0.7	0.8	0.2	0.4	0.5	+0.1	

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 5-4.

TABLE 5-3
Trends in 30-Day Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28
(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change	
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400		
Any Illicit Drug ^a	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	19.8	18.9	20.6	19.9	21.8	+1.9 s	
Any Illicit Drug other than Marijuana ^a	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	8.5	8.6	8.4	7.8	8.3	+0.5	
Marijuana	22.0	20.7	17.9	15.5	13.9	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	15.8	15.7	16.0	16.0	17.0	16.1	18.3	17.7	19.0	+1.3	
Inhalants ^b	0.4	0.6	0.6	0.5	0.6	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	0.2	0.3	0.2	0.4	0.2	0.1	0.1	0.3	0.1	-0.1	
Nitrites ^c	0.5	0.5	0.4	—	0.1	*	0.1	0.2	0.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Hallucinogens ^d	1.3	1.2	1.1	1.1	0.9	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2	1.2	0.9	1.2	0.9	0.8	0.7	0.9	0.9	0.8	1.0	0.9	0.6	1.0	+0.4 s	
LSD	0.9	0.8	0.8	0.8	0.6	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.1	0.2	0.2	0.4	0.2	0.4	0.3	0.3	0.4	+0.2	
Hallucinogens other than LSD ^d	0.6	0.6	0.4	0.5	0.4	0.3	0.5	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.7	0.6	0.8	1.2	0.9	0.8	0.6	0.8	0.7	0.7	0.8	0.6	0.4	0.7	+0.2	
PCP ^e	0.2	0.1	0.3	—	0.2	0.1	0.2	0.2	0.1	*	0.1	0.1	0.2	0.2	*	*	0.1	0.1	0.1	*	*	*	0.1	*	*	0.1	*	0.2	+0.2	
Ecstasy (MDMA) ^f	—	—	—	0.4	0.2	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	0.6	0.7	0.5	0.6	0.6	0.8	0.7	1.0	1.1	+0.1	
Cocaine	8.2	6.0	5.7	3.8	2.4	2.0	1.8	1.4	1.3	1.5	1.2	1.5	1.7	1.9	1.7	2.2	2.2	2.4	2.2	2.2	2.3	2.1	1.9	1.8	1.4	1.5	1.3	1.5	+0.2	
Crack ^g	—	1.0	1.2	0.7	0.4	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2	0.1	0.2	0.1	0.1	+0.1	
Other Cocaine ^h	—	4.8	4.8	3.4	2.1	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	1.9	1.9	2.0	1.7	1.6	1.5	1.4	1.3	1.3	0.0	
Heroin	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.3	+0.2	
Narcotics other than Heroin ^{j,k}	0.9	0.9	0.7	0.7	0.7	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7	2.9	2.9	3.0	3.5	3.2	3.4	3.6	3.2	3.4	2.9	2.7	2.6	-0.1	
Amphetamines, Adjusted ^{l,j}	4.0	3.2	2.7	2.1	1.9	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	2.1	2.2	2.3	2.2	2.5	2.9	3.0	3.2	3.2	0.0	
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.7	1.0	1.0	0.7	0.6	0.7	0.5	0.6	0.3	0.3	0.2	0.3	0.4	0.2	-0.2
Crystal Methamphetamine (Ice) ⁱ	—	—	—	—	—	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.4	0.4	0.6	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.4	+0.1	

↓

(Table continued on next page.)

TABLE 5-3 (cont.)
Trends in 30-Day Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28
(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400	
Sedatives (Barbiturates) ^{j,u}	0.7	0.7	0.7	0.5	0.6	0.5	0.5	0.6	0.6	0.8	0.8	0.9	0.9	1.1	1.3	1.7	1.5	1.5	1.8	1.7	1.5	1.6	1.9	1.2	1.1	1.1	1.1	1.2	—
Sedatives, Adjusted ^{j,m}	0.9	0.8	0.7	0.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^j	0.3	0.2	0.1	*	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{d,j}	1.8	1.6	1.4	1.2	1.1	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8	2.1	2.8	2.4	2.7	2.6	2.3	2.8	2.7	2.8	2.2	2.3	1.9	1.9	0.0
Alcohol ⁿ	75.1	75.4	74.0	72.4	71.2	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.2	68.3	67.0	68.4	68.6	68.7	69.5	68.9	69.4	68.4	68.8	69.5	68.7	-0.8
Been Drunk ^o	—	—	—	—	—	35.4	35.6	34.2	34.3	33.0	33.2	35.6	34.2	37.7	35.7	36.8	37.1	37.8	39.0	39.0	42.1	41.4	40.7	40.5	39.4	39.5	39.1	37.7	-1.4
Flavored Alcoholic Beverage ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	27.6	24.9	25.9	26.7	24.4	24.5	23.8	26.1	25.4	-0.7
Cigarettes	31.1	30.9	28.9	28.6	27.7	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	28.6	27.0	26.2	24.6	23.3	22.4	21.3	19.7	20.0	+0.4
Steroids ^q	—	—	—	0.2	0.1	0.2	0.1	*	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.2	0.3	0.5	0.2	0.1	0.1	0.0

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 5-4.

TABLE 5-4
Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change	
<i>Approximate Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400		
Marijuana ^s	4.1	4.2	3.3	3.2	2.5	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	4.9	5.0	5.0	5.1	5.4	5.3	6.1	5.6	6.2	+0.6	
Any Illicit Drug	0.2	0.1	0.2	0.1	*	0.1	*	0.1	*	0.1	*	*	*	0.1	*	0.1	*	*	0.1	0.1	0.1	*	*	0.1	*	*	*	*	*	0.0
other than Marijuana ^a	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.3	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.0	0.1	+0.1	
Alcohol																														
Daily ^{n,s}	6.1	6.6	6.1	5.5	4.7	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	5.2	5.4	5.6	5.3	5.3	4.6	5.2	5.5	5.1	-0.4	
Been Drunk ^{o,s}	—	—	—	—	—	0.5	0.4	0.4	0.5	0.3	0.4	0.9	0.5	0.9	0.5	0.4	0.6	0.8	0.7	0.5	0.6	0.6	0.5	1.0	0.7	0.7	0.4	0.5	+0.1	
5+ Drinks in a Row in																														
Last 2 Weeks	36.1	36.2	35.2	34.8	34.3	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	37.0	37.6	37.8	37.9	36.7	35.9	36.5	35.5	35.1	-0.4	
Cigarettes																														
Daily	25.2	24.8	22.7	22.4	21.3	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	19.6	18.6	17.3	16.7	15.0	14.8	13.8	12.8	12.1	-0.7	
1/2 Pack+/Day	20.2	19.8	17.7	17.3	16.7	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	12.5	11.9	11.1	10.2	9.3	9.3	7.5	7.6	7.0	-0.6	

Source. The Monitoring the Future study, the University of Michigan.

Note. The illicit drugs not listed here show a daily prevalence of 0.2% or less in all years.

See footnotes on the following page.

Footnotes for Tables 5-1 through 5-4

Notes. Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

The illicit drugs not listed here show a daily prevalence of 0.2% or less in all years.

' * ' indicates a prevalence rate of less than 0.05%.

' — ' indicates data not available.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in four of the five questionnaire forms in 1986–1989; N is four fifths of N indicated. Data were based on five of the six questionnaire forms in 1990–1998; N is five sixths of N indicated. Data were based on three of six questionnaire forms in 1999–2013; N is three sixths of N indicated.

^cThis drug was asked about in one questionnaire form. N is one fifth of N indicated in 1986–1988 and one sixth of N indicated in 1990–1994.

^dIn 2001 the question text was changed on three of the six questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. For tranquilizers, Miltown was replaced with Xanax. Beginning in 2002 the remaining forms were changed to the new wording.

^eThis drug was asked about in one of the five questionnaire forms in 1986–1988; N is one fifth of N indicated. Data were based on one of six questionnaire forms in 1990–2013; N is one sixth of N indicated.

^fThis drug was asked about in two of the five questionnaire forms in 1989; N is two fifths of N indicated. Data were based on two of the six questionnaire forms in 1990–2001; N is two sixths of N indicated. Data were based on three of the six questionnaire forms in 2002–2013; N is three sixths of N indicated.

^gThis drug was asked about in two of the five questionnaire forms in 1987–1989; N is two fifths of N indicated. Data were based on all six questionnaire forms in 1990–2001. Data were based on five of six questionnaire forms in 2002–2013; N is five sixths of N indicated.

^hThis drug was asked about in one of the five questionnaire forms in 1987–1989; N is one fifth of N indicated. Data were based on four of the six questionnaire forms in 1990–2013; N is four sixths of N indicated.

ⁱThis drug was asked about in two of the six questionnaire forms; N is two sixths of N indicated.

^jOnly drug use that was not under a doctor's orders is included here.

^kIn 2002 the question text was changed in three of the six questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is three sixths of N indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003 and beyond.

^lBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^mSedatives, adjusted" data are a combination of barbiturate and methaqualone data.

ⁿIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a drink meant more than just a few sips. Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

^oThis drug was asked about in three of the six questionnaire forms; N is three sixths of N indicated.

^pThis drug was asked about in one of the six questionnaire forms; N is one sixth of N indicated.

^qThis drug was asked about in one of the five questionnaire forms in 1989; N is one fifth of N indicated. Data were based on two of the six questionnaire forms in 1990–2013; N is two sixths of N indicated.

^rThis drug was asked about in two of the six questionnaire forms in 2002–2009; N is two sixths of N indicated. Data were based on three of the six three of the six questionnaire forms. N is three sixths of N indicated.

^sDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.

^tIn 2012 the alcoholic beverage containing caffeine question text was changed to alcoholic beverage mixed with an energy drink. The data in 2011 and 2012 are not comparable due to this question change.

^uIn 2013 the question text was changed on all forms: Tuinal, Nembutal, and Seconal were replaced with Ambien, Lunesta, and Sonata. The data in 2012 and 2013 are not comparable due to this question change.

^vThis drug was asked about in two of the six questionnaire forms in 2011–2012; N is two sixths of N indicated. Data were based on three of the six questionnaire forms in 2013; N is three sixths of N indicated.

^wThis drug was asked about in one of the six questionnaire forms in 2009; N is one sixth of N indicated; Data were based on two of the six questionnaire forms in 2010–2011; N is two sixths of N indicated. Data were based on three of the six questionnaire forms in 2012–2013; N is three sixths of N indicated.

^xThis drug was asked about in two of the six questionnaire forms in 2002–2009; N is two sixths of N indicated; Data were based on three of the six questionnaire forms in 2010–2011; N is three sixths of N indicated. Data were based on two of the six questionnaire forms in 2012–2013; N is two sixths of N indicated.

TABLE 5-5
Trends in Annual and 30-Day Prevalence of an Illicit Drug Use Index ^a
among Respondents of Modal Ages 19–28
Total and by Gender

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change
<i>Percentage who used in last 12 months</i>																													
Any Illicit Drug																													
Total	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	33.3	33.2	34.7	34.0	36.3	+2.3 s
Males	45.3	42.6	39.5	35.7	33.6	30.0	31.4	31.1	32.3	32.1	31.6	31.9	33.6	33.9	34.4	34.9	35.6	36.0	37.0	35.3	35.9	35.4	37.4	35.3	38.1	38.3	37.7	39.2	+1.5
Females	39.0	36.5	33.6	30.5	28.3	24.5	25.8	26.1	25.3	28.1	27.3	27.1	27.1	27.6	28.2	30.1	30.2	31.0	31.4	31.1	29.5	30.7	31.4	32.0	29.9	32.4	31.5	34.3	+2.7 s
Any Illicit Drug other than Marijuana																													
Total	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	17.4	18.5	17.6	17.2	18.1	+0.9
Males	30.4	26.5	23.8	21.0	19.1	16.4	16.3	14.7	16.2	15.4	15.6	16.2	16.7	17.8	17.2	18.9	19.8	21.3	20.4	21.8	20.3	21.1	18.7	21.5	19.9	19.5	21.1	+1.6	
Females	24.0	21.6	19.4	16.2	14.7	12.5	12.2	11.6	10.5	12.0	11.4	12.0	11.0	11.5	12.9	14.1	14.6	17.0	17.1	17.3	16.0	16.7	17.5	16.6	16.5	16.2	15.7	16.1	+0.4
<i>Percentage who used in past 30 days</i>																													
Any Illicit Drug																													
Total	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	19.8	18.9	20.6	19.9	21.8	+1.9 s
Males	29.9	27.1	23.7	21.1	18.8	18.3	17.9	17.4	19.5	18.6	19.0	19.8	20.1	20.0	21.5	21.9	22.8	22.4	23.1	22.0	22.5	22.7	22.8	22.4	23.9	24.5	23.8	26.4	+2.6
Females	22.2	20.2	17.8	15.0	13.5	12.5	12.4	12.9	12.1	13.5	13.3	13.8	13.2	15.0	15.6	16.6	16.3	18.3	16.3	16.4	15.7	16.4	16.9	18.0	15.5	18.2	17.3	18.7	+1.4
Any Illicit Drug other than Marijuana																													
Total	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	8.5	8.6	8.4	7.8	8.3	+0.5
Males	15.2	12.3	10.6	9.1	6.8	6.6	6.5	5.9	7.1	6.8	5.7	6.8	7.1	7.3	7.8	8.1	8.5	9.2	10.6	9.2	10.2	10.0	10.0	8.5	10.0	10.0	9.0	9.6	+0.6
Females	11.0	9.4	8.7	6.2	5.3	4.4	4.7	4.0	3.9	4.8	4.0	4.5	4.4	5.1	5.4	6.3	7.1	7.7	7.1	7.6	6.8	7.7	8.1	8.5	7.6	7.5	7.0	7.4	+0.4
<i>Approximate Weighted N</i>																													
All Respondents																													
Total	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900	4,900	4,600	4,600	4,400	
Males	3,200	3,100	3,000	2,900	3,000	3,000	3,000	3,000	2,900	2,800	2,700	2,800	2,700	2,600	2,400	2,400	2,200	2,200	2,300	2,200	2,100	1,900	2,000	2,000	2,000	1,800	1,900	1,800	
Females	3,700	3,700	3,700	3,700	3,700	3,600	3,700	3,700	3,600	3,600	3,600	3,600	3,500	3,400	3,300	3,400	3,100	3,100	3,400	3,200	3,000	2,900	2,900	2,900	2,800	2,700	2,600		


Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

FIGURE 5-1
ANY ILLICIT DRUG
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-1 (cont.)
ANY ILLICIT DRUG
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	48.1												
1977	51.1												
1978	53.8	55.8											
1979	54.2	54.5											
1980	53.1	54.5	55.3										
1981	52.1	53.4	55.4										
1982	49.4	50.2	51.2	51.7									
1983	47.4	47.4	49.9	48.9									
1984	45.8	45.9	47.3	44.0	44.0								
1985	46.3	45.7	46.3	47.8	45.2								
1986	44.3	42.6	45.8	42.8	39.3	38.4							
1987	41.7	39.5	42.3	37.9	40.1	36.2							
1988	38.5	39.4	38.2	36.6	34.4	32.5	30.5						
1989	35.4	35.7	35.0	31.4	30.5	30.9	28.9						
1990	32.5	32.3	32.7	30.7	29.6	27.4	23.0	23.7					
1991	29.4	28.1	29.9	27.0	25.2	23.9	24.5	23.8					
1992	27.1	29.7	30.0	29.2	26.4	25.3	23.1	21.9					
1993	31.0	30.5	30.2	29.8	25.6	24.6	21.7	22.3					
1994	35.8	32.2	31.6	27.3	25.5	23.6	22.4	22.4	19.5				
1995	39.0	35.6	31.9	28.5	27.3	23.9	21.3	19.8	21.6				
1996	40.2	36.1	33.0	27.6	23.4	23.7	22.7	21.7	21.2				
1997	42.4	36.7	33.5	27.3	25.4	20.7	22.2	21.2	20.3				
1998	41.4	40.6	34.1	27.4	23.9	22.0	19.6	19.3	18.1	20.3			
1999	42.1	40.4	33.3	31.1	24.5	20.8	19.0	17.7	17.7	16.7			
2000	40.9	39.3	36.9	29.6	25.5	21.4	20.3	17.6	19.1	17.2			
2001	41.4	38.4	40.2	31.1	27.4	22.9	21.1	20.2	17.8	15.8			
2002	41.0	39.4	36.7	35.2	27.6	22.9	20.9	—	18.1	18.2			
2003	39.3	38.1	38.3	34.6	27.5	26.3	20.6	—	17.9	15.8	17.8		
2004	38.8	38.0	36.5	34.5	31.6	26.8	22.0	—	18.5	17.5	15.8		
2005	38.4	38.9	36.4	31.9	32.0	24.3	25.2	—	18.2	19.1	15.3		
2006	36.5	36.3	36.0	32.7	28.6	25.7	25.9	—	17.5	16.2	17.2		
2007	35.9	35.2	35.0	34.1	29.3	28.5	22.7	—	17.5	17.4	18.3		
2008	36.6	35.5	36.7	34.4	31.8	30.1	28.2	—	22.1	17.5	17.3	17.9	
2009	36.5	35.5	38.8	34.1	29.6	27.4	27.9	—	20.0	19.1	17.0	16.0	
2010	38.3	32.5	38.1	36.3	31.6	27.1	26.2	—	20.2	16.7	19.1	18.3	
2011	40.0	37.9	37.5	35.4	32.1	29.9	26.2	—	24.2	16.9	17.8	16.8	
2012	39.7	36.2	36.8	35.3	29.7	31.6	25.1	—	21.1	17.6	18.6	18.6	
2013	40.3	37.5	40.9	37.1	34.6	30.8	26.6	—	23.3	18.7	17.7	17.0	16.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-2
ANY ILLICIT DRUG OTHER THAN MARIJUANA
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-2 (cont.)
ANY ILLICIT DRUG OTHER THAN MARIJUANA
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	25.4												
1977	26.0												
1978	27.1	28.6											
1979	28.2	30.2											
1980	30.4	33.3	35.5										
1981	34.0	34.2	37.0										
1982	30.1	32.4	34.2	35.4									
1983	28.4	29.8	33.7	33.2									
1984	28.0	27.5	31.6	29.4	30.2								
1985	27.4	26.9	29.5	33.4	30.3								
1986	25.9	24.7	29.1	29.3	25.5	26.5							
1987	24.1	22.2	25.6	22.6	25.7	23.3							
1988	21.1	21.3	22.8	21.1	21.0	20.4	20.0						
1989	20.0	17.6	19.4	18.8	17.6	18.2	17.4						
1990	17.9	16.5	17.4	17.5	16.6	15.2	12.4	13.8					
1991	16.2	13.8	14.9	14.6	14.4	13.6	13.2	13.1					
1992	14.9	13.4	15.4	14.8	13.4	13.2	11.6	10.7					
1993	17.1	13.5	13.5	12.9	13.0	11.5	9.9	9.5					
1994	18.0	14.6	14.1	12.9	12.0	11.1	10.8	11.5	11.2				
1995	19.4	18.6	15.2	11.5	11.6	10.9	11.0	8.2	10.4				
1996	19.8	17.4	13.7	13.1	10.0	10.7	10.3	10.2	11.4				
1997	20.7	17.6	17.7	12.1	10.7	8.4	11.0	10.8	10.0				
1998	20.2	17.3	15.3	12.9	10.8	8.9	7.8	9.6	8.2	9.3			
1999	20.7	18.7	14.1	14.8	11.6	8.6	8.1	8.3	9.3	7.9			
2000	20.4	19.6	17.0	15.0	12.5	9.9	7.4	7.4	9.3	7.7			
2001	21.6	18.0	20.0	14.1	13.3	11.4	9.9	9.7	8.8	7.3			
2002	20.9	19.6	18.9	17.2	14.6	11.4	10.9	—	9.6	9.7			
2003	19.8	19.9	20.7	20.1	14.5	15.1	11.6	—	9.5	6.7	8.9		
2004	20.5	20.2	21.2	21.2	16.3	14.6	11.8	—	11.0	8.3	9.3		
2005	19.7	20.2	20.5	18.0	19.7	14.2	15.8	—	10.5	9.4	8.4		
2006	19.2	18.1	22.0	19.4	16.9	15.1	15.3	—	10.8	9.8	10.3		
2007	18.5	17.8	19.7	19.1	17.0	16.9	13.0	—	11.0	11.3	10.7		
2008	18.3	16.8	19.5	21.3	19.1	18.0	16.5	—	13.7	11.3	10.7	10.0	
2009	17.0	14.6	22.9	17.6	17.8	14.1	17.2	—	13.3	10.4	9.6	10.3	
2010	17.3	17.2	20.0	20.1	19.5	15.8	14.5	—	12.5	9.3	11.5	10.8	
2011	17.6	17.4	18.2	19.3	17.3	15.8	13.7	—	13.6	9.6	9.8	9.4	
2012	17.0	17.0	17.9	18.8	15.0	17.2	13.7	—	12.5	10.8	11.3	10.2	
2013	17.3	17.3	21.6	18.3	17.4	15.8	15.2	—	13.0	9.6	9.5	8.6	7.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-3a
MARIJUANA
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-3a (cont.)

MARIJUANA

Trends in Annual Prevalence among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	44.5												
1977	47.6												
1978	50.2	52.8											
1979	50.8	51.0											
1980	48.8	49.7	50.1										
1981	46.1	49.0	51.1										
1982	44.3	44.9	45.8	46.0									
1983	42.3	43.0	45.4	43.8									
1984	40.0	41.4	42.1	38.6	38.3								
1985	40.6	40.3	40.9	42.0	39.2								
1986	38.8	39.1	39.6	36.6	34.1	32.5							
1987	36.3	35.8	37.4	33.7	35.4	31.4							
1988	33.1	36.2	33.7	32.0	29.7	26.7	25.4						
1989	29.6	32.2	31.6	27.3	26.2	26.8	24.7						
1990	27.0	28.4	28.2	26.6	24.1	22.6	20.0	19.8					
1991	23.9	25.4	26.8	23.2	21.8	20.9	21.0	19.9					
1992	21.9	26.9	26.9	26.6	23.5	21.2	20.1	17.7					
1993	26.0	27.9	26.1	26.5	22.2	21.3	18.8	19.9					
1994	30.7	29.3	29.2	24.6	22.6	20.1	19.0	18.6	14.5				
1995	34.7	31.8	28.1	25.8	24.4	20.4	18.2	17.2	17.2				
1996	35.8	34.2	30.6	25.8	21.7	20.6	19.5	18.6	16.3				
1997	38.5	34.8	30.6	25.1	23.3	18.0	18.0	16.7	17.5				
1998	37.5	37.2	31.9	25.5	21.2	19.9	16.9	15.8	14.9	17.1			
1999	37.8	37.9	31.5	27.4	21.8	18.2	16.0	14.8	14.7	13.8			
2000	36.5	37.0	33.2	26.9	22.7	18.8	18.4	14.5	13.8	13.7			
2001	37.0	35.4	37.5	28.3	25.0	19.4	17.1	16.7	14.8	12.5			
2002	36.2	36.4	34.3	31.8	24.5	19.4	17.5	—	13.7	14.6			
2003	34.9	35.9	33.1	30.0	24.3	21.2	17.0	—	13.0	13.4	14.0		
2004	34.3	34.5	32.5	27.7	27.6	22.4	16.4	—	13.0	13.9	11.9		
2005	33.6	34.9	32.6	26.8	26.4	19.7	18.9	—	12.9	14.3	11.7		
2006	31.5	33.2	31.1	28.5	24.0	20.9	19.9	—	11.4	11.0	11.6		
2007	31.7	33.1	30.5	29.3	24.7	24.4	18.3	—	10.8	11.6	12.6		
2008	32.4	32.1	33.3	27.4	25.9	23.6	22.3	—	14.2	10.7	11.1	11.7	
2009	32.8	33.2	33.7	29.5	25.2	23.3	22.5	—	12.6	12.2	11.6	10.1	
2010	34.8	30.6	34.0	30.5	25.5	22.3	21.5	—	14.6	12.0	12.7	11.4	
2011	36.4	34.4	34.8	31.8	27.0	25.8	20.9	—	17.7	10.6	11.6	10.8	
2012	36.4	34.0	34.0	30.3	25.6	26.5	19.8	—	14.4	12.5	12.3	12.2	
2013	36.4	35.5	36.7	34.3	28.4	25.2	22.4	—	17.1	14.3	11.9	11.9	12.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-3b
MARIJUANA
Trends in 30-Day Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-3b (cont.)

MARIJUANA

Trends in 30-Day Prevalence

among Respondents of Modal Ages 18 through 55, by Age Group


<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32^a</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976	32.2												
1977	35.4												
1978	37.1	38.0											
1979	36.5	37.5											
1980	33.7	33.9	35.9										
1981	31.6	34.2	35.3										
1982	28.5	28.6	29.1	30.3									
1983	27.0	25.7	29.3	29.7									
1984	25.2	24.6	26.4	25.4	24.9								
1985	25.7	22.8	25.2	26.8	24.8								
1986	23.4	22.9	23.3	23.0	19.9	20.7							
1987	21.0	20.4	21.8	19.6	21.5	20.3							
1988	18.0	20.1	18.5	17.4	17.2	16.1	15.4						
1989	16.7	16.3	15.9	15.6	14.7	14.7	15.0						
1990	14.0	15.2	14.3	13.4	13.4	12.9	11.5	11.5					
1991	13.8	13.2	14.7	13.0	13.0	13.5	12.7	12.1					
1992	11.9	14.1	14.7	12.5	12.6	12.0	12.2	11.3					
1993	15.5	14.6	13.8	13.6	12.4	12.3	11.2	11.7					
1994	19.0	15.3	16.5	13.3	12.9	11.6	11.4	10.8	8.7				
1995	21.2	18.7	15.4	12.2	11.7	10.4	10.8	11.1	11.1				
1996	21.9	19.9	16.4	14.2	12.6	11.0	10.5	10.9	8.8				
1997	23.7	19.9	18.9	14.0	10.5	10.1	9.4	10.0	10.7				
1998	22.8	20.1	17.5	13.8	11.8	10.5	9.0	8.7	9.1	10.5			
1999	23.1	23.1	17.8	15.3	12.0	8.9	9.3	8.5	8.8	8.3			
2000	21.6	22.3	19.8	14.7	12.5	10.7	9.8	7.7	8.3	8.5			
2001	22.4	21.0	22.9	14.9	14.5	10.3	8.3	9.6	8.8	8.3			
2002	21.5	22.2	20.1	17.2	14.8	9.9	9.0	—	8.9	8.1			
2003	21.2	22.5	18.2	18.9	14.5	12.2	8.9	—	7.1	8.2	8.4		
2004	19.9	20.7	18.3	15.6	15.1	12.0	8.5	—	7.8	8.3	6.5		
2005	19.8	18.9	17.9	14.1	15.9	11.9	11.9	—	7.0	8.1	7.2		
2006	18.3	17.5	17.4	16.2	14.0	13.1	10.1	—	6.2	6.7	6.3		
2007	18.8	18.4	18.0	16.2	13.6	13.5	10.4	—	5.8	6.7	6.9		
2008	19.4	17.9	17.8	16.2	13.3	14.2	12.9	—	7.8	6.6	6.4	7.2	
2009	20.6	19.5	20.0	16.0	15.3	13.3	12.1	—	5.9	6.8	7.3	5.9	
2010	21.4	18.0	18.0	17.3	13.6	13.5	11.0	—	8.9	7.1	7.3	6.8	
2011	22.6	20.4	21.9	18.1	15.5	15.0	10.9	—	10.1	6.5	7.3	5.9	
2012	22.9	21.6	19.8	18.0	14.0	14.6	11.5	—	9.1	6.5	6.6	7.3	
2013	22.7	21.8	23.0	20.0	15.8	13.9	13.7	—	10.4	8.2	5.7	7.5	7.6

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-3c
MARIJUANA DAILY
Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-3c (cont.)

MARIJUANA

**Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group**


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	8.2												
1977	9.1												
1978	10.7	10.5											
1979	10.3	10.9											
1980	9.1	8.1	10.9										
1981	7.0	7.9	9.4										
1982	6.3	6.6	6.4	8.1									
1983	5.5	5.2	6.2	6.7									
1984	5.0	4.7	5.3	5.5	6.0								
1985	4.9	4.6	4.5	5.8	6.1								
1986	4.0	3.5	4.1	4.9	3.6	4.8							
1987	3.3	3.4	3.9	4.3	5.0	4.6							
1988	2.7	3.5	3.5	3.1	3.4	3.0	3.2						
1989	2.9	2.8	3.1	3.0	3.3	4.1	3.2						
1990	2.2	2.3	2.5	2.7	2.7	2.4	2.2	2.2					
1991	2.0	2.1	2.4	2.1	2.5	2.6	2.6	2.5					
1992	1.9	1.4	2.6	2.3	2.6	2.5	2.9	2.1					
1993	2.4	2.3	2.3	2.7	2.5	2.3	2.7	2.6					
1994	3.5	3.1	2.9	3.1	2.7	2.2	2.4	2.7	2.3				
1995	4.6	4.7	3.4	3.3	2.3	2.5	2.5	3.1	2.6				
1996	4.9	4.9	3.2	2.3	3.1	2.5	2.2	2.8	2.3				
1997	5.8	5.4	5.3	2.6	2.5	2.7	2.3	2.8	3.5				
1998	5.6	5.2	5.2	3.1	2.4	2.3	2.4	2.8	2.7	3.2			
1999	6.0	6.2	4.6	5.1	3.1	2.8	2.5	2.1	1.9	2.1			
2000	6.0	6.0	5.5	3.8	3.4	2.0	2.2	2.6	2.7	2.6			
2001	5.8	6.1	7.0	4.7	4.6	2.3	2.6	2.9	2.3	1.8			
2002	6.0	6.0	6.0	5.5	2.7	2.5	2.3	—	3.0	3.0			
2003	6.0	6.5	6.0	6.6	3.5	4.0	1.9	—	2.1	2.4	2.6		
2004	5.6	6.0	5.1	5.3	5.5	2.9	2.0	—	2.5	1.8	2.0		
2005	5.0	6.4	4.6	4.5	5.9	3.0	3.9	—	2.1	1.9	2.1		
2006	5.0	5.2	5.3	5.3	5.0	4.3	2.5	—	2.8	2.3	1.4		
2007	5.1	5.1	4.9	5.2	4.1	5.7	3.2	—	1.9	2.3	2.7		
2008	5.4	4.1	6.1	5.4	5.5	4.3	4.8	—	2.2	2.2	2.7	2.0	
2009	5.2	5.8	6.3	5.8	5.1	3.7	5.4	—	1.7	2.1	2.2	2.0	
2010	6.1	6.0	5.1	5.8	4.0	5.3	4.0	—	3.8	2.3	2.2	2.2	
2011	6.6	6.6	6.3	6.9	5.8	4.6	3.7	—	2.7	2.7	2.7	2.3	
2012	6.5	6.2	6.1	5.7	5.1	5.1	4.5	—	3.6	2.6	2.2	2.7	
2013	6.5	6.2	7.8	6.2	5.8	5.1	2.9	—	3.3	2.3	2.4	2.1	2.7

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-4
INHALANTS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32,^b by Age Group



(Figure continued on next page.)

FIGURE 5-4 (cont.)

INHALANTS^a

**Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32,^b
by Age Group**

Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^c
1976	3.0							
1977	3.7							
1978	4.1	2.6						
1979	5.4	2.4						
1980	4.6	2.5	1.8					
1981	4.1	2.2	2.0					
1982	4.5	2.7	1.9	1.9				
1983	4.3	3.0	2.0	1.4				
1984	5.1	2.9	1.7	1.5	0.6			
1985	5.7	3.4	1.8	2.1	0.8			
1986	6.1	3.5	2.6	1.2	1.2	0.8		
1987	6.9	4.2	3.0	1.4	0.9	0.7		
1988	6.5	4.4	2.7	1.0	0.5	0.1	0.5	
1989	5.9	3.7	2.1	1.9	0.5	0.8	0.4	
1990	6.9	4.0	2.3	1.4	0.5	0.7	0.3	0.3
1991	6.6	4.0	2.3	1.0	1.2	0.6	0.6	0.4
1992	6.2	3.5	3.0	1.3	0.6	0.4	0.6	0.6
1993	7.0	3.6	2.8	1.9	0.7	0.8	0.4	0.5
1994	7.7	3.1	3.3	1.9	0.7	0.6	0.6	0.3
1995	8.0	5.0	2.1	2.1	1.7	0.7	0.8	0.9
1996	7.6	4.2	2.7	1.8	0.9	0.7	*	0.2
1997	6.7	4.7	2.8	1.6	1.0	0.5	1.0	0.4
1998	6.2	4.1	2.4	1.1	1.7	0.9	0.1	0.8
1999	5.6	3.1	3.3	3.0	0.4	1.2	0.5	0.3
2000	5.9	3.2	3.0	2.4	1.0	0.9	0.1	0.5
2001	4.5	3.4	2.4	0.9	0.8	1.0	0.7	0.5
2002	4.5	2.8	1.9	1.9	0.8	0.6	0.5	—
2003	3.9	2.2	1.4	0.9	1.3	1.0	0.5	—
2004	4.2	3.1	2.1	1.6	1.2	0.3	0.6	—
2005	5.0	1.5	2.2	1.0	0.4	1.2	1.4	—
2006	4.5	2.4	2.1	0.9	0.5	0.4	0.5	—
2007	3.7	1.8	1.0	0.4	0.4	0.5	0.5	—
2008	3.8	1.8	1.5	1.7	1.1	0.9	0.7	—
2009	3.4	1.5	1.2	0.8	0.5	0.3	0.3	—
2010	3.6	1.9	2.1	1.1	0.6	0.5	0.7	—
2011	3.2	1.1	1.2	0.5	0.7	0.6	0.7	—
2012	2.9	2.1	0.9	1.5	0.7	0.2	0.6	—
2013	2.5	0.4	0.9	0.8	0.2	0.2	0.3	—

Source. The Monitoring the Future study, the University of Michigan.


Notes. * * * indicates a percentage of less than 0.05%. ' — ' indicates data not available.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites. Chapter 5, *Volume I*, shows that such an adjustment would flatten the trend for seniors considerably because the line was adjusted up more in the earlier years, when nitrite use was more prevalent. Questions about nitrite use were dropped from the follow-up questionnaires beginning in 1995.

^bQuestions about the use of inhalants were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

^cBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-5
HALLUCINOGENS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 50,^c by Age Group



(Figure continued on next page.)

FIGURE 5-5 (cont.)
HALLUCINOGENS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 50,^c by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32^b</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>
1976	9.4											
1977	8.8											
1978	9.6	9.5										
1979	9.9	10.9										
1980	9.3	9.7	10.1									
1981	9.0	8.6	10.9									
1982	8.1	9.9	9.3	8.1								
1983	7.3	7.2	7.4	7.4								
1984	6.5	6.0	7.5	5.4	4.7							
1985	6.3	5.1	5.7	4.9	4.7							
1986	6.0	6.3	5.7	4.6	3.0	2.4						
1987	6.4	5.9	5.2	3.7	2.4	2.7						
1988	5.5	5.8	5.8	3.8	2.5	1.3	2.1					
1989	5.6	5.8	4.3	3.8	2.0	1.7	1.4					
1990	5.9	6.3	5.0	4.4	2.3	1.8	1.2	1.0				
1991	5.8	6.2	5.7	4.4	3.2	2.4	1.5	1.3				
1992	5.9	6.7	7.2	4.2	3.7	2.2	1.9	1.2				
1993	7.4	6.9	5.0	4.7	3.0	2.1	1.3	0.9				
1994	7.6	6.7	6.8	4.3	3.0	2.4	1.5	1.0	0.8			
1995	9.3	9.6	6.6	4.9	3.7	2.3	1.9	1.3	0.7			
1996	10.1	10.1	6.2	5.4	3.2	2.2	1.4	0.9	0.5			
1997	9.8	9.6	8.0	5.0	3.7	1.8	1.6	1.6	1.0			
1998	9.0	8.1	6.7	5.2	3.2	2.0	1.4	0.9	0.6	0.8		
1999	9.4	9.4	6.8	5.9	2.7	1.7	1.4	1.3	0.8	0.5		
2000	8.1	8.0	7.4	4.9	3.9	2.6	1.7	0.9	0.5	0.9		
2001	9.1	9.0	8.1	4.6	3.1	1.8	1.7	1.5	0.8	0.2		
2002	6.6	7.3	5.8	5.2	2.8	2.2	2.0	—	0.3	0.7		
2003	5.9	7.7	7.1	5.8	2.8	2.5	1.5	—	0.6	0.5	0.6	
2004	6.2	6.3	6.7	4.4	3.2	2.6	1.4	—	1.0	0.5	0.3	
2005	5.5	6.4	5.3	4.0	4.3	2.1	2.1	—	0.3	0.4	0.1	
2006	4.9	5.8	5.3	4.6	2.1	2.4	1.5	—	0.4	0.1	0.1	
2007	5.4	5.4	4.8	3.5	2.7	2.6	1.3	—	0.4	0.4	0.2	
2008	5.9	5.2	5.5	3.3	3.2	1.7	2.9	—	1.1	0.2	0.1	0.2
2009	4.7	4.7	5.8	3.7	2.8	2.4	2.2	—	0.8	0.5	0.3	0.3
2010	5.5	5.3	5.1	4.7	3.5	2.3	2.1	—	1.8	0.6	0.3	0.2
2011	5.2	4.6	5.3	3.2	3.0	2.4	1.7	—	1.4	0.8	0.7	0.1
2012	4.8	5.3	3.9	3.7	2.6	2.1	2.3	—	1.1	0.2	0.6	0.1
2013	4.5	5.0	4.7	4.1	2.7	2.7	2.6	—	1.2	0.5	0.3	0.1

Source. The Monitoring the Future study, the University of Michigan.


Notes. '—' indicates data not available.

^aUnadjusted for the possible underreporting of PCP.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

^cQuestions about the use of inhalants were not included in the questionnaires for 55-year-olds.

FIGURE 5-6
LSD
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 35,^b by Age Group



(Figure continued on next page.)

FIGURE 5-6 (cont.)

LSD

**Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 35, by Age Group**

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32^a</u>	<u>Age 35^b</u>
1976	6.4								
1977	5.5								
1978	6.3	6.2							
1979	6.6	8.1							
1980	6.5	7.2	7.9						
1981	6.5	6.4	8.0						
1982	6.1	7.7	6.9	6.0					
1983	5.4	5.4	4.9	4.6					
1984	4.7	4.3	5.1	3.1	2.7				
1985	4.4	3.3	3.3	2.9	2.9				
1986	4.5	4.5	4.4	2.7	1.5	1.6			
1987	5.2	4.7	3.7	2.2	1.6	1.8			
1988	4.8	4.9	4.2	2.9	1.6	0.8	1.5		
1989	4.9	4.5	3.2	2.7	1.4	1.1	0.8		
1990	5.4	5.3	4.0	3.5	1.8	1.5	0.8	0.6	
1991	5.2	5.4	5.0	3.8	2.5	1.9	1.0	0.8	
1992	5.6	6.3	6.0	3.5	3.2	1.6	1.4	1.0	
1993	6.8	6.2	4.3	3.5	2.4	1.8	1.0	0.7	
1994	6.9	6.2	5.7	3.2	2.4	1.6	1.0	0.6	0.6
1995	8.4	8.2	5.5	4.1	2.6	1.7	1.4	1.1	0.4
1996	8.8	8.7	4.9	4.6	2.0	1.6	1.0	0.5	0.5
1997	8.4	7.8	5.5	4.0	2.6	1.3	0.8	1.0	0.5
1998	7.6	5.9	4.4	3.5	2.1	1.0	1.0	0.7	0.3
1999	8.1	7.7	4.5	4.3	1.9	1.2	0.8	0.9	0.6
2000	6.6	6.3	4.9	2.6	2.5	1.6	1.0	0.2	0.3
2001	6.6	6.4	4.7	2.5	1.7	1.3	1.1	0.7	0.5
2002	3.5	3.3	1.8	2.2	1.0	0.7	0.9	—	*
2003	1.9	1.9	1.2	1.4	0.6	0.6	0.4	—	*
2004	2.2	1.5	1.4	0.7	0.5	0.3	0.2	—	0.4
2005	1.8	1.5	1.0	0.7	0.7	0.1	0.4	—	0.1
2006	1.7	2.1	1.4	1.1	0.6	0.7	0.2	—	0.1
2007	2.1	1.5	1.3	1.4	0.9	0.6	0.3	—	—
2008	2.7	2.0	2.7	0.9	0.8	0.4	0.5	—	—
2009	1.9	2.2	2.7	1.4	1.0	0.6	0.2	—	—
2010	2.6	2.4	1.5	1.7	1.1	0.6	0.2	—	—
2011	2.7	2.6	2.7	1.5	1.1	0.4	0.9	—	—
2012	2.4	2.5	1.6	1.6	1.3	0.7	0.3	—	—
2013	2.2	2.9	2.9	2.0	1.5	0.7	0.7	—	—


Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a percentage of less than 0.05%. ' — ' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

^bQuestions about LSD use were not included in the questionnaires administered to the 40-, 45-, and 50-year-olds, or the 35-year-olds after 2006.

FIGURE 5-7
HALLUCINOGENS OTHER THAN LSD^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 35, by Age Group



(Figure continued on next page.)

FIGURE 5-7 (cont.)
HALLUCINOGENS OTHER THAN LSD^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 35, by Age Group

Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^b	Age 35 ^c
1976	7.0								
1977	6.9								
1978	7.3	7.1							
1979	6.8	7.3							
1980	6.2	5.4	5.8						
1981	5.6	4.6	6.5						
1982	4.7	6.1	5.2	4.0					
1983	4.1	3.9	4.3	4.2					
1984	3.8	3.2	4.1	3.5	3.0				
1985	3.6	3.2	3.7	2.8	2.7				
1986	3.0	3.6	2.6	2.7	2.0	1.4			
1987	3.2	2.5	2.7	2.4	1.3	1.5			
1988	2.1	2.4	3.2	1.8	1.2	0.7	0.9		
1989	2.2	3.0	2.0	1.9	1.2	0.8	0.9		
1990	2.1	2.6	2.4	1.6	1.0	0.7	0.6	0.5	
1991	2.0	2.6	2.2	1.4	1.3	1.0	0.6	0.8	
1992	1.7	1.9	3.1	1.9	1.5	0.8	0.8	0.4	
1993	2.2	2.8	1.9	2.2	1.4	0.9	0.8	0.3	
1994	3.1	2.2	3.1	2.0	1.3	1.2	0.9	0.5	0.5
1995	3.8	3.9	3.2	1.9	1.8	1.1	0.7	0.5	0.3
1996	4.4	4.4	3.5	2.8	1.6	0.9	0.7	0.6	0.1
1997	4.6	5.1	5.2	2.3	1.6	1.0	1.1	0.8	0.6
1998	4.6	4.8	3.7	3.1	1.8	1.4	0.7	0.4	0.5
1999	4.3	4.2	4.2	3.6	1.5	0.9	0.8	0.6	0.4
2000	4.4	4.6	4.7	3.7	2.1	1.6	1.1	0.6	0.3
2001	5.9	5.5	5.9	3.0	1.9	0.9	1.0	1.1	0.4
2002	5.4	6.5	5.2	4.1	2.5	1.7	1.6	—	0.3
2003	5.4	7.3	6.9	5.5	2.5	2.2	1.3	—	0.6
2004	5.6	6.0	6.3	4.0	3.1	2.4	1.4	—	0.8
2005	5.0	6.2	5.0	3.7	4.0	2.1	1.9	—	0.2
2006	4.6	5.3	4.9	4.2	2.0	2.1	1.4	—	0.4
2007	4.8	5.2	4.7	3.0	2.4	2.5	1.2	—	—
2008	5.0	4.7	4.5	3.0	2.8	1.6	2.6	—	—
2009	4.2	4.1	4.5	3.1	2.4	2.2	2.2	—	—
2010	4.8	4.9	4.6	3.9	3.0	2.1	2.0	—	—
2011	4.3	3.9	4.3	2.6	2.8	2.3	1.3	—	—
2012	4.0	4.4	3.3	3.0	1.9	1.9	2.1	—	—
2013	3.7	4.1	3.7	3.4	2.2	2.4	2.3	—	—

Source. The Monitoring the Future study, the University of Michigan.


Notes. '—' indicates data not available.

^aUnadjusted for the possible underreporting of PCP.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

^cQuestions about the use of hallucinogens other than LSD were not included in the questionnaires administered to the 40-, 45-, and 50-year-olds, or the 35-year-olds after 2006.

FIGURE 5-8
ECSTASY (MDMA)
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32, ^a by Age Group



(Figure continued on next page.)

FIGURE 5-8 (cont.)
ECSTASY (MDMA)
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32, ^a
by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32 ^b</u>
1976								
1977								
1978								
1979								
1980								
1981								
1982								
1983								
1984								
1985								
1986								
1987								
1988								
1989		1.9	2.1	1.3	0.3	1.0	0.1	
1990		2.2	2.0	1.5	1.0	0.7	0.3	0.5
1991		0.6	1.1	1.0	0.6	0.6	0.7	0.2
1992		1.8	1.7	0.4	0.8	0.4	*	0.4
1993		1.1	0.5	1.2	0.5	0.7	*	0.2
1994		0.6	1.4	0.9	0.2	0.4	0.3	0.2
1995		2.2	2.1	1.3	1.2	1.4	0.3	0.2
1996	4.6	3.7	1.9	1.1	1.1	0.7	0.5	0.1
1997	4.0	3.1	3.9	1.3	1.6	0.5	1.4	0.6
1998	3.6	4.0	3.7	2.3	1.8	2.3	*	0.9
1999	5.6	4.9	4.6	3.3	3.4	1.8	0.7	0.8
2000	8.2	9.1	9.8	7.0	6.9	2.6	2.4	0.3
2001	9.2	11.0	10.8	6.8	4.3	4.1	2.6	1.4
2002	7.4	6.3	9.3	8.3	4.4	2.6	2.4	—
2003	4.5	5.3	5.3	5.2	3.4	3.1	1.2	—
2004	4.0	4.2	2.4	3.2	4.0	3.7	0.9	—
2005	3.0	4.1	3.3	2.8	2.3	2.5	2.0	—
2006	4.1	3.8	3.4	3.6	2.0	2.1	2.2	—
2007	4.5	3.1	2.7	2.6	1.9	1.9	1.4	—
2008	4.3	4.7	4.7	2.0	2.7	2.1	2.0	—
2009	4.3	3.4	3.9	3.4	2.7	1.8	1.9	—
2010	4.5	5.0	4.8	4.4	1.6	1.8	1.0	—
2011	5.3	4.8	4.7	3.7	2.5	2.3	2.0	—
2012	3.8	5.8	5.5	4.2	2.6	2.1	2.2	—
2013	4.0	5.0	5.9	4.9	3.3	2.0	1.6	—


Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a percentage of less than 0.05%. ' — ' indicates data not available.

^aQuestions about use of ecstasy (MDMA) were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-9
COCAINE
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-9 (cont.)
COCAINE
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	6.0												
1977	7.2												
1978	9.0	11.8											
1979	12.0	15.0											
1980	12.3	16.3	19.8										
1981	12.4	15.9	20.5										
1982	11.5	16.9	21.6	22.9									
1983	11.4	13.8	21.2	20.8									
1984	11.6	14.6	20.6	20.2	21.1								
1985	13.1	15.4	19.2	23.5	21.6								
1986	12.7	15.9	20.4	22.8	19.7	19.9							
1987	10.3	13.4	16.0	16.2	17.4	15.6							
1988	7.9	10.6	14.1	15.1	15.2	14.2	14.0						
1989	6.5	7.6	11.8	12.0	10.7	12.2	11.6						
1990	5.3	5.6	8.7	9.5	9.9	9.9	8.1	8.9					
1991	3.5	3.8	6.1	7.2	7.4	6.9	6.7	6.8					
1992	3.1	3.7	5.1	6.5	6.6	7.2	6.7	5.7					
1993	3.3	3.2	4.1	4.6	6.3	5.8	4.7	5.1					
1994	3.6	3.2	3.9	4.8	4.2	5.4	6.0	5.5	4.7				
1995	4.0	3.9	4.3	4.5	4.6	4.6	4.5	3.8	4.3				
1996	4.9	3.7	4.2	4.8	3.8	4.3	4.3	5.0	5.1				
1997	5.5	4.5	5.8	4.9	4.3	3.7	4.3	5.1	4.1				
1998	5.7	5.3	6.0	5.2	3.7	3.9	3.7	4.4	4.4	4.5			
1999	6.2	5.7	5.6	6.8	5.0	3.9	3.6	4.1	4.6	4.1			
2000	5.0	5.8	6.3	6.3	4.8	3.6	2.7	3.0	3.9	3.5			
2001	4.8	6.0	7.5	5.4	5.4	4.8	2.8	3.5	3.5	3.0			
2002	5.0	6.5	7.0	6.0	5.6	4.0	4.4	—	3.6	3.7			
2003	4.8	6.3	7.4	8.3	5.4	5.5	4.9	—	2.7	3.1	3.4		
2004	5.3	6.3	8.6	8.4	6.7	5.2	3.6	—	3.3	3.3	3.9		
2005	5.1	6.4	7.5	6.7	8.2	5.7	4.5	—	2.8	2.8	2.9		
2006	5.7	5.7	8.4	6.9	6.6	5.2	4.7	—	2.5	3.0	3.4		
2007	5.2	5.8	7.2	5.8	6.4	5.9	4.1	—	2.0	2.7	3.6		
2008	4.4	5.0	7.3	5.8	6.5	5.3	5.2	—	3.3	2.0	2.7	2.0	
2009	3.4	3.2	6.9	6.9	4.5	4.7	5.6	—	2.5	2.2	3.4	2.6	
2010	2.9	3.4	4.9	5.9	4.8	4.7	4.0	—	2.6	2.0	2.4	1.8	
2011	2.9	3.9	4.3	6.1	6.0	3.4	3.3	—	2.9	1.3	2.1	1.6	
2012	2.7	3.4	3.5	5.4	3.8	4.2	3.3	—	2.3	1.5	1.5	1.8	
2013	2.6	2.6	4.8	4.2	4.4	3.5	3.4	—	3.3	2.2	2.0	1.7	1.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-10
CRACK COCAINE
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-10 (cont.)

CRACK COCAINE

Trends in Annual Prevalence

among Respondents of Modal Ages 18 through 55, by Age Group


<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32^a</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976													
1977													
1978													
1979													
1980													
1981													
1982													
1983													
1984													
1985													
1986	4.1												
1987	3.9	2.7	4.1	3.4	2.4	3.0							
1988	3.1	2.7	2.9	4.0	2.7	3.0	3.2						
1989	3.1	1.8	3.6	3.1	1.9	2.0	2.8						
1990	1.9	1.0	1.6	2.1	2.3	1.3	1.7	1.5					
1991	1.5	0.6	1.3	1.4	1.5	1.6	1.1	1.3					
1992	1.5	1.3	1.3	1.3	1.3	1.6	0.9	1.1					
1993	1.5	1.2	1.1	1.2	1.3	1.7	1.0	0.9					
1994	1.9	1.2	1.1	0.8	1.0	1.5	1.5	1.0	1.0				
1995	2.1	1.0	1.3	1.0	1.1	0.9	1.5	1.0	1.0				
1996	2.1	1.3	1.4	1.2	0.7	0.8	1.0	1.3	1.5				
1997	2.4	1.3	1.2	1.0	0.7	0.6	1.5	0.7	0.8				
1998	2.5	1.3	1.4	1.2	0.4	0.8	0.7	0.9	1.2	1.5			
1999	2.7	2.4	1.2	1.5	0.9	0.8	1.3	0.8	1.1	1.0			
2000	2.2	1.6	1.6	1.2	1.1	0.6	0.5	0.7	0.8	0.5			
2001	2.1	1.7	1.4	1.1	1.0	1.3	0.7	1.0	0.7	0.6			
2002	2.3	1.7	1.0	1.1	0.7	0.6	0.9	—	1.3	1.0			
2003	2.2	1.8	1.2	1.1	0.5	0.6	0.8	—	0.7	0.8	1.1		
2004	2.3	1.6	1.8	1.2	1.1	0.8	0.6	—	0.5	0.8	1.2		
2005	1.9	1.3	1.8	1.1	0.8	1.0	1.2	—	0.7	0.8	0.6		
2006	2.1	1.4	1.5	0.9	1.2	0.5	0.3	—	0.8	0.5	1.1		
2007	1.9	0.8	0.8	1.0	1.5	1.0	0.3	—	0.5	0.3	0.7		
2008	1.6	1.2	0.9	0.9	0.9	0.7	0.5	—	0.5	0.4	0.7	0.4	
2009	1.3	0.8	0.7	0.9	0.6	0.6	0.7	—	0.8	0.4	1.0	0.7	
2010	1.4	0.8	0.3	0.6	0.5	0.4	0.6	—	0.6	0.2	0.4	0.7	
2011	1.0	0.6	0.5	0.6	0.5	0.7	0.6	—	0.1	0.2	0.6	0.5	
2012	1.2	0.6	0.3	0.6	0.4	0.4	0.3	—	0.2	0.3	0.6	0.5	
2013	1.1	0.3	0.7	0.2	0.5	*	0.3	—	0.2	0.1	0.5	0.4	0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-11
HEROIN
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-11 (cont.)

HEROIN

**Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group**


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	0.8												
1977	0.8												
1978	0.8	0.4											
1979	0.5	0.3											
1980	0.5	0.2	0.6										
1981	0.5	0.5	0.4										
1982	0.6	0.2	0.4	0.2									
1983	0.6	0.2	0.3	0.6									
1984	0.5	0.2	0.3	0.2	0.2								
1985	0.6	0.1	0.3	0.2	0.3								
1986	0.5	0.1	0.2	0.1	0.2	0.3							
1987	0.5	0.2	0.3	0.1	0.3	0.3							
1988	0.5	0.1	0.2	0.1	0.1	0.3	0.2						
1989	0.6	0.2	0.2	0.1	0.1	0.2	0.3						
1990	0.5	*	0.1	0.1	0.1	*	*	0.1					
1991	0.4	0.1	0.2	0.2	0.1	0.1	0.2	0.1					
1992	0.6	0.1	0.3	0.2	0.1	0.2	0.1	*					
1993	0.5	0.1	0.2	0.2	0.1	0.2	*	*					
1994	0.6	0.1	0.1	0.1	0.2	0.2	0.3	0.1	0.3				
1995	1.1	0.5	0.4	0.6	0.1	0.1	0.4	0.1	0.2				
1996	1.0	0.7	0.5	0.3	0.3	0.1	0.2	0.3	0.2				
1997	1.2	0.4	0.7	0.1	0.3	0.2	0.3	0.3	0.1				
1998	1.0	1.1	0.4	0.2	0.3	0.1	0.1	0.1	0.1	0.4			
1999	1.1	0.6	0.4	0.5	0.2	0.3	0.1	0.2	0.4	*			
2000	1.5	0.7	0.5	0.4	0.3	0.1	0.1	0.2	0.1	0.1			
2001	0.9	1.0	0.5	0.5	0.3	0.3	0.2	0.0	0.1	0.1			
2002	1.0	0.4	0.1	0.3	0.3	0.1	0.2	—	0.2	0.3			
2003	0.8	0.4	0.4	0.5	0.1	0.6	0.2	—	0.3	0.2	0.2		
2004	0.9	0.4	0.5	0.3	0.2	0.1	0.3	—	0.1	*	0.2		
2005	0.8	0.6	0.7	0.3	0.2	0.4	0.5	—	*	*	*		
2006	0.8	0.5	0.6	0.3	0.4	*	0.1	—	0.3	0.2	0.1		
2007	0.9	0.3	0.5	0.3	0.2	0.2	0.2	—	0.1	0.1	0.1		
2008	0.7	0.6	0.9	0.6	0.3	0.2	0.3	—	0.1	*	*	0.2	
2009	0.7	0.8	0.7	0.7	0.5	0.3	0.3	—	0.4	*	0.1	0.3	
2010	0.9	0.4	0.3	0.8	0.6	0.3	0.2	—	0.3	0.1	*	0.2	
2011	0.8	0.4	0.4	0.6	0.5	0.4	0.3	—	0.0	0.1	0.1	*	
2012	0.6	0.4	0.6	0.3	0.7	0.3	0.4	—	0.1	0.1	0.1	0.1	
2013	0.6	0.5	0.8	0.6	0.8	0.6	0.4	—	0.6	0.2	0.2	0.1	*

Source. The Monitoring the Future study, the University of Michigan.

Notes. '**' indicates a percentage of less than 0.05%. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-12
NARCOTICS OTHER THAN HEROIN ^{a,b}
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-12 (cont.)
NARCOTICS OTHER THAN HEROIN ^{a,b}

Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32</u> ^c	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976	5.7												
1977	6.4												
1978	6.0	4.7											
1979	6.2	4.7											
1980	6.3	5.6	4.9										
1981	5.9	4.9	5.0										
1982	5.3	4.4	3.5	4.4									
1983	5.1	4.2	4.0	3.3									
1984	5.2	3.9	3.3	3.5	2.7								
1985	5.9	3.4	3.8	3.8	3.4								
1986	5.2	4.2	3.8	2.7	2.0	2.7							
1987	5.3	3.7	3.6	2.4	2.5	3.0							
1988	4.6	3.1	3.6	2.3	2.5	1.6	2.2						
1989	4.4	3.0	3.4	2.4	2.4	2.9	2.1						
1990	4.5	3.9	2.7	2.7	2.3	1.5	1.5	1.4					
1991	3.5	3.2	2.4	2.4	2.4	1.8	1.8	1.7					
1992	3.3	2.2	3.4	2.2	2.6	1.7	1.9	1.5					
1993	3.6	2.5	2.9	2.0	2.0	1.4	1.3	1.5					
1994	3.8	2.7	2.9	2.6	1.8	2.1	1.7	1.9	1.7				
1995	4.7	4.7	3.1	2.5	1.8	2.4	1.9	1.8	1.6				
1996	5.4	4.7	2.6	2.9	2.1	2.0	2.1	1.5	1.8				
1997	6.2	4.3	5.0	2.7	2.2	2.0	2.6	2.2	2.0				
1998	6.3	4.3	4.3	3.5	3.0	1.9	1.5	2.0	1.2	1.7			
1999	6.7	5.5	4.2	4.1	3.0	1.8	1.8	2.2	2.1	1.4			
2000	7.0	6.2	5.0	4.2	2.9	2.1	2.5	1.4	2.1	1.7			
2001	6.7	7.0	6.8	4.3	3.7	3.1	2.9	2.9	2.4	1.6			
2002	9.4	8.3	8.9	8.2	6.0	4.3	4.2	—	4.4	3.4			
2003	9.3	9.9	9.6	9.7	6.4	6.7	5.1	—	3.4	2.3	2.8		
2004	9.5	10.4	9.2	9.5	7.9	7.5	5.4	—	4.8	2.9	3.4		
2005	9.0	9.9	10.2	7.6	8.8	6.9	7.8	—	4.3	3.4	3.1		
2006	9.0	8.6	11.5	9.5	8.5	7.0	7.7	—	5.6	4.5	3.5		
2007	9.2	8.2	9.4	9.4	8.5	8.1	6.1	—	3.8	5.8	4.4		
2008	9.1	8.6	8.4	10.5	9.4	8.6	7.1	—	7.4	4.7	3.2	3.3	
2009	9.2	6.4	11.0	8.0	9.8	6.7	8.9	—	6.0	4.7	4.1	4.0	
2010	8.7	8.1	9.2	10.0	10.1	7.8	6.9	—	6.7	4.0	5.0	4.7	
2011	8.7	7.7	7.7	7.8	9.0	7.6	6.7	—	5.8	4.9	4.0	4.2	
2012	7.9	6.5	7.0	7.9	7.1	8.2	6.0	—	6.3	4.7	4.6	4.4	
2013	7.1	7.1	6.9	7.2	6.9	6.8	6.5	—	6.6	4.1	3.0	3.1	2.5

Source. The Monitoring the Future study, the University of Michigan.


Notes. — indicates data not available.

^aIn 2002 the question text was changed on half of the questionnaire forms for 18- to 30-year-olds. The list of examples of narcotics other than heroin was updated. Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2001 data presented here are based on all forms. The 2002 data are based on the changed forms only. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003. Beginning in 2002, data were based on the changed question text for 35- and 40-year-olds.

^bUsed without a doctor's orders.

^cBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-13
AMPHETAMINES^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-13 (cont.)

AMPHETAMINES ^a

**Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group**

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32 ^b</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976	15.8												
1977	16.3												
1978	17.1	18.2											
1979	18.3	21.5											
1980	20.8	23.8	25.5										
1981	26.0	25.5	26.7										
1982	20.3	23.9	22.4	21.8									
1983	17.9	19.7	19.9	18.3									
1984	17.7	15.8	17.4	14.0	14.9								
1985	15.8	14.5	13.0	14.1	12.5								
1986	13.4	11.0	13.0	11.4	8.6	9.1							
1987	12.2	9.1	9.9	7.9	8.3	7.9							
1988	10.9	9.2	8.1	7.6	6.4	5.0	5.5						
1989	10.8	6.9	6.8	5.1	5.5	4.3	5.0						
1990	9.1	6.6	5.5	5.3	4.0	4.3	2.7	3.7					
1991	8.2	4.9	4.9	3.8	3.4	4.0	2.9	3.7					
1992	7.1	5.6	4.3	4.0	2.7	3.5	3.3	2.6					
1993	8.4	5.4	4.8	3.8	2.9	2.6	2.4	2.4					
1994	9.4	5.4	5.3	4.5	3.9	2.9	2.6	2.5	2.4				
1995	9.3	7.2	5.7	3.0	3.5	2.7	2.5	1.5	1.9				
1996	9.5	6.5	4.9	4.1	2.5	2.5	2.6	2.7	1.9				
1997	10.2	5.9	7.3	3.8	3.2	2.0	2.7	2.6	1.7				
1998	10.1	7.5	5.0	4.3	2.9	2.3	1.8	2.6	1.7	1.7			
1999	10.2	7.9	5.0	4.5	3.4	2.6	2.4	1.9	1.9	1.2			
2000	10.5	9.3	6.0	4.8	3.6	2.7	1.4	1.9	1.8	1.4			
2001	10.9	8.7	7.9	5.2	3.6	3.2	1.9	1.5	1.9	1.0			
2002	11.1	9.1	7.1	5.8	3.9	3.3	2.1	—	1.2	1.4			
2003	9.9	8.6	7.5	5.8	3.1	3.6	2.6	—	2.2	1.0	1.4		
2004	10.0	8.5	6.7	7.1	4.6	3.9	2.2	—	1.5	1.1	0.9		
2005	8.6	7.0	6.8	5.0	3.8	2.6	3.0	—	1.2	0.8	0.5		
2006	8.1	6.5	7.6	6.1	4.4	3.3	2.9	—	1.4	1.6	1.4		
2007	7.5	6.7	7.5	5.9	4.2	3.3	2.1	—	1.5	0.8	1.1		
2008	6.8	5.9	6.7	5.7	4.0	4.1	2.6	—	0.7	1.1	0.9	0.6	
2009	6.6	6.2	9.0	5.4	5.3	3.5	2.9	—	1.9	0.9	0.7	1.0	
2010	7.4	8.3	9.0	7.7	5.9	4.3	3.2	—	1.3	0.7	0.8	0.8	
2011	8.2	8.7	8.8	8.8	5.3	3.8	3.2	—	1.4	0.6	1.3	1.0	
2012	7.9	9.3	9.4	8.4	5.8	5.5	4.3	—	1.4	1.0	1.0	0.4	
2013	8.4	8.7	11.1	7.4	6.3	5.1	4.5	—	2.2	1.5	1.2	1.0	0.7


Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aUsed without a doctor's orders.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-14
CRYSTAL METHAMPHETAMINE (ICE)
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32, ^a by Age Group



(Figure continued on next page.)

FIGURE 5-14 (cont.)
CRYSTAL METHAMPHETAMINE (ICE)
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32, ^a
by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32 ^b</u>
1976								
1977								
1978								
1979								
1980								
1981								
1982								
1983								
1984								
1985								
1986								
1987								
1988								
1989								
1990	1.3	0.3	0.7	0.4	0.3	0.5	0.3	0.3
1991	1.4	0.4	0.5	0.4	0.2	*	0.1	0.2
1992	1.3	0.3	0.1	0.7	0.8	0.3	0.4	0.2
1993	1.7	1.4	1.0	0.9	0.1	0.6	0.3	0.2
1994	1.8	1.3	0.4	1.7	0.6	0.3	0.7	0.3
1995	2.4	1.1	2.2	0.8	0.2	0.8	0.6	*
1996	2.8	1.5	0.7	1.4	0.1	0.5	0.2	0.4
1997	2.3	0.7	1.5	1.3	0.4	0.3	0.7	0.7
1998	3.0	2.0	1.1	1.4	0.6	*	*	1.2
1999	1.9	1.4	0.6	1.5	0.8	0.4	*	*
2000	2.2	1.3	1.2	2.3	0.7	0.7	*	*
2001	2.5	1.9	0.9	0.6	1.1	0.8	0.3	0.7
2002	3.0	2.6	1.2	1.6	0.9	0.8	0.7	—
2003	3.0	2.6	1.2	1.6	0.9	0.8	0.7	—
2004	2.0	1.7	1.2	1.1	0.9	1.3	0.1	—
2005	2.3	2.4	1.7	1.0	1.7	1.3	1.7	—
2006	1.9	1.6	1.7	0.5	0.6	1.0	0.9	—
2007	1.6	0.9	0.9	1.8	1.2	0.6	0.7	—
2008	1.1	0.7	0.2	1.4	0.8	0.7	0.3	—
2009	0.9	0.5	0.5	1.3	0.8	0.7	0.3	—
2010	0.9	0.7	0.5	0.3	0.7	*	0.1	—
2011	1.2	0.4	0.5	0.3	0.3	0.9	0.1	—
2012	0.8	0.7	0.5	0.1	0.6	1.0	0.5	—
2013	1.1	0.3	0.2	1.2	1.2	1.0	1.2	—


Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a percentage of less than 0.05%. ' — ' indicates data not available.

^aQuestions about use of ecstasy (MDMA) were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-15
SEDATIVES (BARBITURATES) ^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-15 (cont.)
SEDATIVES (BARBITURATES) ^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group

Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^b	Age 35	Age 40	Age 45	Age 50	Age 55
1976	9.6												
1977	9.3												
1978	8.1	6.4											
1979	7.5	6.9											
1980	6.8	4.5	5.7										
1981	6.6	4.7	5.8										
1982	5.5	4.4	4.1	4.1									
1983	5.2	3.5	3.1	3.7									
1984	4.9	3.5	2.5	2.6	3.3								
1985	4.6	2.0	2.3	3.0	3.4								
1986	4.2	2.2	2.9	2.3	1.8	2.4							
1987	3.6	1.9	2.7	1.5	2.1	2.3							
1988	3.2	2.2	1.9	2.1	1.7	1.2	2.1						
1989	3.3	1.6	1.8	1.8	1.3	1.7	1.4						
1990	3.4	1.7	1.7	2.3	2.2	1.8	1.6	2.2					
1991	3.4	1.8	1.4	2.0	2.5	1.4	1.6	2.2					
1992	2.8	1.7	1.8	1.7	1.5	1.4	2.0	1.7					
1993	3.4	1.9	1.6	1.7	1.8	2.3	1.1	1.5					
1994	4.1	2.3	2.2	1.7	1.1	1.6	1.4	1.2	1.9				
1995	4.7	3.4	2.6	1.4	1.2	1.4	1.7	1.1	1.7				
1996	4.9	3.3	2.4	2.2	1.0	1.7	1.6	1.4	1.6				
1997	5.1	4.0	3.5	1.5	1.5	1.0	1.8	0.8	1.3				
1998	5.5	3.8	3.1	2.4	1.7	1.5	1.0	1.0	1.1	0.9			
1999	5.8	5.0	2.5	3.2	1.8	1.1	1.2	0.9	1.6	1.5			
2000	6.2	4.9	3.9	4.3	2.2	1.4	0.9	0.9	0.9	1.6			
2001	5.7	5.2	4.8	3.4	2.7	2.1	1.3	1.1	1.8	1.4			
2002	6.7	5.8	3.8	4.4	2.9	2.3	1.7	—	1.4	1.0			
2003	6.0	5.2	4.8	3.9	2.5	3.1	1.5	—	1.3	0.7	1.0		
2004	6.5	6.0	4.4	5.0	3.3	2.8	1.9	—	1.2	1.0	1.0		
2005	7.2	5.1	5.0	3.8	4.0	2.8	4.4	—	1.4	1.3	1.4		
2006	6.6	4.3	4.8	4.7	3.3	2.3	3.7	—	1.0	1.2	1.5		
2007	6.2	4.4	4.1	4.6	3.8	4.0	3.2	—	3.8	2.4	2.5		
2008	5.8	5.4	4.9	5.0	4.0	4.1	2.9	—	3.5	3.7	3.5	3.2	
2009	5.2	3.5	5.6	2.9	4.1	2.6	2.9	—	3.1	2.9	2.7	3.0	
2010	4.8	3.0	3.8	3.5	2.6	3.5	2.5	—	3.0	1.9	3.2	3.0	
2011	4.3	2.9	2.8	3.5	4.1	2.7	2.2	—	4.1	2.2	2.0	2.5	
2012	4.5	2.6	2.8	3.1	1.9	2.9	2.2	—	3.1	2.8	2.3	2.3	
2013	4.8	3.1	3.6	3.6	4.2	2.7	2.6	—	2.6	2.7	2.7	1.8	2.2


Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aUsed without a doctor's orders.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-16
TRANQUILIZERS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-16 (cont.)

TRANQUILIZERS ^a

Trends in Annual Prevalence

among Respondents of Modal Ages 18 through 55, by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32 ^b</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976	10.3												
1977	10.8												
1978	9.9	9.4											
1979	9.6	9.8											
1980	8.7	8.8	9.0										
1981	8.0	7.4	7.3										
1982	7.0	5.6	7.2	8.6									
1983	6.9	5.1	5.8	6.6									
1984	6.1	5.4	5.4	5.6	6.7								
1985	6.1	4.4	4.5	6.2	7.1								
1986	5.8	4.2	5.4	5.2	5.4	6.8							
1987	5.5	4.0	5.5	4.1	5.8	6.2							
1988	4.8	3.5	4.5	4.2	4.3	4.8	4.6						
1989	3.8	3.4	3.5	3.8	2.9	4.6	4.1						
1990	3.5	3.0	3.6	3.8	5.0	3.3	3.9	3.8					
1991	3.6	2.7	3.2	4.0	3.9	3.8	4.2	4.1					
1992	2.8	2.2	3.8	3.4	4.5	3.4	3.7	4.1					
1993	3.5	2.1	3.1	3.2	3.7	3.8	2.7	2.7					
1994	3.7	1.9	2.9	3.1	3.3	3.6	3.2	3.8	3.1				
1995	4.4	3.7	3.5	3.0	3.1	3.4	3.5	1.8	3.6				
1996	4.6	3.5	3.7	3.0	2.4	2.9	3.1	3.2	3.6				
1997	4.7	4.7	3.6	2.9	1.9	2.0	4.1	4.1	3.0				
1998	5.5	4.2	4.5	3.7	3.6	2.9	2.4	3.8	2.2	3.0			
1999	5.8	4.1	4.2	4.2	3.5	2.6	2.1	2.7	3.4	2.0			
2000	5.7	5.5	5.3	5.3	3.7	3.0	2.7	2.6	3.2	3.0			
2001	6.9	6.1	7.1	5.4	5.3	3.9	4.2	3.0	4.3	3.7			
2002	7.7	8.8	7.8	6.4	7.0	4.9	5.0	—	4.0	4.2			
2003	6.7	8.0	7.0	7.2	6.3	5.2	4.3	—	3.8	2.2	2.9		
2004	7.3	8.0	8.1	8.3	6.8	5.6	4.8	—	3.6	3.5	3.1		
2005	6.8	6.5	8.5	6.3	7.7	4.3	7.4	—	4.4	3.8	2.9		
2006	6.6	6.1	7.6	6.8	5.6	6.2	6.6	—	4.0	3.5	4.0		
2007	6.2	5.7	7.6	7.5	7.4	7.6	5.0	—	3.5	4.5	3.9		
2008	6.2	7.1	6.3	8.1	6.7	5.7	6.5	—	5.6	3.1	3.8	4.2	
2009	6.3	4.3	7.7	7.1	7.4	5.7	7.2	—	5.0	4.1	2.5	2.9	
2010	5.6	5.2	6.6	7.2	7.2	5.2	6.4	—	4.4	3.7	4.3	3.8	
2011	5.6	5.3	5.2	6.6	7.2	5.2	5.1	—	6.2	3.0	3.8	2.7	
2012	5.3	4.8	4.3	5.9	5.1	6.6	4.1	—	3.1	3.3	4.6	4.3	
2013	4.6	4.8	4.8	6.8	4.6	6.1	5.5	—	4.7	3.8	3.3	4.1	2.6


Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aUsed without a doctor's orders.

^bBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-17
STERIODS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32,^b by Age Group



(Figure continued on next page.)

FIGURE 5-17 (cont.)
STERIODS^a
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 32,^b
by Age Group

<u>Year</u>	<u>Age 18</u>	<u>Ages 19-20</u>	<u>Ages 21-22</u>	<u>Ages 23-24</u>	<u>Ages 25-26</u>	<u>Ages 27-28</u>	<u>Ages 29-30</u>	<u>Ages 31-32^c</u>
1976								
1977								
1978								
1979								
1980								
1981								
1982								
1983								
1984								
1985								
1986								
1987								
1988								
1989	1.9							
1990	1.7							
1991	1.4	0.4	0.3	0.4	0.5	0.8	0.4	0.1
1992	1.1	0.5	0.1	0.7	0.4	*	0.2	*
1993	1.2	*	0.9	0.4	*	0.2	*	*
1994	1.3	0.5	0.6	*	0.2	0.5	*	0.4
1995	1.5	0.4	0.6	0.8	0.2	*	*	0.2
1996	1.4	0.3	0.1	0.3	1.0	*	*	0.5
1997	1.4	0.6	0.2	0.9	0.5	0.2	*	*
1998	1.7	0.8	0.2	0.2	0.3	0.4	*	0.5
1999	1.8	0.5	1.1	0.1	1.0	0.1	*	*
2000	1.7	0.1	0.4	0.3	0.2	0.9	*	0.2
2001	2.4	0.4	0.9	*	*	0.6	0.3	*
2002	2.5	1.0	0.3	0.4	0.3	*	1.1	—
2003	2.1	0.5	0.8	0.2	0.5	0.3	1.1	—
2004	2.5	0.8	0.2	0.5	0.5	0.3	*	—
2005	1.5	0.6	0.4	0.6	0.6	0.3	*	—
2006	1.8	0.4	1.0	*	*	0.1	*	—
2007	1.4	0.6	1.0	0.3	0.5	1.0	*	—
2008	1.5	0.9	0.1	0.3	0.3	0.3	0.3	—
2009	1.5	1.0	0.4	1.3	*	0.6	0.3	—
2010	1.5	0.4	1.9	0.3	0.6	0.7	*	—
2011	1.2	*	0.3	*	0.2	0.3	0.1	—
2012	1.3	0.3	0.5	0.9	0.1	0.1	0.8	—
2013	1.5	0.7	0.3	0.6	0.4	0.5	*	—

Source. The Monitoring the Future study, the University of Michigan.


Notes. ' * ' indicates a percentage of less than 0.05%. ' — ' indicates data not available

^aUsed without a doctor's orders.

^bQuestions about the use of steroids were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^cBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-18a
ALCOHOL
Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-18a (cont.)

ALCOHOL

**Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group**


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	85.7												
1977	87.0												
1978	87.7	89.8											
1979	88.1	90.6											
1980	87.9	89.0	90.2										
1981	87.0	90.6	91.6										
1982	86.8	88.6	91.8	90.0									
1983	87.3	88.5	91.8	91.7									
1984	86.0	88.7	89.1	90.4	88.2								
1985	85.6	88.5	89.8	91.6	89.9								
1986	84.5	88.2	90.1	88.1	88.8	87.8							
1987	85.7	88.2	90.8	89.7	90.5	87.8							
1988	85.3	86.6	89.5	89.7	89.4	87.7	87.2						
1989	82.7	87.5	89.1	88.7	87.5	88.0	86.0						
1990	80.6	85.6	89.6	88.2	87.5	86.4	86.9	84.8					
1991	77.7	84.6	89.0	88.1	87.7	85.3	85.0	83.8					
1992	76.8	81.9	87.9	89.1	86.7	85.6	84.5	85.0					
1993	76.0	80.6	85.9	87.8	87.8	85.7	83.2	83.6					
1994	73.0	78.2	84.4	86.6	86.0	84.5	82.6	83.6	82.5				
1995	73.7	78.3	85.7	87.8	86.7	85.7	83.3	81.8	82.1				
1996	72.5	79.6	84.4	85.7	85.9	85.3	84.7	82.0	83.5				
1997	74.8	79.2	85.1	85.4	86.4	85.9	83.7	83.3	82.3				
1998	74.3	79.7	86.3	84.9	83.8	85.3	84.2	83.2	82.3	77.3			
1999	73.8	79.6	85.5	85.2	85.0	85.4	85.4	85.1	81.0	80.0			
2000	73.2	79.7	86.2	87.2	84.2	82.9	83.7	82.9	81.0	80.3			
2001	73.3	77.6	87.0	86.7	86.3	84.2	84.3	84.4	82.7	81.5			
2002	71.5	78.0	85.8	88.0	88.3	84.7	83.6	—	85.1	80.0			
2003	70.1	75.0	84.3	87.6	86.4	83.6	83.9	—	82.6	81.6	78.9		
2004	70.6	75.2	86.8	87.2	87.9	86.1	83.5	—	86.7	79.8	79.2		
2005	68.6	77.3	84.4	86.6	85.6	85.3	84.8	—	85.8	81.6	80.3		
2006	66.5	77.9	83.6	88.2	86.4	86.9	84.0	—	83.7	80.5	82.8		
2007	66.4	72.9	87.8	87.8	86.1	85.8	85.9	—	84.0	85.2	80.7		
2008	65.5	72.3	88.6	86.6	86.4	84.7	87.8	—	84.3	82.0	80.3	79.0	
2009	66.2	71.4	85.2	89.3	88.2	87.2	84.8	—	83.5	86.6	81.3	79.7	
2010	65.2	68.8	83.4	89.2	86.7	86.6	86.7	—	85.0	86.1	81.1	80.3	
2011	63.5	71.5	82.1	88.3	90.6	86.4	85.1	—	89.0	84.4	80.6	82.1	
2012	63.5	70.3	81.8	85.0	89.4	86.7	84.2	—	87.2	83.0	84.4	80.2	
2013	62.0	68.4	82.8	84.7	87.9	89.6	86.6	—	86.7	83.5	81.5	79.7	76.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-18b
ALCOHOL
Trends in 30-Day Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-18b (cont.)

ALCOHOL

**Trends in 30-Day Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group**


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	68.3												
1977	71.2												
1978	72.1	75.8											
1979	71.8	76.5											
1980	72.0	76.6	78.3										
1981	70.7	77.0	80.5										
1982	69.7	75.7	79.9	77.9									
1983	69.4	73.9	79.3	78.9									
1984	67.2	73.6	78.1	77.6	75.2								
1985	65.9	73.3	75.9	79.7	76.8								
1986	65.3	72.9	77.2	75.7	76.3	73.6							
1987	66.4	72.5	77.2	74.9	77.7	75.0							
1988	63.9	69.6	76.2	75.9	74.1	74.6	72.1						
1989	60.0	69.8	73.8	72.2	72.5	73.9	72.3						
1990	57.1	66.6	74.1	73.6	71.4	70.9	70.2	68.4					
1991	54.0	64.5	75.3	72.4	71.6	69.8	69.6	68.5					
1992	51.3	61.0	72.7	73.0	69.8	69.1	69.2	67.8					
1993	51.0	60.5	71.6	73.1	69.9	68.3	66.2	66.4					
1994	50.1	59.9	70.4	70.1	70.4	69.9	67.0	67.7	65.1				
1995	51.3	59.2	70.4	72.3	71.8	68.0	67.0	67.6	66.8				
1996	50.8	58.1	69.5	69.2	68.5	69.3	68.0	65.5	64.7				
1997	52.7	59.0	69.1	69.3	70.9	70.4	65.8	65.3	65.3				
1998	52.0	59.7	69.4	70.3	66.3	68.7	66.1	65.2	62.9	59.8			
1999	51.0	62.0	69.2	70.2	70.0	70.2	67.4	66.0	64.2	64.2			
2000	50.0	59.1	70.5	71.5	68.7	64.6	65.2	66.7	64.0	63.1			
2001	49.8	59.0	71.8	70.6	68.7	66.5	66.2	67.8	63.7	65.6			
2002	48.6	59.2	71.9	71.9	71.2	67.9	65.4	—	67.3	65.4			
2003	47.5	56.7	69.5	72.7	69.1	67.2	66.5	—	63.7	66.2	62.2		
2004	48.0	56.7	72.4	72.8	72.4	68.8	64.5	—	70.3	63.7	65.7		
2005	47.0	59.0	70.1	71.2	73.0	70.3	65.7	—	68.5	65.1	65.4		
2006	45.3	57.6	69.7	73.8	70.4	72.8	68.7	—	63.3	62.3	66.7		
2007	44.4	54.7	74.5	73.1	73.8	71.9	69.8	—	67.5	66.9	64.1		
2008	43.1	53.8	74.4	74.0	73.9	69.8	73.4	—	65.0	66.3	67.9	63.7	
2009	43.5	52.9	72.9	78.4	75.1	70.7	71.5	—	65.6	71.0	66.5	64.5	
2010	41.2	51.2	71.6	74.6	73.6	72.2	69.1	—	67.7	72.7	67.0	67.1	
2011	40.0	52.3	69.3	75.7	76.4	71.8	69.6	—	71.1	68.6	65.2	68.3	
2012	41.5	54.1	70.5	73.5	76.7	73.4	70.8	—	69.5	68.0	72.0	65.7	
2013	39.2	51.5	70.5	72.7	75.9	73.9	71.1	—	70.0	69.0	67.2	66.7	61.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-18c
ALCOHOL
Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-18c (cont.)

ALCOHOL

**Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group**

<u>Year</u>	<u>Age 18</u>	<u>Ages 19–20</u>	<u>Ages 21–22</u>	<u>Ages 23–24</u>	<u>Ages 25–26</u>	<u>Ages 27–28</u>	<u>Ages 29–30</u>	<u>Ages 31–32^a</u>	<u>Age 35</u>	<u>Age 40</u>	<u>Age 45</u>	<u>Age 50</u>	<u>Age 55</u>
1976	5.6												
1977	6.1												
1978	5.7	7.6											
1979	6.9	7.7											
1980	6.0	7.0	8.4										
1981	6.0	7.2	7.7										
1982	5.7	7.5	7.8	8.2									
1983	5.5	5.3	8.0	8.5									
1984	4.8	5.8	7.7	6.8	7.5								
1985	5.0	5.6	6.4	7.3	7.5								
1986	4.8	5.3	6.3	6.2	5.3	7.3							
1987	4.8	5.7	7.0	6.1	6.9	7.2							
1988	4.2	4.8	7.2	6.2	6.3	5.7	7.6						
1989	4.2	4.7	5.0	5.1	6.0	6.9	5.6						
1990	3.7	4.0	4.9	5.3	4.8	4.9	5.6	6.4					
1991	3.6	3.7	4.9	5.4	4.9	6.2	5.9	5.2					
1992	3.4	3.3	4.4	4.2	6.1	4.4	5.8	6.0					
1993	2.5	3.2	5.1	4.9	5.1	4.7	4.6	5.7					
1994	2.9	3.3	3.9	3.7	3.3	5.4	5.0	4.7	7.2				
1995	3.5	3.1	3.5	4.1	4.4	4.7	4.1	4.9	5.5				
1996	3.7	2.7	5.1	4.8	3.7	3.6	5.1	5.1	7.5				
1997	3.9	4.8	4.6	4.5	5.1	4.2	5.9	5.0	4.8				
1998	3.9	3.6	5.7	3.9	3.4	3.1	3.4	4.8	6.0	6.9			
1999	3.4	4.1	5.9	4.7	5.1	4.3	5.2	4.4	5.2	7.5			
2000	2.9	3.9	5.3	4.2	3.8	3.5	3.9	4.2	5.2	6.5			
2001	3.6	3.6	6.2	4.6	5.0	2.7	4.3	4.7	5.8	7.5			
2002	3.5	3.9	5.6	5.0	5.4	3.7	3.8	—	4.8	6.6			
2003	3.2	3.6	5.7	6.5	4.6	5.1	3.5	—	3.9	7.8	7.8		
2004	2.8	3.7	5.7	5.5	4.3	3.5	3.8	—	6.3	6.6	9.0		
2005	3.1	3.6	6.0	5.8	6.0	4.6	5.8	—	6.1	7.2	8.5		
2006	3.0	4.3	5.9	5.7	5.2	5.8	4.5	—	5.3	7.0	9.5		
2007	3.1	3.4	6.1	6.0	6.5	6.2	5.1	—	8.1	6.1	8.8		
2008	2.8	2.3	5.5	6.3	6.4	6.3	4.8	—	5.4	7.2	9.9	11.0	
2009	2.5	2.5	5.7	6.5	5.9	6.5	7.1	—	6.9	8.5	9.3	9.2	
2010	2.7	1.8	5.4	4.9	5.6	5.7	6.1	—	6.0	7.8	7.2	10.5	
2011	2.1	2.4	6.1	5.2	5.7	7.0	5.3	—	7.6	7.4	7.7	11.3	
2012	2.5	3.0	4.9	6.4	5.9	7.5	7.2	—	6.3	8.3	9.5	10.6	
2013	2.2	2.7	4.9	4.9	6.9	6.5	5.8	—	6.7	8.7	8.4	10.8	10.5

Source. The Monitoring the Future study, the University of Michigan.


Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-18d

ALCOHOL

Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-18d (cont.)

ALCOHOL

Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	37.1												
1977	39.4												
1978	40.3	41.1											
1979	41.2	42.1											
1980	41.2	42.7	40.7										
1981	41.4	43.1	43.6										
1982	40.5	41.7	41.6	37.1									
1983	40.8	40.9	42.3	39.3									
1984	38.7	41.0	40.4	35.1	33.7								
1985	36.7	41.2	40.4	37.3	33.3								
1986	36.8	41.2	40.8	35.8	31.5	30.1							
1987	37.5	37.2	41.0	36.6	33.3	32.2							
1988	34.7	37.3	42.0	37.0	30.7	28.0	26.7						
1989	33.0	36.9	39.3	35.4	31.7	29.8	26.3						
1990	32.2	36.0	38.1	35.5	32.0	28.9	25.2	25.4					
1991	29.8	37.0	40.3	34.4	31.5	28.8	24.3	25.1					
1992	27.9	34.0	39.9	34.9	31.8	29.2	25.7	23.7					
1993	27.5	34.6	40.3	35.0	32.1	29.0	25.1	24.8					
1994	28.2	34.5	40.5	32.9	30.9	28.5	27.5	24.6	21.1				
1995	29.8	31.7	38.5	35.6	28.7	26.9	26.3	24.7	20.0				
1996	30.2	32.7	38.2	36.3	30.0	29.7	24.9	24.3	21.9				
1997	31.3	36.5	40.2	33.4	31.5	29.3	26.5	24.7	22.3				
1998	31.5	34.5	39.7	35.3	31.3	28.9	26.6	22.8	20.4	19.7			
1999	30.8	35.3	40.2	38.1	33.0	32.0	26.9	24.1	21.4	20.5			
2000	30.0	35.3	40.6	37.0	31.5	29.1	24.0	24.1	22.2	18.3			
2001	29.7	36.3	42.4	38.2	33.7	29.2	27.3	24.3	20.6	21.3			
2002	28.6	36.0	40.7	39.4	34.9	28.9	25.8	—	22.9	20.8			
2003	27.9	33.6	39.9	39.3	35.1	31.1	26.4	—	22.4	20.7	20.1		
2004	29.2	35.5	41.7	40.4	36.4	31.3	26.9	—	21.6	20.2	19.2		
2005	27.1	36.3	40.4	39.2	37.7	31.5	29.1	—	23.0	22.2	19.6		
2006	25.4	33.9	42.2	43.2	36.0	32.5	29.1	—	22.5	20.0	19.8		
2007	25.9	31.4	45.8	39.8	38.3	33.4	28.4	—	23.6	20.4	19.4		
2008	24.6	30.7	42.1	42.2	40.0	35.0	31.9	—	24.4	21.9	20.9	20.0	
2009	25.2	28.1	41.2	41.7	39.5	34.2	32.1	—	21.8	25.1	21.8	17.9	
2010	23.2	28.2	39.3	40.1	36.6	35.6	32.6	—	23.0	21.6	22.1	17.8	
2011	21.6	29.8	39.2	39.9	38.7	35.0	30.4	—	25.7	22.2	20.0	19.1	
2012	23.7	29.5	39.1	37.5	36.3	35.1	32.8	—	24.3	22.2	21.0	19.0	
2013	22.1	27.2	40.2	37.7	37.0	33.6	30.9	—	24.4	24.3	20.1	20.3	17.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-19a
CIGARETTES
Trends in 30-Day Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-19a (cont.)
CIGARETTES
Trends in 30-Day Prevalence
among Respondents of Modal Ages 18 through 55, by Age Group


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	38.8												
1977	38.4												
1978	36.7	39.3											
1979	34.4	39.3											
1980	30.5	36.0	37.9										
1981	29.4	34.9	37.5										
1982	30.0	32.1	36.2	36.7									
1983	30.3	32.5	33.5	36.5									
1984	29.3	31.5	32.2	33.6	33.7								
1985	30.1	30.9	32.4	31.9	35.3								
1986	29.6	30.0	32.0	29.9	31.3	32.5							
1987	29.4	30.1	32.4	31.7	28.2	32.3							
1988	28.7	28.4	29.8	29.9	27.3	29.1	28.9						
1989	28.6	27.7	29.4	29.4	29.5	27.2	30.2						
1990	29.4	27.2	28.6	27.8	28.4	26.5	27.8	28.3					
1991	28.3	27.6	28.3	28.5	28.3	28.2	24.4	28.1					
1992	27.8	29.5	29.0	28.4	26.3	27.8	23.8	27.5					
1993	29.9	29.0	29.2	28.1	27.7	25.4	25.8	25.3					
1994	31.2	31.3	28.8	27.0	26.4	25.0	25.5	24.9	24.8				
1995	33.5	33.4	31.8	28.0	25.7	26.8	25.2	25.0	26.1				
1996	34.0	34.0	32.3	30.1	26.8	26.0	23.4	23.8	25.4				
1997	36.5	34.0	32.3	29.1	27.6	24.9	24.6	24.3	22.3				
1998	35.1	33.9	33.7	30.9	29.9	25.6	23.1	22.5	23.6	24.3			
1999	34.6	36.1	33.4	32.4	25.6	22.9	22.7	24.0	22.6	23.5			
2000	31.4	32.2	33.6	29.5	28.2	26.5	21.2	22.7	24.0	23.5			
2001	29.5	32.8	34.0	31.1	28.6	24.2	20.4	21.2	20.4	22.9			
2002	26.7	29.8	32.6	31.9	27.3	24.7	24.4	—	21.9	18.9			
2003	24.4	27.0	30.5	31.0	27.0	26.3	22.0	—	20.1	21.9	20.7		
2004	25.0	27.9	31.3	31.5	29.6	25.9	21.9	—	20.0	20.0	20.2		
2005	23.2	27.5	29.2	29.3	30.7	26.3	23.5	—	19.1	21.4	22.1		
2006	21.6	24.6	27.3	28.1	29.1	26.3	24.4	—	17.7	17.3	18.9		
2007	21.6	22.6	27.8	26.7	27.5	26.6	22.9	—	17.8	18.3	17.6		
2008	20.4	21.8	24.5	26.5	24.5	25.7	24.0	—	20.4	17.8	17.3	18.8	
2009	20.1	21.2	25.2	24.1	22.6	23.9	24.0	—	17.3	16.2	17.8	17.0	
2010	19.2	19.6	22.8	23.0	24.3	22.5	23.9	—	18.3	15.2	18.3	19.9	
2011	18.7	18.5	23.3	22.0	23.4	19.6	20.5	—	19.7	15.1	15.3	16.0	
2012	17.1	16.8	18.9	20.4	20.7	22.0	18.6	—	18.0	12.8	15.9	15.4	
2013	16.3	18.4	20.8	21.4	19.5	20.0	16.5	—	17.8	16.6	13.4	15.3	14.0

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-19b
CIGARETTES
Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-19b (cont.)

CIGARETTES

**Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 55, by Age Group**


Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	28.8												
1977	28.8												
1978	27.5	31.0											
1979	25.4	31.2											
1980	21.3	29.3	31.1										
1981	20.3	26.0	31.4										
1982	21.1	23.9	28.6	30.1									
1983	21.2	24.4	26.0	30.6									
1984	18.7	24.1	25.3	27.8	28.7								
1985	19.5	23.2	25.3	25.1	30.4								
1986	18.7	21.9	24.4	25.2	27.3	27.6							
1987	18.7	22.5	24.2	26.0	23.7	27.9							
1988	18.1	19.5	22.3	24.0	22.9	25.0	25.4						
1989	18.9	18.9	22.5	23.3	25.0	22.9	26.4						
1990	19.1	19.2	20.2	22.2	23.3	22.2	24.2	23.9					
1991	18.5	19.4	20.6	22.5	22.8	23.9	21.0	24.9					
1992	17.2	20.5	21.2	20.9	20.3	21.8	20.3	22.8					
1993	19.0	21.1	20.5	20.1	21.9	20.1	21.7	21.4					
1994	19.4	21.9	21.1	19.9	19.8	20.5	20.9	20.9	22.5				
1995	21.6	22.2	24.0	20.0	19.2	20.9	20.1	21.2	23.0				
1996	22.2	22.5	22.8	22.8	21.1	19.4	18.6	19.8	22.1				
1997	24.6	22.7	21.4	21.5	19.2	17.6	19.7	19.1	18.3				
1998	22.4	23.8	22.8	21.2	21.9	19.5	17.2	17.9	20.4	21.7			
1999	23.1	25.6	24.2	21.4	19.6	16.0	17.2	18.9	19.7	20.9			
2000	20.6	22.7	25.1	21.2	20.1	19.7	15.8	18.1	20.1	20.8			
2001	19.0	21.9	23.6	22.4	20.9	17.2	14.4	16.1	16.4	20.1			
2002	16.9	20.6	23.9	23.5	19.8	18.1	17.4	—	18.2	16.7			
2003	15.8	18.8	20.8	21.5	20.4	19.8	16.4	—	16.3	19.0	19.0		
2004	15.6	18.2	21.5	23.3	22.7	18.2	16.7	—	14.8	16.6	17.8		
2005	13.6	17.6	19.2	20.4	22.5	18.6	18.9	—	14.5	18.5	20.1		
2006	12.2	14.4	17.7	19.5	22.0	20.2	18.3	—	13.5	14.6	16.7		
2007	12.3	12.9	18.3	17.5	19.2	19.3	16.8	—	13.9	15.8	15.4		
2008	11.4	14.3	16.1	17.9	17.4	18.3	17.4	—	16.5	14.7	14.6	16.8	
2009	11.2	12.8	14.9	16.2	15.3	16.5	16.7	—	13.7	12.7	15.6	15.4	
2010	10.7	11.1	15.5	15.3	16.2	16.2	17.3	—	14.3	12.3	16.4	18.0	
2011	10.3	10.2	15.0	13.7	17.0	13.4	14.8	—	15.7	11.8	13.6	14.2	
2012	9.3	9.5	11.5	13.1	14.1	16.0	14.3	—	13.4	10.5	13.8	13.5	
2013	8.5	10.8	12.0	13.1	10.9	13.8	11.3	—	12.4	13.5	11.0	13.2	13.2

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

FIGURE 5-19c
CIGARETTES
Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day
among Respondents of Modal Ages 18 through 55, by Age Group



(Figure continued on next page.)

FIGURE 5-19c (cont.)

CIGARETTES

Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among Respondents of Modal Ages 18 through 55, by Age Group

Year	Age 18	Ages 19–20	Ages 21–22	Ages 23–24	Ages 25–26	Ages 27–28	Ages 29–30	Ages 31–32 ^a	Age 35	Age 40	Age 45	Age 50	Age 55
1976	19.2												
1977	19.4												
1978	18.8	23.8											
1979	16.5	24.6											
1980	14.3	21.9	25.2										
1981	13.5	19.3	25.3										
1982	14.2	18.0	23.0	24.6									
1983	13.8	17.2	19.7	25.1									
1984	12.3	17.2	21.2	22.8	24.1								
1985	12.5	16.6	20.4	20.8	24.8								
1986	11.4	16.2	19.3	21.1	22.0	23.2							
1987	11.4	15.6	19.3	21.6	19.9	23.3							
1988	10.6	13.8	17.3	18.4	18.6	20.6	22.3						
1989	11.2	13.0	16.4	18.6	20.6	19.0	22.0						
1990	11.3	14.3	15.0	17.4	19.6	18.2	20.5	20.8					
1991	10.7	12.7	14.1	17.4	18.2	19.0	16.7	20.8					
1992	10.0	14.5	15.1	15.5	15.8	17.9	17.0	19.3					
1993	10.9	14.5	14.5	15.2	17.4	16.3	17.9	17.8					
1994	11.2	15.0	15.6	15.0	15.0	15.9	16.8	17.8	19.1				
1995	12.4	15.2	18.1	15.3	14.2	16.3	16.5	17.6	19.1				
1996	13.0	14.7	15.7	16.1	15.0	14.8	15.2	16.1	18.5				
1997	14.3	15.4	14.7	16.4	13.2	12.8	15.9	16.1	15.4				
1998	12.6	16.9	16.2	14.5	15.5	14.8	12.2	14.3	16.3	18.7			
1999	13.2	16.3	16.4	14.8	15.0	12.4	13.2	14.8	17.3	17.2			
2000	11.3	14.6	17.2	14.1	14.8	14.7	12.5	12.8	15.7	17.2			
2001	10.3	13.9	15.9	15.8	15.1	12.6	11.4	11.9	13.4	15.9			
2002	9.1	12.8	14.4	15.9	14.1	13.9	14.0	—	13.0	13.6			
2003	8.4	11.7	13.8	15.4	14.0	14.8	12.7	—	12.4	14.9	16.8		
2004	8.0	11.6	12.7	15.2	15.6	12.8	12.5	—	10.9	14.2	15.4		
2005	6.9	10.1	12.1	13.9	13.6	13.1	14.1	—	11.3	16.0	16.4		
2006	5.9	8.8	10.9	12.8	14.0	13.6	13.5	—	10.7	12.2	14.2		
2007	5.7	7.5	10.7	10.6	14.3	13.0	12.6	—	10.5	12.1	12.3		
2008	5.4	7.3	9.8	11.5	10.9	12.0	12.3	—	12.4	11.6	12.2	13.9	
2009	5.0	7.4	9.1	8.6	10.3	11.8	10.5	—	11.1	8.5	13.0	12.2	
2010	4.7	6.7	9.3	9.6	11.3	10.1	10.7	—	10.8	9.1	13.3	14.4	
2011	4.3	4.5	7.9	8.2	9.7	7.6	9.2	—	10.8	8.7	11.0	11.1	
2012	4.0	4.6	7.3	8.2	7.7	10.4	7.8	—	10.8	7.7	10.6	11.2	
2013	3.4	5.4	6.5	8.1	6.4	8.5	7.1	—	8.9	10.2	8.7	10.3	11.1

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available.

^aBeginning in 2002, respondents were followed through age 30 instead of age 32 as in past years.

Chapter 6

ATTITUDES AND BELIEFS ABOUT DRUGS AMONG YOUNG ADULTS

One of the most important theoretical contributions of MTF has been to demonstrate the extent to which attitudes and beliefs about drugs can help explain the use of drugs. Earlier volumes in this monograph series, as well as other publications from the study, have demonstrated that shifts in certain attitudes and beliefs—in particular the degree of risk of harm perceived to be associated with use of a particular drug—are important in explaining changes in actual drug-using behavior. Indeed, on a number of occasions we have accurately predicted such changes in use by using perceived risk as a *leading indicator of use*.⁶⁸ In this chapter, we review trends in these attitudes and beliefs held by young adults since 1980.

PERCEIVED HARMFULNESS OF DRUGS

Figures 6-1 through 6-31 present three separate trend lines for four-year age strata (that is, respondents who are one to four years beyond high school [ages 19–22], five to eight years beyond high school [ages 23–26], and nine to twelve years beyond high school [ages 27–30]). For comparison purposes, data are also included for the high school senior classes, listed as 18-year-olds. Figures 6-1 to 6-3 present trends in the percentages of young adults aged 18 to 30 who perceive a “great risk” of harm associated with three different levels of marijuana use—trying it once or twice (experimental), smoking it occasionally, and smoking it regularly. Subsequent figures do the same for selected levels of use of various other licit and illicit drugs. Table 6-1 provides the tabular information underlying the figures.

For most of the life of the study, these questions were contained in one questionnaire form only, limiting the numbers of follow-up cases. Accordingly, we have used four-

⁶⁸ See also: Bachman, J. G., Johnston, L. D., O'Malley, P. M., & Humphrey, R. H. (1988). Explaining the recent decline in marijuana use: Differentiating the effects of perceived risks, disapproval, and general lifestyle factors. *Journal of Health and Social Behavior*, 29, 92–112; Bachman, J. G., Johnston, L. D., & O'Malley, P. M. (1990). Explaining the recent decline in cocaine use among young adults: Further evidence that perceived risks and disapproval lead to reduced drug use. *Journal of Health and Social Behavior*, 31, 173–184; Bachman, J. G., Johnston, L. D., & O'Malley, P. M. (1998). Explaining recent increases in students' marijuana use: Impacts of perceived risks and disapproval, 1976 through 1996. *American Journal of Public Health*, 88, 887–892; Johnston, L. D. (1981). Characteristics of the daily marijuana user. In R. de Silva, R. L. DuPont, & G. K. Russell (Eds.), *Treating the marijuana-dependent person* (pp. 12–15). New York: The American Council on Marijuana; Johnston, L. D. (1985). The etiology and prevention of substance use: What can we learn from recent historical changes? In C. L. Jones & R. J. Battjes (Eds.), *Etiology of drug abuse: Implications for prevention* (NIDA Research Monograph No. 56, DHHS Publication No. ADM 85 1335, pp. 155–177). Rockville, MD: National Institute on Drug Abuse. Keyes, K.M., Schulenberg, J.E., O'Malley, P.M., Johnston, L.D., Bachman, J.G., Li, G., & Hasin, D. (2011). The social norms of birth cohorts and adolescent marijuana use in the United States, 1976-2007. *Addiction*, 106(10), 1790-1800. doi: 10.1111/j.1360-0443.2011.03485.x

year age bands to increase the available sample size to about 300–600 weighted cases per year for each age band, thereby improving the reliability of the estimates. (The numbers of weighted cases are given at the end of Table 6-1. The actual numbers of respondents are somewhat larger.) Still, these are relatively small sample sizes compared to those available for 8th, 10th, and 12th graders, and thus the change estimates are relatively less stable.

Beginning with 2012 we expanded the numbers of forms from which these data are drawn; this increased the sample sizes from that point forward, thus improving the reliability of both the point estimates and the trend estimates. Because the questions are contained in different numbers of forms for the different drugs, the sample sizes vary between drugs, as is noted in footnotes. We use all forms on which each question exists.

Because of the nature of the MTF design, trend data are available for a longer period for 19- to 22-year-olds (since 1980) than for 23- to 26-year-olds (since 1984) or 27- to 30-year-olds (since 1988). Also displayed in this table are comparison data for 12th graders, shown here as 18-year-olds, from 1980 onward. (See also Table 8-3 in *Volume I* for the longer term trends in 12th graders' levels of perceived risk.) Questions about these attitudes and beliefs are not included in the questionnaires for respondents over age 30 due to the length limitations imposed by using a single questionnaire form for respondents age 35 and older.

- Table 6-1 and Figures 6-1 to 6-31 illustrate considerable differences in the degree of risk young adults associate with various drugs. In general, the results closely parallel the distinctions made by 12th graders.
- ***Marijuana*** is seen as the least risky of the illicit drugs, although sharp distinctions are made between different levels of marijuana use (Figures 6-1 through 6-3). In 2013, experimental use of marijuana was perceived as being of great risk by only 10–12% of all high school graduates ages 19–30, whereas regular use was perceived to carry great risk by a considerably higher percentage (36–40%). In 2007 all four age strata showed declines of three to four percentage points in perceived risk for regular marijuana use; although no one of these declines was statistically significant, taken alone, the consistency across all four groups suggests that the shift was real. Since then the declines have continued, though somewhat erratically; but all four age groups showed a decline between 2007 and 2011 of 7 to 10 percentage points in perceived risk for regular marijuana use, suggesting a possible period effect. That period effect may well have been brought about by the increasing discussion and adoption of medical marijuana use by many states. *In fact, perceived risk associated with regular marijuana use in 2011 was at the lowest level observed since each of the young adult age groups was included in the study—going back to 1980 in the case of 19- to 22-year-olds.* In 2012 the decline halted for all of the groups above age 18, but in 2013 new lows were reached in each of the four age strata. The one year declines in 2013 were over 4 percentage points in three of those strata, making them statistically

significant, while the oldest group declined 3.2 percentage points (not significant). Legalization of recreational use was under discussion in 2013 in addition to the legalization of medical marijuana use.

- In the mid-1980s and early 1990s, fewer of the older age groups attached great risk to marijuana use than did the younger respondents. Indeed, there was a regular negative ordinal relationship between age and perceived risk for some years after 1980, when the first comparisons were available. Although at first this may have looked like an age effect, the MTF design allowed us to recognize it as a cohort effect; the younger cohorts initially perceived marijuana as more dangerous than the older cohorts and persisted in such beliefs as they grew older. Newer cohorts, however, have shown lower levels of perceived risk. Twelfth graders from the class of 2013 are much less likely to perceive regular marijuana use as dangerous, compared with 12th-grade cohorts in the late 1980s and early 1990s. This reflects what we have interpreted as generational forgetting, a phenomenon wherein younger replacement cohorts no longer carry the beliefs—or had the direct or vicarious experience on which those beliefs were based—that the older cohorts had at that age. (This concept appears at other points throughout this chapter, and the implications for prevention are discussed in the last section of this chapter.)

The decline in perceived risk that began in the 1990s was greater in the younger age bands, including grades 8 and 10, and least among the 27- to 30-year-olds. We believe that much of the eventual decline in perceived risk in the older age bands resulted directly from replacement of earlier cohorts by later, less concerned ones. The credibility of this view is strengthened by the 1993–1995 reversal of the relationship between age and perceived risk of regular use. This reversal is consistent with an underlying cohort effect and could not simply reflect an association between age and a regular change in these attitudes. The decline in perceived risk for regular marijuana use ended in a somewhat staggered fashion—among 12th graders in 1999, among 19- to 22-year-olds in 2001, among 23- to 26-year-olds in 2002, and among 27- to 30-year-olds in 2004. This was also indicative of a cohort effect playing out in these attitudes. In 2007 all four age strata showed declines of three to four percentage points in perceived risk for regular marijuana use; although no one of these declines was statistically significant, taken alone, the consistency across all four groups suggests that the shift was real. Since then the declines have continued, though somewhat erratically; but all four age groups showed a decline between 2007 and 2011 of 7 to 10 percentage points in perceived risk for regular marijuana use, suggesting a possible period effect. Young adults ages 19–30 view experimental use of any of the *other illicit drugs* as distinctly more risky than the experimental use of marijuana. About 30–32% of young adults think trying *sedatives (barbiturates)* involves great risk; the corresponding figures are 31–34% for *amphetamines*, 40–45% for *LSD*, 44–47% for *narcotics other than heroin*, 49–52% for *ecstasy (MDMA)*, 47–51% for *cocaine powder*, and 69–73% for *heroin*. Note that two classes of prescription drugs have the lowest levels of perceived risk among this set.

(Perceived risk of tranquilizers is not asked, but likely would rank low as well.) Items about perceived risk of *synthetic marijuana* use were added to the questionnaires in 2012 (Table 6-1). These drugs are sold over the counter in small packets containing plant material that has been sprayed with any of a number of chemicals with chemical structures similar to cannabinoids. The percent seeing great risk in trying synthetic marijuana in the three young adult age bands were 29–31% in 2012 and 29–33% in 2013, reflecting some increase in perceived risk in all four age groups in 2013, though no one stratum's increase was statistically significant.

- The older age groups have been more likely to see *LSD* as dangerous (Figure 6-4). These age distinctions became sharper through about 2001 as perceived risk declined more in the younger age groups—again indicating some important cohort changes in these attitudes, quite likely as a result of generational forgetting of the dangers of LSD. In recent years, there has been a decline in perceived risk in the older age groups, again likely resulting from generational replacement as newer cohorts moved up the age spectrum, which has diminished the age differences. However, there has not been a clear pattern of change in the past several years.
- Questions about perceived risk of *ecstasy (MDMA)* were introduced in the follow-up surveys in 1989, but were not asked of 12th graders until 1997 (due to concerns about introducing the secondary school students to a drug with such an alluring name). At the beginning of the 1990s, all young adult age groups viewed ecstasy as a fairly dangerous drug, even for experimentation. But, again, the different age bands showed diverging trends during the 1990s, with the oldest two age bands continuing to see ecstasy as quite dangerous, but the 19- to 22-year-olds (and very likely the 12th graders, for whom we did not have data until 1997) coming to see it as less so. In 2000, 38% of 12th graders saw great risk in trying ecstasy versus 49% of 27- to 30-year-olds; in 2001, the corresponding figures were 46% and 54%. In fact, three of the four age groups showed appreciable increases in perceived risk for ecstasy in 2001, which led us to predict a decline in use. The increase in perceived risk continued in 2002 in the two youngest age strata, and their use of ecstasy did, indeed, begin to decline—and decline sharply (see chapter 5). Perceived risk continued to rise after 2004 for all age groups, and then began to decline after 2006, with perceived risk of experimenting with ecstasy among 18-year-olds declining fairly steadily from 58% in 2001 to 48% in 2013, quite possibly as a result of generational forgetting (Figure 6-16). In 2012 further declines in the proportions seeing great risk associated with trying ecstasy were observed among all of the young adult age bands (4 percentage points for 19- to 22-year-olds, a statistically significant 11 percentage points for 23- to 26-year-olds, and 7 percentage points for the 27- to 30-year-olds); however, no further change was observed among the 18-year-olds that year, and in 2013 there was some bounce back in the young adult strata partially offsetting the previous year's decline.

- Perceived risk for *salvia* (Table 6-1) also was included for the first time in 2012 in the young adult questionnaires, and the percent seeing great risk in trying salvia ranged from 19% to 23% among the young adults in 2012 and from 20% to 21% in 2013 (Table 6-1).
- Recent years have shown little systematic change in perceived risk for *cocaine* among young adults and not a great deal of difference in this belief among the age groups. A decline in perceived risk began among 19- to 22-year-olds after 1994, among 23- to 26-year-olds after 1997, and among 27- to 30-year-olds after 2001. Young adults have generally reported somewhat higher perceived risk with respect to regular cocaine use than have 12th graders; the differences are smaller for occasional and experimental use.

Among 12th graders and the young adult age groups, the danger associated with using cocaine on a regular basis grew considerably (by 13 and 17 percentage points, respectively) between 1980 and 1986 (Figure 6-9). Interestingly, these changed beliefs did not translate into changed behavior until the perceived risk associated with experimental and occasional use began to rise sharply after 1986. When these two measures rose, a sharp decline in actual use occurred. We hypothesized that respondents saw only these lower levels of use as relevant to them and, therefore, saw themselves as vulnerable only to the dangers of such use. (No one starts out planning to be a heavy user; further, in the early 1980s, cocaine was not believed to be addictive.) Based on this hypothesis, we included the additional question about occasional use in 1986, just in time to capture a sharp increase in perceived risk later that year. This increase occurred largely in response to the growing media frenzy about cocaine—and crack cocaine, in particular—and to the widely publicized, cocaine-related deaths of several public figures (most notably Len Bias, a collegiate basketball star and a leading National Basketball Association draft pick). After stabilizing for a few years, perceived risk began to fall off after about 1991 among 12th graders, but not among the older age groups—again suggesting that lasting cohort differences were emerging, quite possibly as a result of generational forgetting of the dangers of cocaine in the younger age groups. Now, more than 25 years later, none of the young adult age groups has had much exposure to the cocaine epidemic of the mid-1980s, which likely explains why there no longer is much age-related difference in the level of perceived risk, except with regards to regular use, for which 12th graders have a lower level of perceived risk than any of the young adult strata (Figure 6-9).

- Perceived harmfulness of *crack* use has been lowest among 12th graders for some years now (Figures 6-10 through 6-12). High school seniors have been considerably less likely than any of the older age groups to view occasional and regular use of crack cocaine as dangerous, strongly suggesting that an age-effect may have been operating here. Trend data (available since 1987) on the risks perceived to be associated with crack use showed increases in 1987–1990 for all age groups, followed by relatively little change in the older age

strata. During the 1990s, twelfth graders showed decreases in the perceived risk of experimental use of crack—perhaps reflecting the onset of generational forgetting of its dangers—leaving them as perceiving considerably less risk than the older groups. The young adult age groups have shown a staggered decline in this measure, with 19- to 22-year-olds showing a decline after 1994, 23- to 26-year-olds since 1996 and 27- to 30-year-olds after 2001. As a result, the different ages spread out more in their levels of perceived risk of crack use, until declines in the older age groups after about 2002. Given this lack of recent historical or age variation, questions about perceived risk of crack use were dropped from the young adult questionnaires in 2012 to make room for such questions about other drugs.

- Perceived risk for trying *amphetamines* (Figure 6-21) increased in all four age strata very gradually from 1980, when first measured, through 2010, with little difference among them. In 2011 it dropped in all strata and then held level thereafter. There was more difference among the age groups with regard to the risk attached to regular amphetamine use with the older two strata generally seeing greater risk than the younger two strata, and especially the 12th graders. The younger two strata showed an increase in perceived risk during the 1980s and then some fallback in the early 1990s, before stabilizing. The sharp recent decline observed for experimental use was also seen for regular use from 2009 to 2011 among 12th graders and from 2011 through 2012 among all of the young adult strata.
- Perceived risk questions for *Adderall* were added to the young adult questionnaires for the first time in 2012. They showed that perceived risk of using once or twice ranged from 29% to 30% in the three young adult age bands in 2012. In 2013 perceived risk generally rose but only the 5 percentage point increase among 12th graders reached statistical significance (Table 6-1).
- Measures of perceived risk of *crystal methamphetamine (ice)* use were introduced in 1990, and the results show what may be an important reason for its lack of rapid spread. More than half of all 12th graders and young adults perceived it as quite dangerous even to try, perhaps because it was likened to crack in many media accounts. (Both drugs come in crystal form, both are burned and the fumes inhaled, both are stimulants, and both can produce a strong dependence.) There was rather little age-related difference in perceived risk associated with use of crystal methamphetamine in 1990 and 1991, although the two youngest age groups were somewhat higher. But as perceived risk fell considerably among 12th graders (and eventually among 19- to 22-year-olds) and held steady or rose in the oldest two age groups, an age-related difference emerged. Since about 2004, perceived risk has risen some among all of the age strata, narrowing the age-related differences that had emerged for a few years. In 2011 perceived risk for trying this drug stood at 67% among 12th graders and at 73–75% in all of the older strata. Given this lack of variation in recent years and low levels of actual use, these questions

were discontinued in the young adult surveys in 2012 to make room for such questions about other drugs.

- In 2012 perceived risk questions about the use of *bath salts*—over-the-counter synthetic stimulants—were added to the questionnaires. That year fairly high proportions of the young adults saw great risk of harm in even trying bath salts (4649%), but considerably fewer of the 18-year-olds did (33%). Perceived risk increased dramatically in 2013 for bath salts in all four age strata, with increases ranging from 16 to 26 percentage points in a single year. Now even trying bath salts once or twice is seen as dangerous by between 60% and 66% in these age strata—very high levels.
- Young adults have been more cautious than 12th graders about *heroin* use. (See Figures 6-18 through 6-20.) In general, there has been relatively little change over the years in the proportions of all age groups seeing regular heroin use as dangerous, with the great majority of each group (over 85%) consistently holding this viewpoint. However, there has been a long-term gradual slight rise in perceived risk among the young adult strata, but not among the 12th graders. With regard to heroin experimentation, from 1975 to 1986 there had been a downward shift among 12th graders in the proportion seeing great risk associated with trying heroin and some decline among 19- to 22-year-olds. Following this decline (although their data do not extend back as far), young adults showed a gradually increasing caution about heroin use in the latter half of the 1980s—possibly due to the association of heroin injection with the spread of HIV—followed by a leveling through most of the 1990s. In 1996 and 1997, young adults’ perceived risk increased some, as happened among 12th graders (as well as among 8th and 10th graders). These various trends may reflect, respectively, (a) the lesser attention paid to heroin by the media during the late 1970s and early 1980s as cocaine took center stage; (b) the subsequent great increase in attention paid to intravenous heroin use in the latter half of the 1980s due to the recognition of its important role in the spread of HIV/AIDS; (c) the emergence in the 1990s of heroin so pure that people no longer needed to use a needle to administer it; and (d) the subsequent increased attention given to heroin by the media (partly as a result of some overdose deaths by public figures and partly prompted by the emergence of “heroin chic” in the design industry), as well as through an anti-heroin media campaign launched by the Partnership for a Drug-Free America in June 1996. At present, more young adults still see heroin use as dangerous than do 12th graders (Table 6-1).
- Perceived risk questions about *narcotics other than heroin* were first asked of the young adults in 2012; between 43% and 47% of the three age groups saw great risk of harm in experimenting with such drugs. In 2013 there was a significant increase among 12th graders, but no significant changes among the young adult groups (Table 6-1).

- In 2013, only a minority of young adults saw *occasions of heavy drinking* on weekends as dangerous (37–41%), as did a larger proportion of 12th graders (46%). The belief that heavy drinking carries great risk increased over the 1980s in these age groups, rising among 12th graders from 36% in 1980 to 49% in 1992. Among 19- to 22-year-olds, it rose from a low of 30% in 1981 to 42% in 1992; the increases among the older groups were smaller. The increase in this belief may well help to explain the important decline in actual heavy drinking, and may in turn be explained by the media campaigns against drunk driving and the increase in the drinking age in a number of states.⁶⁹ Following a staggered pattern, perceived risk of harmfulness peaked among 18-year-olds in 1992, among 19- to 22-year-olds in 1993, among 23- to 26-year-olds in 1994, and among 27- to 30-year-olds in 1995, suggesting some cohort effect. Since 1995, perceived risk of heavy drinking has not changed much among the 19–30 age groups but has risen slightly among the 18-year-olds.
- Between 1980 and 1991, a gradually increasing proportion of all four age groups viewed drinking *one or two drinks per day* as dangerous (Figure 6-27); but then they all showed a parallel decrease in perceived risk for this behavior through at least 2000. It seems likely that the earlier increase was due to the general rising concern about the consequences of alcohol use, particularly drunk driving, and that the subsequent decline in perceived risk was due to increasing reports of cardiovascular health benefits of light-to-moderate daily alcohol consumption. In recent years there has been little systematic change in this belief in any of the age strata, and there has been little difference by age across the entire 33-plus-year interval, although for about the past ten years the 18-year-olds have been a little more likely to see risk in daily drinking.
- More than four fifths (83–84%) of young adults now perceive regular *pack-a-day cigarette smoking* as entailing high risk. In recent years, 18-year-olds have consistently shown lower perceived risk than young adults have, while 10th graders have been still lower and 8th graders lowest. Clearly, there is an age effect in young people coming to understand the dangers of smoking. Unfortunately, it appears that much of the learning about the risks of smoking happens *after* a great deal of smoking initiation has occurred and many young people have already become addicted. These beliefs about smoking risks have strengthened very gradually in all age groups from senior year forward during the years we have monitored them (see Table 6-1 and Figure 6-30). The

⁶⁹ See O'Malley, P. M., & Johnston, L. D. (1999). Drinking and driving among U.S. high school seniors: 1984–1997. *American Journal of Public Health*, 89, 678–684; O'Malley, P. M., & Johnston, L. D. (2003). Unsafe driving by high school seniors: National trends from 1976 to 2001 in tickets and accidents after use of alcohol, marijuana and other illegal drugs. *Journal of Studies on Alcohol*, 64, 305–312; and O'Malley, P. M., & Johnston, L. D. (2013). Driving after drug use or alcohol use by American high school seniors, 2001–2011. *American Journal of Public Health*, 103(11), 2027–2034. [doi:10.2105/AJPH.2013.301246](https://doi.org/10.2105/AJPH.2013.301246)

parallel changes in these beliefs across the different age groups indicate a period effect, suggesting that all of the age groups responded to common influences in the larger culture. These influences are discussed at length in *Volume I* in chapter 8 on attitudes and beliefs. The rise in perceived risk has stalled some in recent years, with only very slight increases between 2002 and 2013 based on fitting linear trend lines to the data.

- The regular use of *smokeless tobacco* is seen as dangerous by 53–59% of young adults and 42% of 12th graders. These beliefs gradually strengthened until about 2001 in all age groups covered (Table 6-1), particularly among the two older age groups. As with cigarettes, the change appears to reflect a secular trend (period effect) because of its parallel occurrence in all age groups. Perceived risk has not changed among the young adults in any systematic fashion since 2001; these data are based on only one form, so year-to-year fluctuations can appear relatively large. There is a considerable difference across age groups in 2013, with 42% of 18-year-olds seeing great risk in regular smokeless tobacco use compared to 53% to 59% of the three young adult strata.

PERSONAL DISAPPROVAL OF DRUG USE

For most of the life of the study, in one of six questionnaire forms follow-up respondents were asked the same questions asked of 12th graders concerning the extent to which they personally disapprove of various drug-using behaviors among “people (who are 18 or older).” Trends in the answers of young adults in the age bands of 19 to 22, 23 to 26, and 27 to 30 are contained in Table 6-2. Comparison data for 12th graders are also provided for 1980 onward. (See Table 8-6 in *Volume I* for the longer term trends in 12th graders’ levels of disapproval associated with using the various drugs.) As with the perceived risk questions, starting in 2012 the estimates were based on all questionnaire forms on which each disapproval question was located in order to increase sample size and, therefore, reduce sampling error. Each question is footnoted in Table 6-2 to indicate on how many forms it was contained in 2012 and 2013.

- In general, disapproval levels of adult use of the various drugs rank similarly across substances for both 12th graders and young adults. The great majority of young adults disapprove of using, or even experimenting with, all of the *illicit drugs other than marijuana*. For example, 95% or more of young adults in 2013 disapprove of regular use of each of the following drugs: *LSD*, *cocaine*, *heroin*, and *sedatives (barbiturates)*. Fully 80% to 96% of young adults disapprove of experimenting with each of these same drugs. Many of these attitudes differ rather little as a function of age at present; when there is a difference, the younger age groups are usually the least disapproving.

- Even for *marijuana*, about four tenths of young adults now disapprove of experimentation (38–44%), a slight majority (51–55%) disapprove of occasional use, and the great majority (74–76%) disapprove of regular use.

Among drugs measured, marijuana use shows the widest fluctuations in disapproval over time—generally, fluctuations that parallel the changes in perceived risk (though sometimes with a one-year lag, with the change in perceived risk coming first). The most fluctuation has occurred among the younger age groups (Table 6-2). Among 12th graders, disapproval of regular marijuana use increased substantially in the 1980s, peaked in the early 1990s, declined through much of the 1990s, and then leveled around 1998 with little change for some years thereafter. The 19- to 22-year-olds had a quite similar pattern, though the decline continued a year longer, likely due to generational replacement. Among 23- to 26-year-olds, there were some declines starting later in the 1990s, but the declines have been very modest. Since 2007 there has been some decline in disapproval of occasional and regular marijuana use in all four age groups; the pattern is consistent with a secular trend, which would alert us to a possible increase in marijuana use. Indeed, such an increase in use is now occurring among both 12th graders and 19- to 28-year-olds (see Table 2-2).

- In all four age groups from high school seniors to age 30, the great majority has disapproved of even experimenting with *LSD* since 1980, when these data were first available. Beginning around 1990, all age groups decreased some in their disapproval of trying LSD (starting from high levels of disapproval at 90–91%). The decline was steepest among 12th graders, but there was a reversal in this group's disapproval in 1997, and then an increase through 2006. Disapproval in the older age groups declined less and in staggered fashion; this trend has shown some evidence of a reversal among 19- to 22-year-olds and 23- to 26-year-olds since 2001 and 2002, respectively. The pattern again suggests lasting cohort differences in these attitudes. Disapproval levels have fluctuated in recent years, but are generally comparable in 2013 to what they were in 2002. Disapproval of regular LSD use has been near the top of the scale for more than three decades, ranging from 92% to 99%.
- First measured among young adults in 2001, *ecstasy* use was positively associated with age in the early 2000s. In 2001 disapproval of even trying ecstasy was quite high, and from 2001 to 2010, disapproval rose to even higher levels in all age groups, with little systematic change since then. Experimenting with ecstasy is disapproved of by 85% of 12th graders in 2013 and by roughly similar proportions in the upper ages.
- Disapproval of all three levels of *heroin* use (experimental, occasional, and regular use) has remained very high and stable since MTF began. There was one minor exception, however: A little slippage in disapproval of

experimental use occurred among 12th graders from 1991 through 1996 (from 96% to 92%)—a period during which heroin usage rates were rising.

- Disapproval of regular *cocaine* use rose gradually among 19- to 22-year-olds, from 89% in 1981 to 99% in 1990, with little change thereafter (97% in 2013). All three young adult age bands are now near 100%. Disapproval of experimental cocaine use increased during the 1980s, peaking first among 12th graders at 94% in 1991. It then peaked in 1995 among 19- to 22-year-olds (at 94%) and 23- to 26-year-olds (at 92%). Finally, it peaked in 1999 at 90% among 27- to 30-year-olds. All age groups have had some modest falloff in disapproval since those peak levels were attained. Again, the lag in inflection points between the successive age groups suggests some lasting cohort differences in these attitudes. For the last few years, all age groups' disapproval of experimental cocaine use has ranged from 83–94%, with the older strata having somewhat lower disapproval than the younger ones.
- Disapproval of experimenting with *amphetamines* rose in the 1980s as use was falling; thereafter, disapproval leveled at above 80%, with almost no difference among the age strata. For example, trying amphetamines once or twice was disapproved of by 73–74% of 19- to 26-year-olds in 1984, compared to 84% by 1990. After a long period of level disapproval, all strata showed a slight drop in disapproval in 2011, followed by another leveling. Disapproval of regular use started out very high among all age strata in the early 1980s and rose even higher by the early 1990s, where it remained for all age strata until 2011; after that there was a slight decline, but a leveling by 2012. Since 2010 there has been some decline in disapproval of amphetamine use in all four age groups, likely explained by a change in the question wording. Adderall and Ritalin were included in the question for the first time in 2011 as examples of amphetamines. There has been very little difference among the various age strata in either their levels or trends in disapproval.
- Disapproval of experimental use of *sedatives (barbiturates)* has moved very much in parallel with that for amphetamines. Disapproval increased significantly during the 1980s, accompanied by declining use. Disapproval of trying sedatives was at 84–85% in 1984 compared to 89–91% by 1990. Disapproval of sedative use slipped some among 12th graders after 1992 and among 19- to 22-year-olds after 1994, with the 23- to 26-year-olds following suit after 1996, and the 27- to 30-year-old stratum in 2004. This pattern of staggered change again suggests cohort effects, reflecting lasting cohort differences in these attitudes. In recent years a gradual, staggered increase in disapproval of trying sedatives has shown up in all age groups.
- The story for *alcohol* is quite an interesting one, in that changes in the minimum drinking age seem to have led to modest changes in norms for the affected cohorts. Between 1980 and 1992, an increasing proportion of 12th graders favored total abstention; the percent who disapproved of drinking even just once or twice rose from 16% in 1980 to 33% in 1992. (This figure

has fallen back slightly and stands at 27% in 2013.) Among 19- to 22-year-olds there was a modest increase in disapproving of any use between 1985 and 1989 (from 15% to 22%), where it held for some years; it remained at 22% in 2011. For the two oldest age groups, there has been rather little change in these attitudes so far. These differing trends may reflect the fact that during the 1980s, the drinking age was raised in a number of states so that by 1987 it was 21 in all states; this change would have had the greatest effect on 12th graders, who may have incorporated the legal restrictions into their normative structure and, as they entered young adulthood, brought these new norms with them. But the changes may be exhibited only among respondents in the cohorts that were underage after the time that the new law raising the minimum drinking age went into effect.

Disapproval of *daily drinking* (one or two drinks) has not shown any such cohort effects, because all age groups have generally moved in parallel, at similar levels of disapproval. The three youngest age bands (which include 12th graders through 26-year-olds) showed an increase in disapproval of daily drinking up until about 1990 (there was little data yet available on the oldest age group), but disapproval has declined a fair amount in all of the age groups since then. A bit of a gap between 12th graders and young adults opened up between 2004 and 2008, when 12th graders increased their disapproval of daily drinking while young adults did not. This pattern of cross-time change closely parallels what was observed for the perceived risk associated with light daily drinking, discussed previously; the later decline in both variables among the young adults may well be due to widely publicized reports that some cardiovascular benefits may result from having one or two drinks per day.

- There was a considerable increase in disapproval of *occasions of heavy drinking* (also called *binge drinking*) on weekends from the early 1980s for the two youngest age groups, and this continued through 1992 for 12th graders (who then showed some drop-off) and through 1996 among 19- to 22-year-olds (who then also showed some drop-off). As Figure 5-18d illustrates, the prevalence of occasions of heavy drinking declined substantially among 12th graders and 19- to 22-year-olds between 1981 and the early 1990s, as norms became more restrictive. There was little or no change in the older age strata, either in their levels of disapproval or in their rates of occasions of heavy drinking, until the early 2000s, when disapproval began to drop some in both strata.
- Some fluctuations in the disapproval of *cigarette smoking* have occurred over the decades covered by MTF. Twelfth graders showed some increase in disapproval of pack-or-more-a-day smoking between 1982 (69%) and 1992 (74%). Their disapproval then fell through 1997 (to 67%) as their smoking increased; disapproval then increased for several years (to 82% in 2006) before leveling, as smoking declined. In 2013, 83% disapproved of pack-or-more-a-day smoking. The 19- to 22-year-olds showed a similar increase in

disapproval from 66% in 1982 to 81% in 2013. All four age strata showed some upward drift in their level of disapproval of smoking since about 1999 (78–81% in 2013), suggesting a secular change in attitudes during this period.

COHORT DIFFERENCES AND THEIR IMPLICATIONS FOR PREVENTION AND THEORY

An important theoretical point to be made—based on the strong evidence reported here for cohort effects in perceived risk and disapproval of many of the drugs under study—is that among the causes of cohort differences in actual use are lasting cohort differences in these critical *attitudes* and *beliefs*. In other words, *the attitudes and beliefs brought into adulthood from adolescence tend to persevere*.

A second point has to do with the causes of these attitudinal cohort effects. We noted earlier that the older respondents are more likely than the younger ones to see as dangerous the use of marijuana, LSD, heroin, amphetamines, ecstasy, crystal methamphetamine, cocaine, crack, and sedatives (barbiturates). We have offered the framework for a theory of drug epidemics in which direct learning (from personal use) and vicarious learning (from observing use by others in both the immediate and mass media environments) play important roles in changing these key attitudes.⁷⁰ To the extent that the data on perceived risk represent cohort effects (enduring differences between class cohorts), these findings would be consistent with this theoretical perspective. Clearly, use of these particular drugs was greater when the older cohorts were growing up, and public attention and concern regarding the consequences of these drugs were greatest in the 1970s and early to mid-1980s. In the early 1970s, LSD was alleged to cause brain and chromosomal damage, as well as bad trips, flashbacks, and behavior that could prove dangerous. Methamphetamine use was discouraged with the slogan “speed kills.” In addition, there was a serious epidemic of heroin use in the early 1970s. Later cohorts (through the mid-1990s, at least) were not exposed to those experiences. While there may have been a secular trend toward greater perceived risk for drugs in general, in the case of LSD there may have also been an operating cohort effect (with younger cohorts seeing less danger) offsetting the secular trend among 12th graders; the net effect was a decrease in 12th graders’ perceived risk of LSD use after 1980.

This vicarious learning process has a very practical application for national strategy for preventing future epidemics. Because fewer in their immediate social circles and fewer public role models may be using these drugs and exhibiting the adverse consequences of use during certain historical periods, future cohorts of youth may have less opportunity to learn about the adverse consequences of these drugs in the

⁷⁰ Johnston, L. D. (1991). Toward a theory of drug epidemics. In L. Donohew, H. E. Sypher, & W. J. Bukoski (Eds.), *Persuasive communication and drug abuse prevention* (pp. 93–131). Hillsdale, NJ: Lawrence Erlbaum. Available at <http://monitoringthefuture.org/pubs/chapters/ldj1991theory.pdf>.

Monitoring the Future

normal course of growing up. Unless those hazards are convincingly communicated to them in *other ways*—for example, through school prevention programs, by their parents, and through the mass media, including public service advertising—they will become more susceptible to a new epidemic of use of the same or similar drugs.

In *Volume I*, we reported an increase in use of several drugs in 8th, 10th, and 12th grades in 1994 through 1997. This increase suggests that this form of generational forgetting may well have been taking place during those years. For the cohorts that follow such a rise in use, there is once again an increased opportunity for vicarious learning from the adverse experiences of those around them, but by that time, members of affected cohorts have had to learn the hard way what consequences await those who become involved with the various drugs. In the 2000s we have seen drug use subside to some degree, which once again has created the conditions for generational forgetting of the dangers of many of these drugs, as we have been saying in earlier volumes in this monograph series. We are now seeing some softening of attitudes among teens regarding marijuana and ecstasy, and the recent lower levels of perceived risk of LSD use—particularly among 8th graders—suggest a real possibility of future increases in use.

TABLE 6-1
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Percentage saying "great risk"^a

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	(Years Cont.)
Try marijuana once or twice^f	18	10.0	13.0	11.5	12.7	14.7	14.8	15.1	18.4	19.0	23.6	23.1	27.1	24.5	21.9	19.5	16.3	
	19–22	8.3	7.8	9.7	9.7	12.8	11.2	13.0	12.9	16.8	16.9	17.8	19.1	19.7	19.4	18.8	13.3	
	23–26	—	—	—	—	9.6	10.0	12.4	14.5	16.0	14.0	17.7	14.0	15.0	13.0	15.0	15.8	
	27–30	—	—	—	—	—	—	—	—	14.6	16.0	17.0	15.7	15.1	14.0	14.8	16.1	
Smoke marijuana occasionally^f	18	14.7	19.1	18.3	20.6	22.6	24.5	25.0	30.4	31.7	36.5	36.9	40.6	39.6	35.6	30.1	25.6	
	19–22	13.9	14.2	16.9	16.7	21.7	20.6	22.4	23.0	28.7	29.1	30.1	30.2	29.5	30.3	31.3	25.5	
	23–26	—	—	—	—	15.8	16.3	20.9	20.8	26.8	25.3	30.4	26.2	27.4	24.0	25.5	27.7	
	27–30	—	—	—	—	—	—	—	—	24.2	25.7	28.7	27.4	27.5	26.8	28.1	28.3	
Smoke marijuana regularly^f	18	50.4	57.6	60.4	62.8	66.9	70.4	71.3	73.5	77.0	77.5	77.8	78.6	76.5	72.5	65.0	60.8	
	19–22	43.9	47.8	52.4	58.4	62.2	66.8	67.6	69.4	72.4	74.9	73.0	75.0	69.3	69.2	65.0	62.1	
	23–26	—	—	—	—	52.9	57.5	59.4	65.3	68.3	72.1	71.0	70.9	67.3	64.1	63.2	64.2	
	27–30	—	—	—	—	—	—	—	—	67.5	69.1	69.2	67.5	68.8	69.4	65.6	69.2	
Try synthetic marijuana once or twice^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take synthetic marijuana occasionally^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try LSD once or twice^h	18	43.9	45.5	44.9	44.7	45.4	43.5	42.0	44.9	45.7	46.0	44.7	46.6	42.3	39.5	38.8	36.4	
	19–22	44.8	44.4	45.0	44.7	46.0	44.3	47.6	49.4	49.2	49.5	49.3	48.0	45.6	42.4	42.3	40.3	
	23–26	—	—	—	—	48.3	46.9	47.9	51.5	53.7	50.7	52.0	50.1	49.7	49.0	46.8	45.8	
	27–30	—	—	—	—	—	—	—	—	53.3	55.6	54.6	52.5	53.0	51.5	53.5	52.5	
Take LSD regularly^h	18	83.0	83.5	83.5	83.2	83.8	82.9	82.6	83.8	84.2	84.3	84.5	84.3	81.8	79.4	79.1	78.1	
	19–22	83.4	85.3	86.2	86.0	84.5	86.4	87.1	85.6	85.4	85.5	85.8	86.6	87.0	81.3	81.0	80.5	
	23–26	—	—	—	—	89.0	86.6	88.7	90.0	89.2	89.0	88.2	89.1	87.3	85.3	87.5	86.3	
	27–30	—	—	—	—	—	—	—	—	89.1	91.2	92.0	87.1	88.5	89.0	89.2	88.4	
Try PCP once or twice^h	18	—	—	—	—	—	—	—	55.6	58.8	56.6	55.2	51.7	54.8	50.8	51.5	49.1	
	19–22	—	—	—	—	—	—	—	63.6	63.8	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	64.8	63.2	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	65.9	—	—	—	—	—	—	—	
Try ecstasy (MDMA) once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take ecstasy (MDMA) occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try salvia once or twice^{d,g}	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take salvia occasionally^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try cocaine once or twice^h	18	31.3	32.1	32.8	33.0	35.7	34.0	33.5	47.9	51.2	54.9	59.4	59.4	56.8	57.6	57.2	53.7	
	19–22	31.4	30.4	33.3	28.7	33.1	33.2	35.5	45.9	51.9	51.5	58.1	58.7	56.1	60.5	63.8	57.7	
	23–26	—	—	—	—	31.3	31.1	35.9	48.0	47.1	51.3	51.5	50.5	53.5	54.1	56.0	58.7	
	27–30	—	—	—	—	—	—	—	—	45.3	53.0	51.6	52.6	51.8	54.7	53.5	56.4	
Take cocaine occasionally^h	18	—	—	—	—	—	—	54.2	66.8	69.2	71.8	73.9	75.5	75.1	73.3	73.7	70.8	
	19–22	—	—	—	—	—	—	53.8	61.3	67.1	72.6	74.6	72.6	74.9	75.4	78.0	73.4	
	23–26	—	—	—	—	—	—	50.9	62.6	63.2	69.9	69.9	70.3	69.9	72.8	70.3	76.0	
	27–30	—	—	—	—	—	—	—	—	62.6	66.6	66.6	69.1	69.9	69.1	69.9	70.0	
Take cocaine regularly^h	18	69.2	71.2	73.0	74.3	78.8	79.0	82.2	88.5	89.2	90.2	91.1	90.4	90.2	90.1	89.3	87.9	
	19–22	65.2	69.3	71.5	75.2	75.1	82.9	82.0	88.0	90.3	89.1	93.9	93.5	92.9	91.7	92.2	91.5	
	23–26	—	—	—	—	75.6	76.9	83.0	88.9	90.9	91.2	91.2	92.7	89.9	91.9	92.6	93.3	
	27–30	—	—	—	—	—	—	—	—	88.9	92.0	91.4	90.9	92.0	91.6	92.1	91.3	

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Percentage saying "great risk"^a

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change		
Try marijuana once or twice^f	18	15.6	14.9	16.7	15.7	13.7	15.3	16.1	16.1	15.9	16.1	17.8	18.6	17.4	18.5	17.1	15.6	14.8	14.5	-0.3		
	19–22	16.9	14.8	13.4	12.5	14.3	11.9	13.3	17.1	15.3	15.6	14.4	10.8	17.4	13.2	16.8	13.4	12.9	11.8	-1.0		
	23–26	18.5	15.1	16.7	16.4	13.1	13.0	15.1	15.3	13.6	13.0	13.9	13.0	12.5	10.6	12.7	10.5	10.1	9.7	-0.4		
	27–30	16.2	16.1	16.4	16.1	14.4	17.3	16.2	18.0	13.8	14.5	14.5	16.6	11.4	12.3	11.5	12.4	12.5	10.2	-2.2 s		
Smoke marijuana occasionally^f	18	25.9	24.7	24.4	23.9	23.4	23.5	23.2	26.6	25.4	25.8	25.9	27.1	25.8	27.4	24.5	22.7	20.6	19.5	-1.1		
	19–22	25.6	22.0	22.0	19.8	25.8	18.0	21.0	24.1	23.2	24.3	22.1	22.3	23.6	23.1	19.9	19.6	20.6	19.1	-1.5		
	23–26	27.3	26.4	26.8	26.4	24.9	20.5	24.5	22.2	22.7	21.6	22.3	20.2	18.5	18.1	19.3	15.5	17.1	14.4	-2.6 s		
	27–30	28.1	26.0	25.8	25.3	25.8	25.0	30.2	27.9	25.1	24.8	21.8	25.6	21.6	21.7	18.6	19.3	19.7	16.0	-3.6 ss		
Smoke marijuana regularly^f	18	59.9	58.1	58.5	57.4	58.3	57.4	53.0	54.9	54.6	58.0	57.9	54.8	51.7	52.4	46.8	45.7	44.1	39.5	-4.6 s		
	19–22	61.3	60.7	53.4	55.2	58.0	49.6	56.7	57.8	57.2	55.3	54.5	50.4	51.6	46.4	49.8	43.0	43.5	39.4	-4.1 s		
	23–26	62.7	64.1	62.7	60.1	60.3	55.1	53.7	56.7	54.2	53.6	55.9	52.5	52.4	43.0	47.1	39.3	40.1	35.9	-4.1 s		
	27–30	67.3	65.0	63.6	66.1	64.0	61.7	63.5	64.7	59.3	57.0	54.9	51.5	51.2	47.4	48.5	42.2	43.5	40.3	-3.2		
Try synthetic marijuana once or twice^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	23.5	25.9	+2.4		
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	30.6	33.1	+2.5	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.1	29.4	+0.3	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.9	32.7	+3.8	
Take synthetic marijuana occasionally^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	32.7	36.2	+3.5	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	38.5	40.1	+1.6
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	37.3	38.6	+1.3
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.3	41.0	+4.7
Try LSD once or twice^h	18	36.2	34.7	37.4	34.9	34.3	33.2	36.7	36.2	36.2	36.5	36.1	37.0	33.9	37.1	35.6	34.7	33.1	34.9	+1.8		
	19–22	44.4	40.1	38.7	38.1	37.9	37.5	35.3	39.7	39.2	38.7	43.5	40.9	46.5	38.5	40.9	43.5	43.5	40.3	-3.2		
	23–26	46.1	46.6	45.7	49.3	44.9	48.5	45.7	43.8	40.7	39.9	38.1	42.8	43.8	43.0	48.7	44.1	47.2	43.0	-4.2		
	27–30	50.1	52.0	52.0	49.9	46.4	46.7	44.9	47.5	47.2	47.9	44.9	44.6	42.4	41.7	41.5	45.2	45.8	45.2	-0.7		
Take LSD regularly^h	18	77.8	76.6	76.5	76.1	75.9	74.1	73.9	72.3	70.2	69.9	69.3	67.3	63.6	67.8	65.3	65.5	66.8	66.8	0.0		
	19–22	82.4	83.6	78.6	82.2	81.6	79.2	81.1	78.6	78.4	77.8	78.9	77.5	73.9	74.8	72.8	74.4	78.0	76.6	-1.4		
	23–26	84.7	85.6	82.1	85.4	84.1	86.0	85.3	84.3	83.5	80.8	82.0	80.3	80.2	82.0	83.1	81.4	78.9	79.0	+0.1		
	27–30	87.0	87.2	90.5	87.8	85.3	86.9	85.3	87.5	83.9	87.9	82.2	85.7	82.9	80.2	87.0	83.0	83.2	83.8	+0.5		
Try PCP once or twice^h	18	51.0	48.8	46.8	44.8	45.0	46.2	48.3	45.2	47.1	46.6	47.0	48.0	47.4	49.7	52.4	53.9	51.6	53.9	+2.4		
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Try ecstasy (MDMA) once or twice^h	18	—	33.8	34.5	35.0	37.9	45.7	52.2	56.3	57.7	60.1	59.3	58.1	57.0	53.3	50.6	49.0	49.4	47.5	-1.9		
	19–22	46.7	45.5	42.7	37.6	37.9	40.5	46.8	50.1	52.3	53.8	51.0	50.3	51.4	51.4	50.7	49.9	45.9	52.4	+6.4		
	23–26	50.4	50.5	47.7	50.0	46.7	45.7	45.6	45.9	44.9	51.2	46.4	51.4	46.3	46.4	47.5	54.2	43.7	49.1	+5.4		
	27–30	50.6	48.8	50.4	50.9	48.9	53.6	52.0	58.8	49.1	50.2	46.5	51.9	43.5	43.5	52.0	51.3	44.3	51.4	+7.1		
Take ecstasy (MDMA) occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	19–22	—	—	—	—	—	72.5	77.8	81.7	78.3	80.0	82.5	79.3	81.9	79.2	76.2	71.6	76.7	75.3	-1.4		
	23–26	—	—	—	—	—	72.5	71.9	73.6	77.4	77.2	77.0	78.7	78.6	76.2	79.1	76.9	76.6	69.8	-6.8		
	27–30	—	—	—	—	—	75.2	76.5	79.9	76.9	74.7	70.4	72.0	71.3	71.4	69.7	77.8	75.0	76.8	+1.8		
Try salvia once or twice^{dg}	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	39.8	38.7	13.8	12.9	-0.9		
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	22.5	21.4	-1.1		
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	18.6	19.6	+1.0		
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	18.8	20.6	+1.8		
Take salvia occasionally^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	23.1	21.3	-1.8		
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.2	30.6	+1.4		
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	26.6	25.5	-1.1		
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	24.7	25.7	+1.0		
Try cocaine once or twice^h	18	54.2	53.6	54.6	52.1	51.1	50.7	51.2	51.0	50.7	50.5	52.5	51.3	50.3	53.1	52.8	54.0	51.6	54.4	+2.8		
	19–22	61.9	55.5	55.4	52.8	56.7	48.9	55.5	55.0	55.5	55.6	54.0	55.8	56.7	54.9	56.8	56.2	57.0	56.3	-0.7		
	23–26	57.2	63.1	60.2	62.6	63.1	62.4	61.0	55.4	52.1	53.0	52.5	56.9	55.0	56.6	56.7	54.9	60.3	50.9	-9.4 s		
	27–30	53.6	54.6	60.5	61.7	59.9	60.9	58.8	56.4	61.4	56.5	58.1	54.8	56.1	52.0	51.6	54.7	51.8	53.8	+2.0		
Take cocaine occasionally^h	18	72.1	72.4	70.1	70.1	69.5	69.9	68.3	69.1	67.2	66.7	69.8	68.8	67.1	71.4	67.8	69.7	69.0	70.2	+1.2		
	19–22	76.6	76.1	71.2	68.0	72.4	70.0	69.9	70.3	70.2	72.1	71.0	71.5	72.4	67.2	72.9	70.3	78.0	76.5	-1.5		
	23–26	71.3	76.5	74.2	77.8	76.2	74.2	75.4	68.3	74.1	70.4	68.5	70.9	67.2	74.9	71.6	71.6	76.9	75.8	-1.1		
	27–30	67.8	73.8	73.2	75.4	76.5	78.1	74.3	72.6	75.3	76.2	74.6	72.1	73.9	65.4	71.5	71.0	73.2	77.9	+4.7		
Take cocaine regularly^h	18	88.3	87.1	86.3	85.8	86.2	84.1	84.5	83.0	82.2	82.8	84.6	83.3	80.7	84.4	81.7	83.8	82.6	83.3	+0.6		
	19–22	92.2	91.6	88.7	88.5	90.7	85.1	88.3	87.4	87.1	89.2	86.2	86.7	87.0	88.6	87.9	86.3	92.3	91.4	-0.9		
	23–26	90.6	93.2	92.9	92.7	92.9	91.1	91.5	88.5	91.5	88.0	90.9	88.0	86.5	89.2	90.9	88.0	91.2	91.2	0.0		
	27–30	91.6	92.7	93.0	92.4	92.3	94.5	91.2	92.9	91.3	94.0	90.0	89.9	91.1	88.8	92.7	87.2	91.2	91.7	+0.5		

(List of drugs continued.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage saying "great risk" ^a																	
Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	(Years Cont.)
		18	—	—	—	—	—	—	—	—	57.0	62.1	62.9	64.3	60.6	62.4	57.6	58.4	
19–22	—	—	—	—	—	—	—	—	59.4	67.3	68.5	69.4	66.9	65.4	63.5	70.1	61.9		
23–26	—	—	—	—	—	—	—	—	59.1	63.5	69.8	67.3	66.9	67.1	64.2	69.3	64.8		
27–30	—	—	—	—	—	—	—	—	—	66.5	64.9	68.7	66.8	64.3	68.8	65.6	66.4		
Try crack once or twice^h	18	—	—	—	—	—	—	—	70.4	73.2	75.3	80.4	76.5	76.3	73.9	73.8	72.8		
19–22	—	—	—	—	—	—	—	—	75.0	77.3	81.8	82.3	82.7	81.9	83.6	84.3	78.8		
23–26	—	—	—	—	—	—	—	—	70.3	74.0	79.9	81.1	83.9	84.4	81.6	83.2	81.4		
27–30	—	—	—	—	—	—	—	—	—	76.4	76.7	82.6	81.8	79.1	83.6	78.6	81.1		
Take crack occasionally^h	18	—	—	—	—	—	—	—	84.6	84.8	85.6	91.6	90.1	89.3	87.5	89.6	88.6		
19–22	—	—	—	—	—	—	—	—	89.6	91.1	94.1	94.9	95.6	93.4	96.2	96.0	94.2		
23–26	—	—	—	—	—	—	—	—	88.0	89.2	91.5	94.2	95.4	94.1	93.4	94.9	95.5		
27–30	—	—	—	—	—	—	—	—	—	89.6	89.5	95.3	94.4	93.3	93.5	93.0	94.0		
Take crack regularly^h	18	—	—	—	—	—	—	—	45.3	51.7	53.8	53.9	53.6	57.1	53.2	55.4	52.0		
19–22	—	—	—	—	—	—	—	—	44.0	48.6	51.1	54.5	52.7	56.2	49.7	62.0	55.8		
23–26	—	—	—	—	—	—	—	—	41.0	43.6	48.4	48.9	47.4	45.9	45.6	52.5	48.9		
27–30	—	—	—	—	—	—	—	—	—	42.0	45.1	46.2	43.3	42.3	49.9	47.1	48.2		
Try cocaine powder once or twiceⁱ	18	—	—	—	—	—	—	—	56.8	61.9	65.8	71.1	69.8	70.8	68.6	70.6	69.1		
19–22	—	—	—	—	—	—	—	—	58.0	59.0	63.2	70.0	69.9	72.6	70.6	75.4	73.0		
23–26	—	—	—	—	—	—	—	—	50.0	53.2	62.2	63.3	67.0	65.8	64.0	68.8	68.8		
27–30	—	—	—	—	—	—	—	—	—	53.6	52.7	60.9	59.2	61.2	64.3	61.0	65.9		
Take cocaine powder occasionallyⁱ	18	—	—	—	—	—	—	—	81.4	82.9	83.9	90.2	88.9	88.4	87.0	88.6	87.8		
19–22	—	—	—	—	—	—	—	—	86.6	87.6	91.3	92.5	93.8	92.1	94.0	94.9	93.5		
23–26	—	—	—	—	—	—	—	—	82.9	84.1	88.5	92.4	93.8	91.3	92.4	92.8	92.1		
27–30	—	—	—	—	—	—	—	—	—	85.1	86.7	92.7	91.1	91.5	92.5	90.7	92.7		
Take cocaine powder regularlyⁱ	18	52.1	52.9	51.1	50.8	49.8	47.3	45.8	53.6	54.0	53.8	55.4	55.2	50.9	50.7	52.8	50.9		
19–22	57.8	56.8	54.4	52.5	58.7	51.0	55.5	57.9	58.9	59.6	58.3	59.9	59.8	58.9	60.8	58.9			
23–26	—	—	—	—	58.2	59.2	60.8	66.6	65.4	62.3	64.1	62.4	63.7	65.0	63.3	64.1			
27–30	—	—	—	—	—	—	—	—	66.0	69.7	67.5	66.1	66.5	69.3	69.6	66.4			
Try heroin once or twice^g	18	70.9	72.2	69.8	71.8	70.7	69.8	68.2	74.6	73.8	75.5	76.6	74.9	74.2	72.0	72.1	71.0		
19–22	77.5	77.8	73.6	74.5	74.9	73.6	77.2	77.6	77.5	79.8	80.8	80.2	81.6	78.8	79.0	77.9			
23–26	—	—	—	—	81.2	80.7	78.9	84.5	82.4	80.8	83.4	84.4	81.5	82.1	80.8	85.3			
27–30	—	—	—	—	—	—	—	—	86.0	86.8	85.3	84.3	84.9	86.2	86.8	83.1			
Take heroin occasionally^g	18	86.2	87.5	86.0	86.1	87.2	86.0	87.1	88.7	88.8	89.5	90.2	89.6	89.2	88.3	88.0	87.2		
19–22	87.2	89.9	87.5	88.6	86.8	90.2	90.7	90.2	89.6	90.8	91.2	91.5	92.2	89.2	91.2	89.9			
23–26	—	—	—	—	92.0	90.1	90.6	92.8	91.5	91.3	91.0	92.6	91.3	91.6	93.0	93.5			
27–30	—	—	—	—	—	—	—	—	92.7	93.5	93.0	90.7	91.3	92.6	93.8	92.4			
Take heroin regularly^g	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Try narcotics other than heroin once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Take narcotics other than heroin regularly^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Try amphetamines once or twice^{b,h}	18	29.7	26.4	25.3	24.7	25.4	25.2	25.1	29.1	29.6	32.8	32.2	36.3	32.6	31.3	31.4	28.8		
19–22	24.6	24.6	27.8	24.8	26.9	23.9	27.1	27.4	31.7	28.9	35.6	32.8	34.5	33.3	36.3	32.9			
23–26	—	—	—	—	29.6	29.4	29.4	34.1	33.2	32.5	35.3	31.0	32.7	32.6	32.9	34.3			
27–30	—	—	—	—	—	—	—	—	35.2	37.5	36.9	36.5	36.2	34.0	37.5	36.0			
Take amphetamines regularly^{b,h}	18	69.1	66.1	64.7	64.8	67.1	67.2	67.3	69.4	69.8	71.2	71.2	74.1	72.4	69.9	67.0	65.9		
19–22	71.9	69.9	68.3	69.9	68.4	68.5	72.3	72.0	73.9	71.3	74.0	77.1	73.5	73.5	71.6	72.2			
23–26	—	—	—	—	75.8	77.2	75.6	78.2	77.4	76.7	77.8	79.4	76.4	76.2	73.6	80.5			
27–30	—	—	—	—	—	—	—	—	80.6	82.9	83.3	79.4	80.3	79.8	78.4	77.7			
Try crystal methamphetamine (ice)^h	18	—	—	—	—	—	—	—	—	—	—	—	61.6	61.9	57.5	58.3	54.4		
19–22	—	—	—	—	—	—	—	—	—	—	—	—	57.8	58.6	57.7	57.5	61.4		
23–26	—	—	—	—	—	—	—	—	—	—	—	—	56.5	56.0	55.6	52.0	61.0		
27–30	—	—	—	—	—	—	—	—	—	—	—	—	59.6	57.2	52.7	60.3	57.9		

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage saying "great risk" ^a																		2012– 2013 change	
Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .		Age Group	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013
Try crack once or twice^h	18	56.0	54.0	52.2	48.2	48.4	49.4	50.8	47.3	47.8	48.4	47.8	47.3	47.5	48.4	50.2	51.7	52.0	55.6	+3.5	
	19–22	65.2	62.0	59.3	56.1	52.9	54.1	54.1	55.1	56.8	56.6	55.3	51.9	54.9	54.9	53.7	56.6	—	—	—	
	23–26	68.6	64.7	67.3	64.6	63.2	59.8	60.9	58.5	56.4	60.6	54.7	58.4	50.5	50.6	58.4	61.6	—	—	—	
	27–30	66.7	68.5	66.5	65.0	62.9	69.3	67.4	66.0	62.6	61.9	56.8	64.1	56.2	56.2	62.2	60.4	—	—	—	
Take crack occasionally^h	18	71.4	70.3	68.7	67.3	65.8	65.4	65.6	64.0	64.5	63.8	64.8	63.6	65.2	64.7	64.3	66.2	66.5	69.5	+3.0	
	19–22	83.5	79.1	79.1	75.5	74.9	72.3	75.3	75.3	76.0	75.0	72.8	77.7	75.7	75.7	73.6	74.8	—	—	—	
	23–26	85.9	80.8	84.2	81.6	84.0	80.1	82.2	77.1	76.4	78.6	76.8	79.8	75.2	75.2	77.7	82.8	—	—	—	
	27–30	81.3	85.3	81.7	79.8	81.6	84.4	81.5	81.9	82.1	79.5	82.8	79.1	77.3	77.3	80.1	79.6	—	—	—	
Take crack regularly^h	18	88.0	86.2	85.3	85.4	85.3	85.8	84.1	83.2	83.5	83.3	82.8	82.6	83.4	84.0	83.8	83.9	84.0	85.4	+1.4	
	19–22	94.7	93.3	92.8	92.3	91.1	89.6	91.1	93.8	93.3	92.5	90.3	90.3	93.6	93.6	93.1	90.8	—	—	—	
	23–26	96.1	91.4	95.6	94.4	95.6	93.4	94.7	92.2	92.5	93.1	93.3	93.1	91.8	91.8	93.7	94.1	—	—	—	
	27–30	94.3	96.0	94.3	95.2	93.5	96.8	94.2	94.4	94.0	95.2	94.1	93.6	94.1	93.1	93.9	92.6	—	—	—	
Try cocaine powder once or twiceⁱ	18	53.2	51.4	48.5	46.1	47.0	49.0	49.5	46.2	45.4	46.2	45.8	45.1	45.1	46.5	48.2	48.0	48.1	49.9	+1.8	
	19–22	57.1	53.8	53.0	47.9	48.0	47.1	47.9	49.4	48.7	50.2	48.7	46.8	48.3	48.3	44.4	51.3	52.2	51.3	+2.1	
	23–26	57.2	53.6	54.1	53.8	53.2	53.9	52.5	50.8	46.0	53.3	45.8	48.1	44.1	44.2	43.9	47.4	52.5	47.4	-4.3	
	27–30	48.9	49.1	49.8	49.7	52.2	53.3	54.4	56.6	52.5	52.9	49.0	53.6	47.2	47.2	52.1	48.3	53.5	48.3	-1.6	
Take cocaine powder occasionallyⁱ	18	68.8	67.7	65.4	64.2	64.7	63.2	64.4	61.4	61.6	60.8	61.9	59.9	61.6	62.6	62.6	64.2	62.6	65.4	+2.8	
	19–22	77.4	70.7	73.0	69.3	69.3	64.4	68.9	69.3	68.6	68.1	66.4	67.1	68.5	68.5	63.7	64.5	69.4	64.5	+2.1	
	23–26	76.1	72.8	77.0	70.8	76.0	70.5	73.7	67.9	64.6	69.9	66.7	69.9	64.5	64.5	65.5	68.2	73.0	68.2	-6.0 ss	
	27–30	68.2	69.7	68.5	70.1	71.3	73.5	71.9	71.7	71.5	71.7	73.1	69.3	64.9	65.0	68.9	68.8	71.0	68.8	-1.0	
Take cocaine powder regularlyⁱ	18	86.8	86.0	84.1	84.6	85.5	84.4	84.2	82.3	81.7	82.7	82.1	81.5	82.5	83.4	81.8	83.3	83.3	83.9	+0.6	
	19–22	93.8	92.8	91.5	92.4	90.7	89.8	91.0	92.0	91.6	90.7	89.1	89.5	92.3	92.3	90.7	91.0	88.3	90.2	+1.9	
	23–26	94.8	90.8	93.7	93.6	94.2	92.2	93.4	89.1	89.4	91.2	92.9	92.3	90.5	90.5	91.0	93.8	90.6	88.7	-1.9	
	27–30	91.7	93.0	92.3	93.1	91.5	94.0	93.3	94.1	93.1	93.9	92.4	92.5	90.1	90.2	92.1	91.5	92.2	90.9	-1.3	
Try heroin once or twice^g	18	52.5	56.7	57.8	56.0	54.2	55.6	56.0	58.0	56.6	55.2	59.1	58.4	55.5	59.3	58.3	59.1	59.4	61.7	+2.3	
	19–22	61.0	63.9	60.7	63.5	63.2	64.0	63.1	64.6	67.3	66.5	65.0	69.6	67.7	67.3	64.2	66.5	66.8	68.9	+2.0	
	23–26	63.5	67.3	67.3	68.0	70.7	71.9	69.8	70.6	68.5	69.2	67.0	68.3	70.1	69.2	75.6	71.3	74.8	69.2	-5.6 s	
	27–30	66.4	67.9	69.7	70.1	67.4	68.2	70.9	72.3	67.4	74.4	70.8	70.2	70.2	67.6	69.6	69.1	70.4	72.7	+2.3	
Take heroin occasionally^g	18	74.8	76.3	76.9	77.3	74.6	75.9	76.6	78.5	75.7	76.0	79.1	76.2	75.3	79.7	74.8	77.2	78.0	78.2	+0.2	
	19–22	82.1	84.7	80.4	82.5	82.0	83.6	82.2	84.9	85.1	83.8	84.3	85.4	84.5	83.3	81.3	82.9	82.1	85.0	+2.9	
	23–26	82.4	86.5	83.9	88.5	86.6	88.4	90.0	88.3	86.7	87.5	85.2	86.5	88.0	87.8	90.0	88.6	84.2	85.1	+0.9	
	27–30	83.8	85.8	86.6	87.1	86.5	86.4	87.9	87.4	88.6	91.2	88.3	88.5	87.7	87.7	90.1	85.8	86.2	88.6	+2.4	
Take heroin regularly^g	18	89.5	88.9	89.1	89.9	89.2	88.3	88.5	89.3	86.8	87.5	89.7	87.8	86.4	89.9	85.5	87.9	88.6	87.6	-1.0	
	19–22	94.0	93.7	92.4	92.8	94.0	91.3	92.6	93.9	94.3	94.9	94.2	93.6	92.3	92.6	90.8	91.8	93.8	93.5	-0.3	
	23–26	92.7	94.4	93.4	93.7	94.8	95.9	96.3	96.5	96.0	94.8	95.8	93.1	95.7	94.5	97.1	94.2	92.5	95.0	+2.4	
	27–30	92.1	93.8	95.0	93.7	94.2	94.5	95.9	94.9	95.0	97.3	95.3	94.8	95.4	93.9	97.2	94.7	93.6	96.2	+2.6	
Try narcotics other than heroin once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	40.4	39.9	38.4	43.1	+4.8 s
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	47.3	46.1	-1.1
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	45.3	44.1	-1.2
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	43.0	47.3	+4.3
Try narcotics other than heroin regularly^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	74.9	75.5	73.9	75.8	+1.9
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	70.3	74.9	+4.6
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	78.2	75.6	-2.7
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	76.2	75.7	-0.4
Try amphetamines once or twice^{b,h}	18	30.8	31.0	35.3	32.2	32.6	34.7	34.4	36.8	35.7	37.7	39.5	41.3	39.2	41.9	40.6	34.8	34.3	36.3	+2.0	
	19–22	36.8	30.1	31.7	33.7	35.0	34.2	38.1	40.2	36.8	38.3	40.0	38.4	42.1	39.3	40.8	34.7	31.9	33.8	+1.9	
	23–26	34.9	37.8	40.9	41.8	39.9	41.6	38.0	38.3	33.2	39.1	37.0	38.0	40.8	40.7	42.2	31.4	37.8	31.4	-6.3	
	27–30	36.2	34.5	37.6	36.3	39.4	38.5	39.0	40.5	39.2	38.2	39.7	37.4	36.5	36.2	38.5	36.9	35.3	34.0	-1.3	
Take amphetamines regularly^{b,h}	18	66.8	66.0	67.7	66.4	66.3	67.1	64.8	65.6	63.9	67.1	68.1	68.1	65.4	69.0	63.6	58.7	60.0	59.5	-0.5	
	19–22	75.8	72.3	71.9	72.4	73.4	71.1	72.7	75.0	72.4	74.1	72.1	73.8	74.2	74.7	76.9	66.1	69.8	63.9	-5.9	
	23–26	78.5	79.1	77.5	78.7	79.0	77.7	77.9	80.1	75.1	80.1	78.3	77.0	76.5	73.9	80.8	69.7	68.3	64.9	-3.4	
	27–30	75.6	77.4	81.1	82.6	80.8	79.9	79.8	81.5	77.6	78.9	78.9	77.6	78.9	80.1	81.3	75.1	73.5	67.8	-5.7	
Try crystal methamphetamine (ice)^h	18	55.3	54.4	52.7	51.2	51.3	52.7	53.8	51.2	52.4	54.6	59.1	60.2	62.2	63.4	64.9	66.5	67.8	72.2	+4.4 s	
	19–22	61.1	56.4	55.8	50.6	49.2	52.5	56.5	60.0	60.3	63.1	63.5	65.0	70.0	70.0	70.7	74.2	—	—	—	
	23–26	64.1	60.7	58.2	61.3	60.1	59.2	57.7	58.6	55.9	63.9	63.9	66.6	65.6	65.6	70.1	74.6	—	—	—	
	27–30	59.1	59.8	59.9	61.0	59.7	66.4	62.5	66.6	62.8	62.6	64.9	67.9	62.0	62.0	70.2	72.9	—	—	—	

(List of drugs continued.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage saying "great risk" ^a																	
Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	(Years Cont.)
Try bath salts (synthetic stimulants) once or twice ^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try bath salts (synthetic stimulants) occasionally ^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try Adderall once or twice ^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take Adderall occasionally ^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Try sedatives/barbiturates once or twice ^{c,h}	18	30.9	28.4	27.5	27.0	27.4	26.1	25.4	30.9	29.7	32.2	32.4	35.1	32.2	29.2	29.9	26.3		
	19–22	27.6	26.4	30.5	25.4	29.9	25.0	30.7	29.6	32.7	30.5	36.4	33.5	33.5	33.4	35.0	30.5		
	23–26	—	—	—	—	32.2	29.9	30.2	35.5	35.8	32.9	37.9	31.8	33.5	32.8	34.0	34.8		
	27–30	—	—	—	—	—	—	—	—	37.2	38.7	39.0	37.0	38.2	36.5	40.5	36.6		
Take sedatives/barbiturates regularly ^{c,h}	18	72.2	69.9	67.6	67.7	68.5	68.3	67.2	69.4	69.6	70.5	70.2	70.5	70.2	66.1	63.3	61.6		
	19–22	74.0	73.3	72.7	71.3	71.6	71.7	74.5	73.0	74.0	71.7	75.5	75.5	73.6	71.1	69.4	66.4		
	23–26	—	—	—	—	77.4	77.0	74.9	79.9	79.8	76.6	80.5	77.7	76.3	75.0	74.3	77.6		
	27–30	—	—	—	—	—	—	—	—	81.5	83.7	84.0	79.6	78.6	80.2	78.3	77.7		
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ⁱ	18	3.8	4.6	3.5	4.2	4.6	5.0	4.6	6.2	6.0	6.0	8.3	9.1	8.6	8.2	7.6	5.9		
	19–22	3.0	3.4	3.1	2.3	4.7	3.1	5.4	3.5	3.9	5.9	6.1	5.4	5.8	6.6	6.5	4.5		
	23–26	—	—	—	—	5.5	3.0	6.5	6.6	4.2	5.1	5.7	4.4	5.6	3.2	4.5	4.3		
	27–30	—	—	—	—	—	—	—	—	5.0	6.3	4.4	6.6	5.6	4.7	4.1	6.7		
Take one or two drinks nearly every day ⁱ	18	20.3	21.6	21.6	21.6	23.0	24.4	25.1	26.2	27.3	28.5	31.3	32.7	30.6	28.2	27.0	24.8		
	19–22	22.7	22.9	23.2	23.2	25.0	26.3	27.3	26.1	26.5	28.1	30.1	29.1	30.2	28.0	27.5	24.0		
	23–26	—	—	—	—	27.8	27.4	26.9	30.2	29.1	27.8	31.1	30.4	31.6	25.9	26.2	26.1		
	27–30	—	—	—	—	—	—	—	—	27.4	31.7	32.2	31.7	30.9	28.0	27.4	27.2		
Take four or five drinks nearly every day ⁱ	18	65.7	64.5	65.5	66.8	68.4	69.8	66.5	69.7	68.5	69.8	70.9	69.5	70.5	67.8	66.2	62.8		
	19–22	71.2	72.7	73.3	72.7	76.2	74.1	74.0	76.4	72.8	75.7	76.1	75.5	71.8	72.1	70.3	72.5		
	23–26	—	—	—	—	76.7	77.9	80.1	77.2	81.8	76.9	79.7	80.2	78.0	76.7	77.5	75.2		
	27–30	—	—	—	—	—	—	—	—	79.3	81.7	84.7	79.1	79.9	79.1	76.6	82.2		
Have five or more drinks once or twice each weekend ⁱ	18	35.9	36.3	36.0	38.6	41.7	43.0	39.1	41.9	42.6	44.0	47.1	48.6	49.0	48.3	46.5	45.2		
	19–22	34.2	30.1	33.5	36.6	37.9	40.2	34.6	36.7	36.9	42.4	40.6	40.8	41.8	42.4	41.9	39.9		
	23–26	—	—	—	—	38.4	39.7	39.1	39.8	35.8	37.7	40.2	39.3	37.6	36.2	40.2	37.9		
	27–30	—	—	—	—	—	—	—	—	41.0	42.3	44.1	42.2	45.1	42.9	43.2	44.6		
Smoke one or more packs of cigarettes per day ^f	18	63.7	63.3	60.5	61.2	63.8	66.5	66.0	68.6	68.0	67.2	68.2	69.4	69.2	69.5	67.6	65.6		
	19–22	66.5	61.7	64.0	62.1	69.1	71.4	70.4	70.6	71.0	73.4	72.5	77.9	72.6	76.0	71.2	71.6		
	23–26	—	—	—	—	71.1	70.1	75.7	73.6	75.5	71.4	78.5	75.3	76.3	78.4	76.4	76.0		
	27–30	—	—	—	—	—	—	—	—	72.8	75.2	77.8	75.4	77.6	75.0	75.3	75.6		
Use smokeless tobacco regularly ^h	18	—	—	—	—	—	—	25.8	30.0	33.2	32.9	34.2	37.4	35.5	38.9	36.6	33.2		
	19–22	—	—	—	—	—	—	29.7	34.1	31.1	37.1	33.5	38.9	40.1	43.3	37.6	42.3		
	23–26	—	—	—	—	—	—	37.0	38.5	35.8	37.9	40.1	38.9	41.6	44.6	42.9	46.6		
	27–30	—	—	—	—	—	—	—	—	42.8	42.8	43.8	44.3	44.1	47.3	46.3	44.2		
Approximate Weighted N Per Form =	18	3,234	3,604	3,557	3,305	3,262	3,250	3,020	3,315	3,276	2,796	2,553	2,549	2,684	2,759	2,591	2,603		
	19–22	590	585	583	585	579	547	581	570	551	565	552	533	527	480	490	500		
	23–26	—	—	—	—	540	512	545	531	527	498	511	505	518	503	465	446		
	27–30	—	—	—	—	—	—	—	—	513	587	490	486	482	473	443	450		

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Percentage saying "great risk"^a

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change
Try bath salts (synthetic stimulants) once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.2	59.5	+26.4 sss
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	44.5	62.7	+18.2 sss
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	46.7	66.3	+19.6 sss
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	48.7	64.5	+15.9 sss
Take bath salts (synthetic stimulants) occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	45.0	69.9	+24.9 sss
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	52.6	70.1	+17.5 sss
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	54.0	75.3	+21.3 sss
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	55.9	71.5	+15.6 sss
Try Adderall once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.3	31.2	27.2	31.8	+4.6 s
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	30.0	27.7	-2.3
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.9	32.9	+4.1
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.4	32.7	+3.3
Take Adderall occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	41.6	40.8	35.3	38.8	+3.5
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	38.8	39.8	+1.0
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	39.7	44.8	+5.1
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	44.1	45.0	+1.0
Try sedatives/barbiturates once or twice^{c,h}	18	29.1	26.9	29.0	26.1	25.0	25.7	26.2	27.9	24.9	24.7	28.0	27.9	25.9	29.6	28.0	27.8	27.8	29.4	+1.6
	19–22	34.1	31.4	27.7	28.5	30.3	30.0	30.7	32.7	26.7	26.9	28.9	28.1	31.9	26.2	28.7	30.1	32.8	30.5	-2.3
	23–26	35.8	37.3	40.3	39.4	37.0	38.5	34.7	36.5	22.2	29.8	26.3	25.9	28.4	31.1	36.2	28.8	35.9	31.8	-4.1
	27–30	37.2	35.7	36.7	35.2	36.3	40.9	37.3	38.6	31.4	31.7	28.8	28.0	27.8	27.5	27.4	34.4	28.7	31.9	+3.2
Take sedatives/barbiturates regularly^{c,h}	18	60.4	56.8	56.3	54.1	52.3	50.3	49.3	49.6	54.0	54.1	56.8	55.1	50.2	54.7	52.1	52.4	53.9	53.3	-0.6
	19–22	70.7	69.5	65.1	64.7	64.6	61.8	64.5	63.8	60.2	64.4	61.3	63.2	64.0	59.4	64.6	63.6	68.2	64.7	-3.4
	23–26	77.1	75.2	73.9	75.1	73.8	73.1	73.1	72.8	63.9	67.0	67.6	64.8	66.8	64.4	69.6	64.9	71.4	67.6	-3.8
	27–30	74.1	77.1	79.9	80.7	75.5	78.2	75.4	79.0	70.1	75.2	68.0	70.0	70.4	69.0	71.1	71.4	70.7	72.6	+1.9
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)ⁱ	18	7.3	6.7	8.0	8.3	6.4	8.7	7.6	8.4	8.6	8.5	9.3	10.5	10.0	9.4	10.8	9.4	8.7	9.9	+1.2
	19–22	3.3	3.2	4.2	5.7	5.4	4.8	6.6	7.5	5.1	3.8	7.7	5.1	7.9	4.1	6.8	7.2	6.4	5.8	-0.6
	23–26	4.8	4.4	4.4	6.6	3.5	5.5	5.1	5.7	4.7	5.3	5.1	4.8	6.5	5.7	5.5	4.0	3.5	4.3	+0.8
	27–30	4.7	4.0	6.2	5.9	4.7	5.5	3.1	6.9	4.6	7.3	4.2	6.2	3.4	4.1	4.7	6.6	4.8	4.0	-0.8
Take one or two drinks nearly every dayⁱ	18	25.1	24.8	24.3	21.8	21.7	23.4	21.0	20.1	23.0	23.7	25.3	25.1	24.2	23.7	25.4	24.6	23.7	23.1	-0.6
	19–22	23.0	24.2	22.1	23.9	22.1	19.6	22.7	19.8	21.3	22.1	22.0	19.0	24.4	20.6	20.8	20.1	23.1	20.0	-3.1
	23–26	22.0	20.2	21.0	26.0	21.7	23.5	23.4	19.1	22.9	19.9	22.5	21.2	21.0	21.1	20.8	14.4	18.4	18.8	+0.5
	27–30	24.0	24.8	20.8	25.3	22.0	22.7	21.7	21.4	21.8	23.7	20.2	21.5	20.6	18.2	16.9	19.8	17.4	-2.4	
Take four or five drinks nearly every dayⁱ	18	65.6	63.0	62.1	61.1	59.9	60.7	58.8	57.8	59.2	61.8	63.4	61.8	60.8	62.4	61.1	62.3	63.6	62.4	-1.2
	19–22	68.5	71.4	70.4	69.9	69.9	64.5	71.1	66.4	65.3	63.0	66.6	68.8	68.5	67.1	65.6	67.4	69.6	68.7	-0.8
	23–26	72.0	75.1	69.3	72.8	71.7	75.8	74.9	71.1	74.2	71.2	72.4	70.2	70.0	67.8	68.3	69.9	73.1	69.7	-3.5
	27–30	76.1	79.3	75.7	75.1	77.4	72.8	76.2	70.6	72.1	77.5	73.0	76.5	77.1	71.6	71.6	73.8	71.2	68.3	-2.9
Have five or more drinks once or twice each weekendⁱ	18	49.5	43.0	42.8	43.1	42.7	43.6	42.2	43.5	43.6	45.0	47.6	45.8	46.3	48.0	46.3	47.6	48.8	45.8	-3.0
	19–22	40.7	36.6	42.0	37.2	38.9	37.2	37.8	40.4	38.1	37.5	37.2	43.4	41.7	35.2	40.7	40.1	41.6	40.6	-1.0
	23–26	39.1	37.4	41.1	40.2	34.9	39.0	36.8	36.3	37.9	36.8	38.4	39.7	37.0	36.2	35.8	33.6	39.5	40.2	+0.7
	27–30	41.5	40.0	40.2	41.9	37.9	41.6	40.6	42.5	40.5	44.0	39.1	40.4	40.4	40.1	38.6	42.0	41.6	37.2	-4.4
Smoke one or more packs of cigarettes per day^f	18	68.2	68.7	70.8	70.8	73.1	73.3	74.2	72.1	74.0	76.5	77.6	77.3	74.0	74.9	75.0	77.7	78.2	78.2	0.0
	19–22	73.8	76.3	77.2	75.7	77.1	76.6	80.6	77.8	81.1	80.5	80.8	79.3	79.5	80.3	79.7	81.5	82.3	82.8	+0.5
	23–26	76.0	77.6	76.5	80.9	79.7	83.9	85.1	83.6	84.1	81.6	86.4	80.7	83.6	82.0	83.2	84.8	83.1	82.9	-0.2
	27–30	73.0	80.3	80.9	80.7	78.4	82.7	80.6	82.0	81.7	84.1	83.8	84.3	86.6	83.6	89.3	86.6	84.6	84.1	-0.5
Use smokeless tobacco regularly^h	18	37.4	38.6	40.9	41.1	42.2	45.4	42.6	43.3	45.0	43.6	45.9	44.0	42.9	40.8	41.2	42.6	44.3	41.6	-2.7
	19–22	40.9	46.5	47.4	47.0	52.0	48.4	53.6	50.8	49.9	47.6	46.4	48.9	48.7	44.6	45.8	46.0	56.7	52.8	-3.8
	23–26	47.2	46.2	48.4	53.1	49.8	59.8	61.4	58.9	57.8	55.8	59.1	55.3	51.0	52.2	54.2	53.7	59.4	53.5	-5.8
	27–30	43.6	50.2	52.6	53.6	49.9	53.2	56.7	58.2	55.7	58.9	57.5	61.4	61.7	53.6	59.2	62.5	59.6	58.5	-1.1
<i>Approximate Weighted N Per Form =</i>	18	2,449	2,579	2,564	2,306	2,130	2,173	2,198	2,466	2,491	2,512	2,407	2,450	2,389	2,290	2,440	2,408	2,331	2,098	
	19–22	469	464	431	447	424	430	395	402	447	412	411	375	377	393	363	374	345	337	
	23–26	438	420	413	418	400	392	382	401	426	408	361	351	375	345	363	366	323	337	
	27–30	422	434	416	400	377	384	369	380	388	374	358	344	350	337	343	319	335	320	

Source. The Monitoring the Future study, the University of Michigan.

TABLE 6-2
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. Do you disapprove of people (who are 18 or older) doing each of the following?	Age Group	Percentage disapproving ^e																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	
Trying marijuana once or twice^j	18	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8	68.7	69.9	63.3	57.6	56.7	
	19–22	38.2	36.1	37.0	42.0	44.1	46.6	51.6	52.8	55.8	62.4	59.6	60.4	57.8	60.6	63.5	57.1	
	23–26	—	—	—	—	41.2	38.6	42.6	49.1	48.7	52.5	57.5	58.8	55.0	54.6	52.3	51.9	
	27–30	—	—	—	—	—	—	—	—	49.0	50.9	53.8	54.6	51.9	56.8	55.7	57.5	
Smoking marijuana occasionally^j	18	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5	79.4	79.7	75.5	68.9	66.7	
	19–22	49.6	49.1	51.3	56.0	60.4	62.6	66.7	67.2	69.5	77.3	76.3	77.0	74.8	75.8	76.9	70.4	
	23–26	—	—	—	—	54.8	52.8	57.0	64.9	63.4	69.4	73.7	73.3	74.0	71.9	70.9	68.1	
	27–30	—	—	—	—	—	—	—	—	65.3	67.1	68.9	73.0	67.2	72.2	69.4	72.5	
Smoking marijuana regularly^j	18	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0	89.3	90.1	87.6	82.3	81.9	
	19–22	74.3	77.2	80.0	81.8	84.9	86.7	89.2	88.7	89.1	91.2	93.1	91.3	89.5	90.2	90.1	86.8	
	23–26	—	—	—	—	80.6	81.3	83.3	87.4	86.9	90.4	91.0	89.6	90.2	92.1	90.3	90.1	
	27–30	—	—	—	—	—	—	—	—	87.6	87.5	89.7	89.6	87.2	89.4	88.7	91.9	
Trying LSD once or twice^h	18	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8	90.1	88.1	85.9	82.5	81.1	
	19–22	87.4	84.8	85.9	88.4	88.1	89.1	90.4	90.0	90.9	89.3	90.5	88.4	84.6	88.5	86.8	84.2	
	23–26	—	—	—	—	87.3	87.1	88.0	89.9	91.4	91.0	90.7	89.1	88.8	86.9	87.3	87.1	
	27–30	—	—	—	—	—	—	—	—	91.0	87.2	89.7	87.9	85.6	88.8	88.2	87.4	
Taking LSD regularly^h	18	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3	96.4	95.5	95.8	94.3	92.5	
	19–22	98.2	97.4	97.7	97.6	97.6	98.8	98.5	98.0	98.1	97.5	99.1	97.5	97.0	97.8	97.7	96.8	
	23–26	—	—	—	—	99.2	98.0	98.5	99.0	98.0	98.4	98.3	98.4	98.3	98.1	97.7	96.7	
	27–30	—	—	—	—	—	—	—	—	98.8	97.1	98.9	98.9	97.5	98.5	98.7	98.6	
Trying ecstasy (MDMA) once or twice^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Taking ecstasy (MDMA) occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Trying cocaine once or twice^h	18	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5	93.6	93.0	92.7	91.6	90.3	
	19–22	73.0	69.3	69.9	74.1	72.5	77.6	78.9	82.3	85.3	88.8	90.1	91.2	90.6	92.7	93.9	94.2	
	23–26	—	—	—	—	70.2	70.5	72.1	80.0	82.9	85.5	88.3	88.0	87.3	89.2	89.2	91.8	
	27–30	—	—	—	—	—	—	—	—	82.1	81.0	85.5	86.9	83.9	85.7	86.6	86.6	
Taking cocaine regularly^h	18	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7	97.3	96.9	97.5	96.6	96.1	
	19–22	91.6	89.3	91.9	94.6	95.0	96.3	97.0	97.2	97.9	97.4	98.9	97.9	98.4	97.8	98.8	98.2	
	23–26	—	—	—	—	95.7	95.3	97.3	98.1	97.6	98.3	98.4	98.5	98.7	98.4	98.8	97.7	
	27–30	—	—	—	—	—	—	—	—	98.1	97.0	99.3	99.0	97.2	98.7	99.0	98.9	
Trying heroin once or twice^h	18	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1	96.0	94.9	94.4	93.2	92.8	
	19–22	96.3	95.4	95.6	95.2	95.1	96.2	96.8	96.3	97.1	96.4	98.3	95.9	95.9	96.3	96.6	95.6	
	23–26	—	—	—	—	96.7	94.9	96.4	97.1	97.4	96.7	96.8	96.9	96.3	95.4	96.5	95.9	
	27–30	—	—	—	—	—	—	—	—	97.9	95.8	97.5	96.6	94.8	97.3	94.7	96.3	
Taking heroin occasionally^h	18	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7	97.3	96.8	97.0	96.2	95.7	
	19–22	98.6	97.8	98.3	98.3	98.6	98.7	98.3	98.3	98.3	97.9	99.2	98.2	98.1	98.1	98.3	97.7	
	23–26	—	—	—	—	99.2	98.2	98.8	99.1	98.4	98.3	98.1	99.0	98.7	98.4	98.6	97.7	
	27–30	—	—	—	—	—	—	—	—	99.2	97.3	99.0	98.9	97.0	98.9	98.7	98.9	
Taking heroin regularly^h	18	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5	97.8	97.2	97.5	97.1	96.4	
	19–22	99.2	98.5	98.6	98.7	98.7	99.1	98.9	98.6	98.4	98.3	99.5	98.5	98.3	98.4	98.8	98.4	
	23–26	—	—	—	—	99.4	98.8	99.1	99.4	98.7	98.7	98.5	99.3	99.2	98.9	98.8	98.7	
	27–30	—	—	—	—	—	—	—	—	99.4	97.6	99.4	99.0	97.8	99.0	99.4	99.1	

(Table continued on next page.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

	Age Group	Percentage disapproving ^e																		2012–2013 change
		1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Trying marijuana once or twice^j	18	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	55.0	55.6	58.6	55.5	54.8	51.6	51.3	48.8	49.1	+0.3
	19–22	55.4	56.2	55.9	54.0	55.2	49.3	48.7	54.2	48.3	50.3	51.2	47.6	52.7	46.7	50.5	49.0	46.0	44.2	-1.9
	23–26	56.3	54.5	55.3	55.7	54.8	51.2	52.4	47.8	53.4	47.7	47.5	54.6	46.2	44.9	42.5	38.8	40.9	38.1	-2.8
	27–30	54.1	59.0	55.7	52.6	58.0	54.4	56.9	54.9	55.4	52.1	52.0	50.9	49.3	49.3	48.5	46.5	42.7	38.7	-4.0 s
Smoking marijuana occasionally^j	18	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	67.8	69.3	70.2	67.3	65.6	62.0	60.9	59.1	58.9	-0.2
	19–22	68.9	70.2	67.8	66.4	70.7	64.6	62.3	68.0	64.3	67.9	62.6	64.1	63.3	59.8	61.3	61.7	58.2	54.9	-3.3
	23–26	72.5	69.2	70.4	71.1	68.6	67.4	64.0	63.8	69.3	65.6	62.2	68.0	64.5	62.4	59.1	53.1	55.8	51.3	-4.5 s
	27–30	70.5	74.5	72.4	71.5	72.2	70.9	69.1	71.2	69.1	68.2	68.7	67.5	63.7	63.7	62.7	63.7	58.3	55.0	-3.3
Smoking marijuana regularly^j	18	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	82.0	82.2	83.3	79.6	80.3	77.7	77.5	77.8	74.5	-3.3 s
	19–22	87.7	88.1	85.3	84.5	86.6	84.5	82.8	84.8	82.7	84.4	82.5	83.7	83.6	80.8	80.7	78.1	77.0	75.7	-1.4
	23–26	88.9	88.1	87.5	86.1	83.9	86.4	81.7	82.3	87.4	84.3	81.9	85.3	84.3	80.2	78.3	76.4	76.7	73.6	-3.0
	27–30	89.9	92.1	89.2	90.0	89.5	89.3	88.8	87.7	88.6	86.3	86.4	86.8	86.0	84.4	81.7	83.2	77.8	75.9	-2.0
Trying LSD once or twice^h	18	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	87.9	88.0	87.8	85.5	88.2	86.5	86.3	87.2	86.6	-0.5
	19–22	83.0	83.1	80.8	83.2	82.3	81.4	83.7	86.2	85.0	87.6	85.4	88.5	86.5	83.0	86.7	83.3	84.0	83.5	-0.5
	23–26	86.7	87.9	84.1	84.8	80.3	83.0	79.2	80.1	84.0	84.0	84.5	87.6	81.8	85.0	82.6	80.1	83.3	79.7	-3.6
	27–30	88.7	88.7	87.3	86.6	87.2	85.7	82.7	85.6	82.5	82.2	82.0	84.1	82.7	84.5	85.1	85.1	82.4	81.4	-1.0
Taking LSD regularly^h	18	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	95.6	95.9	94.9	93.5	95.3	94.3	94.9	95.2	95.3	+0.1
	19–22	97.0	97.4	96.3	97.0	96.8	96.5	96.9	98.4	97.3	98.9	97.8	97.7	96.8	96.8	96.6	96.5	96.0	96.7	+0.7
	23–26	97.7	96.1	97.6	98.0	97.0	97.1	97.9	96.9	97.1	98.7	97.0	98.4	97.4	98.2	96.5	95.9	97.4	96.1	-1.3
	27–30	98.1	97.5	97.4	97.9	98.6	98.2	98.0	98.2	98.2	97.2	96.7	97.2	97.1	98.6	98.6	97.1	97.3	97.2	-0.2
Trying ecstasy (MDMA) once or twice^h	18	—	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	88.4	89.0	87.8	88.2	88.2	86.3	83.9	87.1	84.9	-2.2
	19–22	—	—	—	—	—	81.5	80.3	87.2	83.5	90.3	87.5	88.5	89.5	89.1	91.4	85.9	87.9	83.9	-4.0
	23–26	—	—	—	—	—	80.6	80.6	80.2	83.1	83.9	83.9	87.4	83.9	85.0	86.9	85.1	85.2	79.9	-5.3
	27–30	—	—	—	—	—	84.2	84.0	86.3	83.2	82.4	82.2	81.8	82.7	83.0	81.9	86.6	83.7	84.5	+0.8
Taking ecstasy (MDMA) occasionally^h	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	19–22	—	—	—	—	—	92.8	91.8	95.6	93.8	96.7	94.0	95.3	94.8	95.2	95.3	91.8	94.4	92.3	-2.1
	23–26	—	—	—	—	—	90.5	91.8	92.1	93.3	94.4	93.7	94.3	94.0	95.4	94.3	92.5	93.3	92.1	-1.2
	27–30	—	—	—	—	—	91.7	93.0	94.3	91.0	92.1	93.4	92.8	94.1	93.6	92.6	94.5	93.5	93.0	-0.6
Trying cocaine once or twice^h	18	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	88.9	89.1	89.6	89.2	90.8	90.5	91.1	91.0	92.3	+1.4
	19–22	92.0	91.7	89.9	90.9	89.9	87.7	87.9	89.3	87.7	92.3	88.2	89.2	85.8	87.8	87.1	90.1	89.7	90.5	+0.7
	23–26	90.7	91.5	89.0	91.3	87.1	90.1	85.8	86.4	87.4	88.3	84.4	87.6	84.5	86.2	86.0	82.7	86.0	85.7	-0.3
	27–30	88.3	89.2	90.3	90.4	89.4	90.3	88.5	91.5	88.0	87.0	85.8	87.7	87.4	88.3	87.3	87.0	85.6	82.5	-3.1
Taking cocaine regularly^h	18	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	96.0	96.1	96.2	94.8	96.5	96.0	96.0	96.8	96.7	-0.1
	19–22	97.9	98.0	97.8	97.6	98.0	97.2	97.0	98.2	98.5	98.7	98.9	99.0	97.6	97.6	97.6	97.2	97.6	97.4	-0.2
	23–26	97.8	96.9	98.5	98.3	97.8	97.5	97.5	97.6	98.1	98.9	97.3	98.1	98.0	98.7	97.6	97.3	98.8	97.8	-1.0
	27–30	98.5	97.9	97.8	98.8	98.7	98.4	97.8	98.8	98.8	97.8	97.2	97.9	97.3	99.0	99.0	98.4	98.5	98.0	-0.5
Trying heroin once or twice^h	18	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	94.3	93.8	94.8	93.3	94.7	93.9	94.3	95.8	95.6	-0.1
	19–22	95.2	95.6	95.1	95.5	94.1	94.2	95.0	96.4	95.9	98.8	95.6	97.6	95.7	95.5	95.8	96.7	95.9	96.3	+0.4
	23–26	96.1	95.2	94.6	96.3	93.1	95.0	94.8	95.0	95.0	96.1	93.7	97.2	95.6	94.9	94.5	95.5	95.7	94.7	-0.9
	27–30	96.0	96.9	95.9	96.7	95.9	96.4	94.4	97.6	94.9	95.6	93.9	96.4	96.2	95.4	96.3	95.7	95.9	94.8	-1.1
Taking heroin occasionally^h	18	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	96.3	96.2	96.8	95.3	96.9	96.2	96.3	97.0	96.9	-0.1
	19–22	97.9	97.8	98.2	97.2	98.0	97.9	97.9	98.3	98.9	99.4	98.2	98.8	97.3	97.9	97.5	97.7	97.4	98.0	+0.5
	23–26	98.7	97.4	97.5	98.5	98.2	97.8	97.5	97.2	98.5	98.3	97.7	98.8	98.3	98.5	97.1	99.0	99.0	98.1	-0.8
	27–30	98.0	98.7	97.6	98.8	98.6	98.4	98.6	98.7	98.1	97.7	97.1	98.1	98.2	98.6	99.3	98.1	97.9	97.7	-0.2
Taking heroin regularly^h	18	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	96.7	96.9	97.1	95.9	97.4	96.4	96.7	97.4	97.4	-0.1
	19–22	98.3	98.1	98.3	98.2	98.5	98.2	98.3	98.8	99.0	99.2	98.9	99.1	98.3	98.1	97.6	97.9	98.3	98.4	+0.1
	23–26	98.9	97.6	98.5	98.7	98.8	98.4	98.3	98.6	98.9	98.9	98.0	99.0	99.1	99.2	97.6	99.3	99.1	98.3	-0.8
	27–30	98.6	98.4	98.1	98.8	98.7	98.7	98.4	99.3	98.8	99.1	97.5	98.2	98.4	99.0	99.3	98.6	98.3	97.9	-0.5

↓
(List of drugs continued.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage disapproving ^e																	
Q. Do you disapprove of people (who are 18 or older) doing each of the following?		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	(Years Cont.)
Trying amphetamines once or twice^{b,h}	18	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3	86.5	86.9	84.2	81.3	82.2		
	19–22	74.5	70.5	68.9	74.0	73.0	75.6	78.9	79.9	81.8	85.3	84.4	83.9	83.8	87.2	88.3	85.0		
	23–26	—	—	—	—	74.2	74.2	74.6	80.3	83.5	83.3	84.1	84.8	83.4	84.8	82.7	86.0		
	27–30	—	—	—	—	—	—	—	—	83.5	81.0	84.3	83.7	80.9	83.5	82.0	83.1		
Taking amphetamines regularly^{b,h}	18	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5	96.0	95.6	96.0	94.1	94.3		
	19–22	94.8	93.3	94.3	93.4	94.9	96.6	96.9	95.1	97.5	96.8	97.5	97.7	96.7	97.3	97.9	96.8		
	23–26	—	—	—	—	96.6	95.9	96.6	97.0	97.2	98.1	97.9	97.7	97.7	98.4	97.7	97.0		
	27–30	—	—	—	—	—	—	—	—	98.1	96.5	98.6	97.8	96.8	97.7	99.0	98.9		
Trying sedatives/barbiturates once or twice^{c,h}	18	83.9	82.4	84.4	83.1	84.1	84.9	86.8	89.6	89.4	89.3	90.5	90.6	90.3	89.7	87.5	87.3		
	19–22	83.5	82.3	83.8	85.1	85.2	86.1	88.3	87.5	90.1	92.0	91.1	90.4	88.8	90.7	91.1	90.5		
	23–26	—	—	—	—	84.0	84.5	84.4	89.8	90.7	89.4	88.8	87.9	88.8	88.5	88.0	89.3		
	27–30	—	—	—	—	—	—	—	—	90.5	88.3	88.4	88.8	86.6	88.9	87.6	88.0		
Taking sedatives/barbiturates regularly^{c,h}	18	95.4	94.2	94.4	95.1	95.1	95.5	94.9	96.4	95.3	95.3	96.4	97.1	96.5	97.0	96.1	95.2		
	19–22	96.6	95.6	97.3	96.5	96.6	98.1	98.0	97.0	97.9	97.7	98.7	98.0	97.9	98.2	98.7	97.7		
	23–26	—	—	—	—	98.4	98.5	97.7	98.6	98.3	98.3	98.5	98.5	98.6	98.5	98.5	97.4		
	27–30	—	—	—	—	—	—	—	—	98.4	97.1	99.1	98.5	97.7	98.4	99.1	99.0		
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)ⁱ	18	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4	29.8	33.0	30.1	28.4	27.3		
	19–22	14.8	14.5	13.9	15.5	15.3	15.4	16.9	16.0	18.4	22.4	17.6	22.2	16.9	20.8	22.2	22.0		
	23–26	—	—	—	—	17.4	16.1	13.2	17.7	13.7	17.5	18.6	19.5	17.4	18.1	17.6	16.5		
	27–30	—	—	—	—	—	—	—	—	19.5	19.1	18.7	18.8	17.9	19.5	18.6	18.2		
Taking one or two drinks nearly every day^j	18	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9	76.5	75.9	77.8	73.1	73.3		
	19–22	67.8	69.7	71.3	73.3	74.3	71.3	77.4	75.3	76.5	80.0	79.7	77.1	76.0	75.0	78.0	74.7		
	23–26	—	—	—	—	71.4	73.7	71.6	72.7	74.6	74.4	77.6	76.9	75.5	74.2	73.3	69.7		
	27–30	—	—	—	—	—	—	—	—	76.0	73.9	73.3	76.1	69.5	73.5	72.4	71.8		
Taking four or five drinks nearly every day^j	18	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9	90.6	90.8	90.6	89.8	88.8		
	19–22	95.2	93.4	94.6	94.6	94.6	94.8	94.9	95.7	94.8	96.1	95.8	96.4	95.5	95.1	96.2	95.5		
	23–26	—	—	—	—	96.2	95.0	95.5	96.9	94.3	95.9	96.9	96.1	95.7	95.7	95.7	95.2		
	27–30	—	—	—	—	—	—	—	—	97.4	94.6	96.1	95.3	94.8	94.8	96.4	96.7		
Having five or more drinks once or twice each weekend^j	18	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9	67.4	70.7	70.1	65.1	66.7		
	19–22	57.1	56.1	58.2	61.0	59.7	59.4	60.3	61.6	64.1	66.3	67.1	62.4	65.6	63.5	68.1	66.0		
	23–26	—	—	—	—	66.2	68.3	66.5	67.5	65.2	63.2	66.9	64.6	69.6	66.8	66.9	65.3		
	27–30	—	—	—	—	—	—	—	—	73.9	71.4	73.1	72.1	68.4	73.4	73.5	73.7		
Smoking one or more packs of cigarettes per day^j	18	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8	71.4	73.5	70.6	69.8	68.2		
	19–22	68.7	68.1	66.3	71.6	69.0	70.5	71.4	72.7	73.8	75.6	73.7	73.2	72.6	72.8	75.3	69.8		
	23–26	—	—	—	—	69.9	68.7	67.5	69.7	66.4	71.1	71.5	77.2	73.6	72.9	70.3	72.2		
	27–30	—	—	—	—	—	—	—	—	72.8	69.4	73.5	71.2	70.7	73.8	72.3	73.9		
<i>Approximate Weighted N Per Form =</i>	18	3,261	3,610	3,651	3,341	3,254	3,265	3,113	3,302	3,311	2,799	2,566	2,547	2,645	2,723	2,588	2,603		
	19–22	588	573	605	579	586	551	605	587	560	567	569	533	530	489	474	465		
	23–26	—	—	—	—	542	535	560	532	538	516	524	495	538	514	475	466		
	27–30	—	—	—	—	—	—	—	—	526	509	513	485	512	462	442	450		

(Table continued on next page.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Percentage disapproving*

Q. Do you disapprove of people (who are 18 or older) doing each of the following?	Age Group	Percentage disapproving*																		2012–2013 change
		1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Trying amphetamines once or twice ^{b,h}	18	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	86.1	86.3	87.3	87.2	88.2	88.1	84.1	83.9	84.9	+1.0
	19–22	84.4	83.3	84.6	84.9	83.8	82.1	81.4	86.3	82.1	88.2	84.9	84.8	86.7	85.4	86.9	80.5	81.3	83.7	+2.4
	23–26	86.4	85.7	83.5	84.5	82.4	83.9	83.5	79.9	81.6	81.3	79.0	85.8	79.7	84.4	84.1	76.5	80.7	77.3	-3.4
	27–30	85.8	86.3	85.9	86.4	84.5	86.0	86.4	84.9	82.4	81.3	81.1	84.5	83.7	82.9	84.3	81.1	81.9	81.5	-0.4
Taking amphetamines regularly ^{b,h}	18	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	94.8	95.3	95.4	94.2	95.6	94.9	92.9	93.9	93.2	-0.8
	19–22	97.2	97.8	96.7	97.5	96.1	97.3	96.4	97.1	97.1	98.4	97.5	98.6	96.2	96.8	96.2	92.1	94.1	94.4	+0.3
	23–26	97.9	97.0	98.0	97.0	97.6	96.8	96.3	97.2	95.9	98.3	96.2	97.6	97.3	98.1	96.8	94.8	95.9	94.6	-1.4
	27–30	98.2	98.1	97.7	98.2	98.5	97.6	97.4	98.1	98.0	97.6	96.4	98.4	97.2	98.1	98.0	97.5	95.8	96.8	+1.0
Trying sedatives/barbiturates once or twice ^{c,h}	18	84.9	86.4	86.0	86.6	85.9	85.9	86.6	87.8	83.7	85.4	85.3	86.5	86.1	87.7	87.6	87.3	88.2	88.9	+0.7
	19–22	89.1	86.6	85.8	86.6	84.2	85.2	84.2	87.7	81.8	86.6	83.4	82.7	82.1	84.7	85.2	85.4	88.0	88.6	+0.6
	23–26	88.3	88.3	87.4	87.3	85.2	86.9	86.8	81.8	80.3	81.6	80.5	84.3	77.7	83.3	80.9	80.6	83.8	84.4	+0.6
	27–30	89.4	88.8	88.4	87.6	87.3	88.5	86.9	89.2	81.8	78.7	80.1	83.5	80.5	82.5	80.3	83.3	83.1	82.6	-0.5
Taking sedatives/barbiturates regularly ^{c,h}	18	94.8	95.3	94.6	94.7	95.2	94.5	94.7	94.4	94.2	95.2	95.1	94.6	94.3	95.8	94.7	95.1	96.1	95.8	-0.3
	19–22	97.9	97.7	97.7	97.3	97.4	96.9	97.8	98.5	96.6	98.3	98.1	98.3	96.7	96.7	96.3	96.7	96.4	96.5	+0.1
	23–26	98.4	97.4	98.5	97.6	97.4	97.0	97.1	97.1	96.1	98.0	96.3	97.8	96.7	98.4	95.7	98.1	97.3	97.2	-0.1
	27–30	98.5	97.9	97.7	98.5	98.1	98.4	97.2	98.4	98.1	96.5	95.6	97.4	97.4	98.4	98.6	97.0	97.7	97.1	-0.6
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor) ⁱ	18	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	26.4	29.0	31.0	29.8	30.6	30.7	28.7	25.4	27.3	+1.8
	19–22	22.0	18.3	21.5	18.3	18.4	16.3	18.3	20.1	20.7	22.3	17.8	17.3	20.5	19.1	23.7	21.6	21.4	19.6	-1.8
	23–26	18.0	15.8	18.6	19.1	19.9	15.9	18.1	13.0	16.3	13.5	14.7	14.9	12.5	16.0	15.4	10.9	14.1	13.5	-0.6
	27–30	16.1	17.4	15.2	15.9	14.8	15.9	18.4	15.4	18.8	16.1	15.0	14.2	11.9	11.5	13.3	11.8	14.7	13.2	-1.5
Taking one or two drinks nearly every day ⁱ	18	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	70.8	72.8	73.3	74.5	70.5	71.5	72.8	70.8	71.9	+1.1
	19–22	73.5	73.2	70.3	67.3	66.7	68.3	63.9	66.9	68.1	64.6	68.2	65.1	65.2	67.4	68.4	71.0	65.7	64.0	-1.6
	23–26	70.6	68.4	70.2	73.4	66.3	66.5	62.7	65.0	61.7	64.4	62.0	62.4	66.4	62.0	62.5	55.7	53.9	54.4	+0.4
	27–30	71.4	71.8	69.8	67.9	65.9	68.9	70.9	63.1	66.7	60.5	62.0	65.8	59.5	63.7	61.4	61.7	55.6	51.3	-4.3
Taking four or five drinks nearly every day ⁱ	18	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	89.4	90.6	90.5	89.8	89.7	88.8	90.8	90.1	90.6	+0.5
	19–22	94.2	93.9	92.4	92.4	92.8	94.2	92.6	92.5	92.2	93.2	92.9	92.9	94.0	93.6	92.2	93.9	91.9	92.1	+0.3
	23–26	96.5	93.8	96.1	95.1	94.3	93.5	93.7	92.6	93.1	94.8	92.9	95.6	94.9	94.6	93.9	94.7	92.8	91.8	-1.0
	27–30	96.4	96.2	95.0	97.2	95.3	96.1	95.4	95.6	96.0	92.8	92.7	95.0	93.9	96.0	94.3	95.8	92.1	92.1	-0.1
Having five or more drinks once or twice each weekend ⁱ	18	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	66.5	68.5	68.8	68.9	67.6	68.8	70.0	70.1	71.6	+1.5
	19–22	69.2	66.5	63.2	63.5	65.1	58.3	57.5	61.9	59.4	60.1	59.3	59.1	63.4	62.3	62.7	65.4	64.7	66.3	+1.7
	23–26	70.9	66.6	69.5	68.1	66.2	66.0	61.2	65.5	60.9	64.5	59.7	62.4	63.0	59.5	61.7	55.9	63.0	63.3	+0.3
	27–30	72.4	73.0	71.1	73.1	73.1	73.0	70.9	71.5	73.8	67.5	67.3	71.5	66.4	65.8	67.5	64.9	63.3	65.0	+1.7
Smoking one or more packs of cigarettes per day ^j	18	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	79.8	81.5	80.7	80.5	81.8	81.0	83.0	83.7	82.6	-1.1
	19–22	72.2	74.3	72.3	70.1	73.1	73.2	73.4	73.4	74.8	81.5	77.2	81.0	80.4	81.8	82.9	83.8	79.5	81.0	+1.5
	23–26	73.0	71.7	73.9	73.8	72.7	77.3	74.8	75.7	76.2	74.8	74.1	76.2	77.9	77.3	77.9	80.3	78.2	77.8	-0.4
	27–30	72.7	74.3	71.7	71.0	78.6	75.2	78.8	76.2	77.6	77.3	73.9	81.1	74.5	80.9	79.6	79.5	79.1	79.9	+0.9
<i>Approximate Weighted N Per Form =</i>	18	2,399	2,601	2,545	2,310	2,150	2,144	2,160	2,442	2,455	2,460	2,377	2,450	2,314	2,233	2,449	2,384	2,301	2,147	
	19–22	480	470	446	449	416	413	402	396	431	378	378	333	365	368	364	340	356	280	
	23–26	449	423	401	397	389	404	346	385	403	374	364	325	335	328	347	309	334	312	
	27–30	430	453	449	429	395	368	359	346	370	367	330	355	339	325	334	306	312	301	

Source. The Monitoring the Future study, the University of Michigan.

Footnotes for Tables 6-1 through 6-2

Notes. Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

The illicit drugs not listed here show a daily prevalence of 0.2% or less in all years.

' * ' indicates a prevalence rate of less than 0.05%.

' — ' indicates data not available.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bIn 2011 the list of examples was changed from upper, pep pills, bennies, and speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

^cIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers.

These changes likely explain the discontinuity in the 2003 and 2004 results.

^dFor 12th graders only: In 2011 the question on perceived risk of using salvia once or twice appeared at the end of a questionnaire form. In 2012 the question was moved to an earlier section of the same form. A question on perceived risk of using salvia occasionally was also added following the question on perceived risk of trying salvia once or twice. These changes likely explain the discontinuity in the 2012 result.

^eAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

^fAge 18 data based on one questionnaire form for all years reported. For ages 19-30 only: Prior to 2012, data based on one questionnaire form. Data based on five questionnaire forms in 2012-2013.


^gAge 18 data based on one questionnaire form for all years reported. For ages 19-30 only: Prior to 2012, data based on one questionnaire form. Data based on two questionnaire forms in 2012-2013.

^hData based on one questionnaire form.

ⁱAge 18 data based on one questionnaire form for all years reported. For ages 19-30 only: Prior to 2012, data based on one questionnaire form. Data based on three questionnaire forms in 2012-2013.


^jAge 18 data based on one questionnaire form for all years reported. For ages 19-30 only: Prior to 2012, data based on one questionnaire form. Data based on four questionnaire forms in 2012-2013.

FIGURE 6-1
Trends in Harmfulness of MARIJUANA Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-2
Trends in Harmfulness of MARIJUANA Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Occasional Use




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-3
Trends in Harmfulness of MARIJUANA Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-4
Trends in Harmfulness of LSD Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-5
Trends in Harmfulness of LSD Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-6
Trends in Harmfulness of PCP Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-7
Trends in Harmfulness of COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-8
Trends in Harmfulness of COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Occasional Use




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-9
Trends in Harmfulness of COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use




Source. The Monitoring to Future study, the University of Michigan.

FIGURE 6-10
Trends in Harmfulness of CRACK COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-11
Trends in Harmfulness of CRACK COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Occasional Use




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-12
Trends in Harmfulness of CRACK COCAINE Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-13
Trends in Harmfulness of COCAINE POWDER Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-14
Trends in Harmfulness of COCAINE POWDER Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Occasional Use




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-15
Trends in Harmfulness of COCAINE POWDER Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-16
Trends in Harmfulness of ECSTASY (MDMA) Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-17
Trends in Harmfulness of ECSTASY (MDMA) Use as Perceived by
Respondents in Modal Age Groups of 19–22, 23–26, and 27–30
Occasional Use




FIGURE 6-18
Trends in Harmfulness of HEROIN Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice






FIGURE 6-19
Trends in Harmfulness of HEROIN Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Occasional Use




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-20
Trends in Harmfulness of HEROIN Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use



Source. The Monitoring the Future study, the University of Michigan.


FIGURE 6-21
Trends in Harmfulness of AMPHETAMINE^a Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice



Source: The Monitoring the Future study, the University of Michigan.

^aIn 2011 the list of examples was changed from upper, pep pills, bennies, and speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.


FIGURE 6-22
Trends in Harmfulness of AMPHETAMINE^a Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use



Source. The Monitoring the Future study, the University of Michigan.


^aIn 2011 the list of examples was changed from upper, pep pills, bennies, and speed to uppers, speed, Adderrall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

FIGURE 6-23
Trends in Harmfulness of CRYSTAL METHAMPHETAMINE (ICE) Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice



Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-24
Trends in Harmfulness of SEDATIVE (BARBITURATE)^a Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers.

These changes likely explain the discontinuity in the 2003 and 2004 results.

FIGURE 6-25
Trends in Harmfulness of SEDATIVE (BARBITURATE)^a Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers.


These changes likely explain the discontinuity in the 2003 and 2004 results.

FIGURE 6-26
Trends in Harmfulness of ALCOHOL Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Trying Once or Twice




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-27
Trends in Harmfulness of ALCOHOL Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Having One or Two Drinks per Day




Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-28
Trends in Harmfulness of ALCOHOL Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Having Four or Five Drinks per Day




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-29
Trends in Harmfulness of BINGE DRINKING as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Having Five or More Drinks Once or Twice Each Weekend




Source: The Monitoring the Future study, the University of Michigan.

FIGURE 6-30
Trends in Harmfulness of TOBACCO Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Smoking One or More Packs of Cigarettes per Day



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 6-31
Trends in Harmfulness of SMOKELESS TOBACCO Use as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30
Regular Use



Source: The Monitoring the Future study, the University of Michigan.

Chapter 7

THE SOCIAL CONTEXT

The social contexts in which individuals place and otherwise find themselves can influence the likelihood of using drugs in a number of ways. The context can provide social modeling and social norms for either use or abstention from use. Through friends and friends' contacts it can also influence the availability of drugs and bring about an awareness of new drugs, including knowledge of their existence and potential for altering mood and consciousness. Since its inception, MTF has measured three important features of the social context: (1) peer groups' norms about drug use, (2) amount of direct exposure to drug use by friends and others, and (3) perceived availability of drugs. All three factors are measured by self-reports and are therefore measures of the perceived context, though evidence suggests that they bear a strong correlation with the actual context. We believe that these three factors exert important influences on substance use, at both the individual (micro) and the aggregate (macro) level.

In *Volume I*, we examined these factors among secondary school students. In this chapter, we do the same for the young adult population ages 19 to 30, whose social contexts typically differ considerably from what they were in high school. Most high school graduates today enter college, many get civilian jobs, and some enter military service. These transitions almost always change the institutional contexts experienced by young adults (e.g., colleges, work organizations, military services, etc.) and therefore the circles of people to whom they are exposed and with whom they develop friendships. They also alter the potential consequences of drug use if it is discovered by authorities in the relevant institution; for example, consequences can be quite severe for those in military service, and we have shown that illicit drug use drops when young people enter the military.⁷¹

Each of the question sets discussed here are contained in only one of the six questionnaire forms, so the case counts are lower than those presented in most chapters in this volume. (Also, in comparison to the secondary school samples covered in *Volume I*, follow-up samples reported here are much smaller.) Therefore, the prevalence and trend estimates are more subject to random fluctuation.

In addition, we include consideration of norms, exposure, and availability where relevant among those age 35 and older. In such cases, the data are based on the one form that is used for all respondents at a particular age.

⁷¹Bachman, J. G., Freedman-Doan, P., O'Malley, P. M., Johnston, L. D., & Segal, D. R. (1999). Changing patterns of drug use among U.S. military recruits before and after enlistment. *American Journal of Public Health, 89*, 672-677. [doi:10.2105/AJPH.89.5.672](https://doi.org/10.2105/AJPH.89.5.672)

PEER NORMS AMONG YOUNG ADULTS (AGES 18–30)

Table 7-1 provides current levels and trends in perceived friends' disapproval of drug use as reported by 12th graders, 19- to 22-year-olds, 23- to 26-year-olds, and 27- to 30-year-olds. (These are the same age groupings used in chapter 6.) Trend data are available since 1980, 1984, and 1988, respectively, for these three 4-year age groupings of young adults.

The results for perceived peer norms are generally quite consistent with those for personal disapproval in the aggregate.⁷² Exceptions are trying marijuana once or twice and smoking one or more packs of cigarettes per day, for which friends' attitudes are consistently reported as more disapproving than respondents' own attitudes (especially in the oldest age band), and heavy weekend drinking, for which friends' attitudes are seen as less disapproving than their own. The question set regarding friends' disapproval employs a shorter list of drug-using behaviors but includes the same answer scale, stated in terms of strength of disapproval associated with different use levels of the various drugs, as the questions on the respondent's own attitudes about those behaviors (discussed in chapter 6). While peer disapproval and personal disapproval questions appear on different questionnaire forms and therefore have different sets of respondents, the forms are distributed randomly in senior year and should leave no systematic sample differences.

Current Perceptions of Close Friends' Attitudes (Ages 18 to 30)

Table 7-1 provides trends for each age band in the proportions of respondents indicating how their close friends would feel about the respondent engaging in various drug-using behaviors. For purposes of simplification, we begin by addressing results across the entire 19- to 30-year age band (tabular data for the entire age band are not presented). Then we distinguish among the three young adult age bands: 19–22, 23–26, and 27–30, along with 18-year-olds. In 2010 questions about friends' disapproval were dropped from the young-adult follow-up questionnaires for all drugs except marijuana, occasions of heavy drinking, and cigarettes. The dropped questions had shown a high degree of redundancy with respondents' reports of their own attitudes in the aggregate, and thus were deleted to make room for other items.

- Generally, the peer norms reported by young adults 1 to 12 years past high school have been quite similar to those reported by 12th graders.
- With regard to *marijuana*, half of young adults (50%) thought their close friends would disapprove of their trying it, while three fifths (60%) thought their close friends would disapprove of occasional use, and about three fourths (78%) thought close friends would disapprove of regular use. Clearly the

⁷²The question reads, "How do you think your close friends feel (or would feel) about you... [smoking marijuana once or twice]?" The answer categories are "don't disapprove," "disapprove," and "strongly disapprove." Percentages discussed are for the last two categories combined.

norms differ as a function of level of marijuana use but for all levels of use they tend to be restrictive for the majority of young adults.

- For each of the *illicit drugs other than marijuana*, 2009 was the last year in which results on peer norms were available. At that time, the great majority of young adults said that their close friends would disapprove of their even *trying* such drugs once or twice; 89% indicated this for *cocaine*, 87% for *LSD*, and 87% for *amphetamines*. (We stopped asking these questions beginning in 2010 to make space for new items on the survey and because the data that they provided on peer norms so closely tracked what their own attitudes were in the aggregate.)
- In 2009 nearly two thirds (63%) of young adults said their close friends would disapprove if they were *daily drinkers*, and 9 out of 10 (91%) thought friends would disapprove if they had *four or five drinks nearly every day*.
- Friends' disapproval of *heavy drinking on weekends* continues to be measured and is distinctly lower than disapproval of daily drinking. In 2013, 49% to 53% of any of the young adult age groups thought that their close friends would disapprove of their having five or more drinks once or twice each weekend. These levels of disapproval are lower than among 18-year-olds (65%).
- Peer disapproval of *cigarette* smoking is high in all four age bands: In 2013, 84% of 12th graders said their friends would disapprove of pack-a-day smoking, as did 86% to 90% of 19- to 30-year-olds.

Trends in Peer Norms (Ages 18 to 30)

Important changes in the social acceptability of drug-using behaviors among both 12th graders' and young adults' peers have occurred since MTF began (see Table 7-1).

- Among 12th graders, the proportion saying their close friends would disapprove of their *trying marijuana* rose from 41% in 1979 to 73% in 1992—a period of substantial decline in use. Friends' disapproval also grew substantially stronger in all of the young adult age bands in the years for which data are available. For example, among 19- to 22-year-olds, the proportion thinking their close friends would disapprove if they even tried marijuana rose from 41% in 1981 to 65% in 1992 (Table 7-1). A similar peak in disapproval occurred for 23- to 26-year-olds in 1992 and 1993, and among 27- to 30-year-olds in 1994 and 1995—66% for both age bands. In all age groups, peer disapproval subsequently declined, though the declines were earliest and greatest among 12th graders. The decline ended in 1997 for 12th graders and began to reverse, but continued through 2002 among 19- to 26-year-olds. There was little systematic change for several years among 19- to 26-year-olds, but more recently—since about 2008—their friends' disapproval

declined appreciably for all three levels of marijuana use. In 2013 all young adult age groups showed a further decline in disapproval of marijuana use; indeed, the declines for the older two age bands were large and statistically significant. For example, the percent of 23- to 26-year-olds and 27- to 30-year-olds saying that their close friends would disapprove of their trying marijuana fell by about 9 percentage points in that one year, possibly reflecting both cohort effects and secular trends. At this point only about half of each age group say that their close friends would disapprove of their trying marijuana, down from between 57-62% as recently as 2008. *Clearly peer norms among young adults, as well as teens, have become more accepting of marijuana use over the past few years.*

Close friends' disapproval of more frequent marijuana use also rose until the early 1990s among 18-year-olds, and then declined between 1992 and 1997. It declined through 1999 among 19- to 22-year-olds and continued to decline among 23- to 30-year-olds through 2003. Recent peak years of friends' disapproval of trying marijuana were 2006 for 12th graders, 2008 for 19- to 26-year-olds, and 2010 for 27- to 30-year-olds. In essence, peer norms have moved in a way consistent with the existence of some lasting cohort differences in these norms, as well as in use. A recent, more formal analysis of age, period, and cohort effects in disapproval came to the same conclusion.⁷³ In 2013 there were declines in friends' disapproval of occasional and regular use among 23- to 30-year-olds, particularly among the older two groups.

- There was a more gradual increase in peer disapproval of *trying* an **amphetamine** for all age groups (18–30) through 1991, followed by a small decline evident among 12th graders through 1997. Between 1997 and 2009—the last year for which data are available—levels of disapproval among 18- to 30-year-olds increased to some extent, though not dramatically. In 2009 (the last year we asked this question of young adults), disapproval levels were at 87% for all four age groups.
- Perceived peer norms for **LSD** were measured from 1980 through 2009 among the follow-up respondents. Through 1991, peer disapproval of trying LSD showed very little change in any of the age bands, but it fell some in the 1990s, especially among 18-year-olds and subsequently among 19- to 22-year-olds. These declines bottomed out in a staggered fashion, beginning with the 12th graders in 1997, which thereafter showed a seven-percentage-point increase in peer disapproval. There was a five-percentage-point increase among 19- to 22-year-olds (since 2000), and a three-percentage-point increase among 23- to 26-year-olds (since 2001)—again suggestive of a cohort effect in these norms. In 2009, the last year we asked, there was almost no difference

⁷³Keyes, K. M., Schulenberg, J. E., O'Malley, P. M., Johnston, L. D., Bachman, J. G., Li, G., & Hasin, D. (2011). The social norms of birth cohorts and adolescent marijuana use in the United States, 1976-2007. *Addiction*, 106(10), 1790-1800. [doi: 10.1111/j.1360-0443.2011.03485.x](https://doi.org/10.1111/j.1360-0443.2011.03485.x)

among the age groups, with 85% to 87% of respondents in each age group saying their friends would disapprove of their trying LSD.

- Perceived peer norms regarding *cocaine* use were measured from 1986 through 2009, after which such questions were dropped. From 1986 to 1994, self-reported cocaine use declined substantially as peer norms in all age bands shifted toward disapproval. For example, by 1994, 95% of the 19- to 22-year-olds thought their close friends would disapprove of their trying cocaine once or twice. After 1994, peer norms against use continued to strengthen a bit in the upper age bands, perhaps through generational replacement, but weakened slightly in the younger age bands, likely reflecting a new cohort effect. By 2009 (the last year we asked) there was little difference by age in peer norms against cocaine use, with 85% to 91% saying their friends would disapprove of their even trying it. By way of contrast, in 1986 that statistic ranged between 71% and 80% among 18- to 26-year-olds.
- Peer norms against *occasions of heavy drinking* on weekends (five or more drinks once or twice each weekend) among the three young adult age groups have tended to be weakest for the 19- to 22-year-old age group, where such behavior is most common, and strongest for the 27- to 30-year-old group. Since 2002, disapproval of such drinking has also been low for the 23- to 26-year-old group relative to the other two age bands. Among 12th graders, friends' attitudes had become somewhat more restrictive between 1981 and 1992 (and respondents' own occasions of heavy drinking declined during that interval), but attitudes have been fairly level since then with just a little upturn in the past couple of years. There was a similar upward trend in peer disapproval among the various young adult age bands that followed a staggered pattern, again likely reflecting a cohort effect in these norms. However, between 1997 and 2000 the 19- to 22-year-old age group themselves became somewhat less disapproving of occasions of heavy drinking on weekends; this was followed by a decline in perceived peer disapproval between 2001 and 2004 among 23- to 26-year-olds, and a decline from 2004 to 2009 among 27- to 30-year-olds. Despite some increases in peer disapproval over the years, this rather extreme form of drinking has the *least* restrictive perceived peer norms of all of the substance-using behaviors measured in MTF; yet, the majority still report peer disapproval. In 2013 the proportions saying that their friends would disapprove of such heavy weekend drinking was 49% to 53% for the three young adult age bands.
- Peer norms against *cigarette smoking* one or more packs per day have strengthened in staggered fashion among 18-year-olds and the young adult age groups in recent years. Between 1998 and 2008, the proportion saying that their close friends would disapprove of their smoking a pack or more of cigarettes per day rose from 69% to 83% among 18-year-olds and from 69% to 86% among 19- to 22-year-olds. The two older strata did not see a comparable change until 2006, when among 23- to 26-year-olds peer disapproval rose from 77% in 2005 to 88% in 2009, before leveling. The

change did not manifest itself among the 27- to 30-year-olds until 2010; their rates of peer disapproval of smoking, which for some years had the highest rates of disapproval for smoking among the four age groups, stayed fairly level after 2000, until there was a 4.4-percentage-point jump in 2010, followed by a leveling. This pattern again suggests some cohort effects in peer norms working their way up the age spectrum. In 2013, very large proportions across the age bands reported that their friends would disapprove of pack-a-day smoking, ranging from 84% among 18-year-olds to 90% among those 27–30 years old.

In the early years of MTF, peer disapproval of smoking a pack or more of cigarettes per day rose among 12th graders from 64% (1975) to 73% (1979). There was little further net change for 13 years through 1992, when friends' disapproval stood at 76%. Between 1992/1993 and 1997/1998, all age groups showed a decrease in perceived peer disapproval of smoking—this time consistent with a secular trend.

ADULTS' EXPOSURE TO DRUG USE THROUGH FRIENDS AND OTHERS (AGES 18–55)

Exposure to drug use is important because it provides both the modeling of the behavior by peers (possibly including direct encouragement to use) and immediate access. Exposure is measured by two sets of questions, each appearing on a (different) single questionnaire form. The first set asks the respondent to estimate what proportion of his or her friends use each drug, while the second set asks, “During the LAST TWELVE MONTHS how often were you around people who were using each of the following to get high or for ‘kicks’?” The same questions are asked of 12th graders, and their results are included here for comparison purposes in Tables 7-2 and 7-3 and Figures 7-1 through 7-24. (Questions about direct exposure to drug use were not included in the questionnaires for 35- to 55-year-olds due to the space limitations imposed by the use of a single questionnaire form at each of these ages.) We continue to deal with four-year age bands for the friends' use measures in order to increase the reliability of the estimates. Ages 35, 40, 45, 50, and 55 are included as one-year age bands, with both half samples from each of those cohorts being surveyed at those modal ages. Starting with age 35, each year has a larger number of cases than single years at the earlier ages because *all* respondents in a cohort at later ages complete the relevant questionnaire form, compared with only one sixth of those at younger ages. At the end of each table is a summary of the weighted number of cases upon which each annual estimate is based. (The actual numbers of cases are somewhat higher.)

Exposure to Drug Use (Ages 18 to 30)

- Relatively high proportions of young adults in all of these age bands have at least *some* friends who use *some illicit drug*; that proportion varies considerably with age, with older respondents reporting that fewer friends use

(Table 7-2). In 2013, illicit drug use by at least some friends is reported by 81% of 12th graders, 73% of 27- to 30-year-olds, 53% of 35-year-olds, 42% of 40-year-olds, 40% of 45-year-olds, 39% of 50-year-olds, and 36% of 55-year-olds. Clearly, the older the respondent, the less likely he or she is to report having friends who use any illicit drugs. The proportions who say that *most or all* of their friends use one or more of the illicit drugs are much lower: 27% for 12th graders, falling to 22% of 19- to 22-year-olds, 14% of 23- to 26-year-olds, 7% of 27- to 30-year-olds, and between 1.0% and 4.7% for the 35-, 40-, 45, 50-, and 55-year-olds—quite a dramatic difference across ages, and one that is consistent with the large differences in their own self-reported current use.

- With regard to *illicit drugs other than marijuana*, taken as a whole, considerably fewer report that *any* of their friends use compared to what is true for marijuana use (see below): 45% for 12th graders, 52% for 19- to 22-year-olds, 45% for 23- to 26-year-olds, 45% for 27- to 30-year-olds, and 19–25% for 35- to 55-year-olds. The proportions saying that *most or all* of their friends use illicit drugs other than marijuana in 2013 are 4%, 7%, 3%, and 0.7%, respectively, for the four youngest age bands, with 1.3% or fewer of respondents in the oldest age groups (35–55) reporting such high proportions of their friends using illicit drugs other than marijuana. Thus, relatively low proportions of these age groups appear to be deeply immersed in a drug culture involving illicit drugs beyond marijuana.
- With respect to individual illicit drugs, exposure among all of the age groups is greatest for *marijuana*, with the percentages saying they have any friends who use at 80% of the 18-year-olds, 76% of the 19- to 22-year-olds, 73% of the 23- to 26-year-olds, 69% of 27- to 30-year-olds, and so on down to 30% of 55-year-olds reporting that at least some of their friends use the drug. The next highest exposures are for *amphetamines* (26% among 12th graders, 38% among 19- to 22-year-olds, 27% among 23- to 26-year-olds, and 25% among 27- to 30-year-olds) and *narcotics other than heroin* (26% among 12th graders, 30% among 19- to 22-year-olds, 26% among 23- to 26-year-olds, and 30% among 27- to 30-year-olds) followed by *cocaine*, *hallucinogens other than LSD*, and *ecstasy (MDMA)*. Because of the dramatic increase in its use during the 1990s and early 2000s, ecstasy surpassed a number of the more traditional drugs, though its use declined sharply in subsequent years. (It should be noted that use of several illicit drugs was not asked of the age groups above 30 due to space limitations in the single questionnaire form used at each of those ages. See Table 7-2.)
- For the remaining illicit drugs, the proportion of young adults reporting that some friends use a given drug is 10% or higher in at least one of the four young age strata for the following drugs: *sedatives (barbiturates)* (14–19%), *LSD* (11–23%), *steroids* (10–17%), *tranquilizers* (9–13%), and *crack* (5–10%). See Table 7-2 for specifics.

- For most illicit drugs, the proportion of young adults having any friends who use them decreases with each higher age band, consistent with the age differentials in self-reported use. The steepest declines occur with *inhalants*—in 2013, 12% of 18-year-olds report that some friends use versus only 3% of 27- to 30-year-olds. (Inhalant use is not asked of the age groups above 30, precisely because of this sharp decline in use with age.) As reported in *Volume I*, the decline with age in inhalant use is actually well under way by 10th grade.
- For some years, *cocaine* showed significantly higher rates of active use among adults compared to 12th graders. That is no longer true, although there is rather little drop-off with age in early adulthood; consequently, there is not a great difference associated with age in having friends who use cocaine (18–20% for all four of the younger age groups). The 35-, 40-, 45-, 50-, and 55-year-olds are asked separately about cocaine powder and crack use; in 2013 far fewer report having friends who use cocaine powder—12% for age 35 and 4% to 7% for the four older groups.
- For *crack*, however, the story is different. Reported friends' use of crack now descends sharply with age, although this was not true in the mid-1980s, when measures of crack use were first included in the surveys. In 2013, 10% of 12th graders report having any friends using crack, versus 8% of 19- to 22-year-olds, 5% of 27- to 30-year-olds, and 1% to 3% of 35- to 55-year-olds.
- The proportion reporting in 2013 that they have any friends who take *heroin* also decreases sharply with age, from 9% among 12th graders to 5% among 27- to 30-year-olds. (Older respondents are not asked this question.)
- At present, the percentage reporting any friends who use *narcotics other than heroin* shows no consistent relationship with age: 26% of 18-year-olds and 23- to 26-year-olds, versus 30% of those 19–22 and 27–30 years old.
- In general, it appears that some respondents who report that their friends use *illicit drugs* are themselves not directly exposed to that use by their friends, judging by the differences in proportions saying they have some friends who use (Table 7-2) and the proportions who say they have been around people who were using during the prior year (Table 7-3 and Figure 7-1).
- With respect to *alcohol* use, the great majority of young adults have at least *some* friends who *get drunk at least once a week*, although this peaks in the early 20s and then drops off gradually with age: 70% of 12th graders, 78% of 19- to 22-year-olds, 83% of 23- to 26-year-olds, 74% of 27- to 30-year-olds, 64% of 35-year-olds, 55% of 40-year-olds, 50% of 45-year-olds, 45% of 50-year-olds, and 35% of 55-year-olds. Given the potential serious consequences of this behavior, these rates are disturbingly high across a wide age range. The proportions who say *most or all* of their friends get drunk once a week differ more substantially by age: 21% of 12th graders and 28% of 19- to 22-year-

olds, declining sharply to 13% of 27- to 30-year-olds and 2% of 55-year-olds. Note in particular how high these rates are among the high school and college-age populations. In terms of having any direct exposure during the prior year to people who were drinking alcohol “to get high or for ‘kicks,’” having some such exposure is almost universal in the three 4-year age groups of young adults: 83%, 89%, and 89%, respectively, as well as among 18-year-olds (85%) (see Table 7-3 and Figure 7-23).

- From ages 18 through 30, about four fifths of respondents (72–82%) have at least a few friends who *smoke cigarettes*, with considerable falloff by age 35. In fact, 11% of the 12th graders and 11% of 19- to 22-year-olds state that *most or all* of their friends smoke. Above those ages, the proportions decline to 7% of 27- to 30-year-olds and 2–5% for those 35 years of age and older. This increase in the segregation of smokers from nonsmokers likely reflects the stratification of young people after high school as a function of educational attainment, which is highly correlated with cigarette smoking. Also, it can be seen in Table 7-2 that there was much less age-related difference in the late 1980s, suggesting that the sharp rise in smoking among high school students during much of the 1990s, followed by a sharp decline in the years since, accentuated the age differentials, and that those differentials remain, reflecting lasting cohort effects.

Trends in Exposure to Drug Use (Ages 18 to 55)

Tables 7-2 and 7-3 also provide *trend* data on the proportions of respondents’ friends using drugs and the proportion of respondents directly exposed to drug use by others. Both of these measures of exposure to use will be discussed in this section. Once again, trends are available for 19- to 22-year-olds since 1980, for 23- to 26-year-olds since 1984, and for 27- to 30-year-olds since 1988. Data for 35-, 40-, 45-, 50-, and 55-year-olds are available on friends’ use since 1994, 1998, 2003, 2008, and 2013, respectively. (Questions about frequency of being around drug users were not included in the questionnaires administered to respondents age 35 and older, so those age bands are not included in Table 7-3 or Figures 7-1 through 7-24. However, they were asked about the proportions of their friends using.) Twelfth-grade data have also been included in these tables for comparison purposes.

Figures 7-1 through 7-24 provide graphic presentation of trends in exposure to use.

- An examination of Table 7-3 and Figures 7-1 through 7-6 shows that exposure to illicit drug use (in the 12 months preceding the survey) declines at each higher age band for *any illicit drug*, *marijuana*, and *any illicit drug other than marijuana*, as well as for nearly all of the specific illicit drugs. In general, these differences replicate across different historical periods, with the exception of *cocaine*, which did not show a decline in exposure with increasing age until after 1996. (Prior to that it showed an increase with increasing age.) These declines reflect age effects (changes with age observed

across multiple cohorts) in both exposure to use and in personal use of most drugs.

- Until 1992, young adults' trends in exposure to use tended to parallel those observed for 12th graders. From 1980 to 1992, that meant a decreasing number of respondents were exposed to **any illicit drug** use (Table 7-3 and Figure 7-1) or reported any such use in their own friendship circle (Table 7-2). After 1992, however, an important *divergence* in trends emerged: 12th graders showed a substantial increase in both friends' use and exposure to use (as well as self-reported use); 19- to 22-year-olds showed a similar rise, but lagged by a few years; 23- to 26-year-olds subsequently showed some rise; while the 27- to 30-year-old age band did not show a rise until 2002. As discussed in earlier chapters, this pattern no doubt reflects the emergence of lasting cohort differences that emerged in secondary school and, driven by generational replacement, continued up the age spectrum as the secondary school students grew older. The age differentials expanded in the 1990s during the relapse phase in the drug epidemic; first observed among the 12th graders, the increases in use then occurred on a staggered basis. The age differentials have diminished considerably during the 2000s as use declined some among the younger age groups but rose among the older ones as the cohort effect worked its way up the age spectrum (see Figure 7-1).
- **Marijuana** showed a very similar pattern of change. It is particularly noteworthy that, while 34% of 19- to 22-year-olds in 1980 said *most or all* of their friends used marijuana, only 8% said the same in 1991. Clearly, the number of friendship groupings in which marijuana use was widespread dropped dramatically over that earlier interval. This measure of friends' use more than doubled to 19% by 1999 during the relapse phase in the larger epidemic, where it remained for a couple of years before falling to 12% by 2008 before increasing to 20% by 2013. Self-reported use (Figure 5-3a) and friends' use both increased significantly among 18-year-olds in 2008, which we interpreted as a turnaround in the marijuana situation. Since 2006, the other adult age strata also have shown some increase in the proportion reporting some friends using marijuana (Figure 7-5 and Table 7-2).
- The proportion of respondents reporting having any friends who use **any illicit drugs other than marijuana** began to decline after 1982 in the two younger age groups spanning 18–22 (for whom we had data at that time; see Figure 7-3). By 1991/1992 there had been a considerable drop in all four age groups (spanning 18–30). This drop appears to be due particularly to decreases in friends' use of **cocaine** and **amphetamines**, although there were decreases for **sedatives (barbiturates)** and **tranquilizers** as well. The levels then began to rise among the 18- to 22-year-olds in the early 1990s, while at the same time declining further for the 23- to 30-year-olds, reflecting cohort effects, opening up a large age-related difference in friends' use in the 1990s and into the early 2000s. The 23- to 26-year-olds showed a later increase in friends' use and the 27- to 30-year-olds showed a still later increase. After 2001 there was some

decline in reported friends' use in the two youngest age strata while friends' use continued to climb in the older two strata. The net effect was to narrow the age differences among the young adult strata considerably. More recently, the gap among the four age bands has narrowed for the proportions saying that they have any friends who use some illicit drug other than marijuana. Since 2004, there has been little change among 18- to 22-year-olds while since 2008 there has been an increase among 23- to 26-year-olds and until 2012 a fairly sharp increase among 27- to 30-year-olds. (the four groups showed mixed results in 2013, however.) Among those ages 35 and older, considerably fewer report having any friends who use (between 19% and 25%). See Table 7-2 for the specifics.

- Between 1986 and about 1992, all four age groups showed a considerable drop in the proportion of respondents with friends who used *cocaine* (Figure 7-11). (Self-reported use declined sharply in the same period.) After that decline, the rates of friends' use peaked in 1998 among 18-year-olds, in 2004 among 19- to 22-year-olds, and in 2006 among 23- to 30-year-olds, suggesting another cohort effect. In 2007 friends' use began a fair decline among 18-year-olds followed by a similar decline among 19- to 22-year-olds in 2009, a decline among 23- to 26-year-olds beginning in 2012, and a decline among 27- to 30-year-olds in 2013.. The story for crack has been fairly similar (Table 7-2).
- There were substantial increases between the early 1990s and about 2000 in the proportion of 18-year-olds and 19- to 22-year-olds reporting that they have friends using *narcotics other than heroin*, and smaller increases among 23- to 30-year-olds, resulting in some considerable age-related differences. After 2002, the proportions of 18-year-olds and 19- to 22-year-olds declined some, while the 23- to 30-year-olds continued to increase in a classic cohort-effect pattern of change, thus narrowing the age differences by 2009. There was a wording change in 2010 that served to increase the rates considerably for all age groups (Figure 7-15). Since then, all except the oldest age band have shown a decline in exposure to use of these drugs.
- The proportions saying that any of their friends use *ecstasy (MDMA)* increased sharply in all age groups between 1992 and 2002, though in a staggered fashion (Table 7-2). Twelfth graders showed the first sharp increase beginning after 1992, 19- to 22-year-olds after 1994, 23- to 26-year-olds after 1996 and 27- to 30-year-olds after 1997. These sharp increases ended among 12th graders in 2001 and among 19- to 30-year-olds a year later. Since those peak levels, the proportions saying that they had any friends using ecstasy have generally declined, corresponding with a decline in self-reported use. In all four age groups, 19–27% now report that any of their friends use ecstasy. The staggered nature of the increases suggests a cohort effect at work, but the simultaneous decline strongly suggests a secular trend, likely due to the heavy media coverage of adverse consequences associated with ecstasy use during that period.

- For all four of the youngest age groups (spanning ages 18–30), the proportions saying that they were often exposed to others drinking *alcohol* declined modestly between 1987 and 1992 (Figure 7-24). The next decade saw rather little change in the four youngest age bands. Since 2002, exposure among the 18- to 22-year-olds has declined slightly, while rising among 23- to 30-year-olds, resulting in rather little differences among the age groups in recent years, but with the two younger groups now slightly higher than the two older groups.
- The age groups above age 30 have consistently been much less likely to report that any of their friends *get drunk* at least once a week, compared with those ages 18 to 26 (Table 7-2). These proportions increased starting at different times: after 1998 among 35-year-olds, after 2004 among 40-year-olds, and after 2005 among 45-year-olds, suggesting somewhat stable cohort differences. The net effect has been to reduce the differences separating those in their 20s from those in later decades in terms of the proportion having any friends who get drunk at least once a week. The rates in 2013 for the four youngest age strata are very high—between 70% among 12th graders and 83% among 23- to 26-year-olds. Among the older age band there has been some increase over the past decade or so in the proportion saying that any of their friends get drunk at least once a week.
- Among 12th graders, the proportion who said *most or all of their friends smoked cigarettes* declined appreciably between 1975 and 1981, the same period in which self-reported use declined. After that, neither measure showed much change until about 1992. Thereafter, substantial increases in both measures occurred. By 1997, fully one third (34%) of 12th graders reported that most or all of their friends smoked cigarettes (up from 21% in 1992); since then, that statistic declined (along with self-reported use) to 14% in 2008, and then a little further to 11% in 2013. Among 19- to 22-year-olds, a decline in friends' use occurred between 1980 (or possibly earlier) and 1985, followed by a leveling through 1994. The percentage saying most or all of their friends smoke increased from 22% in 1994 to 29% in 2000, before beginning to decline, reaching 11% in 2013. Among 23- to 26-year-olds, a downturn was evident between at least 1984 (the first year for which data are available) and 1988, and then reported friends' use leveled (14% in 2013). After 2002, some slight increases occurred, but then a reversal occurred from 2004 to 2012. These staggered changes, until about 1998, illustrate that cohort effects were moving up the age spectrum. Since 1998 (or the earliest year available for the age bands above age 30), the proportion saying that any of their friends smoked cigarettes showed some decline among those above age 30, but little or no change among 23- to 30-year-olds (which contains some of the heavier smoking senior classes of the mid-1990s) until about 2006. All of the age strata show rates in 2013 that are somewhat below where they were in 2008, following a gradual decline in recent years.

Nearly all of these changes in exposure to drug use parallel changes in self-reported use by these age groups. This pattern reinforces the validity of self-report data, because there would presumably be less motivation to distort answers about the proportion of an unnamed set of friends who use a drug than about one's own use. The systematic nature of the patterns of change across age strata (whether in terms of parallel changes consistent with a secular trend, or systematically staggered ones consistent with a cohort-related trend) is also supportive of the data validity.

PERCEIVED AVAILABILITY OF DRUGS AMONG ADULTS (AGES 18–55)

Adults participating in the follow-up surveys receive questions identical to those asked of 12th graders regarding how difficult they think it would be to get each of the various drugs if they wanted them. The questions are contained in only one of the six questionnaire forms used through modal age 30. Data for the young adult follow-up samples, which are grouped into the same four-year age bands used above (19–22, 23–26, 27–30), are presented in Table 7-4, along with data for 12th graders and 35-, 40-, 45-, 50-, and 55-year-olds. Sample sizes are presented at the bottom of the table. The availability question is not asked for all drugs in the adult samples, as may be seen in Table 7-4.

Perceived Availability

Much like 12th graders, substantial proportions of the American adult population have access to various illicit drugs. (We do not ask about access to alcohol and cigarettes because we assume these are readily available to all adults.)

- **Marijuana** is the most available illicit drug in 2013, with 84–87% of the young adult age groups (19–30) group saying it would be “fairly easy” or “very easy” to get. Perceived access decreases somewhat with age after age 30; but even at age 55, more than two thirds of respondents (69%) say they could get marijuana fairly or very easily.
- Though less available than marijuana, **amphetamines** are still fairly available, with 54–59% of young adults and 33–41% of those 35, 40, 45, 50, and 55 years old saying they would be easy to get.
- **Cocaine** is reported as readily available in 2013 by a significant proportion of young adults, with 28–37% saying it would be easy to get. **Powdered cocaine** availability is now highest among those ages 23 and above. **Crack** is available to smaller proportions than powdered cocaine; and again is most available to those ages 23 and above (24–32%). Interestingly, perceived availability for both forms of cocaine tends to rise with age between 19 to 22 years old and 27 to 30 years old.
- About one fifth (22%) of 12th graders say that they could get **heroin** fairly or very easily, though far fewer report having used heroin. Reported availability

is about the same for the 23- to 30-year-olds but is a bit lower for the 19- to 22-year olds (16%) in 2013 at least. (The question is not asked of respondents above age 30.)

- More than half of all young adults (54–64%) in 2013 say that they could get *narcotics other than heroin* fairly easily, as do 47% of 12th graders. (The question is not asked of respondents above age 30.)
- *Sedatives (barbiturates)* are a bit less available than amphetamines to these age groups, with little variation across age up to age 30 (32–45%). (The question is not asked of respondents above age 30.)
- *Tranquilizers* are reported as available in 2013 by considerably fewer respondents (17–28%), which historically was not always the case. (The question is not asked of respondents above age 30.)
- *Ecstasy (MDMA)* is seen as readily available to 29–36% of young adults and 12th graders in 2013. (The question is not asked of respondents above age 30.)
- *Hallucinogens other than LSD* are reported as available by 37% of 12th graders and by slightly fewer (32–34%) of the three young adult strata. (The question is not asked of respondents above age 30.)
- *LSD* is the least available drug, declining by age group from 25% among 12th graders to 17% among 55-year-olds.
- *Crystal methamphetamine (ice)* is perceived to be available by between 15% and 22% in 2013, with availability highest among 27- to 30-year-olds. (The question is not asked of respondents above age 30.)
- *Anabolic steroids* are seen as available by about equal proportions of young adults through to the 27- to 30-year-olds (between 21% and 24%). (The question is not asked of respondents above age 30.) Twelfth graders report a slightly higher availability (29%).

Trends in Perceived Availability

- *Marijuana* has been almost universally available to the adolescent and young adult age groups throughout the historical periods covered by the data. Overall, perceived availability has risen in the five older age groups on whom trend data are available during the 2000s—that is, from 27- through 50-year-olds—but has simultaneously declined among 12th graders and 19- to 23-year olds. The 23- to 26-year-olds have shown little change in the 2000s. As a result there is now less variability by age in availability of marijuana. From the peak year in 1979, perceived availability decreased slightly through 1991 among 12th graders and decreased slightly more from 1980 through 1991 among 19- to 22-year-olds. After the late 1990s, the trends in availability

across the age bands had generally been quite parallel, suggesting secular trends in prevailing conditions that affected availability. Perceived availability is now somewhat higher for the younger age groups (81–87% for 12th graders through 27- to 30-year-olds versus 69–80% for those ages 35 to 55).

- Historic highs in perceived availability of *cocaine* occurred in the 1980s among all three young adult age strata (ages 19–30), reaching highest proportions in 1988 and 1989, at which time the older young adult age strata had higher availability than the younger ages. (From a policy perspective, it is worth noting that in 1987 the perceived availability of cocaine *increased* while use actually dropped sharply.) In the early 1990s, all four groups reported decreased availability by 4–7 percentage points—quite parallel to the drop in numbers of those who had friends who were users and to the decline in personal use. Until about 2000, there was some falloff in perceived cocaine availability in all age strata through age 30—particularly among those ages 23 through 30—and an increasing convergence. Since about 2007 all four age strata have shown considerable declines in reported cocaine availability and these declines continued into 2013 among the three young adult strata. (The question is not asked of respondents above age 30.)
- *Crack* availability peaked in 1988–1989 for all age groups (it was first assessed in 1987) and declined through 1992, with little further change until 1995. Since 1995, crack availability has declined substantially in all of the lower four age strata (ages 18–30). Data on 35-, 40-, 45-, and 50-year-olds are available for shorter intervals, but also show appreciable declines from initial levels.
- In 2013, between 17% and 26% of each age group said they could get *LSD* fairly easily, which contrasts quite dramatically with the mid-1990s, when over 50% of those in the younger age strata said they could get it. Across the decades measured, the trends in LSD availability among young adults have had some parallels to those among 12th graders. For 12th graders, there was a drop of about 10 percentage points in the mid-1970s, and a later drop from 1980 to 1986. The latter drop, at least, was paralleled in the data from 19- to 22-year-olds. After 1986, LSD availability increased considerably in all age bands, reaching its peak levels by 1995 during the relapse phase of the illicit drug epidemic. At that time a considerable age-related difference developed, with availability lower in the older age groups. Since 1995, availability has fallen substantially in all age bands but particularly in the youngest two age bands, narrowing the differences among the age groups. Indeed, the drop-off in availability of LSD to 12th graders and 19- to 22-year-olds was quite sharp in 2002, possibly contributing to the steep decline in use that year. In 2012 all age bands except the 18-year-olds showed some further decline in the availability of LSD, but in 2013 the trends were mixed, although increases were observed in the three young adult age bands as well as among 45-year-olds. Availability among 50-year-olds is the lowest at 17%, and 55-year-olds are not asked the question.

- Since 2001 the general pattern regarding the availability of *hallucinogens other than LSD* has been one of stability. Levels of availability have been more differentiated by age than in prior decades, though these differences have diminished in recent years. (This question is not asked of respondents over age 30.) Generally, the lower the age group, the higher the reported availability, though there has been a considerable convergence. In the early 1980s, there was a fair decline among all age groups in the availability of hallucinogens other than LSD; there was little additional change through 1992. From 1992 to 1995, the three youngest age groups all showed an increase in availability, with 12th graders showing the largest increase. From 1996 to 2000, availability was fairly steady. All age groups showed substantial increases in 2001, undoubtedly due to the changed question wording which added “shrooms,” among other substances to the examples of hallucinogens. (“Shrooms” refer to psilocybin mushrooms.) It appears that the inclusion of “shrooms” introduced a greater variability with age in the availability of hallucinogens other than LSD taken as a class. Availability is now only slightly higher among the 18-year-olds than among young adults.
- *Ecstasy (MDMA)* questions were first introduced in MTF surveys in 1989 and 1990 (and are not asked of those over age 30). Availability rose very substantially in all of these age groups during the 1990s and early 2000s. Among 12th graders, reported availability nearly tripled, from 22% in 1989 to 62% in 2001—the peak year of use for 12th graders. All four age groupings showed sharp increases in 2000 and 2001, with the older age groups continuing to increase through 2002—their peak year. The availability of ecstasy showed considerable declines among the two youngest strata after 2002 and among the 23- to 30-year-olds since about 2008. Reported availability of ecstasy varied little by age in 2013, ranging between 29% and 36% among 18-year-olds and the three young adult strata covered.
- All age groups have shown some gradual, modest decline in *heroin* availability since 1997 or 1998, during which interval there has been rather little variability in heroin availability across the 18-to-30 age range. (The question is not asked of respondents over age 30.) Heroin availability varied within a fairly narrow range from 1980 to 1985, then increased in all age groups through 1990. For the younger ages (18–22) heroin availability rose further through 1995 while in the older two age groups it increased some later in the 1990s. It is clear that heroin was much more available to all of these age groups in the 1990s than it was in the 1980s. This increase in the availability—and purity—of heroin most likely led to the emergence of non-injection forms of heroin use observed during this period. In recent years heroin availability has generally declined modestly among young adults ages 18 to 22 and 27 to 30, but has held fairly steady among 23- to 26-year-olds.
- The availability of *narcotics other than heroin* rose slowly among all age groups from 1980 until recent years, with the exception of a period of considerable stability from 1989 through 1994. (Respondents over age 30 are

not asked this question.) After 1994, the modest increase in availability was accompanied by steadily rising use. Recent years showed a very slight falloff in availability among all age strata except the 27- to 30-year-olds, who continued to show an increase. Note, however, that reported availability jumped in 2010, when new drugs, including Vicodin and OxyContin, were added to the list of examples in the question. For the most part, there has not been a consistent difference by age in the availability of narcotics other than heroin among those ages 18 to 30; the predominant trend has been one of increasing availability over a long period of time in the 23- to 30-year-old segment. The addition of newer drugs, like OxyContin and Vicodin, to the list of examples resulted in some further increase, which suggests that availability climbed considerably more over the past decade or so than the data based on the original question had suggested. Although the proportions of young adults saying that narcotic drugs other than heroin are readily available have been rising since 2008 and continued to rise in 2013, they fell significantly among 12th graders (down 3.9 percentage points to 46.5%) in 2013.

- In general, the age groups above age 30 have reported somewhat lower availability of *amphetamines* than the younger strata, but not dramatically lower. These differential rates of reported availability across the age groups emerged after 1992, when prevalence of use began to rise among 12th graders. In 1982, availability peaked for both 12th graders and 19- to 22-year-olds, after which it fell until 1991, by 14–15 percentage points. Among 23- to 26-year-olds, there was a decline of 14 percentage points between 1984 (when data were first available) and 2005. For 27- to 30-year-olds, reported availability decreased by nine percentage points between 1988 (the first measurement point) and 2005. Decreases also occurred among 35-year-olds in the 2000s but some reversal has been evident in recent years. In 2011 all age strata from age 18 through age 35 showed an increase in perceived availability for amphetamines, statistically significant for those 19–22 and 23–26 years old, followed by little systematic change since. It should be noted that the examples of amphetamines used in the question text were revised in 2011 to include Adderall and Ritalin, while pep pills and bennies were eliminated as examples. Therefore, the sharp rise in reported *availability* of amphetamines in 2011 among young adults may be nothing more than a method artifact resulting from the revision of the examples provided.
- By way of contrast, *crystal methamphetamine* or “*ice*” exhibited an increase in availability in the 1990s, rising for all four age strata from 1991 to 1998/1999, before stabilizing with similar rates of availability from ages 18 to 30. (This question was not asked of those over 30.) All four strata have shown some decline in recent years, starting with the youngest three age strata after 2006 and the 27- to 30-year-olds after 2008. Availability is now lowest for the youngest three age bands (the 23- to 26-year-olds showed a significant 7 percentage point decline in 2013)—a reversal of the situation in the early 1990s—but the differences among the age groups have largely disappeared over the past two years.

- ***Sedatives (barbiturates)*** exhibited a long-term decline in availability over more than two decades, from about 1981 or 1982 through 2003 in the two younger groups—a 20-percentage-point drop among 12th graders and a 23-percentage-point drop among 19- to 22-year-olds. All groups showed increased availability in 2004—no doubt due primarily to a change in the question wording. There has been little further systematic change since then, except for a significant decline of 9 percentage points among 12th graders since 2010 and of 8 percentage points among the 19- to 22-year-olds. Availability is considerably higher in the 27- to 30-year-old group (45%) than in the 18-year-old group (28%).
- ***Tranquilizer*** availability has declined long-term by about four fifths among 12th graders, from 72% in 1975 to 15% in 2013. Since 1980, when data were first collected for 19- to 22-year-olds, tranquilizer availability declined by more than three fourths (from 67% in 1980 to 17% in 2013), such that previous differences in availability between these two groups were eliminated by 1992. The older young adult age groups have also shown a considerable decline in the availability of tranquilizers through 2013 but they still generally have a higher level of availability for tranquilizers than the younger two age bands. For the most part, trend lines for the different age groups have been quite parallel, as has been true for sedatives (barbiturates). Indeed, tranquilizers have shown the most consistent pattern of change in perceived availability since MTF began.
- Data on ***steroid*** availability were first gathered in 1990. There has been some decline in availability in all age groups since about 2000, including a sharper rate of decline in the youngest three age strata after 2007. (This question was not asked of those over 30.) While younger respondents used to report higher levels of availability than those in the older strata, by 2013 there was not much difference among them (from 21% to 29%). In fact, for two of the three young adult age groups, availability of anabolic steroids is now at the lowest point since data were first gathered (and it is at the second lowest level for the third group). Eighteen year olds report the highest level of perceived availability in 2013, due to a significant increase of 3.5 percentage points that year bringing them to 29%.

TABLE 7-1
Trends in Proportions of Respondents Reporting Their Close Friends Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How do you think your close friends feel (or would feel) about you. . .	Age Group	Percentage saying friends disapprove ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Trying marijuana once or twice	18	42.6	46.4	50.3	52.0	54.1	54.7	56.7	58.0	62.9	63.7	70.3	69.7	73.1	66.6	62.7	58.1	55.8
	19–22	41.0	40.6	46.9	47.1	51.6	54.5	55.2	54.7	58.7	63.0	63.6	64.7	64.7	63.4	63.7	58.5	64.3
	23–26	—	—	—	—	47.7	47.0	49.1	53.9	58.2	62.6	61.3	64.5	65.6	65.5	63.2	63.8	61.2
	27–30	—	—	—	—	—	—	—	—	58.6	58.7	61.4	64.6	63.5	64.4	66.3	66.1	65.8
Smoking marijuana occasionally	18	50.6	55.9	57.4	59.9	62.9	64.2	64.4	67.0	72.1	71.1	76.4	75.8	79.2	73.8	69.1	65.4	63.1
	19–22	50.9	49.2	54.0	57.9	59.4	64.6	64.4	65.1	69.8	71.5	74.1	73.9	74.3	73.1	73.0	66.6	71.3
	23–26	—	—	—	—	54.3	56.4	57.1	63.1	68.1	73.2	71.8	72.5	75.3	73.5	72.2	70.7	70.8
	27–30	—	—	—	—	—	—	—	—	67.8	69.4	71.9	73.7	76.0	75.1	76.4	73.8	75.6
Smoking marijuana regularly	18	72.0	75.0	74.7	77.6	79.2	81.0	82.3	82.9	85.5	84.9	86.7	85.9	88.0	83.5	80.6	78.9	76.1
	19–22	70.3	75.2	75.7	79.5	80.0	82.7	83.5	84.8	86.9	87.5	89.1	88.4	89.1	87.6	85.9	83.9	84.5
	23–26	—	—	—	—	77.8	78.4	80.9	82.0	85.8	89.2	88.1	87.9	90.3	89.1	88.8	84.9	89.5
	27–30	—	—	—	—	—	—	—	—	85.4	86.0	88.4	89.2	88.7	88.2	88.9	89.7	89.6
Trying LSD once or twice	18	87.4	86.5	87.8	87.8	87.6	88.6	89.0	87.9	89.5	88.4	87.9	87.9	87.3	83.5	83.4	82.6	80.8
	19–22	87.4	90.5	88.0	89.3	89.3	91.1	90.5	91.8	90.8	91.2	89.1	89.9	87.2	87.7	87.9	84.6	85.3
	23–26	—	—	—	—	87.4	90.8	88.6	89.8	88.9	91.0	90.1	92.4	88.9	87.7	86.3	85.3	88.5
	27–30	—	—	—	—	—	—	—	—	88.8	89.7	92.3	91.1	91.4	89.9	91.2	89.7	89.3
Trying cocaine once or twice	18	—	—	—	—	—	—	79.6	83.9	88.1	88.9	90.5	91.8	92.2	91.1	91.4	91.1	89.2
	19–22	—	—	—	—	—	—	76.4	—	84.8	87.7	89.2	92.3	91.9	92.4	94.7	91.7	91.5
	23–26	—	—	—	—	—	—	70.8	—	81.4	84.5	84.1	86.7	87.4	87.7	87.9	90.4	90.0
	27–30	—	—	—	—	—	—	—	—	81.8	81.1	83.7	83.5	84.4	86.1	87.8	87.5	88.7
Taking cocaine occasionally	18	—	—	—	—	—	—	87.3	89.7	92.1	92.1	94.2	94.7	94.4	93.7	93.9	93.8	92.5
	19–22	—	—	—	—	—	—	84.9	—	91.0	93.8	94.2	95.6	95.9	95.6	97.5	95.6	95.7
	23–26	—	—	—	—	—	—	81.7	—	88.2	91.5	92.4	94.1	93.8	93.5	94.3	94.6	95.4
	27–30	—	—	—	—	—	—	—	—	87.7	89.5	90.0	92.2	92.3	92.8	94.6	94.1	94.6
Trying an amphetamine once or twice	18	78.9	74.4	75.7	76.8	77.0	77.0	79.4	80.0	82.3	84.1	84.2	85.3	85.7	83.2	84.5	81.9	80.6
	19–22	75.8	76.7	75.3	74.3	77.0	79.7	81.5	81.3	83.0	83.5	84.5	86.5	83.8	85.0	87.2	83.1	86.0
	23–26	—	—	—	—	78.4	79.1	76.7	81.7	83.0	85.6	84.3	85.0	83.6	84.2	84.7	87.6	86.5
	27–30	—	—	—	—	—	—	—	—	82.7	84.1	84.9	84.6	84.7	84.1	85.9	85.5	85.6
Taking one or two drinks nearly every day	18	70.5	69.5	71.9	71.7	73.6	75.4	75.9	71.8	74.9	76.4	79.0	76.6	77.9	76.8	75.8	72.6	72.9
	19–22	71.9	72.1	68.6	73.5	71.6	72.2	72.7	70.2	73.9	77.1	73.3	73.7	74.0	71.2	73.0	68.3	68.9
	23–26	—	—	—	—	63.6	66.8	67.7	68.3	69.2	70.8	72.7	72.5	72.1	67.6	71.5	68.2	72.8
	27–30	—	—	—	—	—	—	—	—	71.0	68.0	70.4	71.9	68.8	73.2	70.9	68.8	65.7
Taking four or five drinks nearly every day	18	87.9	86.4	86.6	86.0	86.1	88.2	87.4	85.6	87.1	87.2	88.2	86.4	87.4	87.2	85.2	84.1	82.6
	19–22	93.7	91.7	89.9	91.9	91.7	92.5	91.5	90.8	90.4	92.5	89.9	91.7	92.6	89.6	90.1	88.8	88.1
	23–26	—	—	—	—	90.8	90.2	92.5	92.8	93.7	92.1	92.4	92.4	91.1	93.1	92.1	92.2	92.6
	27–30	—	—	—	—	—	—	—	—	92.8	92.0	92.9	92.7	92.7	93.9	94.0	92.9	91.9
Having five or more drinks once or twice each weekend	18	50.6	50.3	51.2	50.6	51.3	55.9	54.9	52.4	54.0	56.4	59.0	58.1	60.8	58.5	59.1	58.0	57.8
	19–22	53.5	51.7	51.7	53.3	50.8	53.3	47.0	49.4	50.5	56.8	53.1	51.4	53.6	51.9	54.4	55.5	52.1
	23–26	—	—	—	—	53.8	57.3	61.0	57.2	58.8	57.5	55.1	56.8	58.4	57.6	61.4	58.9	58.4
	27–30	—	—	—	—	—	—	—	—	61.9	65.1	66.3	68.2	66.2	66.7	63.7	64.6	61.6
Smoking one or more packs of cigarettes per day	18	74.4	73.8	70.3	72.2	73.9	73.7	76.2	74.2	76.4	74.4	75.3	74.0	76.2	71.8	72.4	69.2	69.3
	19–22	75.6	75.1	75.4	78.5	76.2	79.7	77.7	78.6	80.2	78.4	77.5	78.3	79.0	76.0	73.8	70.9	73.9
	23–26	—	—	—	—	73.9	77.3	80.3	80.5	79.5	80.5	78.5	83.3	82.3	77.4	80.1	78.8	78.3
	27–30	—	—	—	—	—	—	—	—	81.2	80.9	82.9	84.5	83.1	86.8	82.5	83.4	81.9
Approximate Weighted N =	18	2,766	3,120	3,024	2,722	2,721	2,688	2,639	2,815	2,778	2,400	2,184	2,160	2,229	2,220	2,149	2,177	2,030
	19–22	569	597	580	577	582	556	577	595	584	555	559	537	520	510	470	480	471
	23–26	—	—	—	—	510	548	549	540	510	513	516	516	507	481	463	445	436
	27–30	—	—	—	—	—	—	—	—	483	518	479	480	451	451	457	439	439

(Table continued on next page.)

TABLE 7-1 (cont.)
Trends in Proportions of Respondents Reporting Their Close Friends Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How do you think your close friends feel (or would feel) about you. . .	Age Group	Percentage saying friends disapprove ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Trying marijuana once or twice	18	53.0	53.8	55.1	58.1	57.6	54.1	58.4	59.5	60.9	62.3	60.4	60.8	61.4	54.9	53.0	52.9	51.2	-1.8
	19–22	58.4	57.0	56.5	56.0	54.2	53.4	56.5	61.0	57.9	60.5	58.4	62.4	57.0	57.4	52.4	54.6	52.2	-2.4
	23–26	59.3	66.5	62.6	64.6	55.2	53.8	51.4	57.7	55.9	60.7	55.8	62.1	57.1	58.0	55.5	59.3	50.2	-9.1 s
	27–30	65.0	65.4	61.8	63.9	64.9	67.1	61.9	67.2	61.2	64.1	58.2	57.1	55.6	60.5	57.1	56.8	48.2	-8.5 s
Smoking marijuana occasionally	18	59.9	60.4	61.6	63.9	64.3	60.3	64.2	65.0	67.6	68.1	65.8	66.3	68.5	61.8	59.4	59.5	57.6	-1.9
	19–22	65.1	65.1	64.6	61.8	61.0	62.6	63.3	70.1	67.2	68.8	70.6	67.5	65.9	67.1	60.6	60.9	59.0	-1.9
	23–26	68.5	73.6	70.2	70.9	63.9	64.5	61.6	63.5	65.5	71.3	63.8	70.1	66.8	63.4	64.7	69.3	60.9	-8.3 s
	27–30	72.4	74.9	74.5	75.0	74.2	72.9	71.4	76.9	70.4	74.9	66.4	67.0	64.6	68.3	64.9	67.1	59.7	-7.4
Smoking marijuana regularly	18	74.1	74.7	74.5	76.1	77.8	75.3	77.0	77.3	79.5	79.8	78.3	78.0	79.1	73.8	73.3	72.7	71.2	-1.5
	19–22	83.3	81.1	78.2	78.5	80.0	80.5	79.1	84.4	82.2	84.1	83.7	81.4	81.9	81.1	76.3	74.5	75.2	+0.7
	23–26	85.6	87.1	86.8	86.9	83.7	82.8	80.0	79.2	82.7	83.7	81.9	87.1	81.9	83.5	82.7	83.8	77.7	-6.1 s
	27–30	87.8	90.8	89.2	91.6	90.1	87.9	87.2	88.0	87.7	88.2	84.3	84.5	83.4	87.5	83.4	86.1	80.8	-5.3
Trying LSD once or twice ^b	18	79.3	81.7	83.2	84.7	85.5	84.9	87.5	87.3	88.4	89.5	88.4	86.3	87.2	84.5	85.6	85.0	84.9	0.0
	19–22	83.6	81.7	82.0	82.1	85.2	86.9	86.9	88.6	90.5	90.4	90.0	90.0	87.1	—	—	—	—	—
	23–26	85.4	87.6	84.5	85.3	82.8	83.6	79.3	82.4	85.6	89.3	90.4	88.4	88.3	—	—	—	—	—
	27–30	88.5	88.7	88.4	85.6	87.4	86.3	87.1	87.7	86.9	88.5	83.5	85.3	84.6	—	—	—	—	—
Trying cocaine once or twice ^b	18	87.3	88.8	88.7	90.2	89.3	89.1	91.2	87.9	89.0	88.7	89.6	88.7	90.2	89.7	89.7	89.2	89.2	0.0
	19–22	91.8	90.0	91.2	89.4	89.1	91.7	90.6	90.3	90.3	91.2	93.3	90.2	91.2	—	—	—	—	—
	23–26	91.1	92.0	89.6	90.5	88.0	88.5	83.6	84.2	84.6	88.7	91.7	91.0	91.0	—	—	—	—	—
	27–30	89.4	89.3	90.5	90.4	89.3	88.8	89.9	91.8	89.5	92.0	86.4	88.0	84.5	—	—	—	—	—
Taking cocaine occasionally ^b	18	90.8	92.2	91.8	92.8	92.2	92.2	93.0	91.0	92.3	92.4	93.1	92.0	92.7	91.8	92.9	92.8	92.5	-0.4
	19–22	96.6	93.1	95.7	94.7	94.5	95.6	95.1	96.0	95.3	96.1	97.1	95.5	95.6	—	—	—	—	—
	23–26	95.1	95.2	95.2	96.7	94.7	93.2	91.2	90.1	93.0	94.9	95.9	96.6	95.6	—	—	—	—	—
	27–30	94.2	96.1	95.4	95.9	94.2	94.0	95.1	96.3	94.5	95.4	93.2	94.3	94.3	—	—	—	—	—
Trying an amphetamine once or twice ^b	18	80.4	82.6	83.0	84.1	83.8	83.3	85.9	84.7	86.1	86.7	87.3	87.1	87.0	85.8	84.6	83.7	83.5	-0.2
	19–22	84.5	84.0	85.8	81.6	84.5	87.6	87.6	89.4	88.9	89.4	89.1	90.2	87.4	—	—	—	—	—
	23–26	83.3	87.0	85.9	85.1	83.1	83.9	81.5	82.7	86.2	89.9	89.3	89.6	87.2	—	—	—	—	—
	27–30	85.9	85.8	87.2	87.8	86.4	86.0	87.9	88.9	87.5	88.5	82.9	85.3	85.6	—	—	—	—	—
Taking one or two drinks nearly every day ^b	18	71.5	72.3	71.7	71.6	73.4	71.6	74.7	72.8	74.0	73.2	74.5	75.2	75.5	75.0	74.9	74.0	75.4	+1.4
	19–22	73.5	67.3	68.6	66.6	64.9	68.5	64.4	72.4	68.3	68.7	68.4	69.5	68.8	—	—	—	—	—
	23–26	68.1	66.9	66.1	65.4	64.4	61.6	62.1	61.8	62.3	66.1	62.5	63.4	59.4	—	—	—	—	—
	27–30	67.3	66.7	64.3	67.3	67.1	64.0	64.5	65.0	62.8	64.9	59.4	58.9	59.8	—	—	—	—	—
Taking four or five drinks nearly every day ^b	18	82.5	82.8	82.2	82.8	84.4	80.1	83.1	82.9	82.7	83.3	84.8	84.7	84.6	83.4	85.8	84.1	85.8	+1.7
	19–22	90.0	85.9	87.9	86.6	84.6	87.7	86.8	89.8	86.8	89.0	90.7	88.8	89.9	—	—	—	—	—
	23–26	90.7	93.7	89.9	92.5	91.1	88.1	89.3	87.8	89.1	90.8	87.8	93.8	89.1	—	—	—	—	—
	27–30	93.8	92.1	95.3	92.4	91.2	92.7	92.6	92.5	93.4	92.3	91.3	89.0	93.1	—	—	—	—	—
Having five or more drinks once or twice each weekend	18	56.4	55.5	57.6	57.7	57.8	55.6	60.3	59.4	59.9	60.6	60.0	62.1	63.5	62.0	62.2	62.3	65.2	+2.9
	19–22	56.4	52.8	51.8	45.2	47.4	50.4	47.9	52.4	53.2	54.8	54.4	55.2	54.6	47.7	48.7	53.9	53.0	-0.9
	23–26	55.6	60.0	54.5	56.6	56.9	52.9	49.5	49.5	51.9	56.0	51.3	55.3	51.0	51.2	50.7	53.4	48.5	-4.9
	27–30	64.0	63.0	57.7	65.8	58.8	63.3	59.6	64.6	56.9	62.7	56.3	57.3	52.7	52.9	50.6	53.7	52.7	-1.0
Smoking one or more packs of cigarettes per day	18	68.5	69.0	71.2	72.6	74.5	75.7	79.2	78.6	81.1	81.2	81.4	82.5	81.6	81.4	81.6	83.2	84.4	+1.2
	19–22	76.5	69.2	73.9	71.1	74.3	77.3	78.3	82.1	82.7	84.8	87.0	85.5	86.8	85.7	84.8	89.2	87.9	-1.3
	23–26	75.8	76.5	78.0	79.9	77.0	75.4	78.3	77.6	77.4	84.4	82.6	88.2	88.1	88.0	88.2	90.6	85.5	-5.1 s
	27–30	80.5	81.9	82.6	84.0	83.6	86.1	84.0	84.6	82.2	84.1	81.3	83.9	85.0	89.5	88.4	88.1	90.0	+1.9
<i>Approximate Weighted N =</i>	18	2,095	2,037	1,945	1,775	1,862	1,820	2,133	2,208	2,183	2,183	2,161	2,090	2,033	2,101	2,132	2,126	1,916	
	19–22	466	436	430	379	402	361	399	427	395	395	361	370	389	347	364	337	309	
	23–26	419	425	394	398	378	366	363	377	361	344	349	336	322	355	320	329	327	
	27–30	422	440	397	394	374	364	346	408	362	327	330	318	333	322	321	285	303	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '—' indicates data not available.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

^bThese questions were dropped from the questionnaires beginning in 2010.

TABLE 7-2
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take any illicit drug ^b																		
% saying any	18	87.5	85.4	86.3	82.6	81.0	82.4	82.2	81.7	79.1	76.9	71.0	69.1	67.3	71.0	78.3	78.6	80.6
	19–22	90.2	88.0	86.8	85.0	82.3	82.9	80.5	76.7	77.2	78.4	72.7	71.5	66.8	71.7	71.6	71.6	76.2
	23–26	—	—	—	—	83.6	82.7	80.3	80.9	74.4	73.8	65.8	63.0	67.3	64.6	66.7	65.3	64.6
	27–30	—	—	—	—	—	—	—	—	74.8	72.9	69.6	67.1	61.5	60.2	57.1	58.5	59.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	38.1	37.4	39.7	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	32.5	29.8	26.5	23.8	20.9	22.7	21.5	18.6	15.8	15.7	11.6	11.7	12.0	15.5	20.3	21.7	23.8
	19–22	34.9	32.8	28.1	22.4	21.9	18.2	16.2	14.0	13.5	10.9	10.5	8.8	9.0	10.4	14.9	13.1	17.3
	23–26	—	—	—	—	19.6	15.4	16.2	11.7	9.5	9.7	9.5	7.4	6.2	6.4	8.7	7.6	8.8
	27–30	—	—	—	—	—	—	—	—	8.6	6.4	5.9	2.9	5.8	5.0	5.6	6.1	3.6
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.1	1.9	2.0
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take any illicit drug other than marijuana ^b																		
% saying any	18	62.4	63.3	64.7	61.2	61.3	61.8	63.3	62.4	56.5	56.2	50.1	46.3	47.1	48.7	53.7	53.7	54.5
	19–22	67.9	67.8	66.7	65.2	60.8	62.1	61.0	57.3	53.5	60.8	53.4	51.5	45.3	51.4	46.3	46.4	46.5
	23–26	—	—	—	—	63.7	64.0	59.0	61.1	55.1	54.2	47.8	41.8	46.1	42.3	39.4	40.3	32.8
	27–30	—	—	—	—	—	—	—	—	55.9	55.0	49.7	47.2	37.7	38.5	33.9	37.7	36.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	21.4	21.6	22.1
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	11.1	11.9	10.9	11.0	10.3	10.4	10.3	9.2	6.9	7.7	5.1	4.6	5.3	7.1	7.1	7.7	8.9
	19–22	9.8	12.9	11.8	9.8	9.3	8.6	7.6	5.0	5.3	4.0	3.2	2.6	3.3	4.0	4.4	3.5	6.2
	23–26	—	—	—	—	10.6	6.6	8.6	5.2	3.9	4.2	3.4	1.6	1.8	2.8	2.5	1.9	1.9
	27–30	—	—	—	—	—	—	—	—	4.6	3.0	2.8	1.0	1.4	1.5	1.5	1.5	0.9
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.5	0.7
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smoke marijuana																		
% saying any	18	86.4	83.0	84.4	80.3	77.7	79.5	79.2	78.4	75.3	72.5	68.3	65.8	63.1	67.4	75.6	76.1	78.0
	19–22	88.8	86.4	85.2	83.8	81.6	81.1	78.5	75.3	75.1	73.8	67.6	68.0	63.5	67.6	67.4	68.8	74.9
	23–26	—	—	—	—	82.0	80.8	77.7	79.4	71.6	69.8	61.8	59.6	61.3	61.2	62.6	63.2	62.6
	27–30	—	—	—	—	—	—	—	—	71.8	68.2	65.1	62.6	58.0	57.4	52.3	55.7	55.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.9	36.3	36.3
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	31.3	27.7	23.8	21.7	18.3	19.8	18.2	15.8	13.6	13.4	10.1	10.0	10.3	13.9	18.9	20.7	22.2
	19–22	34.1	30.6	25.6	20.6	19.4	16.0	13.3	12.5	12.2	9.0	9.2	8.3	8.2	8.5	13.0	12.5	16.3
	23–26	—	—	—	—	17.0	14.3	13.7	10.4	7.8	8.6	8.3	6.9	5.6	5.6	7.5	6.6	8.2
	27–30	—	—	—	—	—	—	—	—	6.8	4.4	4.0	2.8	5.1	5.2	5.0	5.6	3.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.0	2.5	2.9
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Take any illicit drug ^b																			
% saying any	18	83.4	84.6	82.0	82.0	82.8	81.8	80.7	81.2	79.8	78.8	77.7	80.1	79.2	80.4	81.7	78.9	80.8	+1.9
	19–22	77.2	79.8	77.3	83.1	81.1	78.3	79.4	78.1	78.6	74.7	79.8	77.4	80.3	78.6	78.1	78.8	78.3	-0.5
	23–26	67.0	67.6	67.9	67.8	66.9	73.4	70.8	70.8	74.2	72.2	71.3	72.2	74.5	75.7	80.3	74.2	76.9	+2.7
	27–30	60.9	58.3	59.6	55.6	57.2	61.8	58.6	63.1	63.7	62.3	62.7	66.7	70.0	66.9	69.2	72.3	72.9	+0.6
	35	39.2	38.4	36.3	37.7	39.1	40.9	37.5	37.9	40.0	40.4	42.1	44.9	44.4	45.0	50.8	49.0	52.7	+3.7
	40	—	39.2	38.2	38.0	38.4	36.2	36.5	34.6	36.2	35.4	34.6	35.9	39.0	37.3	36.6	40.3	42.1	+1.8
	45	—	—	—	—	—	—	37.8	38.3	34.3	36.7	38.5	35.9	36.1	37.7	36.2	39.2	39.5	+0.3
	50	—	—	—	—	—	—	—	—	—	—	—	39.3	37.0	36.5	36.0	38.4	39.1	+0.7
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.2	—	—
% saying most or all	18	23.7	25.9	25.5	24.5	25.2	23.1	23.5	23.0	20.2	20.9	21.7	21.3	22.4	25.4	29.1	26.4	26.7	+0.3
	19–22	16.2	16.8	20.6	18.9	20.3	20.2	17.3	14.7	15.8	16.8	14.5	13.7	16.0	17.2	21.8	17.3	22.1	+4.7
	23–26	10.5	9.6	8.4	9.7	10.4	10.3	11.7	9.7	11.1	8.1	8.9	12.7	13.9	10.5	9.1	13.6	13.6	+4.5
	27–30	4.5	5.3	5.7	5.3	7.1	6.9	6.9	3.9	4.7	5.4	6.5	6.3	6.4	6.6	7.1	6.8	6.7	-0.1
	35	3.0	3.1	2.8	3.1	3.2	2.9	3.2	2.8	2.5	2.1	2.2	2.2	2.5	3.7	4.5	3.3	4.7	+1.4
	40	—	2.3	2.0	2.0	1.6	2.2	1.6	1.6	2.1	2.5	2.0	1.3	1.3	2.1	1.9	1.2	2.3	+1.1
	45	—	—	—	—	—	—	2.2	1.5	1.4	1.7	1.3	1.3	1.2	1.5	1.2	1.0	1.5	0.5
	50	—	—	—	—	—	—	—	—	—	—	—	1.4	1.4	1.4	1.8	1.8	1.0	-0.8
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.0	—	—
Take any illicit drug other than marijuana ^b																			
% saying any	18	55.1	55.6	51.2	52.5	55.0	54.3	50.0	51.4	51.3	51.0	49.3	49.4	53.7	49.9	48.9	45.4	-3.5	
	19–22	49.7	53.3	54.8	56.1	60.0	57.2	50.8	53.4	54.9	49.5	52.5	46.4	47.5	52.0	52.0	49.3	52.4	+3.1
	23–26	35.1	35.4	41.1	42.5	42.6	49.4	42.3	47.1	46.6	45.6	42.6	45.9	44.4	52.4	50.5	46.6	45.3	-1.3
	27–30	33.9	34.1	35.2	31.7	33.5	36.0	34.7	35.8	33.1	36.2	34.2	36.4	41.6	40.1	40.9	50.1	44.6	-5.5
	35	19.2	19.3	19.0	17.9	18.7	20.4	18.5	20.2	18.5	18.1	20.7	23.7	20.2	23.9	26.4	25.7	25.2	-0.6
	40	—	20.9	21.0	21.9	21.4	21.0	20.2	18.5	21.0	20.3	20.3	19.8	20.6	18.8	17.4	20.2	18.7	-1.5
	45	—	—	—	—	—	—	23.4	25.1	20.8	22.7	25.0	21.2	20.7	20.9	21.5	22.6	20.9	-1.7
	50	—	—	—	—	—	—	—	—	—	—	—	24.5	24.8	21.7	22.8	22.2	20.1	-2.1
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	19.9	—	—
% saying most or all	18	7.0	8.9	7.4	7.4	7.0	6.1	6.7	7.3	6.7	5.3	6.5	5.3	5.6	7.1	6.5	5.5	4.3	-1.3
	19–22	4.1	4.3	5.1	7.7	8.0	5.7	5.1	3.5	4.8	4.2	3.9	3.4	3.6	4.8	7.4	4.6	6.6	+2.1
	23–26	2.6	2.8	2.2	3.8	3.7	2.8	3.7	3.1	3.2	3.2	1.0	1.5	2.9	3.2	3.1	3.2	3.3	+0.1
	27–30	1.2	0.9	1.3	1.5	2.6	2.3	0.7	0.8	0.9	1.4	2.2	2.5	1.7	1.5	2.5	1.5	0.7	-0.8
	35	0.5	0.7	0.9	1.0	0.9	0.6	0.6	0.4	0.5	0.5	0.6	0.5	0.5	1.0	0.7	0.8	1.3	+0.5
	40	—	0.4	0.8	0.7	0.5	0.3	0.3	0.2	0.5	0.7	0.6	0.7	0.3	*	0.2	0.1	0.2	+0.1
	45	—	—	—	—	—	—	0.7	0.7	0.4	0.9	0.5	0.5	0.3	0.3	0.1	0.2	0.4	+0.2
	50	—	—	—	—	—	—	—	—	—	—	—	0.5	0.4	0.3	0.8	0.4	0.2	-0.1
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.4	—	—
Smoke marijuana																			
% saying any	18	81.4	83.2	80.7	80.5	81.2	79.4	78.9	79.5	77.4	76.4	74.8	78.2	77.2	79.7	80.6	77.7	80.2	+2.5
	19–22	74.7	77.2	73.9	81.2	78.4	77.2	76.5	75.6	75.8	72.0	76.6	74.7	77.7	75.6	74.7	76.8	76.2	-0.5
	23–26	63.5	65.0	64.4	64.8	64.5	68.8	67.7	68.4	70.7	67.6	69.0	67.7	71.7	71.9	77.5	71.5	73.4	+2.0
	27–30	58.3	55.5	57.0	51.7	56.5	59.0	55.8	60.4	60.8	61.0	60.2	64.2	65.2	62.3	65.9	66.6	69.2	+2.6
	35	35.0	34.6	33.3	34.9	35.6	37.4	32.9	34.7	37.2	37.3	38.6	42.1	40.6	41.3	47.4	45.1	48.8	+3.7
	40	—	34.6	32.5	32.3	31.8	31.4	30.7	29.9	30.4	29.4	29.2	29.6	33.6	32.1	32.4	35.8	38.0	+2.2
	45	—	—	—	—	—	—	31.1	29.4	26.3	28.4	30.0	28.6	29.4	32.6	30.3	33.0	34.5	+1.5
	50	—	—	—	—	—	—	—	—	—	—	—	30.1	26.9	28.0	27.9	31.3	33.0	+1.7
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	—	—
% saying most or all	18	22.5	23.8	24.2	23.2	24.0	21.4	21.7	21.1	17.9	19.6	19.2	19.9	20.9	23.6	27.3	25.0	25.7	+0.8
	19–22	16.2	16.4	19.4	16.6	18.5	18.6	16.0	15.0	13.4	15.7	13.4	11.5	14.5	15.4	19.1	16.2	19.7	+3.4
	23–26	9.8	9.0	8.5	8.2	9.0	8.7	9.3	9.8	8.0	10.1	7.9	8.5	12.2	12.3	9.6	8.3	12.8	+4.5
	27–30	3.9	4.8	5.5	4.9	6.3	6.2	6.7	3.5	4.3	5.0	6.6	5.0	5.8	6.3	5.8	5.6	6.6	+1.0
	35	2.9	2.8	2.6	2.8	2.6	2.7	3.1	2.7	2.3	2.0	2.1	1.9	2.3	3.4	4.2	2.9	4.3	+1.5
	40	—	2.1	1.4	1.9	1.2	2.0	1.4	1.6	1.8	2.1	1.6	0.9	1.2	2.1	1.7	1.2	2.1	+0.9
	45	—	—	—	—	—	—	1.9	0.9	1.3	1.1	1.0	1.0	1.1	1.3	1.1	0.9	1.3	+0.4
	50	—	—	—	—	—	—	—	—	—	—	—	1.2	1.2	1.2	1.3	1.5	1.0	+0.4
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.8	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	
Q. How many of your friends would you estimate. . .		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Use inhalants																			
% saying any		18	17.8	16.5	18.4	16.1	19.3	21.2	22.4	24.7	20.8	22.1	20.0	19.2	22.2	23.7	26.5	27.5	27.2
		19–22	11.9	13.2	13.8	12.3	11.7	9.6	10.9	12.7	10.9	11.7	13.0	12.2	12.6	13.8	14.0	14.2	16.2
		23–26	—	—	—	—	7.7	6.7	7.2	6.1	6.2	5.9	6.1	4.4	5.1	6.3	7.0	9.3	5.6
		27–30	—	—	—	—	—	—	—	4.6	3.5	2.9	2.5	3.3	2.9	3.5	4.0	4.1	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.2	0.9	1.3	1.1	1.1	1.5	2.0	1.9	1.2	1.9	1.0	0.7	1.8	1.8	2.0	2.0	2.4
		19–22	0.5	0.4	0.7	0.3	0.5	0.6	0.7	0.7	0.4	0.6	0.2	0.8	0.7	0.7	0.6	1.1	—
		23–26	—	—	—	—	0.6	0.2	0.6	0.1	0.2	0.4	0.4	0.1	*	0.1	0.2	0.7	0.5
		27–30	—	—	—	—	—	—	—	0.3	*	0.2	0.2	*	0.2	*	*	*	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use nitrites																			
% saying any		18	19.0	17.4	17.5	14.5	15.0	15.6	18.0	18.3	13.6	13.3	10.4	8.9	9.0	10.7	10.0	10.7	11.2
		19–22	18.4	16.0	14.2	13.8	8.9	9.9	11.7	13.2	10.2	—	—	—	—	—	—	—	—
		23–26	—	—	—	—	10.8	7.8	8.0	7.9	5.2	—	—	—	—	—	—	—	—
		27–30	—	—	—	—	—	—	—	—	6.6	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.3	1.2	0.9	0.7	1.2	1.0	1.2	1.3	0.7	0.9	0.6	0.4	0.7	0.7	0.8	0.8	0.8
		19–22	0.3	0.4	0.9	0.6	0.6	0.6	0.4	0.4	0.2	—	—	—	—	—	—	—	—
		23–26	—	—	—	—	0.8	0.3	0.4	0.3	0.1	—	—	—	—	—	—	—	—
		27–30	—	—	—	—	—	—	—	—	0.5	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take LSD																			
% saying any		18	28.1	28.5	27.8	24.0	23.9	24.4	24.5	25.3	24.1	25.2	25.0	23.4	28.1	31.3	34.1	36.9	37.9
		19–22	30.9	25.9	26.5	22.6	21.6	18.8	18.7	18.2	19.0	20.1	20.1	22.0	22.2	28.8	23.8	26.9	28.6
		23–26	—	—	—	—	21.5	17.2	15.4	15.9	13.3	14.1	12.3	12.5	15.0	17.2	17.3	21.5	15.3
		27–30	—	—	—	—	—	—	—	—	10.4	7.7	9.1	8.6	10.9	8.7	8.1	12.0	11.6
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.8	2.2	2.4	1.4	2.0	1.5	1.8	1.6	1.5	2.4	1.9	1.7	2.4	3.8	4.2	4.8	5.0
		19–22	1.2	0.8	0.9	1.0	0.6	0.8	0.9	0.6	1.3	0.4	1.2	1.4	1.9	2.1	2.5	2.3	3.8
		23–26	—	—	—	—	0.8	0.5	1.0	0.2	0.6	0.5	0.6	0.2	0.4	0.7	1.1	0.7	0.7
		27–30	—	—	—	—	—	—	—	—	0.3	0.2	0.3	0.3	*	0.3	0.4	0.3	0.4
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	2012–2013	
Q. How many of your friends would you estimate. . .		Age Group	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	change
Use inhalants																				
% saying any		18	27.4	25.9	21.6	23.5	22.2	21.0	17.5	17.9	18.1	18.9	17.9	18.0	18.0	19.0	16.4	12.3	12.1	-0.2
		19–22	13.7	16.2	16.3	13.7	13.7	10.4	10.0	9.5	11.1	11.0	9.6	7.4	6.6	8.3	11.9	8.2	7.3	-0.8
		23–26	7.5	6.2	7.9	6.9	7.5	7.4	7.9	6.2	5.8	5.2	3.7	6.1	6.5	6.0	4.8	5.4	4.1	-1.3
		27–30	3.6	3.8	4.2	3.6	6.0	4.5	3.2	2.6	3.2	3.3	2.8	2.7	3.6	1.7	3.2	3.8	2.9	-0.9
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.9	2.7	1.8	1.4	1.4	1.2	1.1	1.2	2.0	1.2	1.6	1.1	0.9	1.8	1.4	0.9	1.1	+0.2
		19–22	0.7	1.3	0.8	0.6	1.2	0.4	0.6	*	0.1	0.5	0.3	0.6	*	0.6	0.6	0.1	0.9	+0.7
		23–26	0.8	*	0.1	0.7	0.1	0.4	0.3	*	0.1	0.3	*	*	*	0.1	0.1	*	0.3	+0.3
		27–30	*	*	*	*	0.3	0.3	*	*	*	*	*	0.3	0.3	*	0.3	*	0.1	+0.1
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	s
Use nitrites																				
% saying any		18	11.9	12.9	10.9	11.0	11.9	11.2	8.5	9.4	9.1	8.1	7.7	7.3	7.7	—	—	—	—	—
		19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	0.7	1.0	0.7	1.0	0.6	0.8	1.0	1.2	1.0	0.5	0.7	0.5	0.2	—	—	—	—	—
		19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take LSD																				
% saying any		18	36.5	36.8	32.2	31.9	32.2	28.6	21.9	23.5	19.5	18.7	18.3	20.9	21.3	22.3	22.5	21.3	17.7	-3.6 s
		19–22	24.7	29.4	28.2	27.8	28.4	24.0	15.4	15.9	13.9	14.2	15.1	12.5	12.8	16.0	18.0	15.7	23.3	+7.7 s
		23–26	18.2	15.2	18.1	19.3	16.8	15.8	16.1	14.4	12.0	11.7	11.2	9.2	11.0	11.9	10.2	11.5	13.4	+2.0
		27–30	12.3	12.6	13.4	11.8	12.5	13.1	11.4	8.9	6.6	9.1	7.6	8.8	7.6	8.2	7.6	7.7	10.6	+2.9
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	3.7	4.7	3.9	3.1	2.9	1.7	1.9	1.5	1.5	0.8	1.2	1.1	1.1	1.5	1.4	1.3	1.2	-0.1
		19–22	1.4	2.5	1.8	2.1	2.7	1.6	0.8	0.3	0.3	0.2	0.8	0.2	0.3	1.4	0.7	1.0	1.0	-0.1
		23–26	0.6	1.0	1.5	0.9	0.3	0.4	0.8	0.4	0.2	0.3	*	0.1	0.5	0.1	0.6	0.4	0.3	0.0
		27–30	0.4	0.1	0.6	0.4	0.4	0.3	0.1	*	0.3	0.4	0.4	*	0.5	0.2	0.1	*	*	0.0
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																		
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	(Years Cont.)
Take other hallucinogens ^c																				
% saying any		18	28.2	26.3	25.6	22.1	21.3	22.0	22.3	21.7	17.8	18.1	15.9	15.1	17.0	19.3	21.4	23.8	26.4	
		19–22	33.4	25.5	25.1	21.0	20.2	16.6	15.8	15.0	16.1	13.9	15.3	14.2	12.0	15.0	13.8	14.9	17.2	
		23–26	—	—	—	—	20.0	16.7	13.2	13.2	11.7	9.6	8.7	8.5	9.8	9.4	10.3	11.7	10.4	
		27–30	—	—	—	—	—	—	—	—	10.6	7.4	7.1	6.8	7.9	7.1	6.6	7.9	7.5	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	2.2	2.1	1.9	1.6	1.9	1.4	1.3	1.2	0.9	1.4	1.0	0.8	1.0	1.7	2.2	2.2	2.3	
		19–22	1.5	0.9	1.1	1.2	0.7	1.0	0.7	0.6	0.9	0.2	0.5	0.8	0.7	0.9	1.6	1.5	1.0	
		23–26	—	—	—	—	0.8	0.3	0.5	0.3	0.2	0.3	0.8	0.1	0.4	0.7	0.6	0.8	0.1	
		27–30	—	—	—	—	—	—	—	0.2	0.1	0.3	0.2	*	0.2	0.3	0.1	0.2		
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take PCP																				
% saying any		18	22.2	17.2	17.3	14.2	14.2	15.9	16.1	15.5	13.5	14.7	13.0	12.0	12.7	15.6	15.5	18.3	20.3	
		19–22	24.1	15.3	15.3	12.6	9.5	8.9	10.1	9.7	10.1	—	—	—	—	—	—	—	—	
		23–26	—	—	—	—	11.6	6.8	7.4	6.9	5.1	—	—	—	—	—	—	—	—	
		27–30	—	—	—	—	—	—	—	—	6.7	—	—	—	—	—	—	—	—	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	1.6	0.9	0.9	1.1	1.1	1.2	1.2	1.1	0.8	1.2	0.5	0.5	0.9	1.9	1.2	1.2	1.3	
		19–22	0.5	0.3	0.3	0.5	0.7	0.7	0.2	0.1	0.3	—	—	—	—	—	—	—	—	
		23–26	—	—	—	—	0.6	*	0.4	*	0.2	—	—	—	—	—	—	—	—	
		27–30	—	—	—	—	—	—	—	—	0.4	—	—	—	—	—	—	—	—	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take ecstasy (MDMA)																				
% saying any		18	—	—	—	—	—	—	—	—	—	—	12.4	11.9	10.7	12.8	15.9	20.7	24.2	
		19–22	—	—	—	—	—	—	—	—	—	16.3	14.3	12.0	12.9	13.7	11.3	17.2	20.7	
		23–26	—	—	—	—	—	—	—	—	—	7.6	9.0	9.5	11.0	9.8	11.4	11.2	11.3	
		27–30	—	—	—	—	—	—	—	—	—	5.6	6.3	5.4	4.6	6.6	5.8	6.9	10.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	—	—	—	—	—	—	—	—	—	—	2.2	1.7	2.1	1.2	1.7	2.8	3.0	
		19–22	—	—	—	—	—	—	—	—	—	0.4	0.7	0.2	0.7	0.7	0.5	0.5	0.8	
		23–26	—	—	—	—	—	—	—	—	—	0.5	0.2	0.1	0.1	0.5	0.1	0.4	0.1	
		27–30	—	—	—	—	—	—	—	—	—	0.5	0.3	*	0.1	0.3	0.2	0.5	0.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Take other hallucinogens^c																			
% saying any	18	26.3	27.4	22.5	24.0	35.4	33.6	30.1	31.9	31.0	30.1	30.1	29.4	30.5	32.3	31.8	29.5	26.9	-2.7
	19–22	17.2	19.1	18.9	20.9	33.6	33.5	24.8	26.8	25.1	27.8	26.7	21.9	21.8	26.4	26.4	22.6	28.3	+5.6
	23–26	13.0	11.7	9.6	11.3	18.6	22.4	20.2	24.5	18.5	18.9	15.9	21.1	19.6	22.6	16.5	17.5	17.4	-0.1
	27–30	6.8	7.8	9.4	8.0	14.6	14.9	13.5	12.4	9.4	14.9	10.6	16.9	12.1	14.9	13.9	17.1	16.5	-0.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.6	3.1	2.4	2.4	2.9	2.3	2.4	2.6	2.2	1.7	1.7	1.8	1.6	2.0	2.1	2.0	1.6	-0.4
	19–22	1.1	1.7	0.8	2.0	2.3	2.2	1.5	1.1	0.6	0.9	1.0	1.3	0.6	0.9	0.7	0.8	1.0	+0.2
	23–26	0.8	0.7	0.8	0.3	0.6	0.7	1.0	0.8	0.7	1.0	0.4	0.4	1.2	0.2	0.7	0.5	1.2	+0.6
	27–30	0.3	0.2	0.2	0.4	0.6	1.0	0.1	*	0.4	0.4	0.4	0.4	0.6	0.7	0.5	0.3	0.4	+0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take PCP																			
% saying any	18	19.7	20.2	16.8	17.5	19.1	17.2	13.6	11.8	10.1	10.6	9.4	9.4	9.3	—	—	—	—	—
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.4	1.6	1.5	1.7	1.3	1.0	1.5	1.1	1.0	0.5	0.8	0.5	0.5	—	—	—	—	—
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take ecstasy (MDMA)																			
% saying any	18	27.7	24.5	26.7	37.3	41.9	38.0	34.2	28.9	23.1	23.0	23.6	24.7	23.5	25.9	27.5	26.8	25.6	-1.1
	19–22	21.4	26.0	30.7	42.4	43.3	43.4	31.3	27.6	28.3	25.2	21.6	19.3	24.4	20.4	22.0	18.9	27.4	+8.5 ss
	23–26	15.1	13.7	15.2	25.9	29.4	36.8	27.0	31.2	25.3	23.4	16.5	20.8	19.7	20.7	19.5	18.8	19.1	+0.4
	27–30	7.4	8.5	12.4	13.1	17.8	20.6	19.4	20.6	15.6	22.6	15.9	17.8	17.0	12.7	10.6	15.8	13.5	-2.3
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.6	2.5	2.7	4.8	5.2	3.7	2.7	3.2	2.5	1.9	2.1	2.4	2.2	2.1	2.7	2.7	1.8	-0.9
	19–22	1.7	2.0	2.9	4.9	5.8	2.7	1.9	1.9	1.1	0.7	0.8	0.9	1.6	1.3	2.9	1.2	2.0	+0.8
	23–26	0.8	0.8	0.4	2.9	1.7	1.2	2.0	1.1	1.3	0.9	0.3	0.4	0.7	0.6	0.4	1.7	1.0	-0.6
	27–30	0.3	*	0.8	0.4	0.3	0.9	0.5	0.6	0.2	*	0.1	0.9	0.7	0.1	0.5	0.3	0.1	-0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																	
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
																			→ (Years Cont.)
Take cocaine																			
% saying any	18	41.6	40.1	40.7	37.6	38.9	43.8	45.6	43.7	37.7	37.4	31.7	26.8	26.3	24.5	26.1	24.8	28.1	
	19–22	51.0	48.9	49.8	46.5	47.6	45.9	48.3	45.7	42.0	42.7	33.2	29.7	22.8	24.3	21.5	22.0	19.4	
	23–26	—	—	—	—	52.4	53.2	51.6	50.7	47.1	40.8	34.8	29.0	28.8	27.1	22.3	24.4	18.1	
	27–30	—	—	—	—	—	—	—	—	47.9	43.3	38.3	35.7	29.9	27.6	22.6	26.2	20.8	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all	18	6.1	6.3	4.9	5.1	5.1	5.8	6.2	5.1	3.4	3.7	2.1	1.5	1.5	2.1	1.5	2.0	2.2	
	19–22	7.0	8.6	7.8	6.1	6.3	6.1	6.1	3.3	3.5	2.1	1.2	1.1	1.0	0.5	1.5	0.9	1.0	
	23–26	—	—	—	—	9.1	5.3	7.0	4.1	3.1	2.7	2.1	0.6	0.9	0.8	1.0	0.3	0.4	
	27–30	—	—	—	—	—	—	—	—	3.8	2.0	2.3	0.9	1.2	0.8	0.8	0.4	0.4	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take crack																			
% saying any	18	—	—	—	—	—	—	—	27.4	25.4	26.1	19.2	17.6	17.8	17.9	20.0	19.2	21.6	
	19–22	—	—	—	—	—	—	—	23.8	21.8	20.6	14.6	14.3	11.8	13.6	13.8	14.0	9.4	
	23–26	—	—	—	—	—	—	—	26.4	22.4	19.8	14.4	10.8	10.8	8.8	8.8	11.1	8.2	
	27–30	—	—	—	—	—	—	—	—	22.1	18.4	16.6	11.6	10.3	10.2	10.4	10.3	8.6	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.5	5.1	4.4	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all	18	—	—	—	—	—	—	—	2.2	1.1	2.1	0.6	0.6	0.7	0.9	1.0	1.1	0.9	
	19–22	—	—	—	—	—	—	—	0.7	0.8	1.0	0.6	0.2	0.1	0.3	0.4	0.3	0.5	
	23–26	—	—	—	—	—	—	—	0.8	0.9	0.8	0.5	0.1	0.1	0.5	0.2	*	0.3	
	27–30	—	—	—	—	—	—	—	—	1.2	0.9	0.9	0.3	*	0.6	0.3	0.1	0.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.3	0.4	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take cocaine powder																			
% saying any	18	—	—	—	—	—	—	—	—	—	25.3	24.6	19.8	19.7	18.1	20.7	19.2	22.8	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14.2	12.9	15.4	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all	18	—	—	—	—	—	—	—	—	—	2.3	2.5	1.8	2.0	1.6	1.9	1.7	1.9	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.3	0.6	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

TABLE 7-2 (cont.)

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																2012–2013 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013
Take cocaine																			
% saying any	18	28.2	31.2	27.8	27.2	27.1	26.8	23.8	29.3	28.1	29.7	29.7	25.2	24.0	22.9	18.8	18.1	18.8	+0.7
	19–22	22.2	26.8	25.7	24.8	27.4	28.2	25.5	26.2	27.2	26.6	29.4	21.8	21.2	21.8	22.3	15.9	19.5	+3.6
	23–26	19.7	18.7	20.1	20.3	19.4	23.7	21.9	27.4	25.6	24.6	23.1	23.1	23.5	28.0	23.7	21.6	18.9	-2.7
	27–30	21.5	18.6	20.7	16.5	19.7	16.0	17.0	17.0	17.9	19.5	18.6	20.7	22.1	19.2	16.1	21.6	18.4	-3.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.0	3.2	2.9	2.0	1.7	1.7	2.4	2.3	2.3	1.9	2.1	1.2	1.8	1.4	1.0	0.8	1.1	+0.4
	19–22	0.8	1.5	1.1	1.0	1.8	1.0	1.4	0.7	2.1	1.0	1.1	1.3	0.5	0.2	0.5	0.3	0.9	+0.6
	23–26	1.1	0.9	0.5	0.8	1.6	1.0	1.6	1.0	1.5	1.4	0.8	0.6	1.7	0.9	0.4	0.6	0.9	+0.3
	27–30	0.6	0.1	0.4	0.4	0.5	0.6	0.3	0.3	*	*	1.4	0.9	0.7	0.2	0.5	0.6	0.5	-0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take crack																			
% saying any	18	22.2	24.4	19.0	21.4	23.4	21.5	18.7	22.5	22.9	22.3	21.8	19.1	18.8	15.2	12.1	10.4	10.3	-0.1
	19–22	13.1	16.4	15.7	16.5	17.4	18.0	11.8	16.0	14.9	14.5	16.0	12.2	11.3	7.2	8.3	5.1	8.3	+3.2
	23–26	8.3	8.3	8.8	7.9	8.6	10.1	10.4	10.8	10.8	10.0	8.7	9.8	8.5	7.0	6.7	6.5	7.5	+1.1
	27–30	6.3	6.4	8.7	6.0	7.1	6.4	6.5	5.2	8.5	9.1	6.9	5.8	9.5	3.6	4.2	3.6	4.6	+0.9
	35	3.1	2.8	3.2	3.9	2.8	3.2	2.8	3.1	2.6	2.8	3.0	2.8	2.5	3.4	2.1	1.3	2.9	+1.5 s
	40	—	3.8	3.0	2.9	3.5	2.6	2.7	2.6	2.8	2.3	3.1	1.9	1.2	1.5	1.2	1.2	1.2	0.0
	45	—	—	—	—	—	—	3.7	3.3	2.4	3.0	2.9	1.8	2.3	2.2	1.8	1.6	2.1	+0.5
	50	—	—	—	—	—	—	—	—	—	—	—	2.0	1.6	1.8	1.4	1.1	1.2	+0.1
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.1	—
% saying most or all	18	1.1	1.7	1.5	1.4	0.8	0.8	1.4	1.6	1.6	1.0	1.3	1.1	1.1	1.5	0.9	0.8	0.9	+0.1
	19–22	0.3	0.9	0.9	0.5	0.3	0.2	0.4	0.1	1.0	0.8	0.3	0.4	0.3	*	0.3	0.3	0.6	+0.3
	23–26	0.5	0.4	*	0.5	0.3	*	0.3	0.5	0.2	0.7	0.1	0.3	*	0.4	0.1	*	0.3	+0.3
	27–30	0.2	0.1	*	*	*	0.3	0.1	*	0.1	*	0.3	0.6	0.3	*	*	*	0.5	+0.5
	35	*	0.1	0.3	0.5	0.2	0.3	0.3	0.1	0.2	0.2	0.2	*	0.1	0.3	*	0.1	*	-0.1
	40	—	*	0.2	0.2	0.1	*	*	*	0.1	0.2	0.1	0.0	*	*	*	*	0.1	+0.1
	45	—	—	—	—	—	—	0.4	0.3	0.2	0.2	*	*	0.1	*	*	0.1	0.1	0.0
	50	—	—	—	—	—	—	—	—	—	—	—	0.2	0.1	0.2	0.1	*	0.2	+0.2
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.2	—
Take cocaine powder																			
% saying any	18	24.8	22.9	22.0	21.3	20.1	22.4	23.2	25.4	23.2	22.8	22.3	22.6	19.1	17.6	15.9	17.4	15.6	-1.8
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	11.1	10.4	10.0	10.3	9.4	9.4	8.2	9.2	8.3	8.4	9.1	11.4	8.7	10.5	12.8	9.0	11.6	+2.6
	40	—	10.8	8.9	8.8	8.8	8.5	7.6	7.6	8.9	7.3	6.7	6.2	6.5	4.9	4.8	5.2	6.6	+1.5
	45	—	—	—	—	—	—	8.3	8.0	7.0	7.4	8.0	6.7	6.4	5.9	5.8	6.0	6.0	-0.0
	50	—	—	—	—	—	—	—	—	—	—	—	6.0	5.4	5.3	4.9	4.9	4.4	-0.5
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.1	—
% saying most or all	18	2.0	1.9	1.9	1.8	1.5	1.9	1.9	3.3	1.7	1.7	1.8	1.5	1.5	1.0	1.6	1.5	1.2	-0.2
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	0.4	0.4	0.6	0.7	0.4	0.4	0.4	0.2	0.2	0.2	0.4	0.2	0.2	0.4	0.0	0.1	0.7	+0.6 s
	40	—	0.2	0.2	*	0.2	0.1	0.1	0.1	0.2	0.5	0.2	0.2	0.1	*	*	*	*	0.0
	45	—	—	—	—	—	—	0.5	0.5	0.2	0.4	0.1	0.1	*	*	*	0.1	0.3	+0.2
	50	—	—	—	—	—	—	—	—	—	—	—	0.2	0.2	0.2	0.0	*	0.1	+0.1
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.2	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	→	
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	(Years Cont.)
Take heroin																				
% saying any		18	13.0	12.5	13.2	12.0	13.0	14.5	15.3	13.9	12.4	14.0	11.4	11.4	13.2	13.3	14.3	14.5	15.6	
		19-22	11.0	8.1	9.4	7.5	7.1	6.5	8.5	8.5	7.8	6.8	6.5	6.1	4.7	7.0	8.1	10.4	6.7	
		23-26	—	—	—	—	6.1	4.4	4.3	6.5	3.6	5.2	4.2	3.6	3.8	4.5	4.9	5.8	4.0	
		27-30	—	—	—	—	—	—	—	—	3.8	2.8	4.5	2.7	3.1	3.6	4.2	3.6	4.4	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	1.0	0.5	0.7	0.8	0.8	0.9	1.1	0.9	0.7	1.1	0.4	0.4	0.7	1.1	1.0	1.1	0.9	
		19-22	0.3	0.5	0.1	0.2	0.4	0.6	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.4	0.4	0.4	
		23-26	—	—	—	—	0.4	0.2	0.2	*	0.2	0.4	0.2	0.3	0.4	0.1	0.2	0.2	*	
		27-30	—	—	—	—	—	—	—	—	0.2	0.1	0.2	0.2	*	0.2	0.3	*	*	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take other narcotics^d																				
% saying any		18	22.4	23.1	23.9	20.8	21.4	22.8	21.8	23.2	19.2	19.2	17.2	13.7	14.9	16.1	18.5	19.5	21.8	
		19-22	22.8	20.4	21.9	17.9	17.4	16.9	14.6	15.4	14.1	15.0	12.9	14.1	10.8	13.2	10.5	15.9	13.4	
		23-26	—	—	—	—	16.0	14.9	14.0	13.0	10.6	10.8	10.5	8.5	8.4	8.7	8.0	10.5	8.9	
		27-30	—	—	—	—	—	—	—	—	12.1	8.6	9.1	9.3	7.5	8.2	8.0	7.7	9.5	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	1.7	1.5	1.4	1.4	1.6	1.4	1.8	1.4	1.2	1.4	0.9	0.5	1.1	1.2	1.0	1.6	1.5	
		19-22	0.9	0.7	0.6	0.5	0.8	1.0	0.5	0.4	0.9	0.1	0.6	0.4	0.5	0.6	0.6	0.6	0.4	
		23-26	—	—	—	—	0.4	0.3	0.7	*	0.3	0.2	0.2	*	*	*	0.3	0.2	*	
		27-30	—	—	—	—	—	—	—	—	0.3	*	0.2	0.2	0.1	0.2	0.2	*	0.2	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take amphetamines^e																				
% saying any		18	43.9	48.8	50.6	46.1	45.1	43.3	41.8	39.5	33.4	33.5	28.7	24.3	24.3	27.5	28.1	30.3	32.2	
		19-22	54.1	52.2	51.3	49.7	46.1	42.1	38.5	34.5	26.8	29.6	23.3	26.2	19.5	21.0	20.9	21.7	21.6	
		23-26	—	—	—	—	45.6	40.1	33.5	32.1	28.4	23.1	20.6	17.1	15.1	16.8	16.2	18.2	12.5	
		27-30	—	—	—	—	—	—	—	—	26.1	21.6	19.3	17.0	15.3	14.0	13.1	13.7	15.5	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	4.8	6.4	5.4	5.1	4.5	3.4	3.4	2.6	1.9	2.6	1.9	1.3	1.3	2.0	1.8	2.0	2.8	
		19-22	3.8	5.7	4.6	3.8	3.3	2.9	1.3	1.9	1.4	0.7	1.0	0.6	0.9	0.2	1.1	1.2	0.7	
		23-26	—	—	—	—	1.9	1.8	1.7	1.2	0.3	0.6	0.7	0.8	0.4	1.5	0.9	0.5	0.2	
		27-30	—	—	—	—	—	—	—	—	0.6	0.4	0.5	0.5	0.1	0.5	0.5	0.3	0.3	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Take heroin																			
% saying any	18	15.6	16.5	12.7	14.9	13.1	12.9	10.3	12.7	13.1	12.7	12.9	11.2	12.7	12.4	10.2	7.7	8.5	+0.8
	19–22	7.4	9.4	9.7	7.7	8.7	8.9	5.3	7.0	6.4	7.5	9.0	6.4	3.9	5.3	6.2	6.4	4.8	-1.6
	23–26	6.2	5.8	4.8	4.7	5.0	5.2	6.1	2.9	5.1	3.5	4.3	3.1	5.9	6.9	3.9	5.9	4.6	-1.3
	27–30	4.2	3.5	3.8	2.8	4.3	3.9	3.4	3.0	3.8	2.5	3.0	2.1	3.9	3.3	2.6	3.5	4.6	+1.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	0.8	1.3	1.0	1.1	0.9	0.7	0.9	0.9	1.1	0.8	1.4	0.7	0.9	1.3	0.6	0.6	0.6	-0.1
	19–22	0.2	0.5	0.1	0.3	0.6	*	0.3	*	0.3	0.4	0.3	0.6	*	*	0.5	0.1	0.6	+0.4
	23–26	0.7	*	*	0.3	*	0.1	*	*	0.3	0.3	*	*	0.1	0.5	0.1	0.1	0.8	+0.7
	27–30	*	0.1	*	*	*	0.3	*	*	*	*	*	0.3	*	0.3	*	0.1	0.1	+0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take other narcotics^d																			
% saying any	18	22.2	24.8	22.9	23.1	24.0	27.5	21.6	24.6	21.4	23.0	20.7	20.6	21.5	36.3	31.0	28.5	25.8	-2.7
	19–22	13.2	15.2	19.8	23.2	23.0	21.8	21.9	22.6	19.9	17.6	23.7	16.8	15.3	31.4	31.3	25.7	29.5	+3.8
	23–26	9.9	9.4	10.4	11.2	13.5	14.6	18.4	16.8	18.3	17.6	14.2	16.0	19.3	36.7	30.4	27.9	25.6	-2.3
	27–30	7.9	8.3	7.2	8.4	11.2	11.8	11.0	12.0	12.5	13.1	10.6	14.3	14.2	28.4	29.8	32.9	30.4	-2.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.4	2.9	1.8	2.0	2.0	2.1	2.4	2.4	1.9	1.8	2.6	1.3	1.9	3.8	2.6	1.8	1.9	+0.1
	19–22	0.4	0.8	0.4	1.2	1.8	1.3	1.0	0.5	1.0	0.9	0.3	1.3	0.4	1.8	2.1	1.4	2.7	+1.3
	23–26	0.6	0.3	*	0.4	0.5	0.6	0.6	0.5	0.8	0.5	0.3	0.1	0.9	1.6	1.1	1.0	1.6	+0.6
	27–30	*	*	0.2	*	*	0.3	0.1	*	*	0.6	*	0.6	0.6	0.9	1.2	0.1	0.5	+0.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take amphetamines^e																			
% saying any	18	32.7	33.8	30.8	32.9	33.2	34.4	28.1	31.4	28.8	29.0	27.4	27.3	30.0	31.1	31.3	30.5	25.7	-4.8 ss
	19–22	21.1	24.4	25.5	28.4	28.0	28.6	24.0	23.5	25.9	25.4	26.9	19.9	26.6	27.3	29.5	30.5	37.9	+7.4 s
	23–26	14.4	14.1	14.2	14.5	17.5	18.4	18.0	18.8	18.4	19.7	17.6	17.9	21.3	23.8	27.7	26.1	27.0	+0.9
	27–30	12.9	11.0	11.8	11.9	12.9	12.3	12.0	13.5	11.8	12.5	10.0	12.8	16.4	16.4	17.2	22.9	24.7	+1.9
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.4	3.4	2.8	3.1	2.2	2.4	2.1	2.9	2.2	2.0	2.4	1.8	2.0	2.9	2.2	2.4	2.2	-0.2
	19–22	0.7	1.2	0.7	1.7	1.6	1.3	1.2	0.5	0.7	1.1	0.4	1.3	1.6	1.2	4.3	2.0	3.5	+1.5
	23–26	0.8	0.5	0.6	0.3	0.5	0.3	0.7	0.1	0.3	0.7	*	0.1	0.3	0.8	1.3	1.5	1.9	+0.4
	27–30	0.1	0.3	0.6	0.1	0.5	0.9	0.1	*	0.4	0.4	0.4	0.3	0.5	0.3	0.3	0.3	0.1	-0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take sedatives/ barbiturates[†]																			
% saying any		18	30.5	31.1	31.3	28.3	26.6	27.1	25.6	24.3	19.7	20.3	17.4	14.8	16.4	17.8	18.2	17.8	21.6
		19–22	33.2	27.9	27.7	23.6	22.0	17.2	18.8	15.5	14.0	14.1	11.9	12.8	10.7	11.7	9.7	13.3	11.6
		23–26	—	—	—	—	22.2	18.7	16.3	14.1	11.2	10.4	8.9	8.3	8.7	8.2	7.6	9.6	6.9
		27–30	—	—	—	—	—	—	—	—	12.0	8.5	8.8	7.1	6.6	6.7	7.4	7.2	6.7
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	2.6	2.1	1.8	1.7	1.7	1.6	1.4	1.1	1.1	1.4	0.6	0.5	0.6	1.0	1.1	1.4	1.6
		19–22	1.1	1.3	1.0	0.8	0.8	0.5	0.3	0.4	0.8	0.1	0.2	0.3	0.1	0.1	0.3	0.8	0.2
		23–26	—	—	—	—	0.4	0.3	0.3	0.3	0.1	0.2	0.2	0.1	0.1	0.3	0.2	*	*
		27–30	—	—	—	—	—	—	—	—	0.2	*	0.4	0.2	0.2	0.2	*	*	0.3
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take quaaludes																			
% saying any		18	32.5	35.0	35.5	29.7	26.1	26.0	23.5	22.0	17.1	16.6	14.3	12.0	13.1	14.2	14.2	15.5	18.1
		19–22	38.3	36.2	35.4	30.5	24.6	19.9	20.3	16.9	12.5	10.9	10.0	10.6	9.2	10.0	7.8	11.5	10.1
		23–26	—	—	—	—	25.7	21.0	17.4	15.0	12.1	10.3	8.6	5.9	6.4	7.6	7.7	9.0	6.3
		27–30	—	—	—	—	—	—	—	—	11.8	7.9	8.2	7.0	7.1	6.5	6.6	4.5	6.9
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	3.6	3.6	2.6	2.6	1.7	1.3	1.6	1.0	1.0	1.3	0.8	0.5	0.8	1.1	1.1	1.3	1.7
		19–22	1.9	2.7	1.2	1.3	1.2	0.6	0.2	0.4	0.4	0.2	0.6	0.2	0.1	0.1	0.2	0.7	0.1
		23–26	—	—	—	—	0.6	0.3	0.7	0.2	0.2	0.4	0.2	0.1	0.2	0.6	0.2	0.2	*
		27–30	—	—	—	—	—	—	—	—	0.5	0.2	0.2	0.2	*	0.2	*	*	0.2
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take tranquilizers																			
% saying any		18	29.7	29.5	29.9	26.7	26.6	25.8	24.2	23.3	19.9	18.0	14.9	13.5	14.6	15.5	16.5	15.8	18.1
		19–22	37.5	33.9	28.7	22.9	22.0	19.7	20.6	18.0	16.4	14.8	13.4	13.0	11.3	11.9	9.5	13.6	10.5
		23–26	—	—	—	—	29.3	26.3	22.3	20.8	15.5	13.1	14.8	12.1	12.5	11.0	13.4	10.4	10.7
		27–30	—	—	—	—	—	—	—	—	20.1	16.6	16.9	14.9	12.0	12.5	13.9	11.9	11.0
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14.3	12.2	13.1
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.9	1.4	1.1	1.2	1.5	1.2	1.3	1.0	0.7	1.5	0.5	0.4	0.7	0.9	0.9	1.1	1.4
		19–22	0.7	0.9	0.5	0.8	0.3	0.7	0.3	0.6	0.4	0.1	0.4	0.5	0.1	0.1	0.2	0.7	0.7
		23–26	—	—	—	—	0.4	0.3	0.5	*	0.3	0.4	0.2	0.3	0.1	0.4	0.2	*	*
		27–30	—	—	—	—	—	—	—	—	0.5	0.3	0.4	0.2	0.1	0.2	0.4	*	0.2
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.3	0.3
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a																2012–2013 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013
Take sedatives/barbiturates ^f																			
% saying any	18	20.4	22.8	20.9	21.6	22.1	25.3	18.1	25.2	22.3	22.5	20.8	19.8	21.0	23.5	21.1	17.3	15.5	-1.7
	19–22	12.1	14.8	16.0	15.2	18.6	17.1	14.4	18.8	19.6	18.7	20.1	17.8	16.4	19.1	14.5	13.7	19.0	+5.4
	23–26	8.4	7.9	8.3	6.6	11.1	10.9	12.9	16.7	15.7	16.2	16.5	13.4	18.6	17.6	12.2	11.8	14.3	+2.5
	27–30	6.5	6.1	5.7	6.4	7.9	7.4	7.3	11.5	10.5	13.5	12.5	15.2	12.7	15.3	13.7	14.5	16.5	+2.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.1	2.5	1.4	1.7	1.1	1.7	1.9	2.0	1.8	1.3	1.6	1.3	1.3	1.5	1.3	1.5	1.2	-0.3
	19–22	0.7	0.4	0.4	1.0	0.9	0.8	0.7	0.3	0.3	0.2	0.6	0.5	1.0	0.6	0.6	0.3	1.2	+1.0
	23–26	0.8	*	*	0.4	0.4	*	0.2	0.4	0.2	0.5	*	0.4	0.4	0.5	0.3	0.4	0.7	+0.4
	27–30	*	*	0.2	*	0.3	0.6	0.1	*	0.5	0.4	0.6	0.1	0.9	0.4	0.2	*	0.1	+0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take quaaludes																			
% saying any	18	16.1	17.4	15.5	16.2	17.8	18.0	14.2	16.6	13.6	13.4	13.6	11.2	14.3	—	—	—	—	—
	19–22	9.3	10.6	11.4	13.1	14.6	13.0	10.3	8.3	8.2	8.6	8.8	5.9	5.3	—	—	—	—	—
	23–26	6.5	6.6	6.4	4.9	7.7	8.5	8.9	6.5	7.7	5.6	5.6	4.1	8.0	—	—	—	—	—
	27–30	4.9	4.1	5.1	5.0	4.9	6.6	4.3	4.4	3.6	4.9	4.3	5.8	4.5	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.1	2.0	1.4	1.4	1.2	1.2	1.2	1.6	1.3	1.3	1.6	0.8	1.1	—	—	—	—	—
	19–22	0.6	0.5	0.4	0.9	0.8	0.1	0.4	*	0.4	0.2	*	0.2	*	—	—	—	—	—
	23–26	0.8	*	0.2	0.3	0.3	0.1	0.2	0.1	0.2	0.3	0.3	*	0.1	—	—	—	—	—
	27–30	*	*	0.2	0.3	*	0.3	*	*	0.3	0.7	*	0.3	0.5	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take tranquilizers																			
% saying any	18	17.9	19.7	16.4	19.4	18.6	21.2	17.2	18.3	16.9	15.3	15.5	15.0	15.8	16.1	13.9	13.3	11.7	-1.7
	19–22	11.7	13.7	16.2	16.7	21.3	18.1	14.5	12.3	11.5	13.0	17.2	11.6	11.1	11.6	8.2	10.2	12.7	+2.6
	23–26	9.6	8.5	9.8	11.2	12.4	14.9	12.9	15.1	13.1	10.7	12.3	12.6	15.5	13.4	9.9	7.3	9.3	+2.0
	27–30	10.8	12.6	10.4	10.6	9.6	10.6	10.4	9.9	8.5	9.1	12.3	10.3	9.5	9.4	12.6	12.3	-0.3	—
	35	10.8	10.7	11.4	10.8	12.2	12.5	11.4	12.7	12.4	12.2	14.7	16.1	14.8	17.6	17.7	17.9	17.3	-0.6
	40	—	13.7	14.8	15.2	15.1	15.6	15.0	13.6	14.1	16.1	16.0	15.0	15.1	13.6	12.9	15.8	14.5	-1.3
	45	—	—	—	—	—	—	17.3	19.8	15.4	18.3	20.7	17.3	17.5	16.3	16.7	18.8	16.7	-2.1
	50	—	—	—	—	—	—	—	—	—	—	—	19.7	21.0	17.8	19.1	18.1	16.7	-1.4
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17.0	—
% saying most or all	18	0.8	2.3	1.3	2.1	1.3	1.6	1.5	1.7	1.6	1.2	1.8	1.2	1.5	1.4	0.8	0.8	1.0	+0.2
	19–22	0.8	0.6	0.3	0.6	0.9	0.4	0.4	0.3	0.3	0.3	0.3	0.1	0.1	0.6	0.6	0.2	0.6	+0.4
	23–26	1.1	0.1	*	0.5	0.8	0.1	*	0.5	0.7	0.4	*	0.1	0.3	0.2	0.1	0.3	0.3	+0.1
	27–30	*	*	0.4	*	0.4	0.6	0.1	*	0.2	0.2	*	0.1	0.5	*	*	*	0.1	+0.1
	35	0.1	0.2	0.6	0.6	0.2	0.1	0.2	0.3	0.5	0.3	0.5	0.3	0.4	0.6	0.3	0.4	0.4	0.0
	40	—	*	0.4	0.1	0.3	0.2	*	0.2	0.2	0.3	0.3	0.3	0.1	*	*	0.1	*	-0.1
	45	—	—	—	—	—	—	0.3	0.2	0.2	0.1	0.3	*	0.2	0.1	0.1	*	0.2	+0.2
	50	—	—	—	—	—	—	—	—	—	—	—	0.3	0.1	0.1	0.4	0.2	0.1	0.0
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																		
Q. How many of your friends would you estimate...		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	(Years Cont.)
Drink alcoholic beverages																				
% saying any		18	96.1	94.7	95.7	95.5	94.6	94.6	95.6	95.4	95.7	95.1	92.0	91.2	90.5	88.9	90.1	90.9	89.6	
		19-22	96.3	96.7	96.6	97.3	96.8	95.8	96.9	95.6	97.0	97.6	96.1	95.2	93.1	95.1	92.5	94.8	93.7	
		23-26	—	—	—	—	96.8	96.8	96.2	95.9	95.3	95.4	94.7	93.9	95.1	94.4	94.0	94.1	92.7	
		27-30	—	—	—	—	—	—	—	—	96.1	96.0	95.2	94.4	95.6	93.4	93.3	93.3	93.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	89.6	89.9	90.3	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	68.9	67.7	69.7	69.0	66.6	66.0	68.0	71.8	68.1	67.1	60.5	58.6	56.9	57.0	59.6	56.4	56.4	
		19-22	76.6	77.6	75.2	75.1	74.9	71.9	74.2	71.3	73.4	74.1	70.0	71.4	67.4	66.5	68.7	63.9	67.0	
		23-26	—	—	—	—	73.2	74.4	69.5	74.9	68.9	69.8	67.1	69.3	68.8	68.7	70.7	67.0	68.9	
		27-30	—	—	—	—	—	—	—	—	66.7	67.8	62.0	62.7	63.3	61.3	63.2	62.6	64.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	43.8	45.1	49.5	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Get drunk at least once a week																				
% saying any		18	83.1	81.8	83.1	83.9	81.5	82.5	84.7	85.6	84.4	82.8	79.2	79.8	79.9	79.2	81.4	78.9	78.5	
		19-22	80.9	79.9	80.0	80.4	79.8	76.7	82.0	81.1	80.6	80.4	80.1	80.8	76.5	81.1	79.6	83.2	80.9	
		23-26	—	—	—	—	73.1	72.7	73.5	73.7	72.1	73.1	72.2	74.0	73.1	74.3	72.1	73.1	74.5	
		27-30	—	—	—	—	—	—	—	—	66.3	61.8	65.4	65.2	65.5	64.5	62.7	67.1	66.7	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	44.3	43.2	44.9	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	30.1	29.4	29.9	31.0	29.6	29.9	31.8	31.3	29.6	31.1	27.5	29.7	28.6	27.6	28.4	27.4	29.0	
		19-22	21.9	23.3	22.0	20.2	22.7	21.7	20.8	21.3	24.0	22.6	23.6	24.9	22.6	28.8	26.3	28.2	26.0	
		23-26	—	—	—	—	11.4	11.6	12.5	11.9	12.8	12.0	13.9	11.6	14.6	13.2	15.2	15.2	14.0	
		27-30	—	—	—	—	—	—	—	—	5.2	6.3	6.7	6.6	5.9	6.7	6.4	7.9	8.6	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.6	3.6	5.4	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Smoke cigarettes																				
% saying any		18	90.6	88.5	88.3	87.0	86.0	87.0	87.8	88.3	87.7	86.5	84.9	85.7	84.4	84.8	88.1	87.9	88.3	
		19-22	94.4	94.3	93.4	93.1	91.9	91.6	91.1	90.3	89.3	90.0	86.1	86.1	86.7	86.7	86.1	88.8	89.2	
		23-26	—	—	—	—	93.9	95.0	91.6	92.1	89.8	90.1	88.7	89.6	85.6	88.3	86.4	86.8	85.3	
		27-30	—	—	—	—	—	—	—	—	92.6	89.8	90.7	90.4	88.0	85.8	84.8	84.9	85.4	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	72.7	71.7	71.7	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	23.3	22.4	24.1	22.4	19.2	22.8	21.5	21.0	20.2	23.1	21.4	21.8	21.4	25.0	25.3	27.5	30.4	
		19-22	31.8	27.6	25.6	25.2	25.6	22.7	21.9	22.5	19.3	19.9	19.2	20.2	20.3	22.2	21.7	28.4	24.0	
		23-26	—	—	—	—	25.6	22.7	19.7	18.5	16.5	20.5	16.9	18.1	16.0	15.5	16.6	13.9	17.6	
		27-30	—	—	—	—	—	—	—	—	15.8	14.2	11.6	12.9	11.9	14.3	10.9	12.3	10.4	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.9	7.2	9.3	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																	2012–2013 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013		
Drink alcoholic beverages																				
% saying any	18	90.7	91.2	90.2	89.8	89.2	88.0	87.9	87.8	87.2	86.0	85.1	85.2	83.7	83.9	82.6	82.0	82.0	0.0	
	19–22	94.5	94.5	92.8	95.2	93.4	94.5	92.5	90.4	95.0	91.9	94.0	92.6	93.2	90.9	88.9	93.3	92.0	-1.3	
	23–26	95.4	95.5	93.3	94.5	93.1	95.3	92.8	94.9	91.6	93.6	94.7	93.3	95.0	95.3	95.3	92.3	92.5	+0.2	
	27–30	95.1	93.1	94.4	92.7	91.4	92.8	90.5	94.4	93.7	95.6	92.4	91.7	93.9	93.0	92.5	93.4	91.6	-1.8	
	35	89.5	88.1	88.7	89.6	89.3	90.1	87.4	93.4	91.3	90.6	90.5	91.0	90.4	93.3	93.0	92.7	93.2	+0.5	
	40	—	88.4	88.9	90.7	89.6	90.5	89.2	90.5	92.1	90.8	93.0	89.3	92.6	92.1	92.4	91.3	91.9	+0.5	
	45	—	—	—	—	—	—	87.9	90.3	89.8	90.1	89.8	90.5	89.5	90.6	90.8	90.1	91.4	+1.3	
	50	—	—	—	—	—	—	—	—	—	—	—	—	88.9	90.2	89.9	90.4	90.1	89.2	-0.9
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	87.6	—
% saying most or all	18	60.9	61.0	58.2	57.2	59.2	53.7	53.1	53.9	55.3	52.4	52.0	51.6	50.5	51.4	50.3	49.4	46.9	-2.6	
	19–22	63.8	69.4	67.8	70.1	65.4	68.8	63.9	66.4	71.8	65.4	71.1	64.4	69.7	69.1	63.3	66.3	63.2	-3.1	
	23–26	66.6	67.4	63.6	70.8	65.7	73.4	66.0	71.3	69.3	69.2	70.2	76.3	76.9	75.5	79.7	74.3	73.7	-0.6	
	27–30	66.6	62.9	64.4	64.8	64.9	66.3	61.5	69.0	66.2	70.7	65.6	67.1	74.0	72.2	70.9	74.9	72.9	-2.0	
	35	46.6	47.1	46.0	49.1	48.4	52.9	51.6	53.7	55.5	55.2	56.1	55.7	53.2	56.9	61.9	58.7	62.1	+3.5	
	40	—	37.7	41.4	42.5	44.7	44.8	47.2	43.3	47.2	45.9	50.3	48.9	54.5	54.7	54.3	55.9	56.6	+0.6	
	45	—	—	—	—	—	—	38.9	41.7	42.4	45.1	46.6	47.0	45.9	46.7	47.2	53.5	52.0	-1.5	
	50	—	—	—	—	—	—	—	—	—	—	—	—	37.7	39.3	41.9	43.5	45.8	48.2	+2.3
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	39.1	—
Get drunk at least once a week																				
% saying any	18	82.4	81.1	81.5	79.5	79.6	78.3	77.3	79.0	78.7	77.4	75.5	76.2	76.2	73.5	71.9	68.9	69.9	+1.0	
	19–22	79.2	82.3	82.8	82.2	81.9	81.5	81.5	80.5	85.1	81.7	84.4	81.3	82.8	81.2	78.3	83.6	77.7	-5.8	
	23–26	71.9	74.1	71.0	76.5	74.7	81.0	76.4	75.8	80.7	80.9	80.4	79.5	83.0	83.7	83.9	79.7	83.1	+3.4	
	27–30	65.4	65.5	65.9	64.3	64.7	68.9	66.5	73.8	72.4	74.6	72.0	71.7	78.7	78.2	78.3	80.1	74.4	-5.7	
	35	42.9	46.1	44.5	46.9	47.6	48.3	47.9	52.0	50.7	52.6	55.0	56.0	56.0	59.2	63.2	62.4	63.9	+1.5	
	40	—	41.6	40.6	42.2	41.3	42.6	42.9	43.2	48.4	47.2	46.3	48.2	53.7	49.6	48.5	54.9	54.7	-0.2	
	45	—	—	—	—	—	—	41.6	42.2	41.6	40.0	42.7	45.7	45.4	49.1	45.9	50.0	50.5	+0.5	
	50	—	—	—	—	—	—	—	—	—	—	—	—	40.0	38.3	39.6	42.4	42.5	45.0	+2.5
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	35.1	—
% saying most or all	18	30.9	31.7	30.1	32.4	32.7	28.3	27.1	27.6	28.5	27.7	27.0	25.2	24.4	23.7	23.8	21.2	20.7	-0.6	
	19–22	26.6	29.8	29.3	28.1	30.2	31.0	29.6	29.0	31.2	32.9	32.0	28.9	31.4	27.7	27.6	27.2	28.1	+0.9	
	23–26	17.0	16.0	16.8	17.4	19.1	19.2	18.3	24.0	24.0	20.3	22.8	23.1	23.2	24.0	22.6	20.0	23.4	+3.4	
	27–30	7.7	9.3	12.1	9.8	11.7	8.9	13.0	9.4	11.2	13.5	12.2	10.9	17.1	13.7	13.2	13.5	13.2	-0.3	
	35	3.2	4.4	4.9	4.6	4.8	4.5	5.2	5.3	5.3	5.6	6.1	7.3	5.9	7.4	8.4	6.8	8.3	+1.5	
	40	—	2.8	3.0	2.5	2.9	3.8	3.9	3.0	3.6	4.0	3.4	4.8	4.6	4.8	4.8	4.3	4.2	-0.1	
	45	—	—	—	—	—	—	3.6	2.7	2.7	3.1	3.7	4.1	3.2	3.2	3.5	4.3	5.1	+0.8	
	50	—	—	—	—	—	—	—	—	—	—	—	—	3.2	2.7	2.0	2.9	2.5	3.6	+1.1
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.0	—
Smoke cigarettes																				
% saying any	18	89.9	89.5	89.3	87.2	86.8	85.4	83.3	83.7	81.8	81.4	77.1	78.4	79.6	78.0	75.4	74.3	72.1	-2.1	
	19–22	91.3	92.6	91.0	90.9	89.9	89.7	86.5	89.7	89.3	85.8	86.8	84.4	88.3	81.8	79.4	78.2	77.4	-0.8	
	23–26	85.4	88.7	84.1	86.5	86.7	86.4	86.5	87.0	87.3	85.4	84.1	86.8	85.3	87.7	86.5	83.1	80.3	-2.7	
	27–30	84.1	81.1	86.3	85.1	84.9	87.0	82.8	83.5	81.0	84.4	81.7	82.1	84.1	84.6	83.8	85.2	81.6	-3.7	
	35	72.4	71.8	69.9	70.8	69.2	66.6	67.0	67.7	65.5	67.0	64.8	67.6	62.2	65.4	66.1	66.4	63.2	-3.2	
	40	—	70.2	70.0	67.8	64.3	65.5	65.1	62.4	63.8	64.6	59.2	59.7	60.5	57.4	57.4	56.7	59.1	+2.3	
	45	—	—	—	—	—	—	66.1	67.0	62.9	60.9	58.5	56.1	57.7	60.6	58.0	57.4	54.3	-3.1	
	50	—	—	—	—	—	—	—	—	—	—	—	—	62.1	61.3	59.2	55.9	57.4	54.7	-2.7
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	56.5	—
% saying most or all	18	34.4	33.9	31.1	28.2	25.0	23.0	19.6	20.6	16.7	15.8	16.4	13.9	14.1	14.9	14.1	12.2	11.0	-1.3	
	19–22	25.1	28.8	26.8	29.4	27.0	25.7	20.2	20.7	20.4	15.2	17.9	12.9	15.3	16.7	13.7	13.6	10.8	-2.8	
	23–26	17.0	16.8	17.5	17.0	15.5	15.1	18.3	19.8	19.6	13.9	14.7	15.0	13.4	15.0	11.1	10.6	13.5	+2.9	
	27–30	12.1	12.3	13.4	11.7	10.2	12.9	12.2	9.2	12.6	12.6	12.7	10.8	12.4	7.9	7.4	10.0	6.8	-3.2	
	35	7.2	8.0	9.0	6.7	8.8	6.6	6.3	6.9	6.0	6.8	5.7	5.9	6.4	6.8	6.2	5.5	4.9	-0.5	
	40	—	8.1	7.4	6.8	5.7	5.8	5.9	6.0	7.0	5.1	4.7	4.5	3.9	4.0	4.2	2.9	3.8	+0.9	
	45	—	—	—	—	—	—	5.7	5.9	6.1	5.4	4.5	3.7	4.8	5.2	3.8	3.4	4.6	+1.2	
	50	—	—	—	—	—	—	—	—	—	—	—	—	4.0	4.3	4.2	3.6	2.6	2.3	-0.3
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.3	—



(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	→
Q. How many of your friends would you estimate. . .	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	(Years Cont.)
Take steroids																			
% saying any	18	—	—	—	—	—	—	—	—	—	—	25.9	24.7	21.5	19.0	18.1	19.5	17.9	
	19–22	—	—	—	—	—	—	—	—	—	23.4	21.5	22.2	19.7	20.7	16.8	16.6	16.1	
	23–26	—	—	—	—	—	—	—	—	—	15.3	15.0	12.3	14.5	11.1	10.5	12.4	7.3	
	27–30	—	—	—	—	—	—	—	—	—	9.9	10.5	7.5	8.0	8.0	8.0	8.0	10.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all	18	—	—	—	—	—	—	—	—	—	—	1.8	1.0	1.7	0.9	1.2	1.3	0.8	
	19–22	—	—	—	—	—	—	—	—	—	0.2	0.6	*	0.1	0.4	0.2	0.1	*	
	23–26	—	—	—	—	—	—	—	—	—	0.4	*	*	0.2	0.1	0.1	*	*	
	27–30	—	—	—	—	—	—	—	—	—	0.5	*	*	*	0.2	0.1	*	*	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Approximate Weighted N =	18	2,987	3,307	3,303	3,095	2,945	2,971	2,798	2,948	2,961	2,587	2,361	2,339	2,373	2,410	2,337	2,379	2,156	
	19–22	576	592	564	579	543	554	579	572	562	579	556	526	510	468	435	470	469	
	23–26					527	534	546	528	528	506	510	507	516	495	449	456	416	
	27–30									516	507	499	476	478	461	419	450	464	
	35															1,200	1,187	1,187	
	40																		
	45																		
	50																		
	55																		

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

		Percentage saying friends use ^a																	2012– 2013	
Q. How many of your friends would you estimate. . .		Age Group	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	change
Take steroids																				
% saying any		18	18.9	18.3	20.0	19.8	21.7	21.6	21.1	22.8	19.1	19.8	20.1	19.4	19.3	16.4	16.0	18.7	17.4	-1.3
		19–22	16.8	20.0	20.6	18.9	20.0	19.3	17.1	21.4	20.1	21.0	18.3	14.8	16.8	13.8	15.3	12.6	11.1	-1.4
		23–26	13.0	9.2	15.0	12.2	13.6	14.3	12.9	12.4	11.6	13.4	13.8	13.3	12.8	11.7	13.9	10.0	11.6	+1.6
		27–30	9.1	7.0	11.2	9.3	10.7	6.4	11.6	10.1	7.4	7.5	6.7	6.6	12.0	9.2	8.5	11.6	10.0	-1.5
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.7	1.4	0.9	1.9	1.2	1.5	1.5	2.6	1.5	0.9	1.2	1.3	1.5	1.7	1.1	1.8	1.5	-0.3
		19–22	0.1	0.3	0.1	0.3	0.7	0.7	0.4	*	0.1	0.3	0.3	0.3	*	0.7	0.6	0.4	0.7	+0.3
		23–26	0.5	*	0.1	0.3	0.2	0.1	*	0.1	0.3	0.3	*	*	0.7	*	0.1	0.1	0.3	+0.2
		27–30	*	*	*	*	*	0.3	*	*	0.1	*	*	*	0.3	*	*	*	*	0.0
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Approximate Weighted N =		18	2,292	2,313	2,060	1,838	1,923	1,968	2,233	2,271	2,266	2,266	2,253	2,125	2,110	2,195	2,208	2,144	1,973	
		19–22	467	437	426	402	402	375	388	443	395	377	362	375	382	376	353	348	340	
		23–26	419	394	414	387	403	358	362	411	361	336	340	355	311	359	314	330	328	
		27–30	454	428	424	363	359	348	369	396	363	350	324	332	309	340	325	333	284	
		35	1,209	1,067	1,071	1,033	1,005	918	968	985	1,041	953	884	905	974	922	858	877	848	
		40		1,098	1,156	1,144	1,119	1,083	945	1,004	975	951	896	924	905	952	877	852	844	
		45							976	1,074	1,052	1,009	999	904	937	889	887	874	844	
		50												940	1,009	1,016	974	987	840	
		55																	880	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '—' indicates data not available. '*' indicates a prevalence rate of less than 0.05%.

^aAnswer alternatives were: (1) None, (2) A few, (3) Some, (4) Most, (5) All. The any percentage combines categories (2)–(5). The most or all percentage combines categories (4) and (5).

^bFor the young adult sample, any illicit drug includes all of the drugs listed in this table except cigarettes and alcohol. For the 35-, 40-, 45-, and 50-year-olds, any illicit drug includes marijuana, tranquilizers, crack, cocaine powder, and other illicit drugs.

^cIn 2001 the question text was changed from other psychedelics to other hallucinogens, and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^dIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

^eIn 2011 pep pills and bennies were replaced in the list of examples by Adderall and Ritalin. This change likely explains the discontinuity in the 2011 results.

^fIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

TABLE 7-3
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a																	(Years Cont.)
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Any illicit drug^b																			
% saying any	18	84.3	82.7	81.4	79.4	77.9	77.7	75.5	73.9	71.3	68.6	67.6	64.2	61.3	66.1	70.8	75.3	78.0	
	19–22	80.6	81.0	81.5	76.5	76.3	77.4	74.6	72.7	69.5	61.5	60.8	58.9	58.6	58.4	60.7	66.4	67.2	
	23–26	—	—	—	—	68.9	70.2	68.0	62.4	62.7	58.3	54.6	52.1	48.2	49.9	47.1	54.2	50.3	
	27–30	—	—	—	—	—	—	—	—	52.4	50.2	47.0	39.6	41.7	38.9	45.6	42.4	44.9	
% saying often exposed	18	36.3	36.1	31.4	29.8	28.3	27.2	26.3	23.3	20.8	22.0	20.7	18.2	18.0	24.0	29.3	32.3	33.8	
	19–22	34.6	34.0	32.1	24.4	24.4	23.7	21.1	18.9	19.9	16.2	16.4	17.6	21.4	16.1	18.1	23.7	20.4	
	23–26	—	—	—	—	20.7	23.3	18.5	17.4	18.2	13.8	13.7	13.3	12.2	11.1	11.1	12.5	12.8	
	27–30	—	—	—	—	—	—	—	—	13.7	12.0	10.8	8.2	10.5	9.0	12.5	8.5	10.1	
Any illicit drug other than marijuana^b																			
% saying any	18	58.5	62.6	62.5	59.4	59.8	59.3	55.3	51.7	47.8	47.1	45.4	40.0	41.6	42.6	45.3	47.2	49.7	
	19–22	56.9	58.4	61.6	54.9	57.1	53.3	53.4	48.5	46.4	36.5	39.4	33.8	37.1	29.4	33.9	36.8	36.5	
	23–26	—	—	—	—	51.5	51.9	51.5	43.6	42.9	36.8	34.0	30.0	27.3	27.8	24.9	26.8	23.2	
	27–30	—	—	—	—	—	—	—	—	35.8	33.7	31.5	25.8	26.6	24.2	25.8	21.1	21.8	
% saying often exposed	18	14.1	17.1	16.6	14.2	14.6	12.9	12.1	10.2	9.6	10.7	9.2	7.9	7.5	9.6	9.4	11.1	12.1	
	19–22	11.8	15.6	13.5	11.1	10.7	10.2	8.2	8.1	7.5	6.7	4.5	4.4	5.5	4.1	5.1	7.7	3.9	
	23–26	—	—	—	—	9.0	10.4	9.3	8.5	6.7	5.0	5.1	3.5	2.6	3.0	2.2	3.5	3.4	
	27–30	—	—	—	—	—	—	—	—	6.0	4.7	4.1	3.2	3.7	2.4	3.4	2.9	3.4	
Marijuana																			
% saying any	18	82.0	80.2	77.9	76.2	74.4	73.5	72.0	70.4	67.0	64.8	63.4	59.6	56.8	61.0	67.2	72.7	75.6	
	19–22	79.8	79.8	78.7	72.7	74.1	75.5	72.4	70.5	66.3	59.3	57.5	55.0	56.4	55.4	56.8	64.0	64.8	
	23–26	—	—	—	—	65.3	66.0	64.1	59.0	57.6	55.0	50.6	47.9	44.6	45.9	44.4	51.0	47.8	
	27–30	—	—	—	—	—	—	—	—	49.1	47.4	42.1	36.0	38.2	35.3	41.9	38.3	41.8	
% saying often exposed	18	33.8	33.1	28.0	26.1	24.8	24.2	24.0	20.6	17.9	19.5	17.8	16.0	15.6	20.9	27.6	30.7	31.8	
	19–22	32.6	30.5	30.3	21.1	21.9	20.3	18.6	16.4	18.3	14.2	14.7	15.9	19.9	14.7	17.0	22.1	20.3	
	23–26	—	—	—	—	17.5	20.6	14.6	14.8	15.6	11.6	11.2	11.6	10.9	10.4	10.4	11.1	11.5	
	27–30	—	—	—	—	—	—	—	—	10.9	9.8	8.5	6.7	8.9	7.6	10.7	7.4	9.1	
LSD																			
% saying any	18	17.2	17.4	16.1	13.8	12.5	13.2	13.1	12.9	13.4	15.0	14.9	15.7	17.8	21.0	24.2	26.1	27.6	
	19–22	17.4	15.8	16.0	13.5	12.8	12.7	10.8	10.9	12.0	12.0	12.1	13.1	19.3	13.4	16.5	18.6	20.7	
	23–26	—	—	—	—	8.3	9.3	8.8	7.3	6.3	6.7	8.4	8.6	8.8	7.8	8.4	9.9	8.6	
	27–30	—	—	—	—	—	—	—	—	3.6	3.2	3.3	3.6	3.9	4.9	5.3	5.5	4.3	
% saying often exposed	18	1.4	2.0	1.9	1.4	1.5	1.3	1.6	1.8	1.6	2.2	2.6	2.9	3.0	3.9	4.2	6.1	4.7	
	19–22	1.4	1.5	1.4	0.6	0.8	0.7	0.5	1.2	0.6	1.1	1.2	1.0	2.0	1.1	0.4	3.6	1.4	
	23–26	—	—	—	—	0.3	0.4	0.4	0.7	0.6	0.3	0.5	0.2	0.8	0.3	0.5	0.5	0.4	
	27–30	—	—	—	—	—	—	—	—	0.3	0.2	0.5	0.2	0.2	0.5	0.5	0.2	0.2	
Other hallucinogens^c																			
% saying any	18	20.4	17.6	16.8	13.1	12.7	12.5	11.8	10.0	9.0	8.8	9.4	9.4	9.7	12.1	14.0	15.8	16.6	
	19–22	18.3	16.3	16.3	12.5	10.5	11.0	9.2	9.1	7.7	8.4	8.3	8.9	10.6	6.7	8.3	12.8	13.1	
	23–26	—	—	—	—	8.4	8.9	9.1	6.0	5.1	4.8	5.7	5.5	5.1	5.7	5.2	5.5	6.9	
	27–30	—	—	—	—	—	—	—	—	5.0	3.4	3.4	3.4	2.1	3.7	3.4	4.2	3.2	
% saying often exposed	18	2.2	2.0	2.6	1.1	1.7	1.4	1.5	1.2	1.1	1.3	1.2	1.3	1.1	1.9	2.3	2.5	2.7	
	19–22	1.1	0.9	0.9	0.7	0.8	0.8	0.2	0.8	0.3	0.4	0.4	0.5	0.7	0.4	0.2	1.6	0.7	
	23–26	—	—	—	—	0.1	0.3	0.5	0.6	0.8	0.1	0.4	0.4	*	0.2	0.4	0.3	0.3	
	27–30	—	—	—	—	—	—	—	—	0.2	0.4	0.5	0.3	0.1	0.5	0.2	0.3	0.2	

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Any illicit drug^b																			
% saying any	18	78.8	77.2	77.9	76.0	76.5	76.5	73.6	74.3	73.0	73.7	70.8	71.9	74.1	76.0	76.6	76.4	75.4	-1.0
	19–22	65.3	69.1	65.8	64.7	69.7	65.7	68.0	67.6	68.8	67.1	67.4	66.2	69.8	66.0	68.3	70.4	72.1	+1.6
	23–26	55.4	50.6	50.5	55.1	56.4	56.5	57.0	53.5	53.9	56.7	58.3	56.3	57.7	56.3	62.6	67.0	65.2	-1.9
	27–30	41.6	37.5	41.1	40.8	42.2	47.0	46.7	43.3	45.7	48.4	44.1	48.7	42.5	49.3	51.6	58.9	57.2	-1.7
% saying often exposed	18	34.7	33.2	35.6	32.6	33.6	32.6	31.8	30.4	29.9	29.7	27.8	28.6	31.4	33.2	34.6	34.9	32.3	-2.5
	19–22	25.3	24.2	24.0	21.3	26.1	25.2	26.5	26.8	25.2	24.2	22.8	20.1	23.7	26.5	24.8	27.3	24.6	-2.7
	23–26	14.3	14.2	15.0	15.9	16.4	15.9	17.8	15.1	18.7	14.9	18.9	15.4	14.9	18.8	19.4	21.2	20.8	-0.4
	27–30	10.3	8.5	9.6	9.4	10.4	13.8	13.9	10.3	14.5	13.2	9.7	9.7	12.1	13.2	13.6	15.7	18.5	+2.8
Any illicit drug other than marijuana^b																			
% saying any	18	47.9	47.3	46.5	47.2	49.9	49.3	46.3	48.3	45.9	45.4	45.4	43.8	44.3	47.2	46.6	45.0	44.2	-0.8
	19–22	39.4	40.0	36.4	38.1	39.2	38.0	40.2	40.9	41.1	38.5	42.7	38.2	37.1	38.5	38.5	41.8	38.9	-2.8
	23–26	25.6	27.1	28.0	31.0	31.4	31.5	32.2	32.6	32.3	34.5	33.1	31.3	33.0	34.8	39.9	37.8	37.4	-0.4
	27–30	21.4	15.4	19.5	17.2	22.2	23.1	26.1	23.2	27.1	27.4	24.8	27.7	22.8	29.3	33.4	35.2	34.4	-0.7
% saying often exposed	18	11.7	9.9	11.7	10.5	11.9	12.6	10.8	11.4	10.6	11.4	10.8	8.2	9.4	10.2	11.5	11.6	9.3	-2.3
	19–22	7.6	7.0	4.8	6.4	7.8	8.6	5.2	7.9	8.0	6.7	6.9	6.6	6.8	6.6	6.9	10.1	8.1	-2.0
	23–26	3.1	3.1	4.3	3.5	3.4	5.0	5.4	5.4	4.0	5.4	6.7	5.4	3.8	6.4	6.3	7.6	5.8	-1.8
	27–30	3.2	1.0	2.5	1.6	3.7	4.7	4.9	2.4	5.6	4.0	3.4	2.3	3.0	4.8	4.2	4.5	4.8	+0.3
Marijuana																			
% saying any	18	76.8	75.5	75.8	73.8	74.9	74.2	71.4	72.2	70.8	71.4	68.4	69.8	71.8	74.2	74.6	75.1	73.7	-1.5
	19–22	63.4	67.1	63.5	63.9	68.0	64.6	64.8	65.1	66.8	65.4	66.3	64.3	67.5	64.9	65.7	67.6	69.0	+1.5
	23–26	53.1	48.8	48.1	51.8	54.2	53.5	54.4	50.6	49.7	51.9	53.3	54.0	55.5	54.0	57.9	63.9	63.4	-0.4
	27–30	39.1	35.7	38.7	38.8	37.0	44.6	44.1	40.4	42.4	44.1	40.7	44.8	39.8	43.5	46.1	56.0	52.3	-3.7
% saying often exposed	18	32.9	31.4	34.4	30.3	30.8	30.7	30.4	28.0	27.0	27.8	25.1	27.0	29.3	31.3	32.3	32.2	30.8	-1.5
	19–22	23.7	22.8	23.0	20.4	24.5	24.8	24.2	24.5	23.6	23.1	20.1	18.3	22.6	25.2	22.9	24.2	22.6	-1.5
	23–26	12.9	13.6	13.2	15.2	15.6	14.9	16.2	13.7	17.8	12.5	16.2	13.7	13.5	17.0	18.0	19.7	18.3	-1.4
	27–30	8.9	8.1	8.8	8.6	8.4	11.7	11.7	9.6	12.2	11.5	8.2	8.5	12.3	10.8	10.9	13.9	16.0	+2.1
LSD																			
% saying any	18	25.9	23.1	23.6	22.0	21.6	17.2	14.2	12.4	10.8	11.6	12.4	12.1	11.9	14.1	13.5	13.0	13.8	+0.8
	19–22	22.3	21.0	20.1	15.9	15.2	13.6	10.0	8.5	7.2	10.4	6.3	9.2	9.1	9.7	10.1	12.2	10.0	-2.2
	23–26	7.6	9.8	9.4	9.8	11.1	9.3	5.5	4.4	4.7	5.6	4.5	4.8	3.7	5.7	8.9	9.6	8.3	-1.2
	27–30	3.9	3.2	3.7	3.2	4.3	4.8	3.0	4.7	4.0	3.4	3.9	1.7	3.8	4.2	4.1	4.2	4.7	+0.5
% saying often exposed	18	5.1	3.2	4.1	3.3	2.8	2.6	1.8	1.6	1.5	1.9	1.7	0.8	1.3	1.4	1.4	1.6	1.5	-0.1
	19–22	1.8	2.0	1.7	1.4	2.4	0.9	0.2	0.1	0.7	0.7	0.3	0.7	0.3	0.1	0.2	0.6	0.9	+0.3
	23–26	0.2	0.1	0.3	0.2	*	0.3	0.3	*	0.3	*	0.5	0.6	*	0.6	0.3	1.4	0.1	-1.3
	27–30	*	*	0.1	*	*	*	0.3	0.3	0.6	*	0.1	*	0.3	0.5	0.5	0.6	1.0	+0.4
Other hallucinogens^c																			
% saying any	18	17.8	15.9	17.7	16.3	28.1	26.4	25.8	24.8	24.3	23.8	23.5	23.6	22.0	25.0	23.8	22.7	22.3	-0.4
	19–22	15.0	15.0	12.4	11.8	22.8	23.4	18.9	18.7	19.5	17.8	20.2	17.5	17.5	19.6	17.5	17.0	14.6	-2.4
	23–26	5.6	8.7	5.8	8.9	14.8	14.7	11.9	10.1	11.3	10.3	9.8	9.8	9.9	12.5	13.8	13.6	14.6	+1.0
	27–30	2.9	2.6	3.0	3.0	6.4	7.7	6.3	7.9	8.8	7.8	6.8	5.2	7.5	5.0	8.1	7.8	7.2	-0.6
% saying often exposed	18	2.8	1.7	2.7	2.1	3.6	4.5	3.2	3.2	2.6	4.1	3.0	1.9	2.7	2.2	2.5	2.7	2.4	-0.3
	19–22	0.7	0.5	0.6	0.8	2.6	2.4	0.4	0.7	1.2	0.7	0.7	0.9	0.9	1.1	1.3	0.8	*	-0.8
	23–26	0.2	*	*	0.4	0.2	0.4	*	*	0.5	*	0.6	0.7	0.1	0.7	0.3	1.1	0.3	-0.8
	27–30	0.5	*	0.1	*	0.4	*	*	0.3	0.6	*	0.4	0.3	0.3	0.7	0.5	0.8	0.9	+0.1

↓
(List of drugs continued.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

Percentage saying exposed to drug ^a

Age Group	Percentage saying exposed to drug ^a																	(Years Cont.)
	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Cocaine																		
% saying any	18	37.7	36.3	34.9	33.3	35.6	38.3	37.4	34.9	30.2	30.2	27.7	21.3	19.8	19.2	18.8	21.6	25.0
	19–22	37.6	42.3	43.6	36.6	38.9	39.4	41.5	37.0	36.2	26.6	24.0	18.5	19.8	13.5	14.7	14.1	19.3
	23–26	—	—	—	—	38.5	40.6	42.0	34.5	35.9	28.0	24.0	19.9	16.7	14.6	14.3	14.1	12.5
	27–30	—	—	—	—	—	—	—	—	28.9	28.3	24.2	18.6	19.4	16.6	14.3	11.4	12.1
% saying often exposed	18	5.9	6.6	6.6	5.2	6.7	7.1	7.8	5.9	5.1	5.4	4.7	3.4	2.7	2.9	2.5	3.2	4.0
	19–22	5.8	7.6	6.5	4.3	6.5	7.0	5.4	5.2	4.8	4.3	2.2	1.6	1.7	1.7	1.8	1.7	1.2
	23–26	—	—	—	—	5.3	8.5	7.0	6.0	5.4	3.5	2.5	1.7	1.4	1.7	1.0	1.7	1.3
	27–30	—	—	—	—	—	—	—	—	4.4	3.9	2.9	2.2	2.0	1.2	1.5	1.4	1.9
Heroin																		
% saying any	18	7.4	6.6	7.1	5.1	6.0	5.5	6.0	5.8	5.7	6.5	5.4	5.1	5.4	5.7	7.3	7.9	8.6
	19–22	4.4	3.3	4.1	2.9	3.1	4.8	2.9	2.9	2.9	2.5	3.0	2.7	2.0	3.7	3.8	3.6	
	23–26	—	—	—	—	2.3	3.3	3.2	2.9	1.7	2.3	2.3	1.8	1.7	1.5	1.9	2.8	2.9
	27–30	—	—	—	—	—	—	—	—	2.1	1.4	1.5	0.9	1.0	2.0	2.0	1.7	1.5
% saying often exposed	18	0.4	0.6	1.0	0.7	1.1	0.5	1.0	0.9	0.8	1.0	0.5	0.9	0.7	1.1	0.7	1.2	1.6
	19–22	0.2	0.3	0.3	0.1	0.2	0.5	0.2	0.1	0.2	0.1	0.2	0.4	0.6	0.4	0.6	1.2	0.2
	23–26	—	—	—	—	*	0.7	0.3	0.6	0.4	0.3	0.6	0.3	*	*	*	0.2	0.2
	27–30	—	—	—	—	—	—	—	—	0.3	0.3	0.5	0.2	0.2	0.9	0.3	0.6	0.6
Other narcotics ^d																		
% saying any	18	19.6	17.5	18.5	17.3	18.0	18.4	15.6	14.4	14.8	13.8	14.2	11.3	11.1	12.4	14.9	15.5	18.5
	19–22	14.4	14.4	15.2	10.9	12.4	13.7	9.8	12.2	11.2	9.0	9.4	9.2	8.5	6.8	10.1	12.1	11.5
	23–26	—	—	—	—	9.0	12.3	9.2	9.7	7.4	8.0	5.9	8.3	7.0	4.6	6.9	7.8	7.4
	27–30	—	—	—	—	—	—	—	—	6.5	6.5	5.8	5.5	3.7	5.6	5.9	5.7	4.7
% saying often exposed	18	1.7	1.7	2.4	2.2	2.0	1.8	2.1	1.7	1.7	1.7	1.6	1.4	1.3	1.7	1.7	2.1	3.4
	19–22	0.7	0.5	0.5	0.9	0.7	1.0	0.5	0.4	0.9	0.3	0.2	1.0	0.9	0.6	0.8	1.4	0.7
	23–26	—	—	—	—	0.4	0.5	1.3	0.8	0.8	0.5	1.6	0.7	0.1	0.3	0.1	0.1	0.3
	27–30	—	—	—	—	—	—	—	—	0.7	0.5	1.0	0.3	0.8	1.2	0.8	0.8	0.7
Amphetamines																		
% saying any	18	40.8	49.5	50.2	46.1	45.0	41.0	36.5	31.7	27.9	27.4	28.3	23.6	24.5	24.7	28.2	28.1	31.5
	19–22	42.3	48.6	48.4	39.7	41.3	35.9	31.3	26.7	21.2	18.5	19.5	17.4	21.3	15.1	20.3	21.0	22.3
	23–26	—	—	—	—	32.3	30.5	29.1	20.9	18.8	14.0	16.8	14.6	11.8	13.2	11.2	13.0	11.1
	27–30	—	—	—	—	—	—	—	—	15.6	14.3	13.5	10.7	11.4	11.3	11.0	10.6	7.6
% saying often exposed	18	8.3	12.1	12.3	10.1	9.0	6.5	5.8	4.5	4.1	4.7	4.1	3.1	3.0	3.9	4.1	4.5	5.6
	19–22	7.4	9.9	7.7	6.9	5.4	4.4	3.1	3.3	2.2	1.5	1.1	1.9	2.6	1.5	3.3	5.0	1.3
	23–26	—	—	—	—	3.9	3.2	2.2	3.3	1.9	0.7	2.0	1.3	0.2	0.8	0.9	1.6	1.3
	27–30	—	—	—	—	—	—	—	—	2.0	2.0	1.2	0.8	0.8	1.3	0.7	1.6	1.8

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a																2012–2013 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013
Cocaine																			
% saying any	18	25.6	26.6	25.8	24.2	24.5	24.9	24.8	24.4	25.7	28.2	25.2	24.1	20.0	20.0	19.3	17.4	16.7	-0.6
	19–22	18.8	21.6	18.5	19.1	20.6	22.5	18.4	23.6	22.7	22.9	22.5	22.7	18.6	17.8	15.5	18.9	11.5	-7.4 s
	23–26	14.0	16.0	18.2	16.4	16.9	18.3	17.4	18.7	19.2	19.3	19.0	18.2	15.3	14.7	20.5	17.2	14.4	-2.8
	27–30	11.4	8.6	11.6	10.2	11.6	12.2	12.6	13.0	15.8	16.0	14.1	14.8	13.2	11.4	13.1	14.2	15.0	+0.8
% saying often exposed	18	4.2	3.7	4.6	4.6	4.5	5.3	5.0	4.7	4.2	5.3	4.6	3.6	2.6	2.1	2.3	2.8	2.1	-0.7
	19–22	2.4	3.2	1.4	3.8	3.0	4.1	1.6	2.6	4.0	2.6	1.8	2.6	2.6	0.7	1.2	3.2	1.7	-1.5
	23–26	1.8	1.5	2.2	1.8	1.0	2.5	1.9	2.9	1.8	2.0	3.2	1.0	1.0	1.2	1.4	2.0	1.0	-1.0
	27–30	1.6	0.8	1.5	0.3	1.6	2.4	1.7	0.7	2.4	0.8	0.7	1.4	0.8	0.1	1.4	0.8	1.6	+0.8
Heroin																			
% saying any	18	9.1	8.7	8.1	9.1	8.7	8.3	7.3	6.6	7.3	9.0	8.6	6.8	7.3	8.3	6.4	6.0	6.6	+0.6
	19–22	3.7	6.4	3.2	5.2	3.2	5.3	3.4	3.0	3.2	3.7	3.3	4.8	4.3	3.2	4.0	3.6	3.4	-0.3
	23–26	2.7	3.1	2.9	2.6	2.4	3.8	2.0	3.1	2.6	3.5	3.6	1.8	1.8	4.1	4.8	2.4	2.3	-0.1
	27–30	1.3	1.4	1.9	1.9	2.3	2.7	1.3	3.2	2.9	2.3	3.0	2.2	1.9	1.5	3.5	3.0	3.9	+0.9
% saying often exposed	18	1.2	0.9	1.3	1.5	0.7	1.3	1.2	1.2	0.8	1.7	1.1	0.8	0.8	1.0	1.1	1.3	0.7	-0.6
	19–22	0.4	0.7	0.8	0.7	0.8	0.6	0.2	*	0.8	0.1	*	0.6	*	0.4	0.7	1.0	1.0	-0.1
	23–26	0.3	0.5	1.0	*	*	0.8	0.5	0.5	0.3	0.6	0.3	*	*	1.2	0.3	0.2	*	-0.2
	27–30	*	*	0.2	*	*	0.7	0.3	*	0.4	0.3	0.4	0.3	0.6	*	1.2	0.7	1.3	+0.6
Other narcotics^d																			
% saying any	18	20.4	20.7	21.9	21.1	21.6	22.5	21.8	20.3	19.0	18.9	18.9	16.3	16.3	30.3	27.5	27.1	23.9	-4.2 s
	19–22	14.5	15.3	13.9	17.0	18.3	18.7	13.6	14.5	16.8	15.3	12.5	13.2	14.2	27.5	23.7	25.2	19.5	-5.6
	23–26	6.5	8.1	9.4	10.9	12.2	12.0	12.6	12.6	12.4	13.0	14.4	11.2	13.2	25.9	25.3	24.1	22.5	-1.6
	27–30	4.9	3.6	5.2	6.5	9.0	7.9	9.5	8.8	11.6	10.6	9.2	9.1	9.7	23.4	22.7	23.6	24.5	+0.9
% saying often exposed	18	2.5	2.8	3.9	2.9	3.0	3.8	3.0	3.3	2.6	3.4	3.4	2.1	2.7	5.3	5.6	5.7	3.8	-1.9 s
	19–22	1.5	1.7	1.1	2.4	1.6	3.0	1.2	0.8	2.4	1.9	1.7	1.9	1.6	3.3	2.1	1.6	2.2	+0.5
	23–26	0.7	0.5	1.1	0.7	1.0	0.9	1.6	1.4	1.3	1.1	1.8	1.0	1.3	4.4	2.5	3.6	1.5	-2.1
	27–30	0.5	*	0.2	1.1	1.0	0.7	1.2	0.1	1.7	0.7	0.8	0.4	1.4	3.0	3.1	2.3	3.1	+0.8
Amphetamines^e																			
% saying any	18	31.0	29.9	30.1	29.5	31.5	30.6	27.4	27.2	26.4	26.6	23.8	23.3	23.8	23.6	28.0	26.2	25.4	-0.8
	19–22	24.6	24.8	21.2	24.8	23.3	25.5	21.6	23.7	22.2	22.7	22.8	17.6	18.0	19.4	26.0	27.4	26.2	-1.2
	23–26	11.7	14.6	12.3	18.5	18.2	17.9	15.4	18.8	15.6	18.7	16.6	13.7	15.3	15.8	24.2	23.1	21.4	-1.7
	27–30	9.1	6.6	10.4	7.4	11.1	11.5	12.2	11.4	12.2	14.1	10.0	10.3	10.3	12.6	16.4	19.0	19.1	+0.1
% saying often exposed	18	5.2	4.7	6.3	4.4	6.0	6.4	4.9	5.3	4.1	5.6	4.3	3.0	4.3	3.3	6.1	5.7	5.3	-0.5
	19–22	4.1	2.9	2.2	2.4	2.6	5.6	1.7	4.1	3.1	2.9	2.3	2.1	3.0	3.9	3.3	5.5	3.7	-1.8
	23–26	1.4	2.2	1.7	1.4	2.2	0.7	1.3	1.7	1.6	2.6	1.6	1.8	1.1	1.6	3.1	4.1	3.2	-0.9
	27–30	1.0	0.2	1.1	0.4	0.6	1.5	1.0	1.2	1.0	0.8	1.1	0.3	0.7	0.6	1.7	3.0	2.7	-0.3

↓
(List of drugs continued.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

Percentage saying exposed to drug ^a

	Age Group	Percentage saying exposed to drug ^a																	(Years Cont.)
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Sedatives/barbiturates ^b																			
% saying any	18	25.2	25.9	25.7	22.5	21.2	18.9	15.8	13.1	12.4	11.8	13.3	10.0	10.2	11.9	13.0	14.5	15.5	
	19–22	25.6	23.1	21.8	18.3	15.7	14.7	12.8	12.0	8.2	8.3	6.5	7.9	7.3	7.2	7.4	10.1	8.8	
	23–26	—	—	—	—	16.1	13.1	11.0	7.1	7.1	6.6	6.9	5.9	6.5	3.8	4.2	5.7	6.6	
	27–30	—	—	—	—	—	—	—	—	8.0	6.8	5.9	5.4	5.2	5.7	4.5	5.2	3.5	
% saying often exposed	18	3.4	4.0	4.3	3.0	2.7	1.7	2.1	1.5	1.4	1.7	1.7	1.2	1.1	1.6	1.7	2.0	2.9	
	19–22	2.5	2.8	1.1	1.4	0.7	1.3	0.5	0.7	0.7	0.3	0.7	0.4	0.7	0.7	1.3	1.3	0.4	
	23–26	—	—	—	—	0.7	0.9	1.7	0.8	0.6	0.3	1.1	0.3	0.3	*	*	0.2	0.3	
	27–30	—	—	—	—	—	—	—	—	0.7	0.4	0.6	0.2	0.4	1.2	0.2	0.6	0.5	
Tranquilizers ^c																			
% saying any	18	29.1	29.0	26.6	23.5	23.1	23.4	19.6	18.4	18.2	15.1	16.3	14.2	12.7	13.8	16.5	15.7	17.9	
	19–22	29.6	26.9	28.5	19.5	21.2	19.5	16.4	18.5	13.8	12.0	12.7	12.6	11.0	10.0	12.0	11.8	10.7	
	23–26	—	—	—	—	23.1	21.0	16.9	15.9	13.4	12.9	12.0	10.4	9.7	10.9	9.8	10.3	10.1	
	27–30	—	—	—	—	—	—	—	—	15.0	11.6	11.1	9.7	10.3	10.4	9.0	11.2	9.6	
% saying often exposed	18	3.2	4.2	3.5	2.9	2.9	2.2	2.5	2.6	2.2	2.1	1.9	1.4	1.9	1.7	1.8	2.3	3.5	
	19–22	3.2	2.6	1.8	2.1	1.5	1.7	0.9	1.1	1.8	1.0	1.1	1.1	1.5	1.1	1.3	1.5	0.5	
	23–26	—	—	—	—	2.0	1.6	2.6	1.8	1.2	0.8	0.5	1.0	0.6	0.7	0.1	1.1	1.5	
	27–30	—	—	—	—	—	—	—	—	1.4	0.3	1.7	0.8	1.3	1.3	1.0	1.1	0.8	
Alcoholic beverages																			
% saying any	18	94.7	94.0	94.0	94.0	94.0	94.0	94.1	93.9	93.1	92.3	93.6	91.7	90.6	91.8	90.0	91.2	91.5	
	19–22	94.3	93.8	94.5	93.4	94.2	92.7	93.6	94.4	92.5	91.8	92.4	94.0	93.3	92.9	93.7	93.1	93.7	
	23–26	—	—	—	—	90.3	92.7	91.4	90.6	91.1	92.9	91.3	91.0	91.4	90.3	89.5	91.9	89.6	
	27–30	—	—	—	—	—	—	—	—	87.1	88.4	86.2	87.7	87.3	86.6	86.2	89.3	89.2	
% saying often exposed	18	60.2	61.0	59.3	60.2	58.7	59.5	58.0	58.7	56.4	55.5	56.1	54.5	53.1	51.9	54.0	54.0	54.5	
	19–22	59.6	61.2	62.5	56.6	59.3	61.8	59.9	61.4	55.4	53.8	56.0	53.9	56.1	56.8	57.0	56.3	52.3	
	23–26	—	—	—	—	52.1	54.8	51.4	53.0	48.1	50.9	49.7	48.4	45.4	45.4	43.3	47.5	44.8	
	27–30	—	—	—	—	—	—	—	—	39.9	39.5	38.7	38.0	39.9	38.1	39.3	38.0	34.7	
Approximate Weighted N =	18	3,259	3,608	3,645	3,334	3,238	3,252	3,078	3,296	3,300	2,795	2,556	2,525	2,630	2,730	2,581	2,608	2,407	
	19–22	582	574	601	569	578	549	591	582	556	567	567	532	528	489	460	464	485	
	23–26	—	—	—	—	533	532	557	529	531	514	523	494	532	513	471	467	447	
	27–30	—	—	—	—	—	—	—	—	522	507	506	478	502	457	425	452	432	

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Sedatives/barbiturates^f																			
% saying any	18	16.1	16.1	17.1	16.3	17.1	17.7	14.8	21.5	20.4	21.3	18.8	16.7	17.6	18.8	16.2	16.0	15.0	-1.0
	19–22	11.7	13.4	11.6	13.1	13.1	16.0	11.9	17.2	17.8	16.0	16.1	15.2	17.3	16.1	12.2	14.8	10.7	-4.1
	23–26	4.9	8.5	7.1	9.3	9.0	9.8	7.9	15.9	12.5	14.8	13.1	12.4	12.7	13.4	15.2	14.9	10.6	-4.2
	27–30	3.8	2.7	4.1	2.9	5.3	6.0	6.1	9.2	12.4	11.9	10.3	10.1	9.9	11.6	10.4	11.7	10.1	-1.6
% saying often exposed	18	2.5	2.7	3.8	2.7	2.7	4.6	2.8	4.1	3.7	3.9	3.9	2.1	3.4	2.5	3.1	2.9	2.5	-0.4
	19–22	0.9	1.4	0.9	1.6	1.2	1.8	0.8	1.7	2.1	2.5	1.4	2.2	1.9	0.9	1.2	1.7	1.0	-0.8
	23–26	0.8	0.5	0.9	0.7	0.2	0.3	0.4	0.7	1.1	1.1	1.6	1.7	0.7	1.0	1.0	1.5	0.8	-0.6
	27–30	0.2	*	0.6	0.2	0.9	0.4	0.6	0.4	1.7	0.7	1.3	0.4	1.7	0.9	1.3	1.1	2.0	+0.9
Tranquilizers^g																			
% saying any	18	18.9	17.3	18.2	17.7	23.8	22.7	21.0	22.1	20.9	21.8	19.3	19.9	20.0	18.2	17.0	17.6	16.4	-1.2
	19–22	15.6	16.9	14.3	18.5	21.3	23.6	20.0	21.9	20.6	23.1	21.4	20.0	19.6	18.1	16.6	19.0	13.3	-5.6
	23–26	9.4	10.9	10.8	12.3	16.4	20.1	18.7	19.9	20.1	19.9	18.8	18.4	17.5	21.4	19.6	21.0	19.5	-1.6
	27–30	9.6	6.1	8.8	7.6	12.6	13.6	15.3	14.6	18.1	19.2	16.7	16.8	13.5	18.6	16.5	19.5	17.5	-2.0
% saying often exposed	18	3.2	2.8	3.7	3.5	4.9	5.8	4.2	4.1	4.5	5.4	4.9	3.7	3.9	2.8	3.4	3.3	3.4	+0.1
	19–22	1.3	1.6	1.5	1.7	3.1	3.6	2.3	2.7	2.7	3.2	3.0	3.2	2.1	1.7	2.9	2.0	1.7	-0.3
	23–26	0.7	1.1	1.5	1.7	1.3	2.1	1.6	2.0	1.3	2.6	2.4	3.6	1.5	3.2	2.6	2.5	1.6	-0.9
	27–30	1.2	0.2	0.9	0.4	1.6	1.6	1.9	0.8	3.5	2.9	2.6	1.0	2.0	1.7	2.0	2.3	1.8	-0.5
Alcoholic beverages																			
% saying any	18	91.4	92.2	91.8	90.7	90.8	89.5	88.3	87.6	87.4	87.6	86.5	85.7	86.5	85.2	85.0	85.3	84.8	-0.4
	19–22	93.1	91.8	91.0	93.3	94.3	93.7	93.6	92.5	92.7	92.0	91.8	90.5	91.2	86.5	87.5	85.8	82.8	-3.0
	23–26	93.1	89.1	91.5	92.1	90.1	91.9	91.8	92.2	90.0	94.0	94.5	92.0	93.0	91.1	94.2	88.7	88.7	0.0
	27–30	86.4	88.4	88.7	89.8	91.2	89.0	90.0	85.3	92.2	91.8	89.6	94.4	91.0	91.2	92.5	90.5	88.8	-1.7
% saying often exposed	18	53.9	54.5	53.5	50.2	52.7	50.8	49.0	48.2	49.1	47.8	46.4	45.4	46.3	45.8	40.7	43.0	41.7	-1.3
	19–22	54.2	57.9	54.7	54.3	53.4	54.9	55.7	54.3	58.9	55.0	60.7	53.9	53.4	48.5	46.0	50.6	45.3	-5.3
	23–26	49.8	44.6	45.7	49.6	48.8	46.3	50.5	48.3	46.4	57.1	54.2	49.6	53.8	51.3	52.5	55.6	49.3	-6.3
	27–30	37.1	36.6	38.3	34.4	40.0	39.6	40.6	36.8	43.6	47.3	44.3	47.8	45.2	43.0	49.3	50.4	48.1	-2.3
Approximate Weighted N =	18	2,595	2,541	2,312	2,153	2,147	2,162	2,454	2,456	2,469	2,469	2,448	2,332	2,274	2,434	2,372	2,299	2,150	
	19–22	471	445	450	415	412	403	396	432	377	378	333	365	368	364	340	356	281	
	23–26	424	400	398	389	406	345	385	404	374	363	327	333	328	347	308	334	311	
	27–30	455	449	430	395	369	359	347	370	370	330	356	339	324	336	306	312	301	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '—' indicates data not available. ' * ' indicates a prevalence rate of less than 0.05%.

^aAnswer alternatives were: (1) Not at all, (2) Once or twice, (3) Occasionally, (4) Often. The "any" percentage combines categories (2)–(4).

^bThese estimates were derived from responses to the question for the following drugs: marijuana, LSD, other hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), and tranquilizers.

^cIn 2001 the question text was changed from other psychedelics to other hallucinogens, and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^dIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

^eIn 2011 pep pills and bennies were replaced in the list of examples by Adderall and Ritalin. This change likely explains the discontinuity in the 2011 results.

^fIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

^gIn 2001 Xanax was added to the list of examples. This change likely explains the discontinuity in the 2001 results.

TABLE 7-4
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

Percentage saying fairly easy or very easy to get ^a

	Age Group	Percentage saying fairly easy or very easy to get ^a																	(Years Cont.)
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Marijuana	18	89.0	89.2	88.5	86.2	84.6	85.5	85.2	84.8	85.0	84.3	84.4	83.3	82.7	83.0	85.5	88.5	88.7	→
	19–22	95.6	91.1	92.4	89.7	88.3	89.5	87.2	85.9	87.1	87.1	86.2	86.0	87.8	85.6	87.2	87.9	89.3	
	23–26	—	—	—	—	92.5	88.8	88.8	90.3	86.9	88.7	83.3	82.5	83.8	84.6	87.1	86.2	85.3	
	27–30	—	—	—	—	—	—	—	—	89.3	86.0	83.1	83.8	80.7	82.8	80.3	83.3	82.6	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	75.7	75.6	73.0	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Amyl & butyl nitrites	18	—	—	—	—	—	—	—	23.9	25.9	26.8	24.4	22.7	25.9	25.9	26.7	26.0	23.9	→
	19–22	—	—	—	—	—	—	—	22.8	26.0	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	23.1	28.0	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	26.7	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
LSD	18	35.3	35.0	34.2	30.9	30.6	30.5	28.5	31.4	33.3	38.3	40.7	39.5	44.5	49.2	50.8	53.8	51.3	→
	19–22	39.6	38.4	35.1	31.8	32.7	29.6	30.5	29.9	33.9	36.4	36.6	37.8	42.5	44.9	43.7	50.5	50.8	
	23–26	—	—	—	—	32.7	29.1	30.0	27.5	32.7	32.6	30.2	32.8	33.5	33.4	40.1	41.0	43.6	
	27–30	—	—	—	—	—	—	—	—	29.4	29.9	32.3	27.0	30.9	30.5	27.2	35.6	33.6	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.8	32.4	28.4	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Other hallucinogens ^b	18	35.0	32.7	30.6	26.6	26.6	26.1	24.9	25.0	26.2	28.2	28.3	28.0	29.9	33.5	33.8	35.8	33.9	→
	19–22	42.1	37.7	33.5	31.0	28.9	28.7	26.3	27.5	28.7	28.1	28.9	26.6	28.3	29.5	28.6	31.5	31.5	
	23–26	—	—	—	—	31.8	29.6	26.4	25.6	29.6	28.7	27.0	25.7	27.7	25.3	28.3	29.2	32.6	
	27–30	—	—	—	—	—	—	—	—	28.6	29.6	30.8	24.9	24.8	25.4	24.7	29.3	25.9	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
PCP	18	—	—	—	—	—	—	—	22.8	24.9	28.9	27.7	27.6	31.7	31.7	31.4	31.0	30.5	→
	19–22	—	—	—	—	—	—	—	21.7	24.6	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	21.2	27.6	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	24.3	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Ecstasy (MDMA)	18	—	—	—	—	—	—	—	—	—	21.7	22.0	22.1	24.2	28.1	31.2	34.2	36.9	→
	19–22	—	—	—	—	—	—	—	—	—	—	26.6	24.9	27.1	23.9	27.0	29.3	33.4	
	23–26	—	—	—	—	—	—	—	—	—	—	21.4	23.1	26.4	24.0	26.0	27.8	28.7	
	27–30	—	—	—	—	—	—	—	—	—	—	27.1	20.8	22.2	22.8	21.9	27.1	29.3	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		

(Table continued on next page.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, 50, and 55

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

Percentage saying fairly easy or very easy to get ^a

	Age Group	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012–2013 change	
Marijuana	18	89.6	90.4	88.9	88.5	88.5	87.2	87.1	85.8	85.6	84.9	83.9	83.9	81.1	82.1	82.2	81.6	81.4	-0.2	
	19–22	90.6	89.9	87.4	89.6	91.7	88.1	87.7	87.3	88.0	86.8	88.4	87.5	83.0	84.2	82.9	85.4	83.9	-1.6	
	23–26	84.4	87.5	85.9	88.4	87.0	89.1	87.2	88.8	87.0	86.8	87.6	85.3	89.4	83.3	88.3	87.0	87.4	+0.4	
	27–30	84.5	82.1	83.0	81.5	84.8	83.6	81.8	86.0	84.6	87.6	87.8	86.4	88.9	84.6	85.6	85.1	86.8	+1.7	
	35	77.1	76.0	74.9	77.1	75.3	76.5	75.1	75.6	73.8	75.1	75.5	76.4	75.7	75.6	80.4	80.5	80.2	-0.3	
	40	—	73.4	71.7	73.1	70.4	72.1	72.3	68.9	73.6	69.7	71.2	72.5	72.9	73.6	74.6	74.6	78.8	+4.2 s	
	45	—	—	—	—	—	—	—	68.5	69.9	70.1	67.9	70.1	68.1	67.9	73.4	69.8	71.8	73.6	+1.8
	50	—	—	—	—	—	—	—	—	—	—	—	—	64.4	65.8	67.9	65.8	68.9	70.1	+1.2
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	68.8	—
	Amyl & butyl nitrites	18	23.8	25.1	21.4	23.3	22.5	22.3	19.7	20.0	19.7	18.4	18.1	16.9	15.7	—	—	—	—	—
19–22		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
23–26		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
27–30		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
35		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
40		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
45		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
50		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
LSD		18	50.7	48.8	44.7	46.9	44.7	39.6	33.6	33.1	28.6	29.0	28.7	28.5	26.3	25.1	25.1	27.6	24.5	-3.1
	19–22	47.7	51.1	43.8	47.1	42.5	37.9	34.1	30.3	27.7	29.0	23.0	19.7	24.2	26.1	24.8	23.2	26.2	+2.9	
	23–26	39.2	40.4	41.2	40.4	38.3	37.2	34.1	38.5	26.5	30.3	25.2	24.1	26.1	24.2	21.4	19.3	22.8	+3.5	
	27–30	35.2	32.9	35.7	35.6	38.3	32.3	33.5	30.0	29.3	29.7	26.8	28.1	22.5	25.2	26.6	19.1	21.7	+2.6	
	35	32.9	31.2	27.7	32.2	28.7	29.1	29.8	25.6	24.0	28.7	26.6	26.4	26.9	25.5	24.0	23.0	24.1	+1.1	
	40	—	31.1	31.0	28.5	25.7	27.4	25.0	24.4	24.3	23.9	21.5	25.1	22.2	23.3	22.6	21.6	20.1	-1.5	
	45	—	—	—	—	—	—	24.2	27.0	25.4	23.7	23.6	21.1	19.4	23.6	21.3	18.9	23.4	+4.5 s	
	50	—	—	—	—	—	—	—	—	—	—	—	19.0	21.9	18.6	20.3	18.1	17.1	-1.1	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	Other hallucinogens ^b	18	33.9	35.1	29.5	34.5	48.5	47.7	47.2	49.4	45.0	43.9	43.7	42.8	40.5	39.5	38.3	37.8	36.6	-1.2
19–22		33.4	34.1	31.1	33.4	45.9	48.8	45.1	46.9	48.5	41.9	39.3	34.7	38.1	39.1	37.5	36.4	34.1	-2.3	
23–26		31.0	32.4	31.5	28.5	38.3	39.7	39.2	44.4	39.2	41.5	36.8	39.3	39.2	32.3	35.0	32.7	31.8	-0.9	
27–30		28.0	25.2	30.3	25.0	38.6	33.3	35.6	31.2	30.8	32.1	30.0	36.2	32.0	34.7	33.4	31.4	33.3	+1.9	
35		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
40		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
45		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
50		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
55		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
PCP		18	30.0	30.7	26.7	28.8	27.2	25.8	21.9	24.2	23.2	23.1	21.0	20.6	19.2	18.5	17.2	14.2	15.3	+1.1
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	55	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	Ecstasy (MDMA)	18	38.8	38.2	40.1	51.4	61.5	59.1	57.5	47.9	40.3	40.3	40.9	41.9	35.1	36.4	37.1	35.9	35.1	-0.9
19–22		35.6	39.4	43.2	49.9	55.5	59.7	52.1	45.8	43.5	41.2	38.4	34.7	37.1	30.4	37.9	28.3	33.9	+5.6	
23–26		31.1	30.1	34.9	41.8	51.5	52.9	49.3	51.3	46.4	44.6	42.2	41.5	36.8	35.2	34.0	32.2	35.7	+3.5	
27–30		24.3	26.4	30.0	35.5	40.6	41.2	41.0	41.1	38.0	40.5	40.7	42.2	38.0	31.2	33.8	32.8	28.6	-4.2	
35		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
40		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
45		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

↓
(List of drugs continued.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

Percentage saying fairly easy or very easy to get ^a

	Age Group	Percentage saying fairly easy or very easy to get ^a																	(Years Cont.)
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Cocaine	18	47.9	47.5	47.4	43.1	45.0	48.9	51.5	54.2	55.0	58.7	54.5	51.0	52.7	48.5	46.6	47.7	48.1	→
	19–22		56.2	57.1	55.2	56.2	56.9	60.4	65.0	64.9	66.8	61.7	54.3	54.5	49.2	49.9	49.4	44.4	
	23–26	—	—	—	—	63.7	67.2	65.8	69.0	71.7	70.0	65.6	58.0	61.1	53.8	54.4	54.7	50.2	
	27–30	—	—	—	—	—	—	—	—	68.6	68.2	64.0	60.0	63.1	56.8	53.1	57.0	53.0	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Crack	18	—	—	—	—	—	—	—	41.1	42.1	47.0	42.4	39.9	43.5	43.6	40.5	41.9	40.7	
	19–22	—	—	—	—	—	—	—	41.9	47.3	47.2	46.9	42.1	42.1	38.4	41.6	40.7	32.9	
	23–26	—	—	—	—	—	—	—	44.5	53.0	49.9	46.9	42.0	42.6	42.5	42.4	42.3	37.9	
	27–30	—	—	—	—	—	—	—	—	46.5	46.8	46.8	43.1	45.2	45.8	41.1	44.7	39.9	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	49.6	48.2	43.1	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Cocaine powder	18	—	—	—	—	—	—	—	52.9	50.3	53.7	49.0	46.0	48.0	45.4	43.7	43.8	44.4	
	19–22	—	—	—	—	—	—	—	58.7	60.2	61.7	56.5	52.5	48.9	45.7	47.8	45.5	41.3	
	23–26	—	—	—	—	—	—	—	64.9	69.1	60.1	58.6	53.2	56.4	50.5	49.7	49.6	45.9	
	27–30	—	—	—	—	—	—	—	63.5	62.8	57.9	55.8	56.8	55.0	48.9	48.9	52.9	48.4	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	53.9	52.1	46.7	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Heroin	18	21.2	19.2	20.8	19.3	19.9	21.0	22.0	23.7	28.0	31.4	31.9	30.6	34.9	33.7	34.1	35.1	32.2	
	19–22	18.9	19.4	19.3	16.4	17.2	20.8	21.2	24.4	28.5	31.6	30.7	25.3	30.2	30.0	33.2	35.2	29.1	
	23–26	—	—	—	—	18.6	18.1	21.0	22.3	28.4	31.2	28.1	25.6	25.7	25.7	29.2	29.3	32.3	
	27–30	—	—	—	—	—	—	—	—	23.6	27.4	29.5	22.1	25.6	28.5	24.4	30.7	29.5	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Some other narcotic ^c	18	29.4	29.6	30.4	30.0	32.1	33.1	32.2	33.0	35.8	38.3	38.1	34.6	37.1	37.5	38.0	39.8	40.0	
	19–22	32.7	32.4	30.8	31.0	28.7	34.3	32.6	33.8	37.9	37.9	35.6	35.4	35.2	33.5	35.1	38.7	37.3	
	23–26	—	—	—	—	32.8	32.1	33.6	32.2	35.9	36.4	34.7	33.2	33.9	33.1	35.8	32.6	36.7	
	27–30	—	—	—	—	—	—	—	—	31.6	36.2	36.1	29.0	31.8	33.0	34.8	36.9	37.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amphetamines	18	61.3	69.5	70.8	68.5	68.2	66.4	64.3	64.5	63.9	64.3	59.7	57.3	58.8	61.5	62.0	62.8	59.4	
	19–22	71.7	72.6	73.5	69.7	69.1	69.1	63.1	61.8	61.3	62.2	57.7	58.3	56.3	56.0	56.6	60.3	56.9	
	23–26	—	—	—	—	65.8	66.0	64.5	65.3	62.2	60.1	55.8	54.8	54.5	52.6	52.9	56.0	52.8	
	27–30	—	—	—	—	—	—	—	—	54.3	58.6	55.3	54.4	50.4	52.9	48.3	53.7	51.7	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	45.6	43.5	39.1	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

Percentage saying fairly easy or very easy to get ^a

	Age Group	Percentage saying fairly easy or very easy to get ^a																	2012–2013 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	
Cocaine	18	48.5	51.3	47.6	47.8	46.2	44.6	43.3	47.8	44.7	46.5	47.1	42.4	39.4	35.5	30.5	29.8	30.5	+0.7
	19–22	49.7	47.7	52.6	52.1	49.6	47.6	46.7	47.0	50.0	47.4	47.3	44.0	38.5	37.2	39.2	32.9	28.1	-4.7
	23–26	46.9	51.8	45.7	45.0	44.6	47.8	40.8	50.7	48.4	51.2	47.4	45.5	44.0	41.1	37.8	37.4	36.8	-0.6
	27–30	50.4	46.9	50.0	44.6	45.5	46.3	42.9	38.0	43.1	43.2	45.8	50.6	43.6	40.8	44.2	42.3	35.0	-7.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Crack	18	40.6	43.8	41.1	42.6	40.2	38.5	35.3	39.2	39.3	38.8	37.5	35.2	31.9	26.1	24.0	22.0	24.6	+2.6
	19–22	39.9	40.0	40.8	40.2	37.3	35.7	37.5	33.7	34.0	35.2	35.7	31.4	27.3	27.2	27.3	20.6	20.8	+0.2
	23–26	37.2	38.4	35.0	31.9	37.1	33.9	32.8	36.5	35.1	34.0	31.4	33.1	27.4	27.1	25.3	27.6	24.2	-3.5
	27–30	36.5	33.3	38.8	35.9	36.9	33.4	33.7	28.0	34.4	29.6	36.4	36.1	33.1	27.5	28.9	25.2	24.6	-0.6
	35	44.3	45.0	41.6	45.0	41.2	38.9	40.5	36.1	34.2	37.1	35.1	33.2	31.6	30.0	30.4	27.3	28.7	+1.4
	40	—	43.3	44.3	42.0	38.7	39.5	39.0	35.8	38.6	37.1	32.7	35.2	33.2	30.9	30.1	27.9	25.5	-2.4
	45	—	—	—	—	—	—	37.0	40.0	40.6	36.2	37.0	34.2	31.7	36.2	32.3	28.2	32.3	+4.0
Cocaine powder	18	43.3	45.7	43.7	44.6	40.7	40.2	37.4	41.7	41.6	42.5	41.2	38.9	33.9	29.0	26.4	25.1	28.4	+3.3
	19–22	46.0	47.1	45.2	45.2	43.3	43.9	45.5	43.2	44.3	44.2	44.5	39.0	36.1	35.6	35.4	26.0	25.1	-0.8
	23–26	43.6	44.4	44.3	41.8	44.4	40.7	43.4	48.5	45.1	46.4	45.0	41.4	41.6	40.3	37.5	37.0	35.1	-1.9
	27–30	45.1	43.9	46.5	43.9	42.7	42.4	39.7	37.9	40.2	42.7	43.0	47.5	41.3	38.2	38.4	37.0	35.4	-1.6
	35	48.3	47.0	43.4	47.9	43.1	41.7	42.0	39.6	35.8	39.5	37.4	38.6	34.9	35.5	35.3	31.4	35.2	+3.8
	40	—	46.0	46.7	44.7	41.5	41.5	40.7	38.5	40.3	37.8	35.2	36.5	33.9	33.5	31.8	29.5	29.8	+0.3
	45	—	—	—	—	—	—	39.0	40.2	40.6	37.3	38.2	34.1	31.5	37.2	33.2	28.7	34.0	+5.3 s
Heroin	18	33.8	35.6	32.1	33.5	32.3	29.0	27.9	29.6	27.3	27.4	29.7	25.4	27.4	24.1	20.8	19.9	22.1	+2.3
	19–22	31.4	32.1	32.7	29.4	30.2	26.4	26.9	22.6	25.4	25.3	26.5	24.2	19.4	22.0	21.2	19.3	16.0	-3.3
	23–26	30.5	35.1	31.9	25.7	26.6	27.2	25.5	30.9	22.5	28.1	22.2	23.4	23.4	23.1	21.1	22.7	23.1	+0.3
	27–30	30.0	28.3	33.0	29.3	29.9	27.0	27.5	22.0	27.8	25.4	27.5	26.3	25.2	25.2	28.0	23.3	20.9	-2.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Some other narcotic ^c	18	38.9	42.8	40.8	43.9	40.5	44.0	39.3	40.2	39.2	39.6	37.3	34.9	36.1	54.2	50.7	50.4	46.5	-3.9 s
	19–22	38.3	38.9	39.5	41.1	44.1	40.4	40.6	39.4	41.4	38.5	38.3	38.0	35.3	55.2	53.8	52.2	53.5	+1.3
	23–26	35.7	39.9	38.2	38.1	35.8	40.0	40.3	47.7	44.7	45.5	41.7	41.2	42.5	56.2	59.6	58.6	62.1	+3.5
	27–30	35.2	32.2	36.9	32.4	39.4	38.5	38.9	35.8	37.7	39.8	41.3	39.4	43.5	62.3	65.2	59.8	64.4	+4.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines ^d	18	59.8	60.8	58.1	57.1	57.1	57.4	55.0	55.4	51.2	52.9	49.6	47.9	47.1	44.1	47.0	45.4	42.7	-2.7
	19–22	55.5	56.3	57.6	60.2	56.5	53.7	55.1	53.9	56.9	52.3	55.8	49.5	49.8	43.6	52.3	54.4	54.0	-0.4
	23–26	51.2	53.2	49.1	51.1	49.4	48.2	50.3	51.8	51.9	58.0	53.7	46.9	51.0	45.5	55.5	55.6	59.4	+3.8
	27–30	48.1	41.4	48.2	47.6	49.3	45.6	48.7	43.9	45.3	49.2	48.1	45.0	51.1	46.4	49.9	54.6	54.2	-0.3
	35	40.9	39.4	38.5	42.2	39.6	39.2	39.2	35.4	35.4	40.3	40.4	40.6	39.2	37.1	40.4	37.5	40.7	+3.1
	40	—	41.0	41.9	39.4	37.5	39.4	38.7	37.9	41.1	38.4	37.6	39.2	37.2	37.0	34.3	35.8	34.6	-1.2
	45	—	—	—	—	—	—	35.8	39.8	39.3	37.1	38.3	36.8	33.0	39.8	37.0	34.5	39.3	+4.9 s

(List of drugs continued.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a																	(Years Cont.)
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Crystal methamphetamine (ice)	18	—	—	—	—	—	—	—	—	—	—	24.0	24.3	26.0	26.6	25.6	27.0	26.9	
	19–22	—	—	—	—	—	—	—	—	—	—	24.0	21.8	22.5	20.9	24.7	25.5	25.4	
	23–26	—	—	—	—	—	—	—	—	—	—	22.3	20.0	21.3	22.9	24.5	24.7	24.7	
	27–30	—	—	—	—	—	—	—	—	—	—	27.3	19.7	22.0	21.2	21.7	25.8	26.1	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Sedatives/ barbiturates ^d	18	49.1	54.9	55.2	52.5	51.9	51.3	48.3	48.2	47.8	48.4	45.9	42.4	44.0	44.5	43.3	42.3	41.4	
	19–22	59.5	61.1	56.8	54.2	48.1	52.7	46.8	44.6	45.5	47.7	44.2	41.7	43.4	41.9	40.6	42.9	41.1	
	23–26	—	—	—	—	52.7	47.7	46.4	45.9	47.4	44.8	41.6	39.6	42.0	38.8	40.3	42.1	40.6	
	27–30	—	—	—	—	—	—	—	—	43.2	44.5	44.2	38.5	37.8	39.7	37.4	39.9	41.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Tranquilizers	18	59.1	60.8	58.9	55.3	54.5	54.7	51.2	48.6	49.1	45.3	44.7	40.8	40.9	41.1	39.2	37.8	36.0	
	19–22	67.4	62.8	62.0	62.3	52.5	55.6	52.9	50.3	50.0	49.4	45.4	44.8	40.7	40.9	41.0	40.2	37.6	
	23–26	—	—	—	—	60.2	54.3	54.1	56.3	52.8	51.4	47.8	45.1	48.1	43.2	45.9	44.3	42.3	
	27–30	—	—	—	—	—	—	—	—	55.3	54.4	54.9	47.5	47.8	47.4	44.4	44.8	46.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Steroids	18	—	—	—	—	—	—	—	—	—	—	—	46.7	46.8	44.8	42.9	45.5	40.3	
	19–22	—	—	—	—	—	—	—	—	—	—	—	44.1	44.8	46.3	41.7	40.9	41.8	
	23–26	—	—	—	—	—	—	—	—	—	—	—	37.6	35.8	39.3	35.8	37.0	37.4	
	27–30	—	—	—	—	—	—	—	—	—	—	—	36.4	30.6	35.0	31.6	30.5	33.1	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Approximate Weighted N =	18	3,240	3,578	3,602	3,385	3,269	3,274	3,077	3,271	3,231	2,806	2,549	2,476	2,586	2,670	2,526	2,552	2,340	
	19–22	582	601	582	588	559	571	592	581	568	572	571	534	512	480	459	470	467	
	23–26	—	—	—	—	540	541	548	539	526	514	532	511	523	500	463	449	418	
	27–30	—	—	—	—	—	—	—	—	519	513	510	487	475	473	437	446	468	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1,142	1,141	1,146	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

Percentage saying "fairly easy" or "very easy" to get^a

	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a																2012–2013 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012		2013
Crystal methamphetamine (ice)	18	27.6	29.8	27.6	27.8	28.3	28.3	26.1	26.7	27.2	26.7	25.1	23.3	22.3	18.3	17.1	14.5	17.2	+2.7
	19–22	29.3	31.0	31.8	27.4	28.4	31.2	26.5	27.1	28.9	29.1	27.7	24.1	19.2	19.3	19.4	13.5	15.3	+1.8
	23–26	25.8	30.2	28.5	25.8	26.4	25.1	26.4	32.3	27.8	32.3	27.8	27.7	23.1	26.1	18.2	23.5	16.3	-7.1 s
	27–30	25.1	22.6	29.1	25.3	27.6	29.5	30.9	25.5	27.4	31.8	29.7	31.4	27.7	27.6	26.2	24.2	22.3	-2.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Sedatives/barbiturates^o	18	40.0	40.7	37.9	37.4	35.7	36.6	35.3	46.3	44.4	43.8	41.7	38.8	37.9	36.8	32.4	28.7	27.9	-0.7
	19–22	39.8	39.2	42.3	40.6	39.3	40.8	38.4	43.8	47.8	42.6	47.5	43.2	42.6	39.6	38.1	31.6	32.1	+0.5
	23–26	39.1	42.6	39.7	37.6	36.1	36.4	37.8	49.4	48.4	51.4	46.5	43.3	47.7	40.4	41.3	40.1	42.2	+2.1
	27–30	39.1	33.9	38.4	36.1	38.1	34.8	35.6	40.5	42.9	43.3	46.4	44.7	48.5	43.1	42.9	42.3	44.8	+2.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers	18	35.4	36.2	32.7	33.8	33.1	32.9	29.8	30.1	25.7	24.4	23.6	22.4	21.2	18.4	16.8	14.9	15.0	+0.1
	19–22	37.8	36.8	37.1	36.5	34.9	34.6	34.2	29.7	30.1	22.8	28.5	23.3	18.3	20.2	18.6	17.3	17.4	+0.1
	23–26	36.4	39.4	38.3	37.6	38.7	33.7	32.5	36.6	32.9	33.0	31.7	30.3	27.7	21.8	23.0	22.1	18.5	-3.6
	27–30	41.9	39.9	41.5	36.7	42.9	38.1	35.9	30.6	33.5	32.1	32.4	33.1	30.1	30.6	27.1	25.7	28.1	+2.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	18	41.7	44.5	44.6	44.8	44.4	45.5	40.7	42.6	39.7	41.1	40.1	35.2	30.3	27.3	26.1	25.0	28.5	+3.5 s
	19–22	39.2	39.2	40.5	40.3	38.1	41.4	39.4	37.8	37.6	37.1	37.9	33.5	28.7	25.1	24.3	21.2	20.6	-0.7
	23–26	35.5	34.9	37.1	34.0	34.7	33.1	31.1	34.7	31.2	34.2	33.3	30.2	28.6	22.2	29.2	25.6	23.6	-2.0
	27–30	32.5	30.5	34.5	36.2	34.6	33.0	32.6	30.6	32.4	29.7	30.9	31.0	31.9	27.6	27.0	23.9	22.3	-1.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Approximate Weighted N =	18	2,517	2,520	2,215	2,095	1,850	2,138	2,391	2,169	2,161	2,161	2,420	2,276	2,243	2,395	2,337	2,280	2,092	
	19–22	463	433	425	400	398	375	386	441	392	376	362	380	377	377	355	341	342	
	23–26	419	395	415	388	401	362	356	411	359	335	338	355	312	358	313	332	325	
	27–30	459	425	424	365	357	349	368	393	359	347	324	334	305	340	325	334	281	
	35	1,150	1,032	1,022	981	977	890	934	963	1,009	925	863	898	952	895	852	875	844	
	40		1,029	1,093	1,096	1,065	1,037	898	967	928	919	868	881	870	911	850	823	820	
	45							911	1,026	1,005	972	954	851	888	846	852	842	806	
	50												902	975	989	939	958	819	
55																		832	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

'—' indicates data not available.

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, and (5) Very easy.

^bIn 2001 the question text was changed from other psychedelics to other hallucinogens, and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.


^cIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

^dIn 2011 pep pills and bennies were replaced in the list of examples by Adderall and Ritalin. This change likely explains the discontinuity in the 2011 results.

^eIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

FIGURE 7-1
Trends in Exposure to Use of ANY ILLICIT DRUGS
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-2
Trends in Exposure to Use of ANY ILLICIT DRUGS
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Often Exposed



Source: The Monitoring the Future study, the University of Michigan.

FIGURE 7-3
Trends in Exposure to Use of ANY ILLICIT DRUG OTHER THAN MARIJUANA^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source. The Monitoring the Future study, the University of Michigan.

^aThese estimates were derived from responses to the question for the following drugs: marijuana, LSD, other hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), ai

FIGURE 7-4
Trends in Exposure to Use of ANY ILLICIT DRUG OTHER THAN MARIJUANA^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed




Source. The Monitoring the Future study, the University of Michigan.

^aThese estimates were derived from responses to the question for the following drugs: marijuana, LSD, other hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), and

FIGURE 7-5
Trends in Exposure to Use of MARIJUANA
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure



Source: The Monitoring the Future study, the University of Michigan.

FIGURE 7-6
Trends in Exposure to Use of MARIJUANA
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-7
Trends in Exposure to Use of LSD
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-8
Trends in Exposure to Use of LSD
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-9
Trends in Exposure to Use of HALLUCINOGENS OTHER THAN LSD^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2001 the question text was changed from other psychedelics to other hallucinogens, and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

FIGURE 7-10
Trends in Exposure to Use of HALLUCINOGENS OTHER THAN LSD^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2001 the question text was changed from other psychedelics to other hallucinogens, and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

FIGURE 7-11
Trends in Exposure to Use of COCAINE
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Any Exposure



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-12
Trends in Exposure to Use of COCAINE
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-13
Trends in Exposure to Use of HEROIN
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Any Exposure



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-14
Trends in Exposure to Use of HEROIN
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-15
Trends in Exposure to Use of NARCOTICS OTHER THAN HEROIN^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source: The Monitoring the Future study, the University of Michigan.

^aIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

FIGURE 7-16
Trends in Exposure to Use of NARCOTICS OTHER THAN HEROIN^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed




Source: The Monitoring the Future study, the University of Michigan.

^aIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

FIGURE 7-17
Trends in Exposure to Use of AMPHETAMINES^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source: The Monitoring the Future study, the University of Michigan.

^aIn 2011 pep pills and bennies were replaced in the list of examples by Adderall and Ritalin. This change likely explains the discontinuity in the 2011 results.

FIGURE 7-18
Trends in Exposure to Use of AMPHETAMINES^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

^aIn 2011 pep pills and bennies were replaced in the list of examples by Adderall and Ritalin. This change likely explains the discontinuity in the 2011 results.

FIGURE 7-19
Trends in Exposure to Use of SEDATIVES (BARBITURATES)^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

FIGURE 7-20
Trends in Exposure to Use of SEDATIVES (BARBITURATES)^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, rebs, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

FIGURE 7-21
Trends in Exposure to Use of TRANQUILIZERS^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Any Exposure




Source. The Monitoring the Future study, the University of Michigan.

^aIn 2001 Xanax was added to the list of examples. This change likely explains the discontinuity in the 2001 results.

FIGURE 7-22
Trends in Exposure to Use of TRANQUILIZERS^a
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed




Source: The Monitoring the Future study, the University of Michigan.

^aIn 2001 Xanax was added to the list of examples. This change likely explains the discontinuity in the 2001 results.

FIGURE 7-23
Trends in Exposure to Use of ALCOHOL
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30


% Saying Any Exposure



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 7-24
Trends in Exposure to Use of ALCOHOL
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

% Saying Often Exposed



Source. The Monitoring the Future study, the University of Michigan.

Chapter 8

PREVALENCE OF DRUG USE AMONG COLLEGE STUDENTS

College students have often been the harbingers of social and political changes that eventually spread to other segments of the population up and down the age spectrum. The Monitoring the Future (MTF) study tracks multiple forms of substance use among U.S. college students and has done so for a third of a century (34 years). In this process, MTF has been able to document the fluctuations in college substance use as well as some patterns of influence on or by other age groups. This chapter focuses on the prevalence of drug use in 2013 by college students and their age-peers not in college; the next chapter focuses on historical trends in drug use among these two groups.

Definition of college students. College students are defined in this volume as those follow-up respondents one to four years past high school who report that they were taking courses as full-time students in a two- or four-year undergraduate college at the beginning of March of the year in question. Note that students at two-year colleges, such as community colleges, are included. The definition excludes those who are currently enrolled in college part-time and those who previously may have been college students or may have completed college. MTF has been able to generate an unparalleled national sample of college students every year since 1980 by following representative samples of sequential high school classes after they graduate. The graduating class of 1976 was the first such class followed after graduation, and by 1980 the survey included college students one to four years post-high school.

The absence of dropouts in the original high school senior samples has practically no effect on the representativeness of these college samples, as very few dropouts go on to college. One notable limitation of the present design for the purpose of characterizing college students is that it limits the age range of the college sample. For trend estimation purposes, we decided to limit the age band to the most typical one for college attendance, that is, one to four years past high school, which corresponds to modal ages 19 to 22. According to the latest statistics available from the United States Census Bureau,⁷⁴ this age band should encompass about 77% of all undergraduate college students enrolled full-time in 2011, down slightly from the 79% covered in 1989. Although expanding the age band to include an additional two years would cover 79% of all enrolled college students, it would reduce the homogeneity of the college experience by including older classmates, and would limit historical comparability. Special analyses conducted in 1985 and replicated in 1997 indicated that the differences in prevalence-of-use estimates under the two definitions

⁷⁴U.S. Census Bureau, October 2011. Available at: <http://www.census.gov/>.

were extremely small. The annual prevalence of all drugs except cocaine shifted only about one or two tenths of a percentage point. Cocaine, which had the greatest amount of age-related change at that time, would have had an annual prevalence rate only 0.8 percentage-points higher using the six-year age span. A replication of these analyses in 2011 yielded virtually the same results. Thus, for purposes of estimating all prevalence rates except lifetime prevalence, the four- and six-year intervals are nearly interchangeable, suggesting that this limitation is negligible for our purposes of trend estimation.

The MTF panels also include high school graduates one to four years past high school who are not attending college. Having longitudinal data for both groups, a rare feature of the MTF longitudinal design, makes it possible to compare college students' substance use rates with those of their age peers. College students as defined here now constitute well over half (61%) of the entire follow-up sample one to four years past high school. If data from the missing high school dropout segment were available for inclusion as part of the noncollege segment, any difference between the two groups in terms of their substance use would likely be enlarged; therefore, any difference observed here is only an indication of the direction and relative size of difference between the college and the entire noncollege population, not an absolute estimate of the difference.

PREVALENCE OF DRUG USE AMONG COLLEGE STUDENTS VERSUS THEIR NONCOLLEGE PEERS

In 2013, prevalence of use for most illicit drugs among college students is *lower* compared with use among their age peers, but the degree of difference varies considerably by drug, as Tables 8-1 through 8-4 show.

- In 2013, annual prevalence of use of *any illicit drug* was nearly equivalent for college students (39%) and the noncollege respondents (40%). Annual prevalence of using *any illicit drug other than marijuana* was also fairly comparable between the two groups with the college segment at 19% and the noncollege group at 20%.
- A few specific drugs showed roughly equivalent annual prevalence rates for the college student segment vs. noncollege segment: *hallucinogens other than LSD* (3.7% vs. 4.4%), *ecstasy* (5.3% vs. 5.7%), *methamphetamine* (0.4% vs. 0.3%), and *ketamine* (0.9% vs. 0.8%).
- The annual prevalence of *marijuana* use was slightly lower among college students in 2013 (36% vs. 37%), whereas the rate of current *daily marijuana* use was about twice as high among the noncollege group compared to the college students (10.0% vs. 5.1%, see Table 8-4.)

- It is clear that use of a number of *illicit drugs other than marijuana* tends to be distinctly higher among those not in college. (As previously noted, such differences would likely be larger if the noncollege group included high school dropouts.) In fact, several drugs show annual use rates for noncollege respondents in 2013 that are two or more times those for college students, including *synthetic marijuana*, *crack cocaine*, *heroin*, *heroin without a needle*, *Vicodin*, *crystal methamphetamine*, *bath salts (synthetic stimulants)*, and *GHB*.
- In 2013, significant proportions of both the noncollege group (9.6%) and college students (5.4%) reported use of *narcotics other than heroin* without medical supervision in the past year. With respect to specific drugs in this class, *Vicodin* was used by 9.5% of the noncollege group and 4.4% of college students; the corresponding numbers for *OxyContin* were 4.4% and 2.3%.
- *Amphetamines* were the only illicit drugs that now have appreciably higher rates among college students than among their noncollege age-mates. Annual prevalence of amphetamine use among college students was 10.6% in 2013, compared to 8.9% in the noncollege group. Specifically, annual prevalence of *Adderall* use without medical supervision (Table 8-2) was somewhat higher for college students (10.7%) than for noncollege respondents (6.8%), as was the case for the last four years. The higher use by college students is very likely because this amphetamine drug, intended for the treatment of attention deficit hyperactivity disorder (ADHD), is sometimes used by students to stay awake and alert in order to complete course work and to study for exams. Use of *Ritalin*, another stimulant drug prescribed for ADHD, was also slightly higher among college students (3.6%) in 2013 than among their noncollege peer group (2.3%).
- In 2013, college students were modestly higher in annual and 30-day use of *alcohol* than the noncollege group; the difference was largest in the 30-day rate (63% vs. 57%).
- College students also had a higher prevalence of *occasions of heavy drinking* (five or more drinks in a row in the past two weeks)—35% versus 31% among their age peers. Indeed, four in every ten college students (40%) report having *been drunk* in the prior 30 days, compared to 34% of the noncollege respondents. The groups did not differ much in their rates of *daily drinking*, with 3.6% of college students and 4.1% of their age peers reporting such use (Table 8-4). In high school, college-bound students, especially in earlier grades, were far less likely to drink alcohol at any level compared to their non-college-bound peers; thus, the relative and absolute increases in alcohol use among college students in the first few years following high school are quite striking.

- Because of increasing attention paid to the problem of *extreme binge drinking*, in 2005 we introduced a set of questions on the subject into one of the six questionnaire forms used with young adults, including college students. The questions ask respondents about the frequency in the past two weeks of their having 10 or more drinks in a row and 15 or more drinks in a row. The low *Ns* resulting from a single questionnaire form necessitate combining multiple years of data (2005–2013), and even then only 1,876 weighted cases are available from the college student population and 1,112 for their noncollege peers. However, they give us an idea of the prevalence and frequency of these levels of drinking.

About one in eight college students (13%) reported having *10 or more drinks in a row* at least once in the prior two weeks, and one in twenty (5%) reported *15 or more drinks in a row* at least once. The noncollege respondents had essentially identical rates (13% and 5%). While we lack earlier data on these measures to determine whether this type of extreme drinking behavior has changed over time, clearly it is quite high today among both college students and their noncollege age peers. As is discussed below, there is a dramatic gender difference in the prevalence of these behaviors.

- In 2013, 58% of college students reported using *flavored alcoholic beverages* in the prior year, compared to 55% for the noncollege group.
- Among all substances studied, the largest differences for annual, 30-day, and daily prevalence rates between the two groups occur for *cigarette smoking*. For example, the prevalence of daily smoking for college students is 6% versus 20% for noncollege respondents (Table 8-4). Smoking at the rate of a half pack or more per day stands at 2% versus 11% for these two groups, respectively. The 12th-grade data show the college-bound to have much lower smoking rates in high school than the non-college-bound; thus, in contrast to what was true for alcohol use, these substantial differences observed at college age actually preceded college attendance.⁷⁵ The smoking differences would be even greater if dropouts were included in the noncollege group, because dropouts have consistently shown an exceptionally high rate of smoking.⁷⁶

In sum, the noncollege segment is generally more drug-involved than the college student segment. This pattern is a continuation of the high school scenario in which those without college plans are more likely to use drugs. The only substances that

⁷⁵See also Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁷⁶For an analysis showing much higher smoking rates among 8th graders who later dropped out before completing high school, see Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

college students, , are appreciably more likely to use than their noncollege peers are *amphetamines* (including *Adderall* specifically), *Ritalin*, and *alcohol* (particularly *getting drunk* and *binge drinking*). The higher rates of alcohol use among college students emerge only after high school.

GENDER DIFFERENCES IN PREVALENCE OF USE AMONG COLLEGE STUDENTS

Data stratified by gender are provided in Tables 8-1 to 8-4.

- Most gender differences among college students replicate those discussed in chapter 4 for all young adults one to 12 years past high school, and they in turn replicate gender differences among secondary school students. That means that among college students, males have higher annual prevalence rates for most illicit drugs.
- Among college students, annual prevalence rates for use of *any illicit drug* are 44% for males versus 36% for females, and 24% versus 16% for *any illicit drug other than marijuana*. Annual *marijuana* use is higher among college males than females (**40% versus 33%**) and the same is true for the noncollege group (40% versus 35%). **Daily marijuana use is three times as high among male college students compared to female college students (8.7% vs. 2.9%)**; in the noncollege group the gender rates are much **higher and higher** among males (13.0% vs 7.3% among females).
- Most *hallucinogens* show *distinctly* higher annual prevalence rates among male versus female college students. Among the noncollege respondents the differences are in the same direction.
- Among college students, annual use of *narcotics other than heroin* without medical supervision is higher for males (7.4%) than for females (4.1%), as is true in the noncollege group (10.9% versus 8.4%, respectively). The specific narcotic *Vicodin* shows a similar pattern among the college students (7.8% for males versus 2.1% for females), as is true for the noncollege group (annual prevalence rates of 11.0% for males versus 8.0% for females). Note that college females are appreciably less likely to use Vicodin than noncollege females. For *OxyContin* use, males are higher than females in both college and non-college groups in 2014 (4.0% vs. 1.2% in college group and 5.8% vs. 3.0% in non-college group).
- Annual *amphetamine* use is higher among college males (12.4%) than college females (9.4%), and differences are similar in the noncollege segment (10.0% for males and 8.0% for females).

- Annual **Ritalin** use without medical supervision is higher among college males (4.4%) than females (3.0%), and in the noncollege segment males are also higher (3.2% vs. 1.4% for females).
- The annual prevalence of **Adderall** use outside of medical supervision is higher among male college students (one in every seven or eight college males, or 13.3%) than female college students (8.9%), and use is also higher among males than among females in the noncollege segment (7.3% vs. 6.3%, respectively). Again, the higher use of amphetamines among male and female college students compared with male and female noncollege respondents suggests that some college students of both genders are using these drugs to enhance their academic performance. Males report very slightly higher rates than females for **being drunk** in the past 30 days; this is true for both college (41% vs. 40%) and noncollege (35% vs. 33%) groups. Both male and female college students (43% and 30%, respectively) have higher rates of **binge drinking** than their counterparts not in college (34% for males and 30% for females; see Table 8-4).
- **Extreme levels of binge drinking** show a large gender difference in both groups. Among college students, the prevalence of having **10 or more drinks in a row** in the prior two weeks was 7% for college females versus 23% for college males for the years 2005–2013 combined; roughly similar rates were reported by noncollege females (7%) but for noncollege males use was lower (17%) than for college males (23%). The prevalence of having **15 or more drinks in a row** was 1.9% for college females versus 9.4% for college males; very similar rates were reported by noncollege respondents—2.2% for females and 9.5% for males. Put another way, during the prior two weeks nearly one in four college males had 10+ drinks at least once, including one in ten who reported having 15+ drinks at least once.
- **Flavored alcoholic beverages** are only slightly *more* likely to be consumed by females than males in college (58% of females vs. 57% of males reporting past-year use), but the gender difference is considerably larger in the noncollege group (66% of females vs. 47% of males).
- Among college students, 30-day prevalence of **cigarette smoking** is higher for males (18%) than for females (12%) in 2013, and the difference is similar in the noncollege segment (31% and 26%, respectively); rates for both genders are much higher in the noncollege group. **Daily smoking** is reported by more males than females in the college segment (7.0% vs. 4.7%); the rates again are much higher in the noncollege segment, but fairly comparable for the two genders (21% for males and 20% for females). Put another way, among males, daily smoking is about three times as high among the noncollege segment as among college students (21% vs. 7.0%), and among females, more than four times as high (20% vs. 4.7%). Rates of **smoking a half pack or more per day** are 3.3% and 1.8% for males and females among college students, compared with 12% and 11% for the noncollege segment.

In sum, among college student males tend to use most substances, licit and illicit, more than their female counterparts, with the greatest proportional differences occurring for various *hallucinogens*, *daily marijuana use*, and *extreme binge drinking*. Compared with their noncollege peers, college males are also more frequent users of *alcohol* and of *Adderall* (outside of medical supervision), but considerably less likely to use *marijuana daily*.

TABLE 8-1
Lifetime Prevalence of Use for Various Types of Drugs, 2013:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Any Illicit Drug ^a	51.0	57.1	54.3	59.9	48.8	54.6
Any Illicit Drug other than Marijuana ^a	26.7	31.8	30.8	34.7	24.0	29.3
Marijuana	47.7	55.4	51.7	58.7	45.2	52.5
Inhalants ^b	4.3	8.7	5.2	9.6	3.7	8.0
Hallucinogens	7.8	11.7	12.3	16.1	4.9	7.8
LSD	4.4	6.1	7.0	8.6	2.7	4.0
Hallucinogens other than LSD	6.8	10.3	11.4	14.8	3.9	6.3
Ecstasy (MDMA) ^b	8.1	12.0	11.8	13.2	5.7	11.0
Cocaine	5.1	10.4	7.3	11.6	3.8	9.3
Crack ^c	0.7	2.5	0.8	3.8	0.6	1.4
Other Cocaine ^d	5.2	9.3	7.5	10.5	3.8	8.4
Heroin	0.4	1.9	0.6	2.8	0.3	1.1
With a Needle ^e	0.1	0.9	0.3	1.4	*	0.3
Without a Needle ^e	0.8	1.7	1.3	3.0	0.4	0.4
Narcotics other than Heroin ^f	10.8	18.3	13.4	19.5	9.2	17.1
Amphetamines, Adjusted ^{f,g}	15.3	17.1	17.9	19.7	13.7	14.7
Methamphetamine ^e	0.9	2.9	1.0	3.7	0.8	2.2
Crystal Methamphetamine (Ice) ^e	*	1.1	*	1.4	*	0.8
Sedatives (Barbiturates) ^f	5.4	9.8	5.9	10.4	5.0	9.3
Tranquilizers ^f	7.8	11.1	9.5	10.9	6.7	11.3
Alcohol	78.0	79.3	80.7	77.3	76.2	81.1
Been Drunk ^b	66.5	71.0	71.0	68.5	63.6	73.2
Flavored Alcoholic Beverages ^h	67.5	69.9	66.4	70.5	68.1	69.0
Cigarettes	—	—	—	—	—	—
Steroids ^e	0.8	0.3	1.8	0.8	*	*
<i>Approximate Weighted N =</i>	<i>1,090</i>	<i>690</i>	<i>430</i>	<i>330</i>	<i>660</i>	<i>360</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. ' * ' indicates a prevalence rate of less than 0.05%.

' — ' indicates data not available.

See footnotes following Table 8-4.

TABLE 8-2
Annual Prevalence of Use for Various Types of Drugs, 2013:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
by Gender
(Entries are percentages.)

	Total		Males		Females			
	Full-Time		Full-Time		Full-Time			
	College	Others	College	Others	College	Others		
Any Illicit Drug ^a	38.9	39.8	43.8	41.5	35.8	38.2		
Any Illicit Drug other than Marijuana ^a	19.0	20.2	23.5	22.7	16.1	17.9		
Marijuana	35.5	37.2	40.1	40.0	32.6	34.6		
Synthetic Marijuana ^b	2.3	5.1	3.8	5.5	1.3	4.7		
Inhalants ^b	0.5	0.8	0.5	1.2	0.5	0.5		
Hallucinogens	4.5	5.5	7.3	8.0	2.7	3.3		
LSD	2.6	3.3	4.2	5.0	1.6	1.9		
Hallucinogens other than LSD	3.7	4.4	6.4	6.3	1.9	2.8		
Ecstasy (MDMA) ^b	5.3	5.7	7.5	6.3	3.9	5.2		
Salvia ^b	1.0	1.9	0.6	2.5	1.3	1.4		
Cocaine	2.7	5.3	4.2	5.1	1.8	5.4		
Crack ^c	0.3	0.8	0.3	1.1	0.3	0.6		
Other Cocaine ^d	2.8	4.6	4.6	4.5	1.7	4.8		
Heroin	0.3	1.2	0.5	1.7	0.1	0.8		
With a Needle ^e	0.1	*	0.3	*	*	*		
Without a Needle ^e	0.5	1.3	1.1	2.2	0.2	0.4		
Narcotics other than Heroin ^f	5.4	9.6	7.4	10.9	4.1	8.4		
OxyContin ^{b,f}	2.3	4.4	4.0	5.8	1.2	3.0		
Vicodin ^{b,f}	4.4	9.5	7.8	11.0	2.1	8.0		
Amphetamines, Adjusted ^{f,g}	10.6	8.9	12.4	10.0	9.4	8.0		
Ritalin ^{b,f}	3.6	2.3	4.4	3.2	3.0	1.4		
Adderall ^{b,f}	10.7	6.8	13.3	7.3	8.9	6.3		
Methamphetamine ^e	0.4	0.3	0.5	0.7	0.4	*		
Crystal Methamphetamine (Ice) ^e	*	0.7	*	0.5	*	0.8		
Bath Salts (synthetic stimulants) ^b	0.1	0.5	0.2	0.2	*	0.8		
Sedatives (Barbiturates) ^f	2.7	4.4	2.6	4.1	2.8	4.8		
Tranquilizers ^f	4.4	5.4	5.8	5.3	3.6	5.5		
GHB ^e	0.1	0.8	0.3	0.9	*	0.8		
Ketamine ^e	0.9	0.8	1.3	*	0.6	1.5		
Alcohol	75.6	75.0	77.7	73.4	74.2	76.4		
Been Drunk ^b	57.9	58.3	60.3	58.6	56.4	58.2		
Flavored Alcoholic Beverages ^h	57.6	55.0	57.0	46.8	57.9	66.0		
Alcoholic Beverages containing Caffeine ^e	39.1	33.7	41.5	38.6	37.5	29.2		
Cigarettes	23.2	39.3	27.1	42.2	20.7	36.7		
Tobacco using a Hookah ^b	26.1	27.4	28.7	30.2	24.3	24.6		
Small Cigars ^b	19.0	24.1	31.1	31.6	10.8	16.5		
Dissolvable Tobacco ^e	0.2	0.9	0.6	1.2	*	0.6		
Snus ^e	4.8	9.6	9.6	18.0	1.5	2.5		
Steroids ^e	0.8	0.2	1.8	0.3	*	*		
	<i>Approximate Weighted N =</i>		<i>1,090</i>	<i>690</i>	<i>430</i>	<i>330</i>	<i>660</i>	<i>360</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

See footnotes following Table 8-4.

TABLE 8-3
Thirty-Day Prevalence of Use for Various Types of Drugs, 2013:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Any Illicit Drug ^a	22.5	27.2	29.8	31.5	17.9	23.3
Any Illicit Drug other than Marijuana ^a	8.2	10.5	10.2	11.7	7.0	9.5
Marijuana	20.6	25.8	27.8	31.2	15.9	20.7
Inhalants ^b	0.1	0.3	*	0.7	0.1	*
Hallucinogens	1.0	1.7	2.0	2.4	0.4	1.1
LSD	0.4	0.8	0.6	1.6	0.3	0.1
Hallucinogens other than LSD	0.8	1.1	1.7	1.1	0.3	1.1
Ecstasy (MDMA) ^b	0.8	1.4	1.0	1.1	0.6	1.6
Cocaine	0.9	2.6	1.3	2.3	0.6	2.8
Crack ^c	0.3	0.2	0.2	*	0.3	0.4
Other Cocaine ^d	0.9	2.0	1.1	1.9	0.8	2.2
Heroin	0.2	0.3	0.5	0.6	0.1	*
Narcotics other than Heroin ^f	1.5	4.1	2.5	5.0	0.9	3.4
Amphetamines, Adjusted ^{f,g}	5.3	4.1	6.1	5.1	4.7	3.2
Crystal Methamphetamine (Ice) ^e	*	0.2	*	0.5	*	*
Sedatives (Barbiturates) ^f	0.9	1.8	0.4	1.7	1.2	1.9
Tranquilizers ^f	1.2	1.9	1.6	1.9	0.9	1.9
Alcohol	63.1	57.2	65.3	58.6	61.7	56.0
Been Drunk ^b	40.2	33.6	40.7	34.6	39.9	32.7
Flavored Alcoholic Beverages ^h	29.1	27.0	27.1	24.4	30.4	30.5
Cigarettes	14.0	28.1	17.5	30.6	11.9	25.9
Steroids ^e	*	*	*	*	*	*
<i>Approximate Weighted N =</i>	<i>1,090</i>	<i>690</i>	<i>430</i>	<i>330</i>	<i>660</i>	<i>360</i>

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

See footnotes following Table 8-4.

TABLE 8-4
Thirty-Day Prevalence of Dailyⁱ Use for Various Types of Drugs, 2013:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
by Gender

(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Marijuana	5.1	10.0	8.7	13.0	2.9	7.3
Cocaine	*	0.1	*	0.1	*	*
Amphetamines, Adjusted ^{f,g}	*	0.2	*	0.3	0.1	0.1
Alcohol						
Daily	3.6	4.1	5.6	5.6	2.3	2.7
5+ Drinks in a Row in Last 2 Weeks	35.2	31.2	43.3	34.4	30.1	28.4
Cigarettes						
Daily	5.6	20.2	7.0	20.7	4.7	19.7
1/2 Pack+/Day	2.4	11.2	3.3	11.7	1.8	10.7
<hr/>						
<i>Approximate Weighted N =</i>	1,090	690	430	330	660	360

Source. The Monitoring the Future study, the University of Michigan.

Notes. * * * indicates a prevalence rate of less than 0.05%.

See footnotes on the following page.

Footnotes for Tables 8-1 through 8-4

^aUse of any illicit drug includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total *N* in 2013 for college students is approximately 550.

^cThis drug was asked about in five of the six questionnaire forms. Total *N* in 2013 for college students is approximately 900.

^dThis drug was asked about in four of the six questionnaire forms. Total *N* in 2013 for college students is approximately 730.

^eThis drug was asked about in two of the six questionnaire forms. Total *N* in 2013 for college students is approximately 360.

^fOnly drug use that was not under a doctor's orders is included here.

^gBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^hThis drug was asked about in one of the six questionnaire forms. Total *N* in 2013 for college students is approximately 180.

ⁱDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.

Chapter 9

TRENDS IN DRUG USE AMONG COLLEGE STUDENTS

In recent years college students and high school seniors have shown simultaneous increases in marijuana use as well as in the index of any illicit drug use. This secular trend is different from prior trends in which drug use increases among college students either preceded or followed those among younger students. During the 1960–70s epidemic, illicit drug use increased dramatically among U.S. college students, then spread quickly to their noncollege age peers and eventually down the age spectrum to high school and even middle school students. The diffusion process seems to have reversed during the subsequent “relapse” in the 1990s, when use increased first among those in early adolescence and then radiated *up* the age spectrum as those cohorts grew older (a cohort effect). Use subsequently declined among adolescents; this decline, like the preceding increase, radiated up the age spectrum.

Again, we define college students as *follow-up respondents one to four years past high school who report that they were taking courses as full-time students in a two- or four-year undergraduate college at the beginning of March in the year in question.*

For comparison purposes, trend data are provided on the remaining follow-up respondents in this age band, who are also one to four years past high school but do not meet our definition of college students (Figures 9-1 through 9-15c). Because the proportion of an age group in college declines steadily with the number of years beyond high school, this comparison group is slightly older on average than the college-enrolled group.

The proportion of young adult high school graduates one to four years beyond high school who attend college full-time has increased considerably since the MTF follow-ups began. In 2013, about 61% of the weighted number of follow-up respondents one to four years past high school met our definition of college students, compared with only 38% in the 1980 survey. This means, among other things, that the size of our college student sample has risen over the years and, conversely, the size of the noncollege sample of the same age has diminished.

The reader is reminded that the difference between the college group and the noncollege group provides an estimate of the degree to which college students’ usage levels for various substances are above or below other high school graduates in this age band. Were we able to include the high school dropout segment in the calculation

for the noncollege group, many of the differences with the college-enrolled would be accentuated.⁷⁷

For each year, approximately 1,100–1,500 weighted respondents constitute the college student sample (see Table 9-5 for *Ns* per year) and roughly 700–1,700 respondents constitute the noncollege group one to four years beyond high school. Trend comparisons for these two groups are provided in this chapter. The reported results begin with 1980, the first year that enough follow-up surveys had accrued to characterize young people one to four years past high school. The 2013 survey is thus the 34th in the annual series on college students.

Throughout much of the chapter, trends for the 12th-grade samples are included for comparison purposes. It is important to keep in mind that the total 12th-grade samples are shown, and that there are substantial differences in rates of substance use within those samples between the college-bound and those who do not plan to complete college. As shown extensively in [Volume I](#) and in [Occasional Paper 81](#), 12th-grade students expecting to complete college are far less likely to smoke cigarettes and also less likely to use most other substances. So when considering figures that show higher rates of use among all 12th graders (regardless of college expectations) than among college students, it should not be concluded that usage declined after college entrance; the fact is that the college-bound were already lower in usage rates than other 12th graders for almost all substances.

One additional point relevant to interpreting differences over time for those attending college and those not attending college, both in terms of the differences between them and trends over time for either taken separately: the proportion of college students who are female has risen substantially since 1980. In 1980, females constituted about 50% of our college respondents, but by 2013 they constitute 61%. Females thus constitute a declining proportion of the noncollege group. As will be discussed below, we have charted the trends separately for male and female college students to permit an assessment of what effect these changing proportions may have on the overall rates observed for college students.

TRENDS IN PREVALENCE, 1980–2013: COLLEGE STUDENTS VERSUS THOSE NOT ENROLLED IN COLLEGE

- The proportion of college students using *any illicit drug* has continued to rise gradually since 2006, driven primarily by an increase in marijuana use. During the first decade of reportable MTF college student data, between 1980

⁷⁷Panel analyses of samples from the high school classes of 1995–1997, followed for an eight-year period beginning when they were in 8th grade, clearly show that those who dropped out of high school had distinctly higher rates of substance use both before and after they left school. See Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

and 1991, college use of any illicit drug dropped fairly steadily—from 56% to 29%, a decrease of nearly half (Table 9-2 and Figure 9-1). After 1991, annual (and also 30-day) prevalence held fairly steady for a couple of years before beginning to rise, reaching 38% in 1998 and again in 2001—still well below the 1980 peak. There was a modest rise after 2006 (when it was 34%) to an annual prevalence rate of 39% in 2013. Their noncollege peers moved similarly until 2000, when they exhibited a four-percentage-point increase due largely to their sharper increases in marijuana, amphetamine, and tranquilizer use; the noncollege rate remained slightly above the rate for college students for several years; then, in 2007, the noncollege rate dropped about four percentage points, just about eliminating the gap. Twelfth graders showed a parallel trajectory to the other two groups in the decline phase through 1991, followed by a much steeper increase in use through 1997, and then leveling after 1998. Use among high school seniors declined some after 1999 (by about six percentage points), whereas among college students there was less decline. As a result, all three groups had quite similar prevalence rates by 2007. After that, use increased among the seniors but later among the college students, creating some new divergence before they converged in 2013. We believe the divergence among the three groups and subsequent convergence of college students and high school seniors, at least, reflect cohort effects. After 2007 (2006 for college students), all three groups showed some increase in the annual prevalence of any illicit drug use—due largely to a turnaround in their use of marijuana, as described below—but the increase has been greatest and longest among the seniors, likely once again giving rise to a cohort effect. Usage estimates for the noncollege segment have been rather unstable in recent years (see Figure 9-3a), likely due to the smaller sample sizes that comprise that segment.

Figure 9-2 shows that since 1980 the noncollege segment has usually had the highest levels of use of *any illicit drug other than marijuana*. The exception was during most of the 1990s (the relapse phase in the epidemic), when use among 12th graders rose sharply and exceeded the noncollege segment. The noncollege group also showed a slightly lagged increase during that phase and passed the 12th graders in the early 2000s. An increase among college students also occurred after around 1994, but it lagged considerably behind the upturn among 12th graders, reflecting a cohort effect. From 1986 through 2009—a twenty-three year interval—college students exhibited the lowest rates of use of any illicit drug other than marijuana. From 1980 to 1994, use of any illicit drug other than marijuana declined appreciably among college students, with their annual prevalence dropping by nearly two thirds from 32% to 12% (Table 9-2). This generally paralleled the trends for the noncollege group and the 12th graders, indicating a secular trend during that period. All three groups showed some increase in use during the early 1990s; however, the rise in use of illicit drugs other than marijuana was again not as sharp among college students as it was in the other two groups, and it began two years later than among the 12th graders and one year later than among the noncollege group (Figure 9-2). This pattern is more consistent with a cohort

effect. After 1999, use among 12th graders leveled off, whereas the college students and noncollege segment showed a continuing increase. In fact, the college students and noncollege respondents continued to show an increase in their annual prevalence rate from 1998 through 2004, before declining from 2005 through 2007 among the noncollege group and from 2005 through 2008 among the college students. Since 2008 the rate has increased among the college students and declined steadily among those in the noncollege group (at least through 2012), closing the considerable gap observed with both college students and 12th graders. In 2012 all three groups had comparable prevalence rates, at 17%-18%. In 2013, college students and their noncollege age peers showed parallel increases in use (roughly 2 percentage points -- nonsignificant), while use among 12th graders remained unchanged. (The increase in use of any illicit drug other than marijuana among college students, from 15% in 2008 to 19% in 2013, is significant.) Thus by 2013 college student use exceeded that of 12th graders, approached that of the noncollege segment, and reached a new recent peak rate. This increase appears attributable mostly to college students' increased use of *amphetamines* (without a doctor's orders) and of *ecstasy*.

Trends during the 1980s for most individual classes of illicit drugs tended to be similar among the three groups. During the 1990s there was more divergence, with college students usually showing later and lesser increases than 12th graders, and, for some drugs, less increase than their noncollege peers.

- Since 2000, the annual prevalence of *marijuana* use among college students reached a recent high point of 36% in 2001, declined to 30% in 2006, and then increased to 36% by 2013. In an earlier period, from 1981 through 1991, annual prevalence of marijuana use dropped by nearly half from 51% to 27% among college students (Figure 9-3a). Their noncollege peers showed a comparable decline over the same time interval, as did the 12th graders; and the annual prevalence rates for both groups were comparable across that interval. Use among 12th graders rose sharply after 1992, while use among college students and noncollege respondents rose more gradually. From 1991 through 1998, annual prevalence rose by 14 percentage points among 12th graders, compared to 10 percentage points among college students and 7 percentage points among the noncollege group. As a result, the 12th graders came to exhibit the highest rate of marijuana use in the last half of the 1990s. The 12th graders were the first to show a leveling off in marijuana use (in 1998), followed by the college students in 1999 and the noncollege group in 2002. All three groups had very similar rates of use by 2005 after some decline, particularly among the 12th graders. The college students and high school seniors both showed some decline in 2006; then both showed a gradual increase in their marijuana use from 2006 through 2011, with the sharpest increase occurring among the 12th graders, indicating in both cases the end of the gradual improvement in marijuana use seen earlier in the decade. Among college students, annual prevalence was 36% in 2013, up from 30% in 2006. Among the noncollege group the increase since 2007 was more erratic, but in

2013 their annual prevalence reached 40%—slightly above that of the college students.

- **Daily marijuana** use has consistently been highest among the noncollege stratum, next highest among the 12th graders, and lowest among those in college. The differences have been greatest in periods of relatively high use and diminished considerably when use was at its nadir at the beginning of the 1990s. Daily marijuana use has varied widely in all three groups since 1980, when data on college students and their noncollege age peers were first available. The period from 1980 through 1992 saw a large proportional decline in daily use in all three groups with rates falling by half or more. Since 1992 the rates have climbed substantially in all three groups, though there was a period of leveling among high school seniors from 1999 through 2009, among college students from roughly 2003 through 2006, and among the noncollege group from 2003 through 2010. Among college students daily use has risen most recently from 3.5% in 2007 to 5.1% in 2013 (Figure 9-3b). While daily use remained level among the 12th graders since 2011, it has continued to rise gradually among college students and their noncollege peers.
- **Amphetamine** use without medical supervision among college students (Figure 9-11) rose from 2008 (5.7%) through 2012 (11.1%) but did not rise any further in 2013 (10.6%). The 1980s saw a considerable decline of annual prevalence among college students, from 22% in 1981 to 4% in 1991. Proportionately, this was a larger drop than among 12th graders, but fairly parallel to the overall change among the noncollege group. Amphetamine use among college students and their noncollege age peers began to increase in both groups after 1992 and 1993, respectively, through 2001, with a leveling in 2002. During the 1990s and early 2000s, the prevalence rates for amphetamine use in all three groups remained well below the rates observed in the early 1980s. Since 2002 there have been some small nonparallel changes among the three groups, with amphetamine use among college students (who have consistently had the lowest rate of use since the mid-1980s) holding steady through 2008, while use among 12th graders and the noncollege group declined, nearly closing the gap. In 2009, prevalence rates were similar for the college and noncollege groups (7.5% and 7.7%), and slightly lower among 12th graders (6.6%). A recent increase in amphetamine use among just the college students now places them highest in 2013 (10.6%), up from 5.7% in 2008. It seems likely that an increased interest in using these drugs to improve academic performance has contributed to this change. **Adderall** was used by more than three times as many college students (10.7%) as was **Ritalin** (3.6%) in 2013.
- Use of **inhalants** has been very low among college and noncollege respondents since 1980, when rates were first measured (Figure 9-4). Twelfth graders have consistently had higher rates of inhalant use than either of these segments of the young adult population; and as is documented in Volume I, the 8th and 10th graders have had higher levels of use still. There has thus

been an age effect, with use declining with increasing age. All three groups being discussed here have trended in parallel, with an increase in use from around 1981 through 1995, followed by a long decline thereafter. The increase and subsequent decline were substantially more pronounced among 12th graders, opening and then shrinking the gap between them and the two young adult groups. The annual prevalence of inhalant use among college students is now quite low (0.5% in 2013 compared with a high of 4.1% in 1997).

- Annual prevalence of **LSD** remains fairly low among college students in 2013 (Figure 9-6). During the early 1980s, one of the largest proportional declines observed among college students occurred with this drug: annual prevalence fell from 6.3% in 1982 to 2.2% in 1985. After 1989, use in all three groups increased, with the prevalence among college students reaching 6.9% by 1995. After 1995, use fell gradually among college students, their age-mates, and 12th graders until 2001; followed in 2002 by a particularly sharp decrease in all groups. As a result there was a considerable convergence in usage rates, which has remained since then. College students maintained lower levels of use than the other two groups from the mid-1990s through 2007. Use rose slightly in all three groups in 2008, but there has been little consistent difference among them. In 2013 annual prevalence was 2.6% among college students, 3.3% in the noncollege group, and 2.2% among 12th graders.
- Use of **hallucinogens other than LSD** (which primarily involves the use of psilocybin known as mushrooms or “shrooms”) followed a track somewhat parallel to LSD use, at least up until about 2000 (Figure 9-7). That meant declining use in all three groups from the early 1980s through the early 1990s, followed by rising use during the 1990s and then some leveling. But the trends for these other hallucinogen diverged from those for LSD after about 2000, with an increase in their use, including among college students, just before and after the drop off in LSD use in 2002. While the use of LSD was higher than that of the other hallucinogens in the first two decades of the study, its use is now lower. Among college students the annual prevalence for hallucinogens other than LSD was 3.7% in 2013: it was 2.6% for LSD.
- The use of **ecstasy (MDMA)** by college students stands at 5.3% in 2013, up from 2.2% in 2007 (Figure 9-8). College use and use by their noncollege age peers began to rise after 1994 and their rates tracked closely through about 2000 (Figure 9-8). The 12th graders had questions about ecstasy use added in 1996 and their usage rates similarly tracked those of the other two groups through about 2000. After 1997 there was a sharp increase in use in all three groups. The annual prevalence for college students, for example, rose from 2.4% in 1997 to 9.2% in 2001, and use rose even more among the noncollege group. All three groups declined sharply from 2001 to 2004, when annual rates were at 2.2% for college students, 2.7% for 12th graders, and 4.0% for the noncollege segment. Both the college and noncollege groups showed some increase in use by 2013, with annual prevalence rates reaching 5.3% and

5.7%, respectively. While none of these groups has usage rates comparable to what they were in 2001, all three showed some resurgence in use in recent years, though over the past two years use among 12th graders has fallen fairly sharply.

- Ever since data were first available in 1980, college students have had the lowest rates of non-medical *sedative (barbiturate)* use among the three groups (Figure 9-12). At that early date, sedative (barbiturate) use was already quite low among college students (at 2.9%), but it still fell by more than half to 1.3% by 1985. This proportional decline was sharper than among 12th graders and less sharp than among the noncollege respondents, both groups that started at considerably higher levels of use. Annual prevalence remained essentially unchanged between 1985 and 1993 for all three groups. All three then showed a gradual increase in use between 1993 (or 1994 for college students) and 2004 for college students and 2005 for the other two groups. After 2005, declines in use appeared in all three groups through 2011 (2013 for the noncollege stratum), before showing a rise in use since then.
- Figure 9-13 shows that the annual prevalence of non-medical *tranquilizer* use among college students dropped by nearly three fourths in 1980–1994, from 6.9% to 1.8%—a period in which use declined in the other two groups as well. After this long period of decline, tranquilizer use by college students increased gradually, returning to 6.9% by 2003. Use by the noncollege segment and by 12th graders dropped more sharply from 1980 through 1992, eliminating the differences among the three groups. Use rose after 1992 for all, but the noncollege group showed the largest gain after 1999, again creating some differences. By 2002, tranquilizer use was once again at or near its recent high in all three groups. In 2003, however, the noncollege group and the 12th graders showed their first declines in recent years, thus narrowing the differences among the three groups. From 2004 through 2013, all three groups showed fair declines in use.
- The non-medical use of *narcotics other than heroin*⁷⁸ (Figure 9-10a) rose after about the mid 1990s for all three young adult groups. The overall trends in use have been quite parallel to those for sedatives (barbiturates) and tranquilizers. Prior to then, the use of narcotics other than heroin by college students was down to about half by 1994 from what it was in 1980 (2.4% in 1994 vs. 5.1% in 1980) as a result of a fairly gradual decline over that 14-year interval. This trend closely paralleled use among their noncollege counterparts and 12th graders. As with a number of other drugs, use among 12th graders began to rise after 1992, but use among college students did not begin to increase until after 1994, likely due to a cohort effect. In 2003, annual

⁷⁸As discussed in chapters 4 and 5, because the questions about narcotics other than heroin were changed in 2002, the prevalence figures are adjusted estimates. See the earlier discussion for details.

prevalence among college students reached an historic high point of 8.7% before leveling for three years. It then declined some from 8.8% in 2006 to 5.4% by 2013. Use among 12th graders leveled after reaching an historic high of 9.5% in 2004, but it then declined to 7.1% by 2013. The noncollege group emerged after 2000 as the most heavily using group for the first time, as their use kept increasing through 2005, reaching an all-time high of 13%. After that, use in the noncollege group declined to 9.6% in 2013—and remains the highest of the three groups.

- Although data about non-medical *OxyContin* and *Vicodin* use were not collected until 2002 (Figures 9-10b and 9-10c, and Table 8-2), these drugs help to explain past differences between the college and noncollege segments in use of narcotics other than heroin. The noncollege group had annual prevalence rates up to twice that for college students in the use of both drugs when their use was first measured in 2002, but the comparisons among the three groups have changed since

Annual prevalence of non-medical *OxyContin* use among college students rose fairly steadily, from 1.5% in 2002 to 5.0% in 2009, before dropping significantly to 2.3% in 2013 (Figure 9-10c). Use in the noncollege segment rose from 2002 (3.3%) to 2009 (6.2%) before it also fell in 2010 and continues to be lower in 2013 (4.4%); the trend line has been quite uneven, likely due to the limited numbers of cases in this segment. (Questions about *OxyContin* and *Vicodin* are in only three of the six questionnaire forms.) Among 12th graders, *OxyContin* use rose from 4.0% in 2002 to 5.1% in 2010, before declining to 3.6% in 2013. It is clear that *OxyContin* use increased among college students between 2002 and 2009, closing the previously existing gap among the three groups; however, its use has declined sharply among the college students since then, again opening a gap between them and the other two groups.

Vicodin use without medical supervision (Figure 9-10b) showed a somewhat different pattern of change, with annual prevalence among all three groups remaining fairly level—and substantially higher than for *OxyContin*—from 2002 through 2008, and then declining in 2010. As with *OxyContin*, the noncollege segment has consistently had higher *Vicodin* use than the college students. Twelfth-grade *Vicodin* use has been in between. Because of the limited numbers of cases, the trends in use of *Vicodin* and *OxyContin* have generally been uneven in the young adult groups. Since 2010, reported *Vicodin* use has declined fairly sharply in all three groups.

- Over recent years, the rates of *cocaine* use among college students, noncollege peers, and 12th graders (Figure 9-9) have declined to levels below those in the 1990s and far below those in the 1980s. Like the 12th graders, college students showed a relatively stable pattern of cocaine use between 1980 and 1986, when their usage levels (and those of their noncollege age peers) were considerably higher than those observed among 12th graders. This level period was followed by a dramatic drop of nearly nine tenths in annual

prevalence among college students, from 17.1% in 1986 to 2.0% by 1994. Their noncollege counterparts also showed a large but somewhat less dramatic decline, from 18.9% in 1986 to 5.1% in 1994. Because use among college students also dropped more sharply than among 12th graders, there was little or no difference between those two groups in annual prevalence rates for cocaine use between 1990 and 1992; but use then began to rise among 12th graders after 1992, while an increase didn't begin among college students until after 1994, reflecting a cohort effect. Use did not begin to rise until after 1995 among the noncollege segment, which maintained the highest rate of use since then. After 2000 the 12th graders and college students have had similar rates of use and parallel trends, while use in the noncollege stratum has been considerably higher. Since around 2006 all three groups have shown declines in use, with the noncollege group showing the greatest decline. All three groups now have rates of cocaine use below those attained in the relapse phase of the illicit drug epidemic in the 1990s, with the noncollege group showing the greatest decline but still the highest level of use.

- College students have shown some unique shifts in *alcohol* use. Despite different trend patterns among the three groups, college students have exhibited the highest levels and greatest constancy in *occasions of heavy drinking* since they were added to MTF surveys in 1980 (Figure 9-14d). Occasions of heavy drinking are defined as having five or more drinks in a row at least once during the prior two weeks. Over the 33-year interval from 1980 through 2013 college students' rates of such drinking declined 9 percentage points (from 44% to 35%), while noncollege respondents' rates declined 10 percentage points (41% to 31%) and high school seniors' rates declined 19 percentage points (41% to 22%). As can be seen in Figure 9-14d, both the noncollege segment and 12th graders showed fairly substantial declines in the prevalence of occasions of heavy drinking from 1981 through 1990. In contrast, college students showed no decline from 1981 to 1986 and then only a modest decline of five percentage points from 1986 through 1993. Between 1981 (when all three populations were very close in use) and 1992, this measure of heavy drinking dropped by 14 percentage points among 12th graders, by 11 percentage points among the noncollege respondents, but by only 2 percentage points among college students. After 1992, occasions of heavy drinking began to rise among 12th graders, while still declining some among college students—likely reflecting a cohort effect emerging during this period, similar to that observed for a number of illicit drugs—narrowing the gap somewhat. Drinking at that level subsequently began to increase among the noncollege segment after 1995, and by less among college students after 1996—increases that continued into 2001. Between 2001 and 2008, college students held fairly steady in their rates, before showing some decline since 2008, while the noncollege segment held steady from roughly 2003 to 2007, followed by some decline and then a leveling. Meanwhile, among 12th graders, occasions of heavy drinking started a gradual decline after 1998 that continued into 2011, slightly enlarging the difference between them and the

other two groups. Once again there is evidence of cohort effects since the early 1990s, with the inflection points occurring later for the older strata.

Why did college students' heavy drinking show little decline for a decade (1981–1991) compared to their noncollege peers and 12th graders? One possibility is that campuses provided some insulation from the effects of changes in the drinking age laws that took place in many states during that interval. Similarly, entrenched in many college campuses is a culture of binge drinking which has proven impervious to many societal trends (and intervention attempts).⁷⁹ Also, individuals who are under the legal drinking age in college are mixed in with peers who are of legal age to purchase alcohol; this was no longer true in high schools and less true, perhaps, for many of those ages 19 to 22 who are not in college. Finally, much alcohol advertising and promotion was and is directed specifically at the college student population. Nevertheless, it appears that the continuing decline in heavy drinking among 12th graders is beginning to resonate among college students.

Daily drinking among noncollege young adults has shown more change than occasions of heavy drinking among college students (Figure 9-14c). College students' daily drinking estimates—which appeared a little less stable in the 1980s, perhaps due to smaller sample sizes at that time—showed little or no decline between 1980 (6.5%) and 1984 (6.6%), but a considerable decline from 1984 through 1995 (to 3.0%), followed by a period of some increase during and after the relapse phase in the drug epidemic among high school students, reaching 5.0% in 2002. After 2002 their daily drinking dropped to 3.7% in 2004, about where it is in 2013 (3.6%). Twelfth graders showed a somewhat similar pattern of daily drinking with a long period of decline, followed by an earlier reversal beginning in 1994. After 1998, 12th grade daily drinking resumed its decline, reaching 2.1% by 2011 and leveling thereafter. Of the three groups, the noncollege respondents have generally had the highest rate of current daily drinking and the 12th graders the lowest. However, after 2008 daily drinking levels have been fairly comparable between the college students and their noncollege age peers.

- The 30-day prevalence of **cigarette smoking** (Figure 9-15a) among college students has declined dramatically over more than a decade, falling by more than half from a recent high of 31% in 1999 to 14% in 2013, and their *daily* smoking rate has fallen by about two-thirds over the same interval (from 19% to 6%). In the early 1980s, cigarette smoking among U.S. college students declined modestly. Thirty-day prevalence fell from 26% to 22% between 1980 and 1984, remained fairly stable through 1990 (22%), then increased gradually but substantially, reaching 31% by 1999. In 2000 the first evidence of a new decline in smoking among college students began to appear, two

⁷⁹Schulenberg, J. E., & Maggs, J. L. (2002). A developmental perspective on alcohol use and heavy drinking during adolescence and the transition to young adulthood. *Journal of Studies on Alcohol*, Supplement 14, 54–70.

years after smoking had begun to decline among 12th graders—this lag reflecting a cohort effect. The noncollege group, which has consistently had the highest smoking rate of the three groups, showed a fairly consistent decline from 1980 through 1990, an offsetting increase through 2001, and then a considerable decline since then. *Because the noncollege segment has shown a more moderate decline so far, their smoking rate is now much higher than that of the college students.*

While smoking rates have consistently been lower among college students than the noncollege segment, the trend lines for these two groups converged some after 1984, as smoking rates more or less stabilized among college students but continued to decline among young adults not in college (Figure 9-15a). In fact, between 1989 and 1991, use began to rise among college students while continuing to decline among noncollege respondents. Both groups showed fairly parallel increases in smoking between about 1991 and 1999, after which use continued to increase among the noncollege segment, but began to decline among college students, opening up a large difference between them. (Twelfth graders exhibited an increase from 1992 to 1997 and their use has declined significantly since then.) The popularity of Camel cigarettes among the college-bound may help to explain some of the narrowing of the gap between college students and their age peers in the 1990s. The Joe Camel advertising and promotion campaign, which commenced in the late 1980s and ended in the late 1990s, may have succeeded in initiating more college students (particularly males) to smoking than had been the case previously or since.

- For many prescription-type drugs—*amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—differences between college students and their noncollege age peers narrowed over the years, particularly through the early 1990s. Much of this was due to general overall declines in usage rates during the 1980s, but may also reflect the increasing proportion of the age group going to college. After that, the differences between these two groups increased for sedatives (barbiturates) and tranquilizers as use in general grew.

The overall drug use trends among college students parallel the trends among 12th graders, though after the early 1990s they were generally lagged by a few years; still, declines in many drugs from 1980 to 1990 were proportionately larger among 19- to 22-year-olds (both college and noncollege) than among 12th graders. Despite parallel trends in the early 1990s, 12th graders have shown larger, and usually earlier, increases in the use of a number of drugs in the years since; as indicated in *Volume I*, 8th and 10th graders showed increases a year earlier than 12th graders. *Clearly the upsurge, or what we have called a “relapse phase” in the illicit drug epidemic, did not originate on the nation’s college campuses, as did the earlier epidemic. It originated among secondary school students—and the younger ones at that—and was carried up the age spectrum through generational replacement.*

GENDER DIFFERENCES IN TRENDS AMONG COLLEGE STUDENTS

As mentioned earlier, recent decades have seen a gradual rise in the proportion of college students who are female. Females constituted 50% of our 1980 sample of college students compared to 61% of our 2013 sample. Given that substantial gender differences exist in the use of some drugs, we have been concerned that apparent long-term trends in the levels of drug use among college students (and/or among the noncollege group) might actually be attributable to changes in the gender composition of that population. For this reason, in particular, we present separate trend lines for males and females. Gender differences in trends are illustrated in the lower panels of Figures 9-1 through 9-15c. In general, trends in use of the various drugs have been highly parallel for male and female college students, as an examination of the relevant figures will show. The most noteworthy exceptions are mentioned below.

- Certain drug use measures showed a convergence between the genders as use rates declined to low levels in the early 1990s. This was true for the use of *any illicit drug* and *any illicit drug other than marijuana*. Rates for male college students have consistently been higher than those for female college students.
- Before 2000, *marijuana* use is another such example, with some gender convergence in the rates between 1980 and 1991 as overall use declined, and then some gender divergence between 1991 and 1999 as usage rates rose. (Marijuana use has been consistently higher among males than among females.) After 2001, the two genders diverged further, with use among males remaining essentially unchanged through 2008 and use among females decreasing (Figure 9-3a). The divergence continued from 2009 through 2011, this time with college males increasing their marijuana use while use among females held fairly steady until 2012, when it rose among females as well (Figure 9-3b).
- From 1999 to 2005, *LSD* use dropped more steeply among males than among females, offsetting sizeable previous differences in which males had higher use, and bringing the genders close together at very low prevalence rates (Figure 9-6). The small increases in use that have occurred since 2005 have been greater among males.
- Use of *hallucinogens other than LSD* has dropped for both genders since 2002 or 2003, with males remaining consistently higher (Figure 9-7). Since then the gap enlarged again as use rose in the 1990s and early 2000s. Rates of *ecstasy (MDMA)* use have been quite similar for male and female college students since measures were first introduced in 1989, and changes in their usage levels have tracked closely (Figure 9-8). There was some divergence in 2013 with use by males rising to 7.5% and use by females falling to 3.9%. It remains to be seen if this difference is replicated in the future.

- Trends in the non-medical use of *narcotics other than heroin* have generally moved in parallel for both male and female college students (with males almost always higher, except during the nadir in use at the beginning of the 1990s when their rates were equivalent); however, because there was a considerable increase in use after 1993, the gap between the genders widened. Both genders showed a decline in use after 2006, narrowing the gap a bit, at least until use leveled in 2013 (Figure 9-10a).
- After 1986, *cocaine* use, which had been substantially higher among males, dropped more steeply for males than for females in general and among male college students in particular, considerably narrowing the sizable gap between genders (Figure 9-9). Since 1991, both genders have moved in parallel, with males reporting somewhat higher usage rates.
- Non-medical *amphetamine* use (Figure 9-11) also showed some convergence in the 1980s due to a greater decline among males. After 1989 the trends were quite parallel through about 2008, with males generally having a slightly higher annual prevalence rate. Both genders have shown some increase in use since 2008 with some modest divergence between them.
- The gender differences for non-medical *sedative (barbiturate)* and *tranquilizer* use have been modest through most of the life of the study, with college males usually having slightly higher rates than their female counterparts. After 1995 a somewhat larger gap emerged for tranquilizers, again with males being higher. Tranquilizer use by college females peaked in 2003, briefly closing the gender gap, but use by males has consistently been slightly higher since then. Both genders are showing a decline in the use of both drugs in recent years, but particularly for sedatives, which no longer show a gender difference (Figures 9-12 and 9-13).
- Among college students, the *annual* prevalence of *alcohol* use has been virtually identical for the two genders since MTF began (Figure 9-14a). Both have shown a gradual decline over the past thirty years. Prior to 2000, the *30-day* prevalence rate showed a modest difference, with males slightly higher (Figure 9-14b); but that difference disappeared by 2000 as drinking rose some among females. Since then college males have had had slightly higher 30-day rates more years than not. College males have consistently had considerably higher rates of *daily drinking* and *binge drinking* than college females (Figures 9-14c and 9-14d). But since about 2004 the gender gap has been narrowing, with an overall increase in daily drinking by college females and an overall decrease among college males.
- From 1988 through 1994, *occasions of heavy drinking* among college females decreased some (from 37% to 31%); but such drinking among college males declined more, from a high point in 1986 of 58% to 47% in 1995 (Figure 9-14d). From 1998 through 2006 there was some closing of the gender gap in binge drinking, as the rate among college females rose from 31% in

1998 to 34% in 2008, while it actually declined some more, from 52% to 49%, among college males. Since 2007, both genders have shown some decline in binge drinking and a fairly constant gap remains in their rates.

- For the twelve-year interval between 1980 and 1992, the 30-day prevalence of ***cigarette smoking*** was consistently higher among college females than males (Figure 9-15a). However, the gap in 30-day prevalence narrowed because use by female college students declined considerably between 1980 and 1989, while use by male college students did not decline. After 1989, the gap remained quite small and the genders reversed position, with college males catching up to, and passing, females in their rate of smoking by 1994 and then remaining higher thereafter. (A similar reversal had occurred among 12th graders a few years earlier, so the reversal among college students probably reflected a cohort effect.) Both genders exhibited a considerable decrease in 30-day smoking between 1999 and 2011, leaving only a modest difference between them (although the trend line for college males was irregular during this interval). In 2013 the 30-day prevalence rates were 18% and 12% for college males and females, respectively, after a fairly sharp and statistically significant drop between 2011 and 2012 in smoking rates among female college students and a slight increase among males.

While the rise in smoking among college students was longer term and more gradual than in the other two groups, it was nevertheless substantial, rising by nearly half between 1989 (21%) and 1999 (31%). This increase in smoking was sharper among college males than among college females, consistent with the notion that Camel cigarettes' promotion and advertising—which ended in the late 1990s—played a role in the overall increase. As we have reported elsewhere, Camels proved considerably more popular among males, especially among those college-bound and from more educated families.⁸⁰

⁸⁰Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (1999). *Cigarette brand preferences among adolescents* (Monitoring the Future Occasional Paper No. 45). Ann Arbor, MI: Institute for Social Research. Available at: <http://www.monitoringthefuture.org/pubs/occpapers/occ45.pdf>.

TABLE 9-1
Trends in Lifetime Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School

(Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>Approximate Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440
Any Illicit Drug ^a	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	45.5	47.4	49.0	52.9	53.2
Any Illicit Drug other than Marijuana ^a	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5
Marijuana	65.0	63.3	60.5	63.1	59.0	60.6	57.9	55.8	54.3	51.3	49.1	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8
Inhalants ^b	10.2	8.8	10.6	11.0	10.4	10.6	11.0	13.2	12.6	15.0	13.9	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4
Hallucinogens ^c	15.0	12.0	15.0	12.2	12.9	11.4	11.2	10.9	10.2	10.7	11.2	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8
LSD	10.3	8.5	11.5	8.8	9.4	7.4	7.7	8.0	7.5	7.8	9.1	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7
Hallucinogens other than LSD ^c	11.6	9.0	10.6	8.3	9.2	8.1	7.8	6.8	6.2	6.2	6.0	6.0	5.7	5.4	4.4	6.5	6.5	7.5	8.7	8.8
Ecstasy (MDMA) ^d	—	—	—	—	—	—	—	—	—	3.8	3.9	2.0	2.9	2.3	2.1	3.1	4.3	4.6	6.8	8.4
Cocaine	22.0	21.5	22.4	23.1	21.7	22.9	23.3	20.6	15.8	14.6	11.4	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4
Crack ^e	—	—	—	—	—	—	—	3.3	3.4	2.4	1.4	1.5	1.7	1.3	1.0	1.8	1.2	1.4	2.2	2.4
Other Cocaine ^f	—	—	—	—	—	—	—	18.1	14.2	16.0	10.2	9.0	7.6	6.3	4.6	5.2	4.6	5.0	7.4	7.8
Heroin	0.9	0.6	0.5	0.3	0.5	0.4	0.4	0.6	0.3	0.7	0.3	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9
Narcotics other than Heroin ^{g,h}	8.9	8.3	8.1	8.4	8.9	6.3	8.8	7.6	6.3	7.6	6.8	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7
Amphetamines ^{g,i}	29.5	29.4	30.1	27.8	27.8	25.4	22.3	19.8	17.7	14.6	13.2	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9
Methamphetamine ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.1
Crystal Methamphetamine (Ice) ^j	—	—	—	—	—	—	—	—	—	—	1.0	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8
Sedatives (Barbiturates) ^{g,t}	8.1	7.8	8.2	6.6	6.4	4.9	5.4	3.5	3.6	3.2	3.8	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7
Sedatives, Adjusted ^{g,k}	13.7	14.2	14.1	12.2	10.8	9.3	8.0	6.1	4.7	4.1	—	—	—	—	—	—	—	—	—	—
Methaqualone ^g	10.3	10.4	11.1	9.2	9.0	7.2	5.8	4.1	2.2	2.4	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{g,l}	15.2	11.4	11.7	10.8	10.8	9.8	10.7	8.7	8.0	8.0	7.1	6.8	6.9	6.3	4.4	5.4	5.4	6.9	7.7	8.2
Alcohol ^m	94.3	95.2	95.2	95.0	94.2	95.3	94.9	94.1	94.9	93.7	93.1	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0
Been Drunk ⁿ	—	—	—	—	—	—	—	—	—	—	—	79.6	76.8	76.4	74.4	76.6	76.2	77.0	76.8	75.1
Flavored Alcoholic Beverages ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^p	—	—	—	—	—	—	—	—	—	0.4	1.5	1.4	1.7	1.9	0.5	0.8	0.6	1.6	0.9	1.3

→
(Years cont.)

(Table continued on next page.)

TABLE 9-1 (cont.)
Trends in Lifetime Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
(Entries are percentages.)

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
<i>Approximate Weighted N =</i>	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090	
Any Illicit Drug ^a	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	51.4	49.1	49.2	50.5	51.0	+0.4
Any Illicit Drug other than Marijuana ^a	25.8	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	25.6	24.8	24.3	23.8	26.7	+2.9
Marijuana	51.2	51.0	49.5	50.7	49.1	49.1	46.9	47.5	46.8	47.5	46.8	46.6	49.1	47.7	-1.4
Inhalants ^b	12.9	9.6	7.7	9.7	8.5	7.1	7.4	6.3	4.9	6.9	5.5	3.7	5.7	4.3	-1.4
Hallucinogens ^c	14.4	14.8	13.6	14.5	12.0	11.0	10.6	9.1	8.5	8.0	7.8	7.4	7.6	7.8	+0.1
LSD	11.8	12.2	8.6	8.7	5.6	3.7	3.5	3.3	4.3	3.3	4.0	3.7	3.1	4.4	+1.4
Hallucinogens other than LSD ^c	8.2	10.7	11.0	12.8	10.1	10.6	10.1	8.5	8.2	7.8	7.1	6.9	7.2	6.8	-0.4
Ecstasy (MDMA) ^d	13.1	14.7	12.7	12.9	10.2	8.3	6.9	5.4	6.2	6.5	6.2	6.8	8.7	8.1	-0.6
Cocaine	9.1	8.6	8.2	9.2	9.5	8.8	7.7	8.5	7.2	8.1	6.6	5.5	5.2	5.1	-0.1
Crack ^e	2.5	2.0	1.9	3.1	2.0	1.7	2.3	1.3	1.4	1.0	1.2	0.8	0.7	0.7	0.0
Other Cocaine ^f	8.1	8.3	8.6	8.5	9.3	8.1	6.2	8.0	7.1	7.9	6.7	5.4	5.1	5.2	+0.1
Heroin	1.7	1.2	1.0	1.0	0.9	0.5	0.7	0.5	0.7	0.8	0.7	0.6	0.5	0.4	0.0
Narcotics other than Heroin ^{g,h}	8.9	11.0	12.2	14.2	13.8	14.4	14.6	14.1	12.4	14.0	12.2	12.4	10.3	10.8	+0.5
Amphetamines ^{g,i}	12.3	12.4	11.9	12.3	12.7	12.3	10.7	11.2	9.1	11.8	12.1	13.4	14.4	15.3	+1.0
Methamphetamine ^j	5.1	5.3	5.0	5.8	5.2	4.1	2.9	1.9	1.9	1.0	1.1	0.6	0.3	0.9	+0.6
Crystal Methamphetamine (Ice) ^j	1.3	2.3	2.0	2.9	2.2	2.4	1.7	1.3	1.1	0.7	0.8	0.2	0.6	0.0	-0.6
Sedatives (Barbiturates) ^{g,t}	6.9	6.0	5.9	5.7	7.2	8.5	6.3	5.9	6.4	6.0	5.3	3.6	3.5	5.4	—
Sedatives, Adjusted ^{g,k}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{g,l}	8.8	9.7	10.7	11.0	10.6	11.9	10.0	9.1	8.6	9.2	8.1	7.1	6.4	7.8	+1.4
Alcohol ^m	86.6	86.1	86.0	86.2	84.6	86.6	84.7	83.1	85.3	82.6	82.3	80.5	81.0	78.0	-3.0
Been Drunk ⁿ	74.7	76.1	75.1	74.9	73.4	72.9	73.1	71.6	72.5	69.1	70.5	67.9	70.0	66.5	-3.4
Flavored Alcoholic Beverages ^o	—	—	—	—	79.0	84.5	80.9	80.6	78.6	78.1	77.4	76.7	76.6	67.5	-9.1 ^s
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^p	0.6	1.5	1.2	1.2	1.6	1.0	1.9	0.6	1.6	1.3	0.7	1.1	0.4	0.8	+0.4

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 9-5.

TABLE 9-2
Trends in Annual Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
 (Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	(Years cont.)
<i>Approximate Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	
Any Illicit Drug ^a	56.2	55.0	49.5	49.8	45.1	46.3	45.0	40.1	37.4	36.7	33.3	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	
Any Illicit Drug other than Marijuana ^a	32.3	31.7	29.9	29.9	27.2	26.7	25.0	21.3	19.2	16.4	15.2	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	
Synthetic Marijuana ^u	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Marijuana	51.2	51.3	44.7	45.2	40.7	41.7	40.9	37.0	34.6	33.6	29.4	26.5	27.7	27.9	29.3	31.2	33.1	31.6	35.9	35.2	
Inhalants ^b	3.0	2.5	2.5	2.8	2.4	3.1	3.9	3.7	4.1	3.7	3.9	3.5	3.1	3.8	3.0	3.9	3.6	4.1	3.0	3.2	
Hallucinogens ^c	8.5	7.0	8.7	6.5	6.2	5.0	6.0	5.9	5.3	5.1	5.4	6.3	6.8	6.0	6.2	8.2	6.9	7.7	7.2	7.8	
LSD	6.0	4.6	6.3	4.3	3.7	2.2	3.9	4.0	3.6	3.4	4.3	5.1	5.7	5.1	5.2	6.9	5.2	5.0	4.4	5.4	
Hallucinogens other than LSD ^c	5.2	4.7	5.4	3.9	4.1	3.9	3.8	3.1	3.4	3.1	3.0	3.1	2.6	2.7	2.8	4.0	4.1	4.9	4.4	4.5	
Ecstasy (MDMA) ^d	—	—	—	—	—	—	—	—	—	2.3	0.9	2.0	0.8	0.5	2.4	2.8	2.4	3.9	5.5		
Salvia ^v	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Cocaine	16.8	16.0	17.2	17.3	16.3	17.3	17.1	13.7	10.0	8.2	5.6	3.6	3.0	2.7	2.0	3.6	2.9	3.4	4.6	4.6	
Crack ^e	—	—	—	—	—	—	—	2.0	1.4	1.5	0.6	0.5	0.4	0.6	0.5	1.1	0.6	0.4	1.0	0.9	
Other Cocaine ^f	—	—	—	—	—	—	—	10.7	10.6	9.3	5.1	3.2	2.4	2.5	1.8	3.3	2.3	3.0	4.2	4.2	
Heroin	0.4	0.2	0.1	*	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.4	0.3	0.6	0.2	
Narcotics other than Heroin ^{g,h}	5.1	4.3	3.8	3.8	3.8	2.4	4.0	3.1	3.1	3.2	2.9	2.7	2.7	2.5	2.4	3.8	3.1	4.2	4.2	4.3	
OxyContin ^{g,j}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Vicodin ^{g,j}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amphetamines ^{g,i}	22.4	22.2	21.1	17.3	15.7	11.9	10.3	7.2	6.2	4.6	4.5	3.9	3.6	4.2	4.2	5.4	4.2	5.7	5.1	5.8	
Ritalin ^{g,j}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Adderall ^{g,j}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.3	
Crystal Methamphetamine (Ice) ^j	—	—	—	—	—	—	—	—	—	—	0.1	0.1	0.2	0.7	0.8	1.1	0.4	0.8	1.0	0.5	
Bath Salts (synthetic stimulants) ⁿ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Sedatives (Barbiturates) ^{g,t}	2.9	2.8	3.2	2.2	1.9	1.3	2.0	1.2	1.1	1.0	1.4	1.2	1.4	1.5	1.2	2.0	2.3	3.0	2.5	3.2	
Sedatives, Adjusted ^{g,k}	8.3	8.0	8.0	4.5	3.5	2.5	2.6	1.7	1.5	1.0	—	—	—	—	—	—	—	—	—	—	
Methaqualone ^g	7.2	6.5	6.6	3.1	2.5	1.4	1.2	0.8	0.5	0.2	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{g,l}	6.9	4.8	4.7	4.6	3.5	3.6	4.4	3.8	3.1	2.6	3.0	2.4	2.9	2.4	1.8	2.9	2.8	3.8	3.9	3.8	
Rohypnol ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
GHB ^w	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Ketamine ^w	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Alcohol ^m	90.5	92.5	92.2	91.6	90.0	92.0	91.5	90.9	89.6	89.6	89.0	88.3	86.9	85.1	82.7	83.2	83.0	82.4	84.6	83.6	
Been Drunk ⁿ	—	—	—	—	—	—	—	—	—	—	—	69.1	67.3	65.6	63.1	62.1	64.2	66.8	67.0	65.4	
Flavored Alcoholic Beverages ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Alcoholic Beverages containing Caffeine ^{j,s}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Cigarettes	36.2	37.6	34.3	36.1	33.2	35.0	35.3	38.0	36.6	34.2	35.5	35.6	37.3	38.8	37.6	39.3	41.4	43.6	44.3	44.5	
Tobacco Using a Hookah ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Small Cigars ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Snus ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Dissolvable Tobacco ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Steroids ^p	—	—	—	—	—	—	—	—	—	0.4	0.5	0.6	0.2	0.9	0.2	0.4	0.2	0.7	0.2	0.9	

(Table continued on next page.)

TABLE 9-2 (cont.)
Trends in Annual Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
(Entries are percentages.)

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012- 2013 change
<i>Approximate Weighted N =</i>	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090	
Any Illicit Drug ^a	36.1	37.9	37.0	36.5	36.2	36.6	33.9	35.0	35.2	36.0	35.0	36.3	37.3	38.9	+1.7
Any Illicit Drug other than Marijuana ^a	15.6	16.4	16.6	17.9	18.6	18.5	18.1	17.3	15.3	16.9	17.1	16.8	17.1	19.0	+1.9
Synthetic Marijuana ^u	—	—	—	—	—	—	—	—	—	—	—	7.4	5.3	2.3	-3.0 ss
Marijuana	34.0	35.6	34.7	33.7	33.3	33.3	30.2	31.8	32.3	32.8	32.7	33.2	34.9	35.5	+0.6
Inhalants ^b	2.9	2.8	2.0	1.8	2.7	1.8	1.5	1.5	1.1	1.2	1.7	0.9	1.5	0.5	-1.0
Hallucinogens ^c	6.7	7.5	6.3	7.4	5.9	5.0	5.6	4.9	5.1	4.7	4.9	4.1	4.5	4.5	0.0
LSD	4.3	4.0	2.1	1.4	1.2	0.7	1.4	1.3	2.6	2.0	2.1	2.0	1.9	2.6	+0.7
Hallucinogens other than LSD ^c	4.4	5.5	5.8	7.1	5.6	5.0	5.4	4.7	4.4	4.1	4.4	3.4	3.9	3.7	-0.2
Ecstasy (MDMA) ^d	9.1	9.2	6.8	4.4	2.2	2.9	2.6	2.2	3.7	3.1	4.3	4.2	5.8	5.3	-0.5
Salvia ^v	—	—	—	—	—	—	—	—	—	5.8	3.5	3.1	1.5	1.0	-0.5
Cocaine	4.8	4.7	4.8	5.4	6.6	5.7	5.1	5.4	4.4	4.2	3.5	3.3	3.1	2.7	-0.4
Crack ^e	0.9	0.9	0.4	1.3	1.3	0.8	1.0	0.6	0.5	0.3	0.4	0.3	0.3	0.3	+0.0
Other Cocaine ^f	4.1	4.1	5.0	5.1	6.3	5.0	3.8	5.3	4.2	4.2	4.0	3.0	3.0	2.8	-0.1
Heroin	0.5	0.4	0.1	0.2	0.4	0.3	0.3	0.2	0.3	0.4	0.2	0.1	0.1	0.3	+0.2
Narcotics other than Heroin ^{g,h}	4.5	5.7	7.4	8.7	8.2	8.4	8.8	7.7	6.5	7.6	7.2	6.2	5.4	5.4	0.0
OxyContin ^{g,q}	—	—	1.5	2.2	2.5	2.1	3.0	2.8	3.6	5.0	2.3	2.4	1.2	2.3	+1.2
Vicodin ^{g,q}	—	—	6.9	7.5	7.4	9.6	7.6	6.7	6.7	8.4	4.9	5.8	3.8	4.4	+0.6
Amphetamines ^{g,i}	6.6	7.2	7.0	7.1	7.0	6.7	6.0	6.9	5.7	7.5	9.0	9.3	11.1	10.6	-0.6
Ritalin ^{g,q}	—	—	5.7	4.7	4.7	4.2	3.9	3.7	3.2	1.7	1.9	2.3	1.8	3.6	+1.8
Adderall ^{g,q}	—	—	—	—	—	—	—	—	—	10.2	9.0	9.8	9.0	10.7	+1.6
Methamphetamine ^j	1.6	2.4	1.2	2.6	2.9	1.7	1.2	0.4	0.5	0.3	0.4	0.2	0.0	0.4	+0.4
Crystal Methamphetamine (Ice) ^j	0.5	0.6	0.8	0.9	1.1	1.4	0.6	0.7	0.1	0.1	0.5	0.1	0.6	*	-0.6
Bath Salts (synthetic stimulants) ⁿ	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.1	-0.2
Sedatives (Barbiturates) ^{g,t}	3.7	3.8	3.7	4.1	4.2	3.9	3.4	3.6	3.7	3.1	2.5	1.7	2.2	2.7	—
Sedatives, Adjusted ^{g,k}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{g,l}	4.2	5.1	6.7	6.9	6.7	6.4	5.8	5.5	5.0	5.4	4.9	4.2	3.4	4.4	+1.0
Rohypnol ^j	—	—	0.7	0.4	0.3	0.1	0.2	0.1	0.3	*	—	—	—	—	—
GHB ^w	—	—	0.6	0.3	0.7	0.4	*	0.1	0.2	*	0.1	0.1	*	0.1	+0.1
Ketamine ^w	—	—	1.3	1.0	1.5	0.5	0.9	0.2	0.4	0.1	0.7	0.6	0.4	0.9	+0.5
Alcohol ^m	83.2	83.0	82.9	81.7	81.2	83.0	82.1	80.9	82.1	79.4	78.6	77.4	79.2	75.6	-3.6 s
Been Drunk ⁿ	64.7	68.8	66.0	64.7	67.1	64.2	66.2	64.8	66.8	61.5	63.8	60.1	61.5	57.9	-3.5
Flavored Alcoholic Beverages ^o	—	—	—	—	63.2	67.0	63.5	62.6	65.0	66.1	60.3	63.0	58.1	57.6	-0.6
Alcoholic Beverages containing Caffeine ^j	—	—	—	—	—	—	—	—	—	—	—	33.6	33.8	39.1	+5.4
Cigarettes	41.3	39.0	38.3	35.2	36.7	36.0	30.9	30.7	30.0	29.9	28.1	25.8	23.4	23.2	-0.2
Tobacco Using a Hookah ^j	—	—	—	—	—	—	—	—	—	—	—	27.9	25.7	26.1	+0.3
Small Cigars ^j	—	—	—	—	—	—	—	—	—	—	—	23.6	20.3	19.0	-1.3
Snus ^j	—	—	—	—	—	—	—	—	—	—	—	6.5	4.7	4.8	+0.1
Dissolvable Tobacco ^j	—	—	—	—	—	—	—	—	—	—	—	*	0.3	0.2	0.0
Steroids ^p	0.1	0.6	0.5	0.3	0.6	0.5	0.8	0.6	0.1	0.7	0.3	0.2	0.3	0.8	+0.5

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 9-5.

TABLE 9-3
Trends in 30-Day Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
(Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	(Years cont.)
<i>Approximate Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	
Any Illicit Drug ^a	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	
Any Illicit Drug other than Marijuana ^a	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	
Marijuana	34.0	33.2	26.8	26.2	23.0	23.6	22.3	20.3	16.8	16.3	14.0	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	
Inhalants ^b	1.5	0.9	0.8	0.7	0.7	1.0	1.1	0.9	1.3	0.8	1.0	0.9	1.1	1.3	0.6	1.6	0.8	0.7	0.6	1.5	
Hallucinogens ^c	2.7	2.3	2.6	1.8	1.8	1.3	2.2	2.0	1.7	2.3	1.4	1.2	2.3	2.5	2.1	3.3	1.9	2.1	2.1	2.0	
LSD	1.4	1.4	1.7	0.9	0.8	0.7	1.4	1.4	1.1	1.4	1.1	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	
Hallucinogens other than LSD ^c	1.9	1.2	1.4	1.0	1.2	0.7	1.2	0.8	0.8	1.1	0.8	0.6	0.7	1.1	0.8	1.6	1.2	1.2	0.7	1.2	
Ecstasy (MDMA) ^d	—	—	—	—	—	—	—	—	—	0.3	0.6	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	
Cocaine	6.9	7.3	7.9	6.5	7.6	6.9	7.0	4.6	4.2	2.8	1.2	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	
Crack ^e	—	—	—	—	—	—	1.3	0.4	0.5	0.2	0.1	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	
Other Cocaine ^f	—	—	—	—	—	—	—	3.5	3.2	3.2	1.0	1.0	0.9	0.6	0.3	0.8	0.6	1.3	1.5	1.0	
Heroin	0.3	*	*	*	*	*	*	0.1	0.1	0.1	*	0.1	*	*	*	0.1	*	0.2	0.1	0.1	
Narcotics other than Heroin ^{g,h}	1.8	1.1	0.9	1.1	1.4	0.7	0.6	0.8	0.8	0.7	0.5	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	
Amphetamines ^{g,i}	13.4	12.3	9.9	7.0	5.5	4.2	3.7	2.3	1.8	1.3	1.4	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	
Methamphetamine ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.2	
Crystal Methamphetamine (Ice) ^j	—	—	—	—	—	—	—	—	—	—	*	*	*	0.3	0.5	0.3	0.1	0.2	0.3	*	
Sedatives (Barbiturates) ^{g,t}	0.9	0.8	1.0	0.5	0.7	0.4	0.6	0.5	0.5	0.2	0.2	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	
Sedatives, Adjusted ^{g,k}	3.8	3.4	2.5	1.1	1.0	0.7	0.6	0.6	0.6	0.2	—	—	—	—	—	—	—	—	—	—	
Methaqualone ^g	3.1	3.0	1.9	0.7	0.5	0.3	0.1	0.2	0.1	0.0	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{g,l}	2.0	1.4	1.4	1.2	1.1	1.4	1.9	1.0	1.1	0.8	0.5	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	
Alcohol ^m	81.8	81.9	82.8	80.3	79.1	80.3	79.7	78.4	77.0	76.2	74.5	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	
Been Drunk ⁿ	—	—	—	—	—	—	—	—	—	—	—	—	45.0	45.0	43.8	42.8	37.9	40.3	46.4	44.3	44.6
Flavored Alcoholic Beverages ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Cigarettes	25.8	25.9	24.4	24.7	21.5	22.4	22.4	24.0	22.6	21.1	21.5	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	
Steroids ^p	—	—	—	—	—	—	—	—	—	*	0.2	0.3	0.2	0.2	0.2	0.1	*	0.2	0.2	0.4	

(Table continued on next page.)

TABLE 9-3 (cont.)
Trends in 30-Day Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
(Entries are percentages.)

	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	2012– 2013 change
<i>Approximate Weighted N =</i>	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090	
Any Illicit Drug ^a	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	20.7	19.2	21.4	22.3	22.5	+0.2
Any Illicit Drug other than Marijuana ^a	6.9	7.5	7.8	8.2	9.1	8.2	8.2	8.1	7.3	8.4	8.1	8.2	7.8	8.2	+0.4
Marijuana	20.0	20.2	19.7	19.3	18.9	17.1	16.7	16.8	17.0	18.5	17.5	19.4	20.5	20.6	+0.2
Inhalants ^b	0.9	0.4	0.7	0.4	0.4	0.3	0.4	0.1	0.4	0.1	0.5	0.3	0.2	0.1	-0.2
Hallucinogens ^c	1.4	1.8	1.2	1.8	1.3	1.2	0.9	1.3	1.7	1.0	1.4	1.2	1.1	1.0	-0.1
LSD	0.9	1.0	0.2	0.2	0.2	0.1	0.3	0.3	0.8	0.3	0.7	0.5	0.4	0.4	-0.1
Hallucinogens other than LSD ^c	0.8	0.8	1.1	1.7	1.2	1.1	0.7	1.1	1.3	0.8	1.2	0.8	0.7	0.8	+0.1
Ecstasy (MDMA) ^d	2.5	1.5	0.7	1.0	0.7	0.8	0.6	0.4	0.6	0.5	1.0	0.7	1.4	0.8	-0.6
Cocaine	1.4	1.9	1.6	1.9	2.4	1.8	1.8	1.7	1.2	1.3	1.0	1.2	1.1	0.9	-0.3
Crack ^e	0.3	0.1	0.3	0.4	0.4	0.1	*	0.1	0.1	0.1	0.1	0.1	*	0.3	+0.3
Other Cocaine ^f	0.9	1.5	1.4	1.9	2.2	1.8	1.3	1.6	1.1	1.2	1.0	1.2	1.3	0.9	-0.3
Heroin	0.2	0.1	*	*	0.1	0.1	0.2	0.1	*	0.1	*	*	0.1	0.2	+0.1
Narcotics other than Heroin ^{g,h}	1.7	1.7	3.2	2.3	3.0	3.1	3.1	2.2	2.3	2.7	2.3	2.1	2.2	1.5	-0.7
Amphetamines ^{g,i}	2.9	3.3	3.0	3.1	3.2	2.9	2.5	3.1	2.8	3.4	4.1	4.5	4.6	5.3	+0.6
Methamphetamine ^j	0.2	0.5	0.2	0.6	0.2	0.1	0.2	0.1	*	0.1	*	*	*	*	0.0
Crystal Methamphetamine (Ice) ^j	*	0.1	*	0.3	0.1	0.2	*	0.1	*	*	0.2	*	0.3	*	-0.3
Sedatives (Barbiturates) ^{g,t}	1.1	1.5	1.7	1.7	1.5	1.3	1.3	1.4	1.4	1.2	0.6	0.8	0.8	0.9	—
Sedatives, Adjusted ^{g,k}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{g,l}	2.0	1.5	3.0	2.8	2.7	2.2	2.1	1.8	1.6	2.2	1.3	1.6	1.1	1.2	+0.1
Alcohol ^m	67.4	67.0	68.9	66.2	67.7	67.9	65.4	66.6	69.0	65.8	65.0	63.5	67.7	63.1	-4.6 s
Been Drunk ⁿ	43.9	44.7	44.4	40.4	47.4	43.1	47.6	46.8	45.3	42.4	43.6	39.9	40.1	40.2	+0.2
Flavored Alcoholic Beverages ^o	—	—	—	—	34.0	30.9	26.2	27.5	35.8	32.3	31.5	29.5	31.3	29.1	-2.2
Cigarettes	28.2	25.7	26.7	22.5	24.3	23.8	19.2	19.9	17.9	17.9	16.4	15.2	12.5	14.0	+1.5
Steroids ^p	*	0.3	*	0.1	*	*	*	0.1	*	0.2	*	0.2	*	*	0.0

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 9-5.

TABLE 9-4
Trends in 30-Day Prevalence of Daily^r Use of Various Types of Drugs
among College Students 1 to 4 Years beyond High School

(Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	(Years cont.)
<i>Approximate Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	
Marijuana	7.2	5.6	4.2	3.8	3.6	3.1	2.1	2.3	1.8	2.6	1.7	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	
Cocaine	0.2	*	0.3	0.1	0.4	0.1	0.1	0.1	0.1	*	*	*	*	*	0.1	*	*	*	*	*	*
Amphetamines ^g	0.5	0.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines, Adjusted ^{g,i}	—	—	0.3	0.2	0.2	*	0.1	0.1	*	*	*	0.1	*	0.1	0.1	0.1	*	0.2	0.1	0.1	
Alcohol ^m																					
Daily	6.5	5.5	6.1	6.1	6.6	5.0	4.6	6.0	4.9	4.0	3.8	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	
Been Drunk ⁿ	—	—	—	—	—	—	—	—	—	—	—	0.5	0.2	0.3	0.8	0.5	0.1	1.3	0.8	1.0	
5+ Drinks in a Row in Last 2 Weeks	43.9	43.6	44.0	43.1	45.4	44.6	45.0	42.8	43.2	41.7	41.0	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	
Cigarettes																					
Daily	18.3	17.1	16.2	15.3	14.7	14.2	12.7	13.9	12.4	12.2	12.1	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	
1/2 Pack+/Day	12.7	11.9	10.5	9.6	10.2	9.4	8.3	8.2	7.3	6.7	8.2	8.0	8.9	8.9	8.0	10.2	8.5	9.1	11.3	11.0	

(Table continued on next page.)

TABLE 9-4 (cont.)
Trends in 30-Day Prevalence of Daily^r Use of Various Types of Drugs
among College Students 1 to 4 Years beyond High School

(Entries are percentages.)

	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
<i>Approximate Weighted N =</i>	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090	
Marijuana	4.6	4.5	4.1	4.7	4.5	4.0	4.3	3.5	3.9	4.9	4.4	4.7	4.8	5.1	+0.4
Cocaine	*	*	*	*	*	0.1	0.1	*	*	*	*	*	*	*	0.0
Amphetamines ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines, Adjusted ^{g,i}	0.1	0.2	0.1	0.3	0.2	0.2	0.4	0.1	0.2	0.1	*	0.2	0.1	*	-0.1
Alcohol ^m															
Daily	3.6	4.7	5.0	4.3	3.7	4.6	4.8	4.3	4.0	4.3	3.6	3.8	3.9	3.6	-0.3
Been Drunk ⁿ	0.7	0.5	0.8	1.1	0.8	0.5	0.6	0.7	0.5	0.7	0.3	1.3	0.4	0.5	+0.1
5+ Drinks in a Row in Last 2 Weeks	39.3	40.9	40.1	38.5	41.7	40.1	40.2	41.1	40.0	36.9	37.0	36.1	37.4	35.2	-2.1
Cigarettes															
Daily	17.8	15.0	15.9	13.8	13.8	12.4	9.2	9.3	9.2	8.0	7.6	7.3	5.2	5.6	+0.4
1/2 Pack+/Day	10.1	7.8	7.9	7.6	6.8	6.7	4.9	4.3	4.3	3.8	3.9	2.5	2.4	2.4	0.0

Source. The Monitoring the Future study, the University of Michigan.

See footnotes following Table 9-5.

TABLE 9-5
Trends in Lifetime, Annual, and 30-Day Prevalence of an Illicit Drug Use Index^a
among College Students 1 to 4 Years beyond High School, by Gender

→
(Years cont.)

	<u>1980</u> ⁱ	<u>1981</u> ⁱ	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	
Percentage who used in lifetime																			
Any Illicit Drug																			
Total	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	47.4	49.0		
Males	71.0	67.5	68.1	71.3	66.4	69.8	64.7	63.5	56.0	56.5	52.5	51.3	50.8	45.7	49.5	47.3	50.3	52.1	
Females	67.5	66.3	61.5	63.0	59.2	61.6	59.4	57.4	60.2	54.9	55.1	49.7	47.1	46.0	42.6	44.3	45.6	46.7	
Any Illicit Drug other than Marijuana																			
Total	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	
Males	42.8	39.8	45.1	44.6	40.9	42.1	38.2	37.2	31.8	30.6	26.2	27.6	26.3	24.3	24.6	26.6	25.0	27.3	
Females	41.6	42.6	34.7	39.2	36.4	38.3	37.0	34.6	34.6	30.4	30.1	24.3	26.1	24.3	20.1	22.9	21.2	22.2	
Percentage who used in last 12 months																			
Any Illicit Drug																			
Total	56.2	55.0	49.5	49.8	45.1	46.3	45.0	40.1	37.4	36.7	33.3	29.2	30.6	30.6	31.4	33.5	34.2	34.1	
Males	58.9	56.2	54.6	53.4	48.4	50.9	49.8	43.3	37.0	38.2	34.2	30.2	32.8	32.6	33.9	36.1	36.6	38.3	
Females	53.3	54.0	44.9	46.7	41.9	42.7	41.1	37.7	37.6	35.4	32.5	28.4	28.7	29.1	29.5	31.7	32.7	31.1	
Any Illicit Drug other than Marijuana																			
Total	32.3	31.7	29.9	29.9	27.2	26.7	25.0	21.3	19.2	16.4	15.2	13.2	13.1	12.5	12.2	15.9	12.8	15.8	
Males	33.7	32.8	33.4	33.5	29.2	29.7	28.6	23.5	19.4	18.7	15.7	14.4	13.8	15.0	14.9	19.5	15.1	18.1	
Females	31.1	30.8	26.9	26.8	25.2	24.4	22.1	19.6	19.0	14.6	14.8	12.1	12.6	10.5	10.2	13.3	11.3	14.1	
Percentage who used in last 30 days																			
Any Illicit Drug																			
Total	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	
Males	42.9	40.6	37.7	33.8	30.4	29.9	31.0	24.0	18.8	20.0	18.2	16.0	18.0	16.0	20.5	23.7	20.6	23.4	
Females	34.0	34.8	25.6	25.5	23.7	23.2	21.7	21.1	18.3	16.7	12.7	14.6	14.5	14.5	12.7	15.7	15.8	16.2	
Any Illicit Drug other than Marijuana																			
Total	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	
Males	22.8	18.6	20.2	16.0	16.1	12.6	14.4	9.0	8.2	8.0	4.9	4.8	5.1	7.3	6.2	8.8	6.1	7.8	
Females	18.7	18.5	14.2	12.1	11.5	11.2	9.3	8.5	8.8	6.0	4.0	3.9	4.2	3.8	3.4	4.5	3.4	6.1	
Approximate Weighted N																			
All Respondents																			
Total	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	
Males	520	530	550	550	540	490	540	520	560	580	620	640	680	660	590	610	560	630	
Females	520	600	610	620	570	600	650	700	750	720	780	770	810	830	820	840	890	860	

(Table continued on next page.)

TABLE 9-5 (cont.)
Trends in Lifetime, Annual, and 30-Day Prevalence of an Illicit Drug Use Index^a
among College Students 1 to 4 Years beyond High School, by Gender

	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2012– 2013 change
<i>Percentage who used in lifetime</i>																	
Any Illicit Drug																	
Total	52.9	53.2	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	51.4	49.1	49.2	50.5	51.0	+0.4
Males	54.4	58.4	54.4	53.9	54.3	54.1	54.9	54.2	55.0	52.3	50.7	53.2	53.5	52.3	52.4	54.3	+1.9
Females	52.0	49.6	53.2	53.5	50.2	53.7	50.6	51.3	47.8	49.4	48.8	50.2	46.2	47.3	49.2	48.8	-0.4
Any Illicit Drug other than Marijuana																	
Total	24.8	25.5	25.8	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	25.6	24.8	24.3	23.8	26.7	+2.9
Males	27.3	29.4	28.9	27.0	30.4	27.6	31.1	29.0	29.2	26.5	25.2	29.9	27.8	27.8	26.0	30.8	+4.9
Females	23.3	22.8	23.5	25.9	24.6	27.5	26.2	25.1	24.4	24.6	21.0	22.7	22.8	22.1	22.2	24.0	+1.7
<i>Percentage who used in last 12 months</i>																	
Any Illicit Drug																	
Total	37.8	36.9	36.1	37.9	37.0	36.5	36.2	36.6	33.9	35.0	35.2	36.0	35.0	36.3	37.3	38.9	+1.7
Males	40.1	42.5	38.0	38.8	39.5	39.2	40.9	40.7	39.2	38.0	38.7	37.6	40.3	41.2	39.5	43.8	+4.3
Females	36.4	33.2	34.7	37.3	35.4	34.8	33.4	34.2	30.6	33.1	32.9	35.0	31.6	33.2	35.7	35.8	+0.1
Any Illicit Drug other than Marijuana																	
Total	14.0	15.4	15.6	16.4	16.6	17.9	18.6	18.5	18.1	17.3	15.3	16.9	17.1	16.8	17.1	19.0	+1.9
Males	17.0	19.0	18.6	17.2	19.2	19.3	22.1	21.1	22.6	19.0	17.8	19.7	20.3	20.1	19.6	23.5	+3.9
Females	12.1	12.8	13.5	15.8	15.0	17.1	16.5	16.9	15.2	16.3	13.7	15.0	15.1	14.7	15.4	16.1	+0.7
<i>Percentage who used in last 30 days</i>																	
Any Illicit Drug																	
Total	19.7	21.6	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	20.7	19.2	21.4	22.3	22.5	+0.2
Males	23.1	26.7	24.0	25.0	25.1	22.8	26.1	22.9	23.4	22.7	23.1	23.4	25.9	27.0	27.0	29.8	+2.8
Females	17.6	18.1	19.6	19.8	19.3	20.5	18.4	17.5	16.6	17.1	16.2	19.0	15.0	17.9	19.1	17.9	-1.2
Any Illicit Drug other than Marijuana																	
Total	6.1	6.4	6.9	7.5	7.8	8.2		8.2	8.2	8.1	7.3	8.4	8.1	8.2	7.8	8.2	+0.4
Males	8.6	7.5	8.2	9.0	8.4	8.1	11.3	10.3	10.3	9.5	9.6	9.0	10.4	10.6	9.2	10.2	+0.9
Females	4.6	5.6	6.0	6.4	7.4	8.3	7.8	7.0	6.9	7.2	5.8	8.0	6.7	6.7	6.8	7.0	+0.2
<i>Approximate Weighted N</i>																	
All Respondents																	
Total	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	1,320	1,260	1,230	1,150	1,090	
Males	570	590	560	540	490	480	520	500	500	470	510	530	500	480	480	430	
Females	880	850	790	800	770	790	880	860	780	770	760	790	760	750	670	660	

Source. The Monitoring the Future study, the University of Michigan.
 See footnotes on the following page.

Footnotes for Tables 9-1 through 9-5

Notes. Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '—' indicates data not available. '* *' indicates a prevalence rate of less than 0.05%.

^aAny illicit drug includes use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in four of the five questionnaire forms in 1980–1989, in five of the six forms in 1990–1998, and in three of the six forms in 1999–2012. Total N in 2012 is approximately 580.

^cIn 2001 the question text was changed on three of the six questionnaire forms. Other psychedelics was changed to other hallucinogens, and shrooms was added to the list of examples. Beginning in 2002 the remaining forms were changed to the new wording.

^dThis drug was asked about in two of the five questionnaire forms in 1989, in two of the six questionnaire forms in 1990–2001, and in three of the six questionnaire forms in 2002–2012. Total N in 2012 is approximately 580.

^eThis drug was asked about in one of the five questionnaire forms for annual use only in 1986, two of the five questionnaire forms in 1987–1989, in all six questionnaire forms in 1990–2001, and in five of the six questionnaire forms in 2002–2012. Total N in 2012 is approximately 960.

^fThis drug was asked about in one of the five questionnaire forms in 1987–1989 and in four of six questionnaire forms in 1990–2012. Total N in 2012 is approximately 770.

^gOnly drug use that was not under a doctor's orders is included here.

^hIn 2002 the question text was changed on three of the six questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is three sixths of N indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003 and beyond.

ⁱRevised questions about amphetamine use were introduced in 1982 to more completely exclude inappropriate reporting of nonprescription amphetamines.

^jThis drug was asked about in two of the six questionnaire forms. Total N in 2012 is approximately 380. Questions about Rohypnol use were dropped from the questionnaires beginning in 2010.

^kSedatives, adjusted data are a combination of barbiturate and methaqualone data.

^lIn 2001 the question text was changed on three of the six questionnaire forms. Miltown was replaced with Xanax in the list of examples. Beginning in 2002 the remaining forms were changed to the new wording.

^mIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a drink meant more than just a few sips. Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

ⁿThis drug was asked about in three of the six questionnaire forms. Total N in 2012 is approximately 580.

^oThis drug was asked about in one of the six questionnaire forms. Total N in 2012 is approximately 190.

^pThis drug was asked about in one of the five questionnaire forms in 1989 and in two of the six questionnaire forms in 1990–2012. Total N in 2012 is approximately 380.

^qThis drug was asked about in two of the six questionnaire forms through 2010 and in three of the six questionnaire forms beginning in 2012. Total N in 2012 is approximately 580.

^rDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, measured as actual daily use, and 5+ drinks, measured as having five or more drinks in a row in the last two weeks.

^sIn 2012 the alcoholic beverage containing caffeine question text was changed to alcoholic beverage mixed with an energy drink. The data in 2011 and 2012 are not comparable due to this question change.


^tIn 2013 the question text was changed on all forms: Tuinal, Nembutal, and Seconal were replaced with Ambien, Lunesta, and Sonata. The data in 2012 and 2013 are not comparable due to this question change.

^uThis drug was asked about in two of the six questionnaire forms in 2011–2012; N is two sixths of N indicated. Data were based on three of the six questionnaire forms in 2013; N is three sixths of N indicated.


^vThis drug was asked about in one of the six questionnaire forms in 2009; N is one sixth of N indicated; Data were based on two of the six questionnaire forms in 2010–2011; N is two sixths of N indicated. Data were based on three of the six questionnaire forms in 2012–2013; N is three sixths of N indicated.

^wThis drug was asked about in two of the six questionnaire forms in 2002–2009; N is two sixths of N indicated; Data were based on three of the six questionnaire forms in 2010–2011; N is three sixths of N indicated. Data were based on two of the six questionnaire forms in 2012–2013; N is two sixths of N indicated.

FIGURE 9-1
ANY ILLICIT DRUG
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




ANY ILLICIT DRUG
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-2
ANY ILLICIT DRUG OTHER THAN MARIJUANA
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



ANY ILLICIT DRUG OTHER THAN MARIJUANA
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-3a
MARIJUANA


**Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



MARIJUANA

**Trends in Annual Prevalence
among Male vs. Female College Students**



Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-3b

MARIJUANA


Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School

(Twelfth graders included for comparison.)



MARIJUANA

Trends in 30-Day Prevalence of Daily Use among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-4
INHALANTS^a


Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School

(Twelfth graders included for comparison.)



INHALANTS^a

Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites.

FIGURE 9-5
HALLUCINOGENS^a
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



HALLUCINOGENS^a
Trends in Annual Prevalence
among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.


^aUnadjusted for the possible underreporting of PCP.

FIGURE 9-6

LSD


**Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



LSD


**Trends in Annual Prevalence
among Male vs. Female College Students**




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-7
HALLUCINOGENS OTHER THAN LSD
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




HALLUCINOGENS OTHER THAN LSD
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-8
ECSTASY (MDMA)
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



ECSTASY (MDMA)
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.
 Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-9

COCAINE


**Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



COCAINE


**Trends in Annual Prevalence
among Male vs. Female College Students**




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-10a
NARCOTICS OTHER THAN HEROIN ^{a,b}
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



NARCOTICS OTHER THAN HEROIN ^{a,b}
Trends in Annual Prevalence
among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUsed without a doctor's orders.


^bIn 2002 the question text was changed on half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only. In 2003 the remaining forms were changed to the new wording.

FIGURE 9-10b

VICODIN^a

Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School

(Twelfth graders included for comparison.)



Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUsed without a doctor's orders.

FIGURE 9-10c
OXYCONTIN^a

**Trends in Annual Prevalence among College Students vs. Others
 1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)




Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUsed without a doctor's orders.

FIGURE 9-11
AMPHETAMINES^a
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



AMPHETAMINES^a
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUsed without a doctor's orders.

FIGURE 9-12
SEDATIVES (BARBITURATES) ^a
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



SEDATIVES (BARBITURATES) ^a
Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUsed without a doctor's orders.

FIGURE 9-13
TRANQUILIZERS^a
Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



TRANQUILIZERS^a
Trends in Annual Prevalence
among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.


^aUsed without a doctor's orders.

FIGURE 9-14a

ALCOHOL


**Trends in Annual Prevalence among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



ALCOHOL


**Trends in Annual Prevalence
among Male vs. Female College Students**




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14b
ALCOHOL
Trends in 30-Day Prevalence among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



ALCOHOL
Trends in 30-Day Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.
 Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14c

ALCOHOL


Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School

(Twelfth graders included for comparison.)



ALCOHOL

Trends in 30-Day Prevalence of Daily Use among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.


Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14d

ALCOHOL


**Trends in 2-Week Prevalence of 5 or More Drinks in a Row
among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



ALCOHOL

**Trends in 2-Week Prevalence of 5 or More Drinks in a Row
among Male vs. Female College Students**




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-15a
CIGARETTES


**Trends in 30-Day Prevalence among College Students vs. Others
1 to 4 Years beyond High School**

(Twelfth graders included for comparison.)



CIGARETTES

**Trends in 30-Day Prevalence
among Male vs. Female College Students**




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-15b
CIGARETTES


Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School

(Twelfth graders included for comparison.)



CIGARETTES

Trends in 30-Day Prevalence of Daily Use among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.


FIGURE 9-15c
CIGARETTES

Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among College Students vs. Others 1 to 4 Years beyond High School
(Twelfth graders included for comparison.)



CIGARETTES

Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.
Note. Others refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

INDEX

A

Adderall

- cohort effect, 16, 25-26
- harmfulness, perceived, 241, table 6-1
- prevalence, 94, table 4-3
 - college enrollment, 366, 368, 370, table 8-2
 - gender, 30, 158, 369
 - population density, 103
 - race/ethnicity, 38
- trends, 46, table 2-2, table 5-2
 - college enrollment, 380, table 9-2

alcohol

- daily use
 - college enrollment, 29, 366, 385, table 8-4
 - table 9-4, figure 9-14c
 - disapproval, 247, table 6-2
 - friends' disapproval, table 7-1
 - gender, 100, table 4-1, table 4-5, table 8-4,
 - figure 9-14c
 - harmfulness, perceived, table 6-1, figure 6-27
 - population density, 103, 174, table 4-5
 - prevalence, 42, 97, table 4-1, table 4-5,
 - figure 4-20b
 - region, 166, table 4-5
 - trends, table 2-4, 150, 160, 385, table 5-4,
 - figure 5-18c, figure 9-14c
- disapproval, 246, table 6-2
- drunk, getting
 - college enrollment, 366, tables 8-1-8-3,
 - tables 9-1-9-4
 - friends' use, 299, table 7-2
 - gender, tables 4-1-4-4, tables 8-1-8-3
 - harmfulness, perceived, figure 6-29
 - prevalence, 368, tables 4-1-4-4, tables 8-1-8-3
 - race/ethnicity, 39
- exposure to use (*see also* friends' use), table 7-3,
 - figures 7-23-7-24
- flavored alcoholic beverages
 - college enrollment, 367, 369, tables 8-1-8-3
 - gender, 369, tables 4-1-4-5, tables 8-1-8-3
 - population density, tables 4-2-4-5
 - prevalence, 369, tables 4-1-4-4, tables 8-1-8-3
 - region, tables 4-2-4-4
 - trends, tables 2-1-2-3, tables 5-1-5-3, tables 9-1-9-3

friends' use, 303, table 7-2

harmfulness, perceived, table 6-1, figures 6-26-6-29

heavy drinking

- college enrollment, 366, 370, 384, figure 9-14d
- disapproval, table 6-2
- friends' use, 299, table 7-2
- gender, 100, tables 4-1-4-5, 388, figure 9-14d
- harmfulness, perceived, table 6-1, figure 6-28
- population density, 174, 103, tables 4-2-4-5
- prevalence, 30, 42, 91, 97, 368, 370, tables 4-1-4-5, table 8-4, figure 4-20b
- race/ethnicity, 38
- region, 102, 166, tables 4-2-4-5
- trends, 149-150, 160, 166, 384, figure 5-18d
- prevalence, 39, 40, 42, 91, 97, tables 4-1-4-5,
 - figure 4-20a
- college enrollment, 368, tables 8-1-8-4
- gender, 30, 32, tables 4-1-4-4, 369, 370,
 - 388, tables 8-1-8-4
- population density, 103, table 4-2-4-5
- race/ethnicity, 38
- region, 102, tables 4-2-4-5
- trends, 30-32, 141, 149, 154, tables 2-1-2-4,
 - tables 5-1-5-4, figures 5-18a-5-18d
- college enrollment, 31, 384-385, tables 9-1-9-4, figures 9-14a-9-14b
- friends' use, 303, table 7-2
- gender, 32, 159, 388, figures 9-14a-9-14b
- population density, 173
- region, 166

amphetamines

- availability, perceived, 304, 308, table 7-4
- cohort effect, 15-16, 24, 29, 30, 248
- daily use, table 8-4, table 9-4
- disapproval, 246, table 6-2
- exposure to use, 298, table 7-3, figures 7-17-7-18
- friends' disapproval, perceived, 294, 295, table 6-2, table 7-1
- friends' use, 301, table 7-2
- harmfulness, perceived, 238, 241, table 6-1,
 - figures 6-21-6-22
- prevalence, 24, 40, 41, 93, 94, tables 4-1-4-4,
 - figure 4-4
- college enrollment, 29, 365, 366, 368, tables 8-1-8-4

- gender, 100, 368, tables 4-1-4-4, tables 8-1-8-4
- population density, tables 4-2-4-4
- region, tables 4-2-4-4
- trends, 142, 144, 147, table 2-1-2-3, tables 5-1-5-3, tables 6-1-6-2, tables 7-1-7-4, tables 9-1-9-4, figure 5-13
- college enrollment, 379, 380, 386, 388, figure 9-11
- gender, 157, 388, figure 9-11
- population density, 103, 171
- race/ethnicity, 38, 39
- region, 161, 163
- amyl nitrites (*See* nitrites)
- anabolic steroids (*See* steroids)
- androstenedione
 - cohort effect, 26
 - prevalence, 27
 - gender, 27
- any illicit drug
 - cohort effect, 7, 8-9
 - exposure to use, 300, 301, table 7-3, figure 7-1-7-2
 - friends' use, table 7-2
 - prevalence, 9, 91, 92, figure 4-1
 - college enrollment, 365, 368, tables 8-1-8-3
 - gender, 100, 368, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - region, 101, tables 4-2-4-4
 - trends, 141, 142, 144, tables 2-1-2-3, tables 5-1-5-5, tables 9-1-9-3, figure 2-1, figure 5-1
 - college enrollment, 28, 377, 387, figure 9-1
 - gender, 154, table 5-5, 387, table 9-5, figure 9-1
 - population density, 167
 - region, 161
- any illicit drug other than marijuana
 - cohort effect, 8, 12, 29
 - exposure to use (*see also* friends' use), 300, tables 7-2-7-3, figure 7-3-7-4
 - friends' use, table 7-2
 - prevalence, 91, 93, figure 4-2
 - college enrollment, 365, 368, tables 8-1-8-3
 - gender, 100, 368, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - region, 101, tables 4-2-4-4
 - trends, 142, 144, tables 5-1-5-5, tables 2-1-2-3, tables 7-2-7-3, figure 5-2, figures 7-3-7-4
 - college enrollment, 378, 387, tables 9-1-9-3, table 9-5, figure 9-2

- gender, 154, 387, table 5-5, table 9-5, figure 9-2
- population density, 168
- region, 161-162

B

barbiturates (*See* sedatives)

bath salts

- harmfulness, perceived, 242, table 6-1
- prevalence, 8, 46, 93, 172
 - college enrollment, 366, table 8-2
 - gender, 30, 100, table 4-3
 - population density, 103, table 4-3
 - region, 164, table 4-3
- trends, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 158
 - perceived risk, 158
 - population density, 172
 - region, 164

bidis, table 2-2

butyl nitrites (*See* nitrites)

C

cigarettes

- cohort effect, 9, 35, 98

daily use

- college enrollment, 369, table 8-4, table 9-4, figures 9-15b-9-15c
- disapproval, table 6-2
- friends' disapproval, perceived, 294, 296, table 7-1
- gender, 369, table 4-5, table 8-4, figures 9-15b-9-15c
- harmfulness, perceived, 243, table 6-1, figure 6-30
- population density, 103, table 4-5
- prevalence, 42, 45, 98, table 4-5, figure 4-21
- region, table 4-5
- trends, 34, 42, 45, 243, 385, table 2-4, table 5-4, table 9-4, figures 5-19b-5-19c
- disapproval, 247, table 6-2
- friends' disapproval, perceived, 294, 296, table 7-1
- friends' use, 300, 303, table 7-2
- harmfulness, perceived, table 6-1, figure 6-30
- prevalence, 40, 45, 75, 80, 98, figure 4-21
 - college enrollment, 367, 369, tables 8-1-8-4
 - gender, 101, 369, tables 4-1-4-5, tables 8-1-8-4
 - population density, 104, tables 4-2-4-5
 - race/ethnicity, 38
 - region, 102, tables 4-2-4-5
- trends, 9, 32-37, 141, 151-154, tables 2-1-2-4, tables 5-1-5-4, tables 7-1-7-2, figures 5-19a-5-19c

- college enrollment, 36, 385, 389, tables 9-1-9-4, figures 9-15a-9-15c
 - gender, 36-37, 160, 389, figures 9-15a-9-15c
 - population density, 174
 - race/ethnicity, 38
 - region, 167
 - club drugs (*See* GHB; ketamine; Rohypnol)
 - cocaine
 - availability, perceived, 20, 42, 304, 306, table 7-4
 - cohort effect, 19-20, 24, 30, 248
 - disapproval, 244, 246, table 6-2
 - exposure to use (*see also* friends' use), 299, 300, table 7-3, figures 7-11-7-12
 - friends' disapproval, perceived, 294, 296, table 7-1
 - friends' use, 296, 299, 300, 302, table 7-2
 - harmfulness, perceived, 240, table 6-1, figures 6-7-6-9
 - prevalence, 24, 40, 41, 42, 44, 91, 93, 95, figure 4-7
 - college enrollment, tables 8-1-8-4
 - gender, 100, tables 4-1-4-4, tables 8-1-8-4
 - population density, 103, tables 4-2-4-4
 - region, tables 4-2-4-4
 - trends, 19-20, 29, 42, 142, 144-145, tables 2-1-2-3, tables 5-1-5-3, figure 5-9
 - college enrollment, 383-384, 388, tables 9-1-9-4, figure 9-9
 - gender, 30, 156, 388, figure 9-9
 - population density, 171
 - race/ethnicity, 39
 - region, 161, 163
 - See also* cocaine powder; crack cocaine
 - cocaine, other (*See* cocaine powder)
 - cocaine powder
 - availability, perceived, 304, table 7-4
 - cohort effect, 20, 30
 - friends' use, 299, table 7-2
 - harmfulness, perceived, 238, table 6-1, figures 6-13-6-15
 - prevalence, 93, figure 4-9
 - college enrollment, tables 8-1-8-3
 - gender, 30, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - region, tables 4-2-4-4
 - trends, 41, 142, tables 2-1-2-3, tables 5-1-5-3
 - college enrollment, tables 9-1-9-3
 - codeine, 22
 - cough/cold medicines, 56
 - crack cocaine
 - availability, perceived, 42, 304, 306, table 7-4
 - cohort effect, 18-19, 30, 248
 - friends' use, 298, 299, 302, table 7-2
 - harmfulness, perceived, 240-241, table 6-1, figures 6-10-6-12
 - prevalence, 40, 41, 42, 44, 93, 96, figure 4-8
 - college enrollment, 366, tables 8-1-8-3
 - gender, 30, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, tables 4-2-4-4
 - race/ethnicity, 35
 - region, tables 4-2-4-4
 - trends, 142, 144, 145, tables 2-1-2-3, tables 5-1-5-3, figure 5-10
 - college enrollment, tables 9-1-9-3
 - gender, 156
 - population density, 171
 - region, 161, 163
 - creatine
 - prevalence, 27
 - gender, 27
 - crystal methamphetamine (ice)
 - availability, perceived, 305, 308, table 7-4
 - cohort effect, 17
 - harmfulness, perceived, 241, 248, table 6-1, figure 6-23
 - prevalence, 93, 94-95, figure 4-6
 - college enrollment, 366, tables 8-1-8-3
 - gender, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - race/ethnicity, 38-39
 - region, tables 4-2-4-4
 - trends, 144, 147, table 2-1-2-3, tables 5-1-5-3, figure 5-14
 - college enrollment, tables 9-1-9-3
 - gender, 158
 - population density, 172
 - race/ethnicity, 38-39
 - region, 164
- D**
- Dextromethorphan, 28
 - diet pills, 27, 28
- E**
- ecstasy (MDMA)
 - availability, perceived, 305, 307, table 7-4
 - cohort effect, 14-15, 248
 - disapproval, 245, table 6-2
 - exposure to use (*see also* friends' use), 298
 - friends' use, 302, table 7-2
 - harmfulness, perceived, 238, 239, table 6-1, figures 6-16-6-17
 - prevalence, 25, 40, 42, 96, figure 4-17
 - college enrollment, 365, tables 8-1-8-3
 - gender, 100, tables 4-1-4-4, tables 8-1-8-3

- population density, 103, tables 4-2-4-4
- race/ethnicity, 38
- region, 102, tables 4-2-4-4
- trends, 25, 146, tables 2-1-2-3, tables 5-1-5-3, figure 5-8
- college enrollment, 379, 381-382, 387, tables 9-1-9-3, figure 9-8
- gender, 155, 387, figure 9-8
- population density, 170-171
- region, 162-163

exposure to drug use (*see also* friends' use), table 7-3

F

flavored alcoholic beverages (*See* alcohol)

G

GHB

- cohort effect, 25
- prevalence, 25
 - college enrollment, 366, table 8-2
 - gender, 100, table 4-3, table 8-2
 - population density, 103, table 4-3
 - region, table 4-3
- trends, 46, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 159
 - region, 166

H

hallucinogens

- prevalence, 41, 93, figure 4-10
 - college enrollment, 28, 368, tables 8-1-8-3
 - gender, 100, 368, 370, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - race/ethnicity, 38
 - region, 102, tables 4-2-4-4
- trends, 38, 142, 144, tables 2-1-2-3, tables 5-1-5-3, tables 9-1-9-3, figure 5-5
- college enrollment, 28, figure 9-5
- gender, figure 9-5

hallucinogens other than LSD

- availability, perceived, 305, 307, table 7-4
- exposure to use (*see also* friends' use), 298, table 7-3, figures 7-9-7-10
- friends' use, 298, table 7-2
- prevalence, 40, 41, 93, figure 4-12
 - college enrollment, 365, tables 8-1-8-3
 - gender, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, tables 4-2-4-4
 - race/ethnicity, 38
 - region, 102, tables 4-2-4-4
- trends, tables 2-1-2-3, tables 5-1-5-3, figure 5-7

- college enrollment, 381, 387, tables 9-1-9-3, figure 9-7
- gender, 155, 387, figure 9-7
- population density, 169

hashish (*See* marijuana)

heroin (including with and without a needle)

- availability, perceived, 304-305, table 7-4
- cohort effect, 20-22, 248
- disapproval, 244, 245-246, table 6-2
- exposure to use (*see also* friends' use), table 7-3, figures 7-13-7-14
- friends' use, 299, table 7-2
- harmfulness, perceived, 238, 242, table 6-1, figures 6-18-6-20
- prevalence, 20, 40, 41, 93, figure 4-19
 - college enrollment, 366, tables 8-1-8-3
 - gender, 30, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, tables 4-2-4-4
 - race/ethnicity, 38
 - region, tables 4-2-4-4
- trends, 46, tables 2-1-2-3, tables 5-1-5-3, figure 5-11
 - college enrollment, tables 9-1-9-3
 - gender, 156
 - population density, 172-173
 - region, 165

hydrocodone, 22

I

ice (*See* crystal methamphetamine)

inhalants, 17, 24, 25

- cohort effect
- friends' use, 299, table 7-2
- harmfulness, perceived, 17
- prevalence, 39, 40, 44, 93-94, figure 4-13
 - college enrollment, tables 8-1-8-3
 - gender, 30, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, tables 4-2-4-4
 - race/ethnicity, 38
 - region, tables 4-2-4-4
- trends, 40, 145-146, tables 2-1-2-3, tables 5-1-5-3, figure 5-4
- college enrollment, 380-381, tables 9-1-9-3, figure 9-4
- gender, 159, figure 9-4
- population density, 173
- region, 163

K

ketamine (Special K)

- prevalence, 25
- college enrollment, 365, table 8-2

- gender, 100, table 4-3, table 8-2
- population density, table 4-3
- region, table 4-3
- trends, 25, 46-47, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 159
 - region, 166
- kreteks, table 2-2

L

look-alike drugs, 28

LSD

- availability, perceived, 305, 306, table 7-4
- cohort effect, 13-14, 24-25, 248
- disapproval, 244, 245, table 6-2
- exposure to use (*see also* friends' use), 298, table 7-3, figures 7-7-7-8
- 7-1
 - friends' disapproval, perceived, 294, 295, table 7-1
 - friends' use, 298, table 7-2
 - harmfulness, perceived, 238, 239, table 6-1, figures 6-4-6-5
 - prevalence, 24-25, 40, 93, figure 4-11
 - college enrollment, 365, tables 8-1-8-3
 - gender, 30, 100, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - race/ethnicity, 38
 - region, 102, tables 4-2-4-4
 - trends, 46, tables 2-1-2-3, 144, tables 5-1-5-3, figure 5-6
 - college enrollment, 381, 387, tables 9-1-9-3, figure 9-6
 - gender, 155, 387, figure 9-6
 - population density, 169
 - region, 162

M

marijuana (hashish)

- availability, perceived, 304, 305, table 7-4
- cohort effect, 8-12, 153, 248
- daily use
 - college enrollment, 365, 368, 370, 380, 387, table 8-4, table 9-4, figure 9-3b
 - gender, 29, 100, table 4-5, 155, 368, 370, 387, table 8-4, figure 9-3b
 - population density, 168, table 4-5
 - prevalence, 29, 41, 92, table 8-4
 - region, table 4-5
 - trends, 44, 141, 143, 153, 380, table 2-4, table 5-4, table 9-4, figure 5-3c
- disapproval, 245, table 6-2
- exposure to use (*see also* friends' use), 298, 300, 301, table 7-3, figures 7-5-7-6

- friends' disapproval, perceived, 293, 294-295, table 7-1
- friends' use, 301, table 7-2
- harmfulness, perceived, 237-238, table 6-1, figures 6-1-6-3
- prevalence, 24-25, 40, 41, 91, 92, figure 4-3
 - college enrollment, 28, 365, 368, tables 8-1-8-3
 - gender, 29, 100, 368, tables 4-1-4-4, tables 8-1-8-4
 - population density, 103, tables 4-2-4-4
 - race/ethnicity, 38-39
 - region, 101, tables 4-2-4-4
- trends, 42, 141, 142, 143, 144, tables 2-1-2-4, tables 5-1-5-4, figure 5-3a-5-3c
- college enrollment, 29, 379-380, tables 9-1-9-3, figure 9-3a
- gender, 154, figure 9-3a
- population density, 168
- race/ethnicity, 38
- region, 161-162

MDMA (*See* ecstasy)

methamphetamine

- cohort effect, 17, 248
- prevalence, 40, 94, figure 4-5
 - college enrollment, 365, tables 8-1-8-2
 - gender, 30, tables 4-1-4-4, tables 8-1-8-2
 - population density, 103, tables 4-2-4-4
 - region, tables 4-2-4-4
- trends, 147, tables 2-1-2-3, tables 5-1-5-3
- college enrollment, tables 9-1-9-3
- gender, 158
- population density, 172
- region, 164

methaqualone

- availability, perceived, 24
- prevalence, 24, 172
- trends, 144, tables 2-1-2-3, tables 5-1-5-3
- college enrollment, tables 9-1-9-3
- population density, 172

N

narcotics other than heroin

- availability, perceived, 305, 307
- harmfulness, perceived, 238, 242, table 6-1
- cohort effect, 22, 25
- exposure to use (*see also* friends' use), 298, figures 7-15-7-16
- friends' use, 299, 302
- prevalence, 41, 93, 95, figure 4-15
 - college enrollment, 366, 368, tables 8-1-8-3
 - gender, 100, 368, tables 4-1-4-2, tables 8-1-8-3
 - population density, table 4-2
 - race/ethnicity, 38

- region, table 4-2
- trends, 22, 46, 142, 147-148, tables 2-1-2-3, tables 5-1-5-3, figure 5-12
- college enrollment, 29, 382, 388, tables 9-1-9-3, figure 9-10a
- gender, trends 156-157, 388, figure 9-10a
- population density, 173
- region, 165

nitrites (*See also* inhalants)

- availability, perceived, table 7-4
- friends' use, table 7-2
- prevalence, 18
- trends, 18, tables 2-1-2-3, tables 5-1-5-3

O

OxyContin

- prevalence, 22-23, 25
 - college enrollment, 366, 368, table 8-2
 - gender, 30, 100, 368, table 4-3, table 8-2
 - population density, table 4-3
 - race/ethnicity, 38
 - region, table 4-3
- trends, 22-23, 46, 148, table 2-2, table 5-2
- college enrollment, 383, table 9-2, figure 9-10c
- gender, 157
- population density, 173
- region, 165

P

PCP

- availability, perceived, table 7-4
- cohort effect, 20
- friends' use, table 7-2
- harmfulness, perceived, table 6-1, figure 6-6
- prevalence
 - gender, tables 4-1-4-4
 - population density, tables 4-2-4-4
 - region, tables 4-2-4-4
- trends, tables 2-1-2-3, tables 5-1-5-3

Percocet, 22

powder cocaine (*See* cocaine powder)

Provigil

- cohort effect, 26
- prevalence, 97
- trends, 46, table 2-2, table 5-2

psychedelics (*See* hallucinogens)

psychotherapeutic drugs, 24

Q

Quaaludes, table 7-2

R

Ritalin

- cohort effect, 16
- prevalence, 94
 - college enrollment, 366, 368, 369, table 8-2
 - gender, 30, table 4-3, 369, table 8-2
 - population density, table 4-3
 - race/ethnicity, 38
 - region, table 4-3
- trends, table 2-2, table 5-2
 - college enrollment, 380, table 9-2
 - gender, 157
 - race/ethnicity, 39

Rohypnol

- prevalence, 25, 40
 - gender, 30, 159
 - region, 166
- trends, 40, 46-47, tables 2-1-2-3, table 5-2
- college enrollment, table 9-2

S

salvia

- harmfulness, perceived, 240, table 6-1
- prevalence, 26, 96-97
 - college enrollment, table 8-2
 - gender, table 4-3, table 8-2
 - population density, 103, table 4-3
 - race/ethnicity, 38
 - region, table 4-3
- trends, 26, 146-147, table 2-2, table 5-2
- college enrollment, table 9-2
- gender, 155-156
- population density, 170
- region, 162

sedatives (barbiturates)

- availability, perceived, 305, 309, table 7-4
- cohort effect, 25, 248
- disapproval, 244, 246, table 6-2
- exposure to use (*see also* friends' use), table 7-3, figures 7-19-7-20
- friends' use, 298, 301, table 7-2
- harmfulness, perceived, 238, table 6-1, figures 6-24-6-25
- prevalence, 23, 41, 93, 95, figure 4-14
 - college enrollment, tables 8-1-8-3
 - gender, tables 4-1-4-4, tables 8-1-8-3
 - population density, tables 4-2-4-4
 - race/ethnicity, 38, 39
 - region, tables 4-2-4-4
- trends, 46, 47, 142, 149, tables 2-1-2-3, tables 5-1-5-3, figure 5-15
- college enrollment, 29, 382, 386, 388, tables 9-1-9-3, figure 9-12
- gender, 158, 388, figure 9-12

- population density, 172
 - region, 164
- smokeless tobacco
 - harmfulness, perceived, 244, table 6-1, figure 6-31
 - prevalence, 99
 - gender, 40, 101, tables 4-1-4-2, tables 4-4-4-5
 - population density, 104, table 4-2, tables 4-4-4-5
 - race/ethnicity, 38
 - region, 102, table 4-2, tables 4-4-4-5
 - trends, 45, tables 2-1-2-4
- stay-awake pills, 27
- steroids
 - availability, perceived, 305, 309, table 7-4
 - friends' use, 298, table 7-2
 - prevalence, 40, 99, 149, figure 4-18
 - college enrollment, tables 8-1-8-3
 - gender, 26, 27, 100, 101, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - region, tables 4-2-4-4
 - trends, 149, tables 2-1-2-3, tables 5-1-5-3, figure 5-17
 - college enrollment, tables 9-1-9-3
- synthetic marijuana
 - harmfulness, perceived, 239, table 6-1
 - prevalence, 12, 92-93, 366, table 4-3
 - college enrollment, table 8-2
 - gender, 100, table 8-2
 - population density, 103
 - region, 101
 - trends, 46, 141, 144, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 155
 - population density, 168
 - region, 162

T

- tobacco (*See* cigarettes; smokeless tobacco)
- tranquilizers
 - availability, perceived, 305, 309, table 7-4
 - cohort effects, 23, 25
 - exposure to use (*see also* friends' use), table 7-3, figures 7-21-7-22
 - friends' use, 298, 301, table 7-2
 - prevalence, 40, 41, 93, 95, figure 4-16
 - college enrollment, 29, tables 8-1-8-3
 - gender, tables 4-1-4-4, tables 8-1-8-3
 - population density, 103, tables 4-2-4-4
 - race/ethnicity, 38, 39
 - region, tables 4-2-4-4
 - trends, 46, 47, 142, 144, 149, tables 2-1-2-3, tables 5-1-5-3, figure 5-16

- college enrollment, 382, 386, 388, tables 9-1-9-3, figure 9-13
- gender, 158, 388, figure 9-13
- population density, 172
- race/ethnicity, 38, 39
- region, 165

V

- Valium, 23
- Vicodin
 - cohort effect 22, 25
 - prevalence
 - college enrollment, 366, 368, table 8-2
 - gender, 30, 100, 368, table 4-3, table 8-2
 - population density, table 4-3
 - race/ethnicity, 38
 - region, table 4-3
 - trends, 148, table 2-2, table 5-2
 - college enrollment, 383, table 9-2, figure 9-10b
 - gender, 157
 - population density, 173
 - region, 165-166

X

- Xanax, 23



**Monitoring the Future website:
<http://www.monitoringthefuture.org>**

