

NATIONAL INSTITUTE ON DRUG ABUSE

Monitoring the Future

National Survey Results on Drug Use, 1975–2004

Volume II

College Students & Adults Ages 19–45 2004

National Institutes of Health
U.S. Department of Health and Human Services

MONITORING THE FUTURE
NATIONAL SURVEY RESULTS ON DRUG USE, 1975-2004

Volume II
College Students and Adults Ages 19-45

by

Lloyd D. Johnston, Ph.D.
Patrick M. O'Malley, Ph.D.
Jerald G. Bachman, Ph.D.
John E. Schulenberg, Ph.D.

The University of Michigan
Institute for Social Research

National Institute on Drug Abuse
6001 Executive Boulevard
Bethesda, Maryland 20892

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
National Institutes of Health

2005

This publication was written by the principal investigators and staff of the Monitoring the Future project, at the Institute for Social Research, the University of Michigan, under Research Grants No. R01 DA 01411 and R01 DA 016575 from the National Institute on Drug Abuse.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the NIH.

Public Domain Notice

All material appearing in this volume is in the public domain and may be reproduced or copied without permission from the Institute or the authors. Citation of the source is appreciated.

Recommended Citation

Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2005). *Monitoring the Future national survey results on drug use, 1975-2004: Volume II, College students and adults ages 19-45* (NIH Publication No. 05-5728). Bethesda, MD: National Institute on Drug Abuse.

ABBREVIATED CONTENTS*

Detailed Contents.....	v
List of Tables	ix
List of Figures.....	xiii
<i>Chapter 1</i> Introduction	1
<i>Chapter 2</i> Key Findings: An Overview and Integration Across Five Populations ...	5
<i>Chapter 3</i> Study Design and Procedures	55
<i>Chapter 4</i> Prevalence of Drug Use in Early and Middle Adulthood	75
<i>Chapter 5</i> Trends in Drug Use in Early to Middle Adulthood	119
<i>Chapter 6</i> Attitudes and Beliefs About Drugs Among Young Adults	173
<i>Chapter 7</i> The Social Milieu for Young Adults	189
<i>Chapter 8</i> Prevalence of Drug Use Among College Students	225
<i>Chapter 9</i> Trends in Drug Use Among College Students	235
Index	273

*See next page for Detailed Contents.

DETAILED CONTENTS

	Page
<i>Chapter 1</i> Introduction	1
Surveys of College Students.....	1
Surveys of Young Adults and Those Aged 35, 40, and 45.....	2
General Purposes of the Research.....	2
<i>Chapter 2</i> Key Findings: An Overview and Integration Across Five Populations	5
Trends in Illicit Drug Use.....	6
College-Noncollege Differences in Illicit Drug Use.....	19
Male-Female Differences in Illicit Drug Use.....	20
Trends in Alcohol Use.....	21
College-Noncollege Differences in Alcohol Use.....	22
Male-Female Differences in Alcohol Use.....	23
Trends in Cigarette Smoking.....	23
Age and Cohort-Related Differences in Cigarette Smoking.....	25
College-Noncollege Differences in Cigarette Smoking.....	26
Male-Female Differences in Cigarette Smoking.....	26
Racial/Ethnic Comparisons.....	27
Drug Use in Eighth Grade.....	28
Drug Use by Age 45.....	30
Summary and Conclusions.....	31
<i>Chapter 3</i> Study Design and Procedures	55
Research Design and Procedures for the Surveys of Seniors.....	55
The Population Under Study.....	56
The Omission of Dropouts.....	56
Sampling Procedures.....	56
Questionnaire Administration.....	57
Questionnaire Format.....	57
Research Design and Procedures for the Surveys of Lower Grades.....	57
Mode of Administration.....	58
Questionnaire Forms and Sample Proportions.....	59
Research Design and Procedures for the Follow-Up Surveys of Seniors.....	60
Follow-Up Procedures.....	60
Follow-Up Questionnaire Format.....	61
Representativeness and Sample Accuracy.....	61
School Participation.....	61
Student Participation.....	63
Sampling Accuracy of the Estimates.....	64
Panel Retention.....	64

DETAILED CONTENTS (continued)

	Page
The Problem of Panel Attrition.....	64
Response Rates Attained.....	65
The Impact of Panel Attrition on Research Results.....	66
Effects on Relational Analyses.....	68
Validity of the Measures of Self-Reported Drug Use.....	68
Consistency and the Measurement of Trends.....	69

EARLY AND MIDDLE ADULTHOOD

Chapter 4 Prevalence of Drug Use in Early and Middle Adulthood.....	75
A Note on Adjusted Lifetime Prevalence Estimates.....	76
Prevalence of Drug Use as a Function of Age.....	77
Prevalence Comparisons for Subgroups of Young Adults.....	83
Gender Differences.....	83
Regional Differences.....	84
Population Density Differences.....	85
Chapter 5 Trends in Drug Use in Early to Middle Adulthood.....	119
Trends in Prevalence: Early and Middle Adulthood.....	120
Trends for Important Subgroups of Young Adults.....	128
Gender Differences in Trends.....	128
Regional Differences in Trends.....	133
Population Density Differences in Trends.....	136
Chapter 6 Attitudes and Beliefs About Drugs Among Young Adults.....	173
Perceived Harmfulness of Drugs.....	173
Personal Disapproval of Drug Use.....	177
A Further Commentary: Cohort Differences and Their Implications for Prevention and Theory.....	180
Chapter 7 The Social Milieu for Young Adults.....	189
Peer Norms as Perceived by Young Adults.....	189
Current Perceptions of Friends' Attitudes.....	190

DETAILED CONTENTS (continued)

	Page
Trends in Peer Norms	190
Exposure to Drug Use by Friends and Others	192
Exposure to Drug Use Among Adults	192
Trends in Exposure to Drug Use Among Young Adults	194
Perceived Availability of Drugs by Young Adults	197
Perceived Availability.....	197
Trends in Perceived Availability	198

COLLEGE STUDENTS

<i>Chapter 8</i> Prevalence of Drug Use Among College Students	225
Prevalence of Drug Use: College Students Versus Those Not Enrolled in College.....	226
Gender Differences in Prevalence of Use Among College Students.....	228
<i>Chapter 9</i> Trends in Drug Use Among College Students	235
Trends in Prevalence 1980-2004: College Students Versus Those Not Enrolled in College	236
Gender Differences in Trends Among College Students.....	242
<i>Index</i>	273

LIST OF TABLES

	Page
Table 2-1. Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)	35
Table 2-2. Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28).....	42
Table 2-3. Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28).....	48
Table 2-4. Trends in Thirty-Day Prevalence of <u>Daily</u> Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)	53
Table 3-1. Sample Sizes and Response Rates	71
Table 4-1. Prevalence of Use of Various Types of Drugs by Gender, 2004, Among Respondents of Modal Ages 19-30.....	87
Table 4-2. Lifetime Prevalence of Use of Various Types of Drugs by Subgroups, 2004, Among Respondents of Modal Ages 19-30.....	89
Table 4-3. Annual Prevalence of Use of Various Types of Drugs by Subgroups, 2004, Among Respondents of Modal Ages 19-30.....	91
Table 4-4. Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups, 2004, Among Respondents of Modal Ages 19-30.....	94
Table 4-5. Thirty-Day Prevalence of <u>Daily</u> Use of Various Types of Drugs by Subgroups, 2004, Among Respondents of Modal Ages 19-30.....	96
Table 5-1. Trends in Lifetime Prevalence of Various Types of Drugs Among Respondents of Modal Ages 19-28	141
Table 5-2. Trends in Annual Prevalence of Various Types of Drugs Among Respondents of Modal Ages 19-28	143
Table 5-3. Trends in Thirty-Day Prevalence of Various Types of Drugs Among Respondents of Modal Ages 19-28.....	144
Table 5-4. Trends in Thirty-Day Prevalence of <u>Daily</u> Use of Various Types of Drugs Among Respondents of Modal Ages 19-28.....	145

LIST OF TABLES (continued)

	Page
Table 5-5. Trends in Annual and Thirty-Day Prevalence of an Illicit Drug Use Index Among Respondents of Modal Ages 19-28.....	146
Table 6-1. Trends in Perceived Harmfulness of Drugs: High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30.....	182
Table 6-2. Trends in Proportions Disapproving of Drug Use: High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30.....	186
Table 7-1. Trends in Proportions of Friends Who Disapprove of Drug Use: High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30.....	201
Table 7-2. Trends in Proportions of Friends Using Drugs: High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45.....	203
Table 7-3. Trends in Exposure to Drug Use: High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30.....	215
Table 7-4. Trends in Reported Availability of Drugs: High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45.....	219
Table 8-1. Lifetime Prevalence of Use for Various Types of Drugs, 2004: Full-Time College Students vs. Others Among Respondents 1-4 Years Beyond High School.....	230
Table 8-2. Annual Prevalence of Use for Various Types of Drugs, 2004: Full-Time College Students vs. Others Among Respondents 1-4 Years Beyond High School.....	231
Table 8-3. Thirty-Day Prevalence of Use for Various Types of Drugs, 2004: Full-Time College Students vs. Others Among Respondents 1-4 Years Beyond High School.....	232
Table 8-4. Thirty-Day Prevalence of <u>Daily</u> Use for Various Types of Drugs, 2004: Full-Time College Students vs. Others Among Respondents 1-4 Years Beyond High School.....	233
Table 8-5. Lifetime, Annual, and Thirty-Day Prevalence of an Illicit Drug Use Index, 2004: Full-Time College Students vs. Others Among Respondents 1-4 Years Beyond High School.....	234

LIST OF TABLES (continued)

	Page
Table 9-1. Trends in Lifetime Prevalence of Various Types of Drugs Among College Students 1-4 Years Beyond High School	244
Table 9-2. Trends in Annual Prevalence of Various Types of Drugs Among College Students 1-4 Years Beyond High School	246
Table 9-3. Trends in Thirty-Day Prevalence of Various Types of Drugs Among College Students 1-4 Years Beyond High School	247
Table 9-4. Trends in Thirty-Day Prevalence of <u>Daily</u> Use of Various Types of Drugs Among College Students 1-4 Years Beyond High School	248
Table 9-5. Trends in Lifetime, Annual, and Thirty-Day Prevalence of an Illicit Drug Use Index Among College Students 1-4 Years Beyond High School, by Gender	249

LIST OF FIGURES

	Page
Figure 2-1. Trends in Annual Prevalence of an Illicit Drug Use Index Across Five Populations.....	54
Figure 3-1. Schools Included in One Year’s Data Collection: Eighth, Tenth and Twelfth Grades.....	72
Figure 3-2. School Response Rates	73
Figures 4-1–4-19. Lifetime, Annual, and Thirty-Day Prevalence Among High School Seniors and Adults Through Age 45, 2004, by Age Group	
4-1. Any Illicit Drug.....	97
4-2. Any Illicit Drug Other Than Marijuana	98
4-3. Marijuana	99
4-4. Amphetamines	100
4-5. Methamphetamine.....	101
4-6. Cocaine	102
4-7. Crack Cocaine.....	103
4-8. Other Cocaine	104
4-9. Hallucinogens	105
4-10. LSD	106
4-11. Hallucinogens Other Than LSD.....	107
4-12. Inhalants.....	108
4-13. Sedatives (Barbiturates).....	109
4-14. Narcotics Other Than Heroin.....	110
4-15. Tranquilizers	111
4-16. MDMA.....	112
4-17. Crystal Methamphetamine (“Ice”).....	113
4-18. Steroids	114
4-19. Heroin	115
Figure 4-20a. Alcohol: Lifetime, Annual, and Thirty-Day Prevalence Among High School Seniors and Adults Through Age 45, 2004, by Age Group.....	116
Figure 4-20b. Alcohol: Two-Week Prevalence of Five or More Drinks in a Row and Thirty-Day Prevalence of Daily Use Among High School Seniors and Adults Through Age 45, 2004, by Age Group.....	117
Figure 4-21. Cigarettes: Annual, Thirty-Day, Daily, and Half-Pack-a-Day Prevalence Among High School Seniors and Adults Through Age 45, 2004, by Age Group.....	118

LIST OF FIGURES (continued)

Page

Figure 5-1.	Any Illicit Drug: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	147
Figure 5-2.	Any Illicit Drug Other Than Marijuana: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group...	148
Figure 5-3a.	Marijuana: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	149
Figure 5-3b.	Marijuana: Trends in Thirty-Day Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	150
Figure 5-3c.	Marijuana: Trends in Thirty-Day Prevalence of <u>Daily</u> Use Among High School Seniors and Adults Through Age 45, by Age Group.....	151
Figure 5-4.	Inhalants: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	152
Figure 5-5.	Hallucinogens: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	153
Figure 5-6.	LSD: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	154
Figure 5-7.	Hallucinogens Other Than LSD: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	155
Figure 5-8.	MDMA: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	156
Figure 5-9.	Cocaine: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	157
Figure 5-10.	Crack Cocaine: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	158
Figure 5-11.	Heroin: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	159
Figure 5-12.	Narcotics Other Than Heroin: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	160

LIST OF FIGURES (continued)

		Page
Figure 5-13.	Amphetamines: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	161
Figure 5-14.	Ice: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	162
Figure 5-15.	Sedatives (Barbiturates): Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	163
Figure 5-16.	Tranquilizers: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	164
Figure 5-17.	Steroids: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	165
Figure 5-18a.	Alcohol: Trends in Annual Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	166
Figure 5-18b.	Alcohol: Trends in Thirty-Day Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	167
Figure 5-18c.	Alcohol: Trends in Thirty-Day Prevalence of <u>Daily</u> Use Among High School Seniors and Adults Through Age 45, by Age Group.....	168
Figure 5-18d.	Alcohol: Trends in Two-Week Prevalence of Having Five or More Drinks in a Row at Least Once Among High School Seniors and Adults Through Age 45, by Age Group	169
Figure 5-19a.	Cigarettes: Trends in Thirty-Day Prevalence Among High School Seniors and Adults Through Age 45, by Age Group.....	170
Figure 5-19b.	Cigarettes: Trends in Thirty-Day Prevalence of <u>Daily</u> Use Among High School Seniors and Adults Through Age 45, by Age Group.....	171
Figure 5-19c.	Cigarettes: Trends in Thirty-Day Prevalence of Smoking a Half-Pack or More Daily Among High School Seniors and Adults Through Age 45, by Age Group.....	172
Figure 9-1.	Any Illicit Drug: Trends in Annual Prevalence Among College Students vs. Others and Among Male vs. Female College Students.....	250

LIST OF FIGURES (continued)

Page

Figure 9-2.	Any Illicit Drug Other Than Marijuana: Trends in Annual Prevalence Among College Students vs. Others and Among Male vs. Female College Students	251
Figure 9-3.	Marijuana: Trends in Prevalence Among College Students vs. Others and Among Male vs. Female College Students	
a.	Annual Prevalence	252
b.	Thirty-Day Prevalence of <u>Daily</u> Use.....	253
Figures 9-4–9-13.	Trends in Annual Prevalence Among College Students vs. Others and Among Male vs. Female College Students	
9-4.	Inhalants	254
9-5.	Hallucinogens	255
9-6.	LSD	256
9-7.	Hallucinogens Other Than LSD.....	257
9-8.	MDMA (Ecstasy).....	258
9-9.	Cocaine	259
9-10a.	Narcotics Other Than Heroin.....	260
9-10b.	Vicodin.....	261
9-10c.	OxyContin.....	262
9-11.	Amphetamines	263
9-12.	Sedatives (Barbiturates)	264
9-13.	Tranquilizers	265
Figure 9-14.	Alcohol: Trends in Prevalence Among College Students vs. Others and Among Male vs. Female College Students	
a.	Annual Prevalence	266
b.	Thirty-Day Prevalence	267
c.	Thirty-Day Prevalence of <u>Daily</u> Use.....	268
d.	Two-Week Prevalence of Five or More Drinks in a Row	269
Figure 9-15.	Cigarettes: Trends in Prevalence Among College Students vs. Others and Among Male vs. Female College Students	
a.	Thirty-Day Prevalence	270
b.	Thirty-Day Prevalence of <u>Daily</u> Use.....	271
c.	Thirty-Day Prevalence of Smoking a Half-Pack or More per Day	272

Chapter 1

INTRODUCTION

This volume—the second in a two-volume set from the Monitoring the Future study—provides findings on the substance use and related behaviors of several segments of the adult population. It also contains findings on attitudes and beliefs about drugs, as well as on several particularly salient dimensions of their social environments. Volume I presents similar findings for American secondary students in grades 8, 10, and 12.

One important segment covered here is the population of American college students; a second is their age peers who are not attending college. Also covered in this volume are young adult high school graduates ages 19 to 30 (including the college students), as well as high school graduates at ages 35, 40, and 45.

Monitoring the Future is a long-term research program conducted at the University of Michigan's Institute for Social Research under a series of investigator-initiated research grants from the National Institute on Drug Abuse. Now in its 30th year, it comprises, in part, ongoing series of annual nationally representative surveys of 12th (begun in 1975) and of 8th- and 10th-grade students (begun in 1991).

Over the years, *follow-up surveys* have been conducted of representative samples of the previous participants from each high school senior class. The present volume presents data from the 1977 through 2004 follow-up surveys of the graduating high school classes of 1976 through 2003 as these respondents have progressed into adulthood—through age 45 for the oldest respondents.

To permit this volume to stand alone, we have repeated some material from Volume I. Specifically, chapter 2 in this volume is the same as chapter 2 in Volume I; it provides an integrated overview of the key findings presented in both volumes. Chapter 3, “Study Design and Procedures,” is also the same as chapter 3, Volume I. The reader already familiar with Volume I may wish to skip over these chapters. Otherwise, the content of the two volumes does not overlap.

SURVEYS OF COLLEGE STUDENTS

The follow-up samples in Monitoring the Future have provided very good coverage of the national college student population for almost a quarter of a century—since 1980. College students tend to be a difficult population to study. They generally were not well covered in normal household surveys, which typically excluded dormitories, fraternities, and sororities. Further, institution-based samples of college students must be quite large in order to attain accurate national representation because of the great heterogeneity in the types of student populations served in those institutions. Obtaining good samples and high response rates within many institutions may also pose difficulties. The current study, which in essence draws the college sample in senior year of high school, has considerable advantages for generating a broadly representative sample of college students who

emerge from each graduating cohort (and it does so at very low cost). Further, its “before,” “during,” and “after” college measurement design permits examination of the many changes associated with the college experience. Moreover, it has comparable panel data on the high school graduates who are not attending college, a segment that is important not only in its own right (having been called the “forgotten half”) but also for purposes of providing a comparison group for the college students.

As defined here, the college student population comprises all full-time students, one to four years post-high school, enrolled in a two- or four-year college in March during the year of the survey. More is said about this sample definition in chapters 3 and 8. Results on the prevalence of drug use among college students in 2004 are reported in chapter 8, and results on the trends in substance use among college students over the past 25 national surveys are reported in chapter 9. Both chapters also report data on the portion of the follow-up samples who are in the same age band as the college students but who are not enrolled in college.

SURVEYS OF YOUNG ADULTS AND THOSE AGED 35, 40, AND 45

A “young adult” sample, on which we also report here, comprises representative samples from each graduating class from 1992 to 2003, all surveyed in 2004. Since 18 is the modal age of 12th graders, the young adults covered here correspond to modal ages 19 through 30. (The college students are included as a part of this young adult sample.) The study design calls for annual follow-up surveys of each class cohort (though each individual participates in a follow-up survey every two years) through age 30, after which surveys occur at five-year intervals beginning at age 35. In 2004 the graduating classes of 1977, 1982, and 1987 were sent the “age 45,” “age 40,” and “age 35” questionnaires, respectively. Data were collected from 45-year-old respondents for the first time in 2003.

In this volume, we have re-weighted respondent data to adjust for the effects of panel attrition on measures such as drug use. We are less able to adjust for the absence of high school dropouts who were not included in the original high school senior sample. Because nearly all college students have completed high school, the omission of dropouts should have almost no effect on the college student estimates, but this omission does affect the estimates for entire age groups. Therefore, the reader is cautioned that the omission of about 15% of each cohort who drop out of high school will make the drug use estimates given here for the various young adult age bands somewhat low for the age group as a whole. The proportional effect may be greatest for some of the most dangerous drugs, such as heroin and crack, and also for cigarettes—the use of which is highly correlated with educational aspirations and attainment.

GENERAL PURPOSES OF THE RESEARCH

The research purposes of the Monitoring the Future study are extensive and can be sketched only briefly here.¹ One major purpose is to serve a social monitoring or social indicator function, intended

¹For a more complete listing and discussion of the study’s many objectives, see Johnston, L. D., O’Malley, P. M., Schulenberg, J. E., & Bachman, J. G. (2001). *The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2001*. (Monitoring the Future Occasional Paper No. 52). Ann Arbor, MI: Institute for Social Research. 139 pp. It is available on-line at www.monitoringthefuture.org/pubs.html#papers.

to characterize accurately the levels and trends in certain behaviors, attitudes, beliefs, and conditions in the population. Social indicators can have important agenda-setting functions for society, and they are useful for gauging progress with national goals and providing an indication of the impacts of major historical events or social changes. Another purpose of the study is to develop knowledge that increases our understanding of how and why changes in these behaviors, attitudes, beliefs, and so on are taking place. (In health-related disciplines, such work is usually labeled *epidemiology*.) These two broad purposes are addressed in the current series of volumes.

The many additional purposes for the research are addressed through other types of publications and professional products. These include helping to determine what types of young people are at greatest risk for developing various patterns of drug abuse; gaining a better understanding of the lifestyles and value orientations associated with various patterns of drug use, and monitoring how those orientations are shifting over time; determining the immediate and more general aspects of the social environment that are associated with drug use and abuse; and determining how drug use is affected by major transitions into and out of social environments (such as military service, civilian employment, college, unemployment) or social roles (marriage, pregnancy, parenthood). We also seek to determine the life course of the various drug-using behaviors during this period of development; distinguish such “age effects” from cohort and period effects that are influencing drug use; determine the effects of social legislation on various types of substance use; and determine the changing connotations of drug use and changing patterns of multiple drug use among youth. We believe that differentiating between period, age, and cohort effects on use of various types of substances has been a particularly important contribution of the project; its cohort-sequential research design is especially well suited to allow such differentiation. In fact, a number of important cohort effects that emerged in the 1990s in terms of both use and attitudes about use will be featured in this volume.

Readers interested in publications dealing with any of these other areas are invited to visit the study’s Web site at www.monitoringthefuture.org. A complete listing of all publications from the study is available there, as well as abstracts and/or complete manuscripts for many of those publications. Complete text of press releases from the study is also provided. Additional information may be requested through the Web site or by writing the authors at the Institute for Social Research, the University of Michigan, Ann Arbor, Michigan, 48106-1248.

Chapter 2

KEY FINDINGS: AN OVERVIEW AND INTEGRATION ACROSS FIVE POPULATIONS

Monitoring the Future, which is now in its 30th year, has become one of the nation's most relied-upon sources of information on changes taking place in licit and illicit psychoactive drug use among American adolescents, college students, young adults and, more recently, middle-aged adults. During the last three decades, the study has tracked and reported the use of an ever-growing array of such substances in these populations.

This annual series of monographs, written by the study's investigators and published by its sponsor—the National Institute on Drug Abuse—is one of the major vehicles by which the epidemiological findings from the study are reported. The present two-volume monograph reports findings from the inception of the study in 1975 through 2004. (A companion series of annual reports provides a much briefer, advance synopsis of the key findings from the latest surveys of secondary school students.²)

Monitoring the Future has conducted in-school surveys of nationally representative samples of (a) high school seniors each year since 1975 and (b) 8th- and 10th-grade students each year since 1991. In addition, beginning with the class of 1976, follow-up surveys have been conducted by mail on representative sub-samples of the respondents from each previously-participating 12th-grade class; follow-up surveys of all those classes continue into young adulthood and beyond.

A number of important findings have been summarized and integrated in this chapter to provide the reader with an overview of the key results. Because so many populations, drugs, and prevalence intervals are discussed here, a single integrative set of tables (Tables 2-1 through 2-4) showing the 1991-2004 trends for all drugs on all five populations (8th-grade students, 10th-grade students, 12th-grade students, full-time college students ages 19-22, and all young adults through age 28 who are high school graduates) is included in this chapter. (Note that the young adult group includes the college student population.) Volume II contains additional data on older age bands: specifically, ages 35, 40, and 45.

²Johnston, L. D., O'Malley P. M., Bachman, J. G., & Schulenberg J. E. (2005). *Monitoring the Future national results on adolescent drug use: Overview of key findings, 2004*. (NIH Publication No. 05-5726). Bethesda, MD: National Institute on Drug Abuse. (Also available on the Web at <http://monitoringthefuture.org>.)

TRENDS IN ILLICIT DRUG USE

Early in the 1990s we noted an increase in use of several illicit drugs among secondary students and some important changes among the students in terms of certain key attitudes and beliefs related to drug use. In the volume reporting 1992 survey results, we noted the beginning of such reversals in both use and attitudes among 8th graders, the youngest respondents surveyed in this study, and also a reversal in attitudes among the 12th graders. Specifically, the proportions seeing great risk in using drugs began to decline, as did the proportions saying they disapproved of use. As we suggested then, those reversals indeed presaged “an end to the improvements in the drug situation that the nation may be taking for granted.” The use of illicit drugs rose sharply in all three grade levels after 1992 as negative attitudes and beliefs about drug use continued to erode. This pattern continued for several years.

In 1997, for the first time in six years, illicit drug use finally began to decline among 8th graders. Use of marijuana continued to rise among the 10th and 12th graders, although their use of several other drugs leveled off and relevant attitudes and beliefs also began to reverse in many cases. In 1998, illicit drug use continued a gradual decline among 8th graders and started to decline at 10th and 12th grades. In 1999, 2000, and 2001, the decline continued for 8th graders while use held fairly level among 10th and 12th graders. In 2002 and 2003, use by 8th and 10th graders decreased significantly and use by 12th graders dropped, but by less than a statistically significant amount. Nonsignificant declines continued for all three grades in 2004. As we have noted previously, the fact that use continued to decline steadily, albeit slowly, among the 8th graders suggested there would be an eventual further decline at the upper grades. We are now seeing those declines.

As will be illustrated in the discussion of specific drugs, the increase in use of many drugs during the 1990s among secondary school students, combined with fairly level rates of use among college students and young adults, resulted in some unusual reversals in the usage rates by age. Figure 2-1 illustrates the point. In the early years of the epidemic, illicit drug use rates clearly were higher in the college-age group (and eventually the young adults) than they were among secondary school students. But by the late 1990s, the highest rates of active use (i.e., annual or 30-day prevalence) tended to be found in the late secondary school years. In fact, in 1996 and 1997 both 10th and 12th graders actually had higher prevalence rates for illicit drug use than either college students or young adults. This changed somewhat after 2001, when use in the older age groups rose, as the heavier-using cohorts of adolescents began to comprise the college student and young adult populations. At the same time, use among the secondary students was declining.

- In 2004 the rank order for annual prevalence of using *any illicit drug* was 12th graders (39%), college students (36%), 19- to 28-year-olds (34%), 10th graders (31%), and 8th graders (15%). With respect to using *any illicit drug other than marijuana* in the past 12 months, the rank order was as follows: 12th graders (21%), college students and 19- to 28-year-olds (both at 19%), 10th graders (14%), and finally 8th graders (8%). As can be seen by this divergence of trends for the different age groups, something other than a simple secular trend in drug use was taking place; specifically, important cohort differences were emerging.

- From the early 1990s until 1997, *marijuana* use rose sharply among secondary school students and their use of a number of *other illicit drugs* also rose, though more gradually. We have called this period a “relapse phase” in the longer-term epidemic. An increase in marijuana use also occurred among American college students, largely reflecting “generational replacement,” wherein earlier graduating high school class cohorts were replaced in the college population by more recent ones who were more drug-experienced before they left high school—in other words, as the result of a cohort effect. A resurgence in illicit drug use spreading *up* the age spectrum is a reversal of the way the epidemic spread several decades earlier. In the 1960s the epidemic began on the nation’s college campuses, and then the behavior diffused downward in age to high school students and eventually to junior high school students. This time the increases began in middle schools and radiated up the age spectrum. The graduating class cohorts in the middle and late 1990s were bringing with them the pattern of heavier drug use that emerged among them in the early 1990s while they were in secondary school.

The increases in use of *marijuana* and of *other illicit drugs* treated as a class were substantially larger, in both proportional and absolute terms, in the three secondary school grades than in either the college or young adult populations. In fact, at present there still is only a modest, albeit continuing, increase in illicit drug use in the young adult population of 19- to 28-year-olds. From 1991 through 1997, their annual prevalence of use of *any illicit drug* held remarkably stable at the same time that adolescent use rose appreciably. As we have said in the past, we believe that as generational replacement continues to occur we will likely see some increase in use of illicit drugs by the young adults. As would be expected, that happened sooner and more sharply among the college students, whose annual prevalence of marijuana use peaked a year later than among 12th graders and whose 30-day prevalence peaked two years later. Their use of *any illicit drug other than marijuana* continued to rise through 2004, whereas use by 12th graders peaked in 1997 at 21%. (It is still at 21% in 2004.) Indeed, the rates on this index are also still rising among 19- to 28-year-olds, even though it has been level for some years among high school seniors and has actually been declining among younger students.

Again, these diverging trends across the different age groups clearly show that changes during the 1990s reflected some important cohort effects—lasting differences between class cohorts—rather than broad secular trends, which would appear simultaneously in all of the age groups covered by the study. During all of the previous 18 years of the study, the use of most drugs moved in parallel across most age groups, indicating that secular change was prevailing.

- A somewhat parallel finding occurred for *cigarette* smoking, in that college students showed a sharp increase in smoking, beginning in 1995, which no doubt reflected a generational replacement effect as the heavier-smoking cohorts of adolescents entered college. This has been a more typical pattern of change for cigarettes, however, since differences in cigarette smoking rates among class cohorts tend to remain through the life course and also tend to account for much of the overall change in use observed at any given age.

In the early 1990s, cigarette smoking among 8th and 10th graders rose by about 50%—a particularly sharp and worrisome rise to which this study called widespread attention. Smoking also had been rising among high school seniors since 1992. The increase in current smoking ended among 8th and 10th graders in 1996, among 12th graders in 1997, and among college students in 1999. The appreciable decline in the smoking rate that began among the 8th graders now appears to be radiating up the age spectrum as they get older. (Their 30-day prevalence rate has fallen from 21% in 1996 to 9% in 2004.) Among the young adult stratum there has been little evidence yet of a decline in current smoking. The rate is still about the same as it was in 1995 (29%); but with time we expect their current smoking also will drop as the cohort effect works its way up the age bands. In fact, the current smoking rate among young adults already has slipped by a percentage point or two in the past few years, and smoking among college students has been falling for several years.

- ***Marijuana*** use, which had been rising sharply in all three grades of secondary school during the early to mid-1990s, began to turn downward in 1997 among 8th graders and then did the same in 1998 among 10th and 12th graders. Its decline has been rather modest, however, perhaps due in part to effects of the public debates over medical use of marijuana. The 8th and 12th graders showed a continuation of this decline in 2000; but in 2001, use remained level in all three grades. Since 2001, all three grades have shown significant declines in their annual prevalence of marijuana use, with the proportional decline greatest among the 8th graders.

Earlier in the 1990s, the annual prevalence of marijuana use (i.e., the percent reporting any use during the prior 12 months) tripled among 8th graders (from 6% in 1991 to 18% in 1996), more than doubled among 10th graders (from 15% in 1992 to 35% in 1997), and grew by three-quarters among 12th graders (from 22% in 1992 to 39% in 1997). Among college students, however, the increase in marijuana use, presumably largely due to a “generational replacement effect,” was much more gradual. Annual prevalence rose by about one-third from 27% in 1991 to 36% in 1998, before beginning to level. Among young adults there so far has been even less change, from 24% in 1991 to 29% in 2001, with no decline yet, though we expect one to begin soon.

Daily marijuana use rose substantially among secondary school and college students between 1992 and 2000 but somewhat less so among young adults (see Table 2-4). Daily use began a slow decline after 1999 at 8th grade, after 2001 at 10th grade, and after 2003 at 12th grade, consistent with a cohort-effect pattern. It appears that the college student and young adult rates at least leveled in 2004, and may have begun to reverse. However, none of the one-year changes in 2004 reached statistical significance. Prevalence of daily marijuana use has been highest among these five groups since the mid-1990s and was slow to decline even though annual and 30-day prevalence had been dropping. Still, the rate for seniors, for example, at 5.6% in 2004, is only about half the 10.7% peak figure reached in 1978, at the height of the illicit drug epidemic.

The amount of risk associated with using marijuana fell during the earlier period of increased use in the late 1970s, and perceptions of risk fell again during the more recent

resurgence of use in the 1990s. Indeed, at 12th grade, perceived risk began to decline a year *before* use began to rise in the upturn of the 1990s, making perceived risk a leading indicator of change in use. (The same may have happened in 8th grade, as well, but we do not have data starting early enough to check that possibility.) The decline in perceived risk halted after 1997 in 8th and 10th grade, and annual prevalence began to decline a year or two later. Again, perceived risk was a leading indicator of change in use, as it has proven to be for a number of drugs. As is discussed in Volume I, chapter 8, on attitudes and beliefs toward drugs, these attitudes themselves show evidence of cohort effects over the past decade and a half.

Personal disapproval of marijuana use slipped considerably among 8th graders between 1991 and 1996 and among 10th and 12th graders between 1992 and 1997. For example, the proportions of 8th, 10th, and 12th graders who said they disapproved of trying marijuana once or twice fell by 17, 21, and 19 percentage points, respectively, over those intervals of decline. There has since been some increase in disapproval among 8th and 10th graders and, beginning more recently, among 12th graders. Over the past two years, both perceived risk and disapproval of marijuana use have risen in all three grades, suggesting that the recent downturn in use can be expected to continue.

- Among seniors, the proportions using *any illicit drug other than marijuana* in the past year rose from a low of 15% in 1992 to 21% in 1997 (and is at 21% in 2004). (This recent peak was substantially below the 34% peak rate reached more than two decades earlier, in 1981.) In fact, all of the younger groups showed significant increases (though not as large in proportional terms as for marijuana). Use of any illicit drug other than marijuana began to increase in 1992 among 8th graders, in 1993 among 10th and 12th graders, and in 1995 among college students—again reflecting evidence of a cohort effect. Use peaked in 1996 among 8th and 10th graders and by 1997 among 12th graders, but it has yet to peak among the college students and young adults. The 8th graders have shown some gradual decline in their use of the other illicit drugs, treated as a class, since 1996; the decline among 10th graders paused after 1998 and did not resume until after 2001. Twelfth graders also showed some decline after 2001, but the decline appears to have halted in 2004. However, among college students and young adults, the proportions using any illicit drug other than marijuana are still rising, likely due to a continuing cohort effect.
- Between 1989 and 1992 we noted an increase among high school seniors, college students, and young adults in their use of *LSD*, a drug quite popular in the late 1960s and early 1970s. In 1992 the newly added populations (8th and 10th graders) were also showing an increase in LSD use; and for several more years, modest increases persisted in all five populations. Use of LSD among college students and young adults was the first to peak, in 1995. Use in all three grades of secondary school peaked a year later. After those peak years in the mid-1990s, there was a gradual decline in LSD use across the board, followed by a sharp decline after 2001 in all five populations. Overall, the pattern of change for LSD seems more consistent with secular change, rather than a cohort effect. The different age groups moved in parallel for the most part, likely in response to

historical events in the environment, including a sharp reduction in LSD availability after 2001.

Prior to the significant increase in LSD use among seniors in 1993, there was a significant 4.3-percentage-point decline between 1991 and 1992 in the proportion seeing great risk associated with trying LSD. (Once again this belief proved a leading indicator of change in use.) The decline in perceived risk continued through 1997 and halted in 1998. The proportion of seniors disapproving of LSD use also began to decline in 1992 and continued through 1996.

Because LSD was one of the earliest drugs to be popularly used in the overall American drug epidemic, there is a distinct possibility that young people—particularly the youngest cohorts, like the 8th graders—were not as concerned about the risks of use. They had less opportunity to learn vicariously about the consequences of use by observing others around them or to learn from intense media coverage of the issue, which occurred some years earlier. We were concerned that this type of “generational forgetting” of the dangers of a drug, which occurs as a result of generational replacement, could set the stage for a whole new epidemic of use. In fact, perceived harmfulness of LSD began to decline after 1991 among seniors. The measures for risk and disapproval were not introduced for 8th and 10th graders until 1993; both measures dropped until 1997 or 1998, after which perceived risk and disapproval leveled and then declined some. In 2004, 12th graders’ personal disapproval of trying LSD increased significantly. Because the decline in use in the last few years generally has *not* been accompanied by expected changes in these attitudes and beliefs, we suspected that some displacement by another drug might have been taking place, at least through 2001. The most logical candidate is *ecstasy*, which, like LSD, is used for its hallucinogenic effects; ecstasy was popular in the club scene and was very much on the rise through 2001. After 2001 a sharp decline in the reported availability of LSD, observed in all five populations, very likely played a role in the sharp decline in use among all of them.

- Questions about the use of *ecstasy* (*MDMA*) have been included in the follow-up surveys of college students and young adults since 1989; however, because of our concern about stimulating interest in an attractive-sounding and little-known drug, these questions were not added to the secondary school surveys until 1996. From 1989 to 1994, the annual prevalence rates tended to be quite low in the older age groups for whom we had data, but in 1995 there was a substantial increase (from 0.5% to 2.4% among college students, and from 0.7% to 1.6% among young adults generally).

When data were first gathered on secondary school students in 1996, the 10th and 12th graders showed higher rates of annual use (both 4.6%) than the college students (2.8%). Ecstasy use then fell steadily at all three grades of secondary school between 1996 and 1998, though it did not fall in the older age groups. Between 1998 and 2001, use rose sharply in all five populations. In fact, annual prevalence more than doubled in that three-year period among 12th graders, college students, and young adults and nearly doubled in the lower grades. In 2000 even the 8th graders showed a significant increase in use. Among young adults, the increase in use occurred primarily among those under age 29.

In 2002, use declined for all five age groups, but only the 10th graders' change was significant. Use decreased again in 2003 for all five populations, and only the drop among college students was not significant. There was some decline among all age groups in 2004; the largest declines were among the college students and young adults, suggesting that cohort effects and a secular trend may have been at work. Once again, among the secondary students this decline in use was predicted by an increase in perceived risk in 2001—an increase that continued through 2004. The rates of annual prevalence in 2004 for ecstasy were 1.7%, 2.4%, and 4.0% among 8th, 10th, and 12th graders, respectively; 2.2% among college students; and 3.5% among all young adults. The annual prevalence fell by half in 2004 alone among the college students, and all five groups are at rates that range from one-half to three-quarters lower than their recent peaks in 2001.

All five populations have been moving fairly synchronously since 1999, which appears to reflect an important secular trend and suggests that events in the social environment are reaching everyone. We believe that relevant events include increasing media coverage of people suffering adverse outcomes due to their ecstasy use, increasing dissemination of the scientific evidence on effects produced by the National Institute on Drug Abuse, and an anti-ecstasy media campaign by the Partnership for a Drug-Free America and the Office of National Drug Control Policy, initiated in 2002.

The quite dramatic increase in reported availability of ecstasy through 2001 was substantiated by law enforcement data on ecstasy seizures. Of the 12th graders surveyed in 1991, only 22% thought they could get ecstasy fairly easily, but a decade later (in 2001) 62% thought that they could. Since 2001, however, ecstasy availability has been decreasing in all three grades, possibly due in part to the steep decline in the number of users, who serve as supply points for others. See Figure 8-6 in Volume I, chapter 8, for a graphic presentation of the trends in use, availability, and perceived risk.

- In the decade between 1982 and 1992, annual prevalence rates for *amphetamine* use (other than that ordered by a physician) among seniors fell by nearly two-thirds, from 20.3% to 7.1%. Rates among college students fell even more over the same interval, from 21.1% to 3.6%. During the relapse phase in the overall drug epidemic, annual use increased by about half among 8th and 10th graders between 1991 and 1996, and there were increases among 12th graders and college students between 1992 and 1996. After 1996 the age groups diverged, with amphetamine use declining gradually among 8th and 10th graders and continuing to rise among 12th graders, college students, and young adults. The decline in the lower two grades continued through 2004, while the rise among the 12th graders and college students finally halted by 2003. The 12th graders have since shown some decline in their annual prevalence of amphetamine use, while the college students have held steady and the young adults appear still to be rising in their usage rate. This is another drug for which a cohort effect seems to best explain the pattern of change across the several age groups.

The increase in non-medical use of amphetamines (and a concurrent decrease in disapproval) that began among seniors in 1993 followed a sharp drop in perceived risk a

year earlier (which, as we have said, often serves as a leading indicator). Following a period of decline, perceived risk among seniors generally drifted up from 1995 through 2003; and disapproval also drifted up from 1996 through 2004. However, amphetamine use among seniors held relatively steady, with just a slight rise from 1997 through 2002 and with a small decline since then.

- Use of the specific amphetamine **Ritalin** showed a distinct increase around 1997—with annual prevalence among 12th graders going from 0.1% in 1992 to 2.8% in 1997—and then stayed level for a few years. (See Volume I, Appendix E, Table E-2.³) Because of its increasing importance, a differently structured question was introduced for Ritalin use in 2001. This new question, which we prefer to the original, does not use a prior branching question and produced somewhat higher prevalence rates. Results from the new question show Ritalin use to have been steady or declining slightly in recent years in all five populations (see Table 2-2).
- **Methamphetamine** questions were introduced in 1999 because of rising concern about its use; but a modest decline in use actually was observed among all five populations through 2002, with the exception of young adults, whose use has held steady. All groups except the young adults now have lower rates of annual prevalence in 2004 than they did in 1999. The only one-year change in 2004 that reached significance was for 8th graders, whose annual prevalence dropped from 2.5% in 2003 to 1.5% in 2004.
- We have had questions for a longer time—since 1990—about the use of **ice** (a crystallized form of methamphetamine that can be smoked, much like crack). The use of ice increased between the early and the late 1990s among 12th graders, college students, and young adults (the only populations asked about their use of ice). The estimates are a bit “bouncy” due to the relatively small sample sizes asked about this drug, but it appears that ice use has held fairly steady since 1998 among 12th graders and college students, while it may have risen some among young adults generally.
- **Inhalants** are defined as fumes or gases that are inhaled to get high, and they include common household substances such as glues, aerosols, butane, and solvents. In the early 1990s, there was a troublesome increase in inhalant use among secondary school students, followed by a reversal after 1995. One class of inhalants, **amyl and butyl nitrites**, became somewhat popular in the late 1970s, but their use has been almost eliminated. For example, the annual prevalence rate among 12th-grade students was 6.5% in 1979 but only 0.8% in 2004.
- **Crack cocaine** use spread rapidly from the early to the mid-1980s. Still, among high school seniors, the overall prevalence of crack leveled in 1987 at a relatively low prevalence rate (3.9% annual prevalence), even though crack use had continued to spread to new communities. Clearly, it had quickly attained a reputation as a dangerous drug, and by the time of our first measurement of perceived risk in 1987, it was seen as the

³As is discussed in Appendix E of Volume I, the absolute prevalence rates for Ritalin probably were higher than these statistics indicate, but the trend story likely is quite accurate. See Table 2-2 for more accurate estimates of the absolute prevalence rates in recent years; these estimates are based on a new question that does not require the respondent to first indicate some amphetamine use before asking about his or her Ritalin use.

most dangerous of all of the drugs. Annual prevalence dropped sharply in the next few years, reaching 1.5% by 1991, where it remained through 1993. Perceived risk began what turned out to be a long and substantial decline after 1990. Use began to rise gradually after 1993, when it was 1.5%, to 2.7% by 1999, before finally declining in 2000 and then leveling.

Among 8th and 10th graders, crack use had risen gradually in the 1990s: from 0.7% in 1991 to 2.1% by 1998 among 8th graders, and from 0.9% in 1992 to 2.5% in 1998 among 10th graders. Use among 12th graders peaked a year later, in 1999, at 2.7% and among young adults at 1.4%. Since those peak years, crack use has gradually declined in the lower two grades, held fairly steady in 12th grade and among young adults, and possibly increased some among college students. In general, the prevalence rates for this drug are relatively low—between 1.3% and 2.3% in all five groups. Among high school seniors, the group with the highest prevalence rate, annual crack prevalence among the college-bound is considerably lower than among those not bound for college (1.0% for college-bound versus 4.9% for noncollege-bound, in 2004).

We believe that the particularly intense and early media coverage of the hazards of crack cocaine likely had the effect of “capping” an epidemic early by deterring many would-be users and by motivating many experimenters to desist use. As has been mentioned, when we first measured crack use in 1987, it had the highest level of perceived risk of any illicit drug. Also, it did not turn out to be “instantly addicting” upon first-time use, as had been reported widely. While 3.9% of seniors in 2004 reported ever having tried crack, only 1.0% reported use in the past month, indicating that about three-quarters of those who tried crack did not establish a pattern of continued use.

In 1993 the levels of perceived risk and disapproval associated with crack dropped in all three grade levels, foretelling the rise in use that occurred in all three grades between 1994 and 1998. Because more than a decade had passed since the 1986 media frenzy about crack use, it is quite possible that “generational forgetting” of the risks of that drug was occurring. Indeed, perceived risk of crack use had been eroding steadily at all grade levels since 1991 (or 1992 in the case of the 12th graders) through 2000. Since then there has not been much systematic change in risk or disapproval of crack, though any change has been in an upward direction.

- Use of *cocaine*⁴ in general began to decline a year earlier than crack, probably because crack was still in the process of diffusing to new parts of the country, being still quite new. Between 1986 and 1987 the annual prevalence rate for cocaine dropped dramatically, by about one-fifth in all three populations then being studied—seniors, college students, and young adults. The decline occurred when young people finally began to view experimental and occasional use—the type of use in which they are most likely to engage—as more dangerous. This change first began to occur in 1987, probably partly because the hazards of cocaine use received extensive media coverage during the

⁴Unless otherwise specified, all references to “cocaine” refer to the use of cocaine in any form, including crack.

preceding year, but almost surely in part because of the highly publicized cocaine-related deaths in 1986 of sports stars Len Bias and Don Rogers. By 1992 the annual prevalence of cocaine use had fallen by about two-thirds among the three populations for which long-term data are available (12th graders, college students, and young adults).

During the 1990s, however, cocaine use in all five age populations increased some, both beginning and ending in a staggered pattern by age. Use rose among 8th graders from 1991 to 1998, among 10th and 12th graders from 1992 to 1999, among college students from 1994 to 2004, and among young adults from 1996 through 2004. (Note that a turnaround has yet to occur in the two older groups and they are currently at their highest level since 1990.) So, again, there is evidence of a cohort effect at work.

The story regarding attitudes and beliefs is informative. Having risen substantially after 1986, the perceived risk of using cocaine actually showed some (nonsignificant) decline in 1992 among seniors. In 1993, perceived risk for *cocaine other than crack* fell sharply in all grades and disapproval began to decline in all grades, though not as sharply as perceived risk. The decline in perceived risk had virtually ended by 1995 among 8th graders, by 1998 among 10th graders, and by 2001 among 12th graders, suggesting that there was a cohort effect at work in this important belief. Disapproval declined between 1991 and 1996 among 8th graders, before leveling; it also declined in 1992 through 1998 among 10th and 12th graders, with the exception of an increase for 12th graders in 1995. These changes foretold a subsequent leveling of use at each grade level.

Through 1989, there was no decline in perceived availability of cocaine among 12th graders; in fact, it rose steadily from 1983 to 1989, suggesting that availability played no role in bringing about the substantial downturn in use after 1986. After 1989, however, perceived availability fell some among seniors; that decline may be explained by the greatly reduced proportions of seniors who said they have any friends who use, because friendship circles are an important part of the supply system. From 1992 through 1998 or 1999, there was rather little change in reports of availability of powder cocaine in the three grades, but since 1999 there has been some continuing falloff. Perceived availability of powder cocaine has not changed much since 2001 in the upper grades but has been declining among the 8th graders.

As with all the illicit drugs, lifetime cocaine prevalence climbs with age; in 2004 it reached 38% among those aged 45. Unlike all of the other illicit drugs, active use of cocaine—i.e., annual prevalence or monthly prevalence—holds fairly steady after high school (and, until recent years, increases in use after high school) rather than declining. (See Figure 4-5 in Volume II.) Nearly all of the other illicit drugs show a decline in active use with age.

- **PCP** use fell sharply among high school seniors between 1979 and 1982, from an annual prevalence of 7.0% to 2.2%. It reached a low point of 1.2% in 1988, rose some in the 1990s to 2.6% in 1996, and declined to 0.7% by 2004. For the young adults, the annual prevalence rate rose very slightly from 0.2% in 1996 to 0.6% in 2001 before declining to 0.1% in 2004.

- Looking at the long-term trends, we see that the annual prevalence of *heroin* use among 12th graders fell by half between 1975 (1.0%) and 1979 (0.5%). It then stabilized for 15 years, through 1994. Heroin use was also stable in the early 1990s among the other four populations covered here. Then, in 1994 in the case of the 8th graders, and in 1995 in the case of all other groups, there was a sudden uptick in use, with rates doubling or tripling in one or two years. The new higher levels of heroin use remained among all five populations for the rest of the decade. In 2000, however, there was a significant decrease in use among 8th graders (from 1.4% in 1999 to 1.1% in 2000) and a significant increase in use among seniors (from 1.1% in 1999 to 1.5% in 2000). The increase among seniors was due entirely to an increase in non-injection use. Use of heroin declined significantly among 10th and 12th graders in 2001, as did their use of heroin without a needle. In 2002 little change took place among the secondary school students, but young adults showed a significant decline in their reported heroin use. A significant decline in use of heroin overall, as well as use of heroin without a needle, occurred among 10th graders in 2003. In sum, all age groups have rates of heroin use in 2004 that are below their recent peaks.

Two factors very likely contributed to the upturn in heroin use in the 1990s. One is a long-term decline in the perceived risk of harm, probably due to “generational forgetting,” because it had been a long time since the country had experienced a heroin epidemic. The second factor, not unrelated to the first, is that in the 1990s the greatly increased purity of heroin allowed it to be used by means other than injection. This may have lowered an important psychological barrier for some potential users by making heroin use less aversive and by making it seem less addictive as well as safer, because avoiding injection reduces the likelihood of transmission of HIV, hepatitis, or other serious blood-borne diseases. By introducing some new questions on heroin use in 1995, we were able to show that significant proportions of past-year users in all five populations were indeed taking heroin by means other than injection. (See Table 2-2 and chapter 4 of Volume I for details.)

The risk perceived to be associated with heroin fell for more than a decade after the study began, with 60% of the 1975 seniors seeing a great risk of trying heroin once or twice and only 46% of the 1986 seniors saying the same. Between 1986 and 1991, perceived risk rose some, from 46% to 55%, undoubtedly reflecting the newly recognized threat of HIV infection associated with heroin injection. After 1991, however, perceived risk fell again (to 51% by 1995), this time perhaps reflecting the fact that the newer heroin available on the street could be administered by methods other than injection because it was so much purer. Between 1996 and 1998, perceived risk among seniors rose—perhaps as the result of an anti-heroin campaign launched by the Partnership for a Drug-Free America in June 1996, as well as the visibility of heroin-related deaths of some celebrities in the entertainment and fashion design worlds. The perceived risk of trying heroin decreased among seniors in 1999, however, foretelling a significant increase in their use of the drug in 2000. In 2001, as the perceived risk of trying heroin increased slightly, seniors’ use declined significantly. Since 2001, perceived risk and use have remained fairly level.

Questions about the degree of risk perceived to be associated with heroin use were first introduced into the questionnaires for 8th and 10th graders in 1995. The questions asked

specifically and only about use “without using a needle” because we thought this was the form of heroin use of greatest concern at that point. (Similar questions were asked of 12th graders, as well, in one of the six questionnaire forms.) In general, perceived risk in all three grades for use without a needle rose in 1996 and 1997, before leveling.

- The use of *narcotics other than heroin* is reported only for high school seniors and older populations because we believe that younger students are not accurately discriminating among the drugs that should be included or excluded from this general class. Use had been declining gradually over most of the first half of the study in these groups. Seniors had an annual prevalence rate in 1977 of 6.4%, which fell to 3.3% by 1992. But from about 1992 through 2001, all of the older age groups showed a continuing increase, reaching peak levels of use in 2003 or 2004. The specific drugs in this class are listed in Table E-4 in Appendix E of Volume I, which shows that *Vicodin*, *codeine*, *Percocet*, and *OxyContin* are among the drugs most commonly mentioned by high school seniors in recent years. They also account for much of the increase in the general class, though there has also been an increase in the reported use of other substances in the class.
- In 2002, free-standing questions were included for two drugs in this class—*Vicodin* and *OxyContin*—and the observed prevalence rates suggest that they very likely help to account for the upturn in the use of the general class of narcotics other than heroin. We found that in 2003 Vicodin had attained surprisingly high prevalence rates in the five populations under study here—an annual prevalence of 2.8% in 8th grade, 7.2% in 10th grade, 10.5% in 12th grade, 7.5% among college students, and 8.6% among young adults. Lower rates were found for OxyContin, but considering that it is a highly addictive narcotic drug, the rates are not inconsequential—1.7%, 3.6%, 4.5%, 2.2%, and 2.6% annual prevalence in the same five populations, respectively. Both of these narcotic drugs also had shown some (not statistically significant) increases in all five populations in 2003. In 2004, use of OxyContin leveled among 8th and 10th graders but continued to increase some among 12th graders, college students, and young adults. Use of Vicodin leveled or decreased in 2004 among all populations under study except the young adults, among whom it increased slightly. Because OxyContin has received considerable adverse publicity in recent years, it is possible that perceived risk (which we do not measure) will increase. But because its use appears to have originated in several fairly delimited geographic areas, it seems likely that OxyContin will be diffusing to new communities for some time to come; this could delay its turnaround overall, as seems to have happened earlier for crack and ecstasy.
- A long, substantial decline, which began after 1977, occurred for *tranquilizer* use among high school seniors. By 1992, annual prevalence reached 2.8%, down from 11% in 1977. After 1992, use increased significantly (as has been true with most of the drugs), reaching 5.8% in 1999. Use of this drug continued to rise through 2002 to 7.7% (although because the question was revised slightly in 2001 to include Xanax as an example of a tranquilizer, part of the increase may be artifactual). In 2003 there was a significant decline in use among seniors, to 6.7%. In 2004 the rate stood at 7.3%. Reported tranquilizer use also exhibited some modest increase among 8th graders, from 1.8% in 1991 to 3.3% in 1996, before declining a bit to 2.6% in 1998 with little change since

then. As with a number of other drugs, the downturn in use began considerably earlier among the 8th graders than among their older counterparts. Among 10th graders, annual prevalence remained stable between 1991 and 1994, at around 3.3%, and increased significantly to 5.6% in 2000. Their use began to decline after 2001, reaching 5.1% in 2004. After a period of stability, college student use also showed an increase between 1994 and 2003, more than tripling their rate of use. For the young adult sample, after a long period of decline, annual prevalence increased appreciably between 1997 and 2004, more than doubling. Most of the reported tranquilizer use in recent years has involved *Valium* and *Xanax*. (See Table E-3 in Appendix E of Volume I.)

- The long-term gradual decline in *sedative (barbiturate)* use, which began at least as early as 1975, when the study began, halted in 1992. Use among 12th graders then rose to 6.7% in 2002—still well below the peak rate of 10.7% in 1975. Their use finally leveled in 2003. The 2004 annual prevalence of this class of drugs is lower among young adults (4.4%) and college students (4.2%) than among seniors (6.5%). Use among college students began to rise a few years later than it did among 12th graders, no doubt reflecting the impact of generational replacement in the form of a cohort effect. (Data are not included here for 8th and 10th grades, again because we believe that the younger students have more problems with proper classification of the relevant drugs.) Among young adults, use increased since the early 1990s, rising from 1.6% in 1992 to 4.4% in 2004.
- *Methaqualone*, another sedative drug, has shown a trend pattern quite different from barbiturates. Methaqualone use rose among seniors from 1975 to 1981, when annual prevalence reached 7.6%. Its use then fell very sharply, declining to 0.2% by 1993, before rising significantly during the general drug resurgence in the 1990s, to 1.1% by 1996. Use then leveled before decreasing significantly to 0.3% in 2000, but it is now up a bit to 0.8% in 2004. Use also fell among all young adults and among college students, who had annual prevalence rates of only 0.3% and 0.2%, respectively, by 1989—the last year they were asked about this drug. In the late 1980s, shrinking availability may well have played a role in this drop, as legal manufacture and distribution of the drug ceased. Because of its very low usage rates, only the 12th graders are now asked about use of this drug.
- It should be noted that during much of the 1990s and into the 2000s we were seeing a virtually uninterrupted increase among high school seniors, college students, and young adults generally in the use of nearly all illicit drugs that are central nervous system depressants. These include *sedatives (barbiturates)*, *tranquilizers*, and *narcotics other than heroin*. All of these drugs tended to fall from favor from the mid-1970s through the early 1990s, but many made a comeback after the early 1990s. The resurgence now seems to be leveling off.
- To summarize, for some years five classes of illicitly used drugs—*marijuana*, *amphetamines*, *cocaine*, *LSD*, and *inhalants*—had an impact on appreciable proportions of young Americans in their late teens and 20s. In 2004, high school seniors showed

annual prevalence rates of 34.3%, 10.0%, 5.3%, 2.2%, and 4.2%, respectively. Among college students in 2004, the comparable annual prevalence rates are 33.3%, 7.0%, 6.6%, 1.2%, and 2.7%; and for all young adults the rates are 29.2%, 6.2%, 7.1%, 0.9%, and 1.7%. Because **LSD** use has fallen so precipitously since 2001 in all five populations, it really no longer ranks as one of the major drugs of abuse, whereas **narcotics other than heroin** have become quite important due to their long-term rise that began in the 1990s. The narcotics other than heroin now have annual prevalence rates of 9.5% at 12th grade, 8.2% among college students, and 9.0% among young adults. **Tranquilizers** also have become more important due to a similar rise in use, with prevalence rates of 7.3%, 6.7%, and 7.4%, respectively, in the same three populations.

- Joining this set of long-established, more prevalent drugs was **MDMA (ecstasy)** for a period of time. However, it has shown a considerable drop in use and in 2004 has annual prevalence rates of 1.7%, 2.4%, 4.0%, 2.2%, and 3.5% among 8th, 10th, and 12 graders and college students and young adults, respectively.
- In 8th grade, **inhalants** rank second only to marijuana among the illicitly used drugs in terms of annual prevalence, and they actually rank first on lifetime use. Because of their importance among the younger adolescents, a new index of illicit drug use including inhalants was introduced in Tables 2-1 through 2-3. The use of inhalants reflects a form of illicit, psychoactive drug use; its inclusion makes relatively little difference in the illicit drug index prevalence rates for the older age groups but considerable difference for the younger ones. For example, in 2004 the proportion of 8th graders reporting any illicit drug use in their lifetime, exclusive of inhalants, was 22%, whereas including inhalants raised the figure to 30%.
- Several additional classes of drugs have been added to the study's coverage in recent years, and they are all discussed in chapter 4 of Volume I. These include **ketamine**, **GHB**, and **Rohypnol**, so-called "club drugs" (in addition to LSD and ecstasy). In general, these drugs have not attained high prevalence rates among 8th, 10th, or 12th graders: the 2004 annual prevalence rates for **ketamine** are 0.9%, 1.3%, and 1.9%, respectively; for **GHB**, 0.7%, 0.8%, and 2.0%; and for **Rohypnol**, 0.6% and 0.7% for 8th and 10th grade (the Rohypnol question for 12th graders was changed in 2002 and in 2004 stands at 1.6%). There was little change in the use of any of them this year except for significant declines in 10th graders' use of GHB and ketamine. The two narcotic drugs added to our coverage in 2002—**OxyContin** and **Vicodin**—show higher prevalence rates, as stated earlier.
- Two new substances used primarily by males to develop their physique and physical strength were added to the question set in 2001. One is **androstenedione**, a precursor to anabolic steroid, which could be purchased over the counter until early 2005. Among males, where use is heavily concentrated, the 2004 annual prevalence rate is relatively high, at 1.2%, 1.6%, and 3.7% in grades 8, 10, and 12. (Among females, the rates are 0.5%, 0.5%, and 0.6%.) As is discussed in chapter 10 of Volume I, the proportion of young males who report past-year use of either **androstenedione** or **steroids** is appreciable. The peak rate was observed in 2001, when the "andro" question was first introduced; 1 in every 12 or 13 boys (8.0%) in 12th grade indicated using one or both of

these drugs in the prior year. The rate has fallen some in all three grades since then, and in 2004 it was 5.3% among 12th-grade boys.

- Another physique-enhancing substance that is not a drug, but rather a type of protein supplement, is *creatine*. Because we thought its use often was combined with the use of steroids and androstenedione, we included a question on it in 2001 and found prevalence of use to be very high. Among boys, who again are the primary users, the 2004 annual prevalence for creatine is 3.3%, 9.8%, and 15.9%, in grades 8, 10, and 12. In other words, one in every six 12th-grade boys had used creatine in the prior year. (For girls, the rates are 0.6%, 0.9%, and 1.0%.)
- For some years, the study has contained a set of questions about the use of nonprescription stimulants, including stay-awake pills, diet pills, and the so-called “look-alikes.” The annual prevalence among 12th graders of over-the-counter *stay-awake pills*, which usually contain caffeine as their active ingredient, nearly doubled between 1982 and 1990, increasing from 12% to 23%. After 1990 this statistic fell, reaching 12% by 2004. Earlier decreases also occurred among the college-aged young adult population (ages 19 to 22), in which annual prevalence was 26% in 1989 and declined to 12% in 2003—its lowest level since 1986. The *look-alikes* also have shown some falloff in recent years. Among high school seniors, annual prevalence decreased slightly from 6.8% in 1995 to 5.0% in 1999, increased to 7.1% in 2001, and then decreased to 5.0% by 2004; among young adults aged 19 to 22, use also declined from 6.0% in 1995 to 1.9% in 2004. Among high schools seniors, annual prevalence of over-the-counter *diet pills* declined between 1986 and 1995, from 15% to 10%, increased to 15% by 2002, then declined to 11% in 2004. (Among 12th-grade girls in 2004, 22% had tried diet pills by the end of senior year, 14% used them in the past year, and 7% used them in just the past 30 days.) Among young adults aged 19 to 22 there also had been an earlier decline from 1986 to 1995, with annual prevalence moving from 17% to 7%. Use then rose to 15% by 2004. Use of these over-the-counter drugs by high school seniors is covered in chapter 10 of Volume I.

College-Noncollege Differences in Illicit Drug Use

- For analytic purposes, American college students are defined here as those respondents one to four years past high school who were actively enrolled full-time in a two- or four-year college. For most categories of drugs, college students have rates of use that are *below* those of their age peers, including *any illicit drug other than marijuana*, *hallucinogens*, *LSD* specifically, *ecstasy*, *cocaine*, *crack cocaine* specifically, *narcotics other than heroin*, *OxyContin*, and *Vicodin* specifically, *amphetamines*, *methamphetamine*, *ice*, *sedatives (barbiturates)*, and *tranquilizers*. College students show *higher* rates of use than their age mates for a few drugs—*Ritalin*, *Rohypnol*, *GHB*, *ketamine*, and *inhalants*. Their high rate of Ritalin use may derive from its use for staying awake while studying. For a few categories of drugs—including *any illicit drug*, *marijuana*, and *heroin*—they also show annual usage rates that are about *average* for all high school graduates their age. College students are about average on an index of any

illicit drug use likely because they have very similar rates of use of marijuana, which largely drives the index.

- Although college-bound seniors have below-average rates of use on *all* of the illicit drugs while they are in high school, these students' eventual use of some illicit drugs attain parity with, or even come to exceed, the rates of those who do not attend college. As results from the study published in two recent books have shown, this college effect of "catching up" is largely explainable in terms of differential rates of leaving the parental home after high school graduation and of getting married. College students are more likely than their age peers to have left the parental home and its constraining influences and less likely to have entered marriage, with its constraining influences.⁵
- In general, the trends since 1980 in illicit substance use among American college students have paralleled those of their age peers not in college. Most drugs showed a period of substantial decline in use sometime after 1980. Further, all young adult high school graduates through age 28, as well as college students taken separately, showed trends highly parallel for the most part to the trends among high school seniors until about 1992. After 1992 a number of drugs showed an increase in use among seniors (as well as 8th and 10th graders) but not among college students and young adults for some period of time.

This divergence, combined with the fact that the upturn began first among the 8th graders (in 1992), suggests that cohort effects were emerging for illicit drug use, as we discussed earlier. In fact, as those heavier-using cohorts of high school seniors entered the college years, we saw a lagged increase in the use of several drugs in college. For example, annual prevalence reached a low point among 12th graders in 1992 for a number of drugs (e.g., *cocaine*, *amphetamines*, *sedatives [barbiturates]*, *tranquilizers*, *other narcotics*, and *any illicit drug other than marijuana*) before rising thereafter; among college students, those same drugs reached a low two years later in 1994, and then began to rise gradually. Then, in 1998, as *marijuana* use was declining in the three grades of secondary school, we saw a sharp increase in use among college students. The evidence for cohort effects resulting from generational replacement is impressive, as well as consistent with our earlier predictions.

Male-Female Differences in Illicit Drug Use

- Regarding gender differences in the three older populations (high school seniors, college students, and young adults), males are more likely to use *most illicit drugs*, and the differences tend to be largest at the higher frequency levels. *Daily marijuana use* among high school seniors in 2004, for example, is reported by 7.7% of males versus 3.1% of females; among all adults (aged 19 to 30 years) by 6.3% of males versus 3.3% of females; and among college students, specifically, by 6.8% of males versus 3.1% of females.

⁵Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates.

- In the 8th- and 10th-grade samples there are fewer and smaller gender differences in the use of drugs—perhaps because girls tend to date and then emulate older boys, who are in age groups considerably more likely to use drugs. While the rate of using *marijuana* in the past year is slightly higher for males, the rate for the use of *any illicit drug other than marijuana* tends to be slightly higher for females. There is little male-female difference in 8th and 10th grades in the use of *LSD*, *ecstasy*, *cocaine*, *crack*, *heroin*, *Ritalin*, *Rohypnol*, and *GHB*. *Inhalant*, *amphetamine*, and *tranquilizer* use are slightly higher among females.

TRENDS IN ALCOHOL USE

- Several findings about *alcohol* use in these age groups are noteworthy. First, despite the fact that it is illegal for virtually all secondary school students and most college students to purchase alcoholic beverages, experience with alcohol is widespread among them. Alcohol has been tried by 44% of 8th graders, 64% of 10th graders, 77% of 12th graders, and 85% of college students; and active use is also widespread. Most important, perhaps, is the occurrence of *occasions of heavy drinking*—measured by the percent reporting five or more drinks in a row at least once in the prior two-week period. Among 8th graders this statistic stands at 11%, among 10th graders at 22%, among 12th graders at 29%, and among college students at 42%. After people pass their early 20s, this behavior recedes somewhat with age, reflected by the 37% rate found in the entire young adult sample and the 27% rate found specifically among 29- to 30-year-olds.
- Alcohol use did not increase as use of other illicit drugs decreased among seniors from the late 1970s to the early 1990s, although it was common to hear such a “displacement hypothesis” asserted. This study demonstrates that the opposite seems to be true. After 1980, when illicit drug use was declining, the *monthly prevalence of alcohol use* among seniors also declined gradually, but substantially, from 72% in 1980 to 51% in 1992. *Daily alcohol use* declined by half over the same interval, from a peak of 6.9% in 1979 to 3.4% in 1992; and the prevalence of drinking *five or more drinks in a row* during the prior two-week interval fell from 41% in 1983 to 28% in 1993—nearly a one-third decline. When illicit drug use rose again in the 1990s, there was evidence that alcohol use (particularly binge drinking) was rising some as well—albeit not nearly as sharply as did marijuana use. In the late 1990s, as illicit drug use leveled in secondary schools and began a gradual decline, similar trends were observed for alcohol. So, the evidence suggests that alcohol moves much more in concert with illicit drugs, rather than in the opposite direction.

College-Noncollege Differences in Alcohol Use

- The data from college students show a quite different pattern of change in relation to alcohol use than that of 12th graders or noncollege respondents of the same age. (See Figure 9-14 in Volume II.) From 1980 to 1993, college students showed considerably less drop-off in monthly prevalence of *alcohol* use (82% to 70%) than did high school seniors (72% to 51%). *Occasions of heavy drinking* also declined less among college students from 1980 to 1993, from 44% to 40%, compared to a decline from 41% to 28% among high school seniors. Among their noncollege age-mates, the decline was from 41% to 34%. Thus, because both their noncollege age-mates and high school students were showing greater declines, the college students stood out as having maintained a high rate of binge (or party) drinking. Since 1993, this behavior has changed little among college students (42% in 2004, slightly above the 40% rate observed in 1993), while their noncollege age-mates also increased by two percentage points, to 34%; high school seniors' binge drinking increased to 32% in 1998 but then decreased to 29% by 2004. Still, college students continue to stand out as having a relatively high rate of binge drinking.

Because the college-bound seniors in high school are consistently less likely to report occasions of heavy drinking than the noncollege-bound students, the higher rates of such drinking in college indicate that they catch up to and pass their peers in binge drinking after high school graduation. We have shown that this differential change after high school is largely attributable to the fact that college students are more likely to leave the parental home in the four years after graduating from high school than their noncollege peers, and also less likely to get married.⁶ A recent publication from the study also shows that membership in a fraternity or sorority tends to increase heavy episodic drinking and marijuana use.⁷

- Since 1980, college students have generally had *daily drinking* rates that were slightly lower than their age peers, suggesting that they were more likely to confine their drinking to weekends, when they tend to drink a lot. College men have much higher rates of daily drinking than college women (7.2% versus 1.8% in 2004). This gender difference also exists in the noncollege group (9.0% versus 3.1% in 2004).
- Comparisons between college students and their noncollege peers, in terms of binge drinking, have typically shown that college students are more likely to engage in this activity.
- The rate of daily drinking fell considerably among the noncollege group, from 8.3% in 1980 to 3.2% in 1994, but by 2000 had risen to 5.8%, where it stands in 2004. Daily

⁶See Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁷McCabe, S. E., Schulenberg, J. E., Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Kloska, D. D. (2005). Selection and socialization effects of fraternities and sororities on U.S. college student substance use: A multi-cohort national longitudinal study. *Addiction, 100*, 512-524.

drinking by the college group also dropped in approximately the same time period, from 6.5% in 1980 to 3.0% in 1995, and then increased to 4.5% in 1997. It is at 3.7% in 2004.

Male-Female Differences in Alcohol Use

- Given that the physiological impacts of five drinks are considerably greater for the typical young female versus the typical young male, it is not surprising that we find substantial gender differences in the prevalence of *occasions of heavy drinking*. Among high school seniors the rates in 2004 are 24% for females versus 34% for males; this difference generally has been diminishing very gradually since the study began. (In 1975 there was a 23-percentage-point difference between them, versus a 10-point difference in 2004.)
- Among college students and young adults generally, there also are substantial gender differences in alcohol use, with college males drinking the most. For example, 49% of college males report having *five or more drinks in a row* over the previous two weeks versus 38% of college females. There has not been a great deal of change in this gender difference since 1980.

TRENDS IN CIGARETTE SMOKING

Quite a number of very important findings about *cigarette smoking* among American adolescents and young adults have emerged during the life of the study, and one of the study's more important contributions to the long-term health of the nation's people has been to document these trends. Despite the demonstrated health risks associated with smoking, young people continued to establish regular cigarette habits during late adolescence in sizeable and, during the first half of the 1990s, growing proportions. In fact, since the study began in 1975, cigarettes have consistently remained the class of abusable substances most frequently used on a daily basis by high school students.

- During most of the 1980s, when smoking rates were falling steadily among adults, we reported that smoking among adolescents was not declining. Then the situation went from bad to worse. Among 8th and 10th graders, the *current (past 30-day) smoking rate* increased by about half between 1991 (when their use was first measured) and 1996; and among 12th graders, the current smoking rate rose by nearly one-third between 1992 and 1997. This study played an important role in bringing these disturbing increases in adolescent smoking to public attention during those years.

Fortunately, there have been some important declines in current smoking since 1996 in the case of 8th and 10th graders and since 1997 in the case of 12th graders. In fact, the declines have more than offset the increases observed earlier in the 1990s. In 2004, 9% of 8th graders (down from 14% in 1991 and 21% in 1996) reported smoking one or more cigarettes in the prior 30 days—a decline of more than one-half from the recent peak. Some 16% of 10th graders were current smokers in 2004 (down from 21% in 1991 and 30% in 1996), representing a 47% drop from the recent peak rate. And in 2004, 25% of

12th graders were current smokers (versus 28% in 1991 and 37% in 1997). This represents a one-third drop from the recent peak. The decline in the smoking rate among 12th graders halted in 2004, even though the decline continued in the lower grades, albeit at a much decelerated rate. Despite these very important recent improvements, at present one-quarter of American young people are current smokers by the time they complete high school; and other research consistently shows that smoking rates are substantially higher among those who drop out before graduating. Perhaps most important at present is that the improvement appears to be drawing to an end.

Among college students the peak rate in current smoking was not reached until 1999 (31%), but since then there has been a fair decline (to 23% in 2003 and 24% in 2004). The young adults 19 to 28 years old have shown little change in rates of current smoking since 1996 (when their rate was 30%); it is 29% in 2004. Note that none of the three older groups—high school seniors, college students, and young adults—showed any decline in their smoking rates from 2003 to 2004. All three showed nonsignificant increases.

- **Daily smoking** rates also increased by about half among 8th graders (from a low of 7.0% in 1992 to 10.4% in 1996) and 10th graders (from a low of 12.3% in 1992 to 18.3% in 1996), while daily smoking among 12th graders increased by 43% (from a low of 17.2% in 1992 to 24.6% in 1997).⁸ In 1997 we saw the first evidence of a change in the situation, as daily smoking rates declined among 8th graders and leveled among 10th graders. There was a significant decline in 10th and 12th graders' daily smoking rates by 1998. All three grades have been continuing to decline in use through 2004, with significant declines among all three grades in 2004. Among college students daily smoking increased by nearly half from 1994 (13%) through 1999 (19%)—reflecting the cohort replacement effect of the heavier-smoking senior classes—before a turnaround began in 2000, decreasing the levels of use to 14% by 2003, where it remained in 2004.
- The dangers perceived to be associated with **pack-a-day smoking** differ greatly by grade level and seem to be unrealistically low at all grade levels. Currently, nearly three-quarters of the seniors (74%) report that pack-a-day smokers run a great risk of harming themselves physically or in other ways, but only 62% of the 8th graders say the same. All three grades showed a decrease in perceived risk between 1993 and 1995, as use was rising rapidly, but a slightly larger and offsetting increase between 1995 and 2000, presaging the subsequent downturn in smoking. Between 2000 and 2003, perceived risk remained relatively level in all grades. In 2004, perceived risk increased significantly among 8th and 10th graders and showed a nonsignificant increase among 12th graders. This is a hopeful sign for the possibility of further declines in the smoking rates among young people, in light of the deceleration of the declines between 2002 and 2004.
- Disapproval of cigarette smoking was in decline longer: from 1991 through 1996 among 8th and 10th graders, and from 1992 to 1996 among 12th graders. Since then there has been a fairly steady increase in disapproval of cigarette smoking in all three grades. Undoubtedly the heavy media coverage of the tobacco issue (the proposed settlement

⁸For high school seniors, during a much earlier period (from 1977 to 1981), there had been a substantial decline in daily smoking, a leveling for nearly a decade (through 1990), and a slight decline in 1991 and 1992.

with the state attorneys general, the congressional debate, the eventual state settlements, etc.) had an important influence on these attitudes and beliefs. However, that coverage diminished considerably in 1998, raising the question of whether these changes in youth attitudes would continue. It may well be, of course, that the removal of certain kinds of cigarette advertising and promotion, combined with national- and state-level anti-smoking campaigns and more recent significant increases in cigarette prices, have served to sustain and prolong these changes. This study has shown that there have been important changes in reported recall of anti-smoking ads as a result of both state and national campaigns.⁹

Age and Cohort-Related Differences in Cigarette Smoking

- Initiation of smoking occurs most often in grades 6 through 9 (i.e., at modal ages 11-12 to 14-15), although according to the 2004 8th graders, 11% had already initiated smoking below grade 6. The initiation rate trails off considerably by 12th grade, although a number of the light smokers in 12th grade make the transition to heavy smoking in the first two years after high school. Analyses presented in this volume and elsewhere have shown that cigarette smoking evidences a clear “cohort effect.” That is, if a class (or birth) cohort establishes an unusually high rate of smoking at an early age relative to other cohorts, the rate is likely to remain high throughout the life cycle relative to that of other birth cohorts at equivalent ages.
- As we reported in the “Other Findings From the Study” chapter in the 1986 volume in this series, some 53% of the half-pack-a-day (or more) smokers in senior year said that they had tried to quit smoking and found they could not. Of those who had been daily smokers in 12th grade, nearly three-quarters still were daily smokers seven to nine years later (based on the 1985 follow-up survey), despite the fact that in high school only 5% of them thought they would “definitely” be smoking five years hence. A more recent analysis, based on the 1995 follow-up survey, showed similar results. Nearly two-thirds (63%) of those who had been daily smokers in the 12th grade were still daily smokers seven to nine years later, although in high school only 3% of them had thought they would “definitely” be smoking five years hence. Clearly, the smoking habit is established at an early age, it is difficult to break for those young people who have it, and young people greatly overrate their own ability to quit. Additional data from the 8th- and 10th-grade students show us that younger children are even more likely than older ones to underestimate seriously the dangers of smoking.
- The surveys of 8th and 10th graders also show that cigarettes are almost universally available to teens. Nearly two-thirds (60%) of 8th graders and four-fifths (81%) of 10th graders say that cigarettes are “fairly easy” or “very easy” for them to get, if they want them. Until 1997 there had been little change in reported availability since these questions were first asked in 1992. Over the last eight years, however, perceived availability of cigarettes decreased significantly for 8th and 10th graders, quite likely

⁹Johnston, L. D., Terry-McElrath, Y. M., O'Malley, P. M., & Wakefield, M. (2005). Trends in recall and appraisal of anti-smoking advertising among American youth: National survey results, 1997-2001. *Prevention Science*, 6, 1-19.

reflecting the impact of new regulations and related enforcement efforts aimed at reducing the sale of cigarettes to children. (Twelfth graders are not asked this question.)¹⁰

College-Noncollege Differences in Cigarette Smoking

- A striking difference in smoking rates has long existed between college-bound and noncollege-bound high school seniors. For example, in 2004, smoking a half-pack or more per day is about three times as prevalent among the noncollege-bound seniors (16.0% versus 5.6%). Among respondents of college age (one to four years past high school), those not in college show the same dramatically higher rate of smoking than those who are in college, with half-pack-a-day smoking standing at 19.2% versus 6.8%, respectively. Clearly, the differences precede college attendance.
- In the first half of the 1990s, smoking rose some among college students and their same-age peers, although the increases were not as steep for either group as they were among high school seniors. But in 1998 and 1999, while smoking was declining among secondary school students at all grades, smoking increased significantly for college students, no doubt reflecting the cohort effect from earlier, heavier-smoking classes of high school seniors moving into the older age groups. Between 1991 and 1999, the 30-day prevalence of cigarette smoking by college students rose from 23% to 31%, or by about one-third, and daily smoking rose from 14% to 19%, also by about one-third. The year 2000 showed, for the first time in several years, a decline in college student smoking, one which continued with a significant decline, to 23%, in 2003, though it did not appear to continue into 2004 (24%). (Because of the smaller numbers of cases in the college student samples, the trend lines are not always as smooth as they are for most of the other groups discussed here.) Some decline has been observed among their noncollege-aged peers, but only since 2001.

Male-Female Differences in Cigarette Smoking

- In the 1970s, high school senior females caught up to and passed senior males in rates of *current smoking*. Both genders then showed a decline in use followed by a long, fairly level period, with use by females consistently higher, but with the gender difference diminishing. In the early 1990s, there was another crossover—rates rose among males and declined among females. Both genders in 12th grade showed increasing use between 1992 and 1997 and then a decline in use through 2003. In 2004, current smoking among males continued to decline (to 25% 30-day prevalence); use among the females bumped up 2 percentage points to 24% (following a 3 percentage-point drop the year before), but their smoking rate still remained slightly below that for males.

Among college students, females had slightly higher probabilities of being daily smokers from 1980 through 1994—although this long-standing gender difference was not true among their age peers not in college. However, there was a crossover from 1994 through 2001, with males being higher—an echo of the crossover among seniors in 1991. Since

¹⁰For a more detailed examination of recent changes in youth access to cigarettes, see Johnston, L. D., O'Malley, P. M., & Terry-McElrath, Y. M. (2004). Methods, locations, and ease of cigarette access for American youth, 1997-2002. *American Journal of Preventive Medicine*, 27, 267-276.

about 2001 there has been little consistent gender difference in smoking among college students.

RACIAL/ETHNIC COMPARISONS

The three largest ethnic groups in the population—Whites, African Americans, and Hispanics taken as a group—are examined here for 8th, 10th, and 12th graders. (Sample size limitations simply do not allow finer subgroup breakdowns unless many years are combined. Separate publications from the study have done just that.) A number of interesting findings emerge from the comparison of these three groups, and the reader is referred to chapters 4 and 5 of Volume I for a full discussion of them and to Appendix D for a tabular documentation of them across all drugs.¹¹ The trends for these three subgroups are also presented graphically in an occasional paper available on-line.¹²

- African American seniors have consistently shown lower usage rates of most drugs, licit and illicit, than White seniors; this also is true at the lower grade levels where few have yet dropped out of school. The differences are quite large for some drugs at all three grade levels: these include *inhalants*, *LSD*, *hallucinogens other than LSD*, *ecstasy*, *powder cocaine*, *amphetamines*, and *tranquilizers*.
- African American students have a much lower 30-day prevalence rate of *cigarette smoking* than White students (10% versus 28% in senior year, in 2004) because their smoking rate continued to decline after 1983, while the rate for White students stabilized for some years. (Smoking rates had been rising among White seniors and African American seniors after 1992, but by 1998 there was a leveling, and since then a reversal, in both groups in all grades.) The White students showed a continuing decline in 2004 for all three grades, while the decline among African American students halted for all three grades.
- In 12th grade, *occasions of heavy drinking* are much less likely to be reported by African American students (11%) than by White students (33%) or Hispanic students (26%).

¹¹Periodically we publish comparisons that contain a number of the smaller racial/ethnic groups in the population, based on data combined for a number of contiguous years in order to attain adequate sample sizes. The first was Bachman, J. G., Wallace, J. M. Jr., O'Malley, P. M., Johnston, L. D., Kurth, C. L., & Neighbors, H. W. (1991). Racial/ethnic differences in smoking, drinking, and illicit drug use among American high school seniors, 1976-1989. *American Journal of Public Health*, 81, 372-377. More recent articles are: Wallace, J. M., Jr., Bachman J. G., O'Malley, P. M., Johnston, L. D., Schulenberg, J. E., & Cooper, S. M. (2002). Tobacco, alcohol and illicit drug use: Racial and ethnic differences among U.S. high school seniors, 1976-2000. *Public Health Reports*, 117 (Supplement 1), S67-S75; Wallace, J. M., Jr., Bachman J. G., O'Malley, P. M., Schulenberg, J. E., Cooper, S. M., & Johnston, L. D. (2003). Gender and ethnic differences in smoking, drinking, and illicit drug use among American 8th, 10th, and 12th grade students, 1976-2000. *Addictions*, 98, 225-234; and Delva, J., Wallace, J. M., Jr., O'Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2005). The epidemiology of alcohol, marijuana, and cocaine use among Mexican American, Puerto Rican, Cuban American, and other Latin American 8th-grade students in the United States: 1991-2002. *American Journal of Public Health*, 95, 696-702.

¹²Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2005). *Demographic subgroup trends for various licit and illicit drugs, 1975-2004*. (Monitoring the Future Occasional Paper No. 61) [On-line]. Ann Arbor, MI: Institute for Social Research. c. 411 pp. Available: www.monitoringthefuture.org.

- In 12th grade, of the three racial/ethnic groups, Whites tend to have the highest rates of use on a number of drugs, including *marijuana*, *inhalants*, *hallucinogens*, *LSD* specifically, *hallucinogens other than LSD*, *ecstasy*, *narcotics other than heroin*, *amphetamines*, *sedatives (barbiturates)*, *tranquilizers*, *alcohol*, *getting drunk*, *cigarettes*, and *smokeless tobacco*.
- However, Hispanics have the highest usage rate in 12th grade for a number of the most dangerous drugs, for example, *heroin with a needle*, *crack*, and *crystal methamphetamine (ice)*. Further, in 8th grade, Hispanics have the highest rates not only for these drugs, but for many of the others, as well. For example, in 8th grade, the annual prevalence of *marijuana* for Hispanics is 17%, versus 12% for Whites and 13% for African Americans; the two-week prevalence of *binge drinking* is 16% for Hispanics, 11% for Whites, and 10% for African Americans. In other words, Hispanics have the highest rates of use for many drugs in 8th grade, but not in 12th, which suggests that their considerably higher dropout rate (compared to Whites and African Americans) may change their relative ranking by 12th grade.
- With regard to trends, seniors in all three racial/ethnic groups exhibited a decline in *cocaine* use from 1986 through 1992, although the decline was less steep among African American seniors because their earlier increase in use was not as large as the increase among White and Hispanic students.
- For virtually *all of the illicit drugs*, the three groups have tended to trend in parallel. Because White seniors had achieved the highest level of use on a number of drugs—including *amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—they also had the largest declines; African Americans have had the lowest rates and, therefore, the smallest declines.
- The important racial/ethnic differences in *cigarette smoking* noted earlier among high school seniors have emerged during the life of the study. The three groups were fairly similar in their smoking rates during the mid-1970s, and all three mirrored the general decline in smoking from 1977 through 1981. Over the 11-year interval from 1981 through 1992, however, smoking rates declined very little, if at all, for Whites and Hispanics; but the rates for African Americans continued to decline steadily. As a result, by 1992 the daily smoking rate of African Americans was one-fifth that of Whites. Subsequently, all three ethnic groups of 12th graders exhibited fairly parallel trends in smoking.

DRUG USE IN EIGHTH GRADE

It may be useful to focus specifically on the youngest age group in the study—the 8th graders, most of whom are 13 or 14 years old—because the exceptional levels of both licit and illicit drug use that they already have attained help illustrate the nation’s urgent need to continue to address the substance abuse problems among its young.

- By 8th grade in 2004, 44% of youngsters report having tried *alcohol* (more than just a few sips), and one in five (20%) say they have already been *drunk* at least once.
- More than a quarter of the 8th graders (28%) have tried *cigarettes*, and one in eleven (9.2%) say they have smoked in the prior month. Shocking to most adults is the fact that only 62% of 8th graders recognize that there is great risk associated with being a pack-a-day smoker. While an increasing proportion will recognize the risk by 12th grade, for many this is too late, because by then they will have developed a hard-to-break smoking habit.
- *Smokeless tobacco* has been tried by 15% of male 8th graders, is used currently by 6.4% of them, and is used daily by 1.7%. (Rates are much higher among males than among females.)
- Among 8th graders, one in six (17%) have used *inhalants*, and 1 in 24 (4.5%) say they have used them in just the past month. This is the only class of drugs for which use is substantially higher in 8th grade than in 10th or 12th grade.
- *Marijuana* has been tried by nearly one in every six 8th graders (16%) and has been used in the prior month by almost 1 in every 16 (6.4%).
- A surprisingly large number of 8th graders (7.5%) say they have tried prescription-type *amphetamines* without medical instruction; 2.3% say they have used them in the prior 30 days.
- Relatively few 8th graders say they have tried most of the *other illicit drugs*. (This is consistent with the retrospective reports from seniors concerning the grades in which they first used the various drugs.) But the proportions having at least some experience with them is not inconsequential because a 3.3% prevalence rate, for example, on average represents one child in every 30-student classroom. The 2004 8th-grade proportions reporting any lifetime experience with the other illicit drugs are *tranquilizers* (4.0%), *hallucinogens other than LSD* (3.0%), *ecstasy* (2.8%), *cocaine other than crack* (2.6%), *methamphetamine* (2.5%), *crack* (2.4%), *steroids* (1.9% overall, and 2.3% among males), *LSD* (1.8%), *heroin* (1.6%), and *Rohypnol* (1.0%).
- In total, 12% of all 8th graders in 2004—one in every eight—have tried *some illicit drug other than marijuana* (excluding inhalants). Put another way, in an average 30-student classroom of 8th graders, about three or four students have used some drug other than marijuana and nearly six have used marijuana.
- The very large number of students who have already begun use of the so-called “gateway drugs” (*tobacco, alcohol, inhalants, and marijuana*) suggests that a substantial number of today’s 8th-grade students are already at risk of proceeding further to such drugs as LSD, cocaine, amphetamines, and heroin.

DRUG USE BY AGE 45

Because we have now followed up graduating high school seniors into their 40s, we can characterize the drug-using history of today's 45-year-olds. This is important not only because it characterizes how use by these respondents has developed over more than two decades since they left high school, but also because many of them are now themselves the parents of adolescents. Their active use of substances may serve as role modeling for their children, and their own past experience may complicate their communications with their children regarding drugs. The level of use they have attained is truly impressive. (See chapter 4 of Volume II for greater detail and discussion.)

- Among 45-year-old high school graduates in 2004, we estimate that about four-fifths (81%) have tried *marijuana* and that over two-thirds (71%) have tried an *illicit drug other than marijuana* (estimates are adjusted to correct for panel attrition, as described in chapter 4 of Volume II).

Their current behavior is far less extreme than those statistics would imply, however. “Only” one in eight (12%) indicates using marijuana in the last 12 months, while 1 in 11 (9%) affirms use of any other illicit drug in that time period. (Their past-month prevalence rates are lower still—6.5% and 3.7%, respectively.) About 1 in 50 45-year-olds (2.0%) is a *current daily marijuana* user, though a great many more have been so at some time in the past.

- Quite high proportions of the 45-year-old respondents have had some experience during their lifetime with several of the specific illicit drugs other than marijuana. These include *amphetamines* (51%), *cocaine* in any form (44%), *non-crack forms of cocaine* (37%), *tranquilizers* (38%), *hallucinogens* of any type (33%), *narcotics other than heroin* (28%), *sedatives (barbiturates)* (29%), *LSD* specifically (20%), and *other hallucinogens* (19%). In sum, today's adults in their mid-40s are a very drug-experienced segment of the population, as might be expected from the fact that they graduated from high school near the peak of the drug epidemic. To repeat, 81% have tried marijuana and 71% have tried some illicit drug other than marijuana.
- Among the illicit drugs other than marijuana that have been used in just the past year by this age group (outside of medical regimen) are *cocaine* (3.9% annual prevalence), *narcotics other than heroin* (3.4%), and *tranquilizers* and *non-crack forms of cocaine* (both at 3.1%). There is very little active use being reported by our respondents at this age of *LSD*, *other hallucinogens*, *amphetamines*, *crack*, or *heroin*. (Of course, we would not expect *heavy* heroin or crack users to have remained in the panel studies.)
- *Alcohol* consumption is relatively high at this age, with 66% indicating that they consumed at least one alcoholic drink in the prior 30 days, 9.0% indicating current *daily drinking* (defined as drinking on 20 or more occasions in the prior 30 days), and 19% indicating *occasional heavy drinking* (defined as five or more drinks on at least one occasion in the prior two weeks). The rate of occasional heavy drinking is much lower than was exhibited by this cohort when they were of high school and college ages.

- One in five (20%) 45-year-old high school graduates currently smokes *cigarettes*. Almost all of those are current *daily smokers*.

SUMMARY AND CONCLUSIONS

We can summarize the findings on trends as follows: over more than a decade—from the late 1970s to the early 1990s—there were very appreciable declines in use of a number of *illicit drugs* among 12th-grade students and even larger declines in their use among American college students and young adults. These substantial improvements—which seem largely explainable in terms of changes in attitudes about drug use, beliefs about the risks of drug use, and peer norms against drug use—have some extremely important policy implications. One is that these various substance-using behaviors among American young people are malleable—they *can* be changed. It has been done before. The second is that demand-side factors appear to have been pivotal in bringing about most of those changes. The reported levels of marijuana availability, as reported by high school seniors, have held fairly steady throughout the life of the study. (Moreover, both abstainers and quitters rank availability and price very low on their list of reasons for not using.) And, in fact, the perceived availability of cocaine actually was rising during the beginning of the sharp decline in cocaine and crack use, which occurred when the risks associated with that drug rose sharply. (See the last section of chapter 9, Volume I, for more examples and further discussion of this point.)

However, improvements surely are not inevitable; and, when they occur, they should not be taken for granted. Relapse is always possible and, indeed, just such a “relapse” in the longer-term epidemic occurred during the early to mid-1990s, as the country let down its guard on many fronts. (See chapter 8 of Volume I for a more detailed discussion of this point.)

In 1992, 8th graders exhibited a significant increase in annual use of *marijuana*, *cocaine*, *LSD*, and *hallucinogens other than LSD*, as well as an increase in *inhalant* use. (In fact, all five populations showed some increase in *LSD* use, continuing a longer-term trend for college students and young adults.) Further, the attitudes and beliefs of seniors regarding drug use began to soften.

In 1993, use of several drugs began to rise among 10th and 12th graders, as well, fulfilling our earlier predictions that we had made based on their eroding beliefs about the dangers of drugs and their attitudes about drug use. Increases occurred in a number of the so-called “gateway drugs”—*marijuana*, *cigarettes*, and *inhalants*—increases that we argued boded ill for the use of later drugs in the usual sequence of drug use involvement. Indeed, the proportion of students reporting the use of *any illicit drug other than marijuana* rose steadily after 1991 among 8th and 10th graders and after 1992 among 12th graders. (This proportion increased by more than half among 8th graders, with annual prevalence rising from 8.4% in 1991 to 13.1% in 1996.) The softening attitudes about *crack* and *other forms of cocaine* also provided a basis for concern—the use of both increased fairly steadily through 1998.

Over the years, this study has demonstrated that changes in perceived risk and disapproval have been important causes of change in the use of a number of drugs. These beliefs and attitudes

surely are influenced by the amount and nature of public attention paid to the drug issue in the historical period during which young people are growing up. A substantial decline in attention to this issue in the early 1990s very likely helps to explain why the increases in perceived risk and disapproval among students ceased and began to backslide. News coverage of the drug issue plummeted between 1989 and 1993 (although it made a considerable comeback as surveys—including this one—began to document that the problem was worsening again), and the media's *pro bono* placement of ads from the Partnership for a Drug-Free America also fell considerably. (During that period the 12th graders in this study showed a steady decline in their recalled exposure to such ads and in the judged impact of such ads on their own drug-taking behavior.)

Also, the deterioration in the drug abuse situation first began among our youngest cohorts—perhaps because they had not had the same opportunities for vicarious learning from the adverse drug experiences of people around them and people they learn about through the media. Clearly, there was a danger that, as the drug epidemic subsided in the 1980s and early 1990s, newer cohorts would have far less opportunity to learn through informal means about the dangers of drugs—that what we have called a “generational forgetting” of those risks would occur through a process of generational replacement of older, more drug-savvy cohorts with newer, more naive ones. If true, this suggests that as drug use subsides, as it did by the early 1990s, the nation must redouble its efforts to ensure that such naive cohorts learn these lessons about the dangers of drugs through more formal means—from schools, parents, and focused messages in the media, for example—and that this more formalized prevention effort be *institutionalized* so that it will endure for the long term. Clearly, for the foreseeable future, American young people will be aware of the psychoactive potential of a host of drugs and will continue to have access to them. That means that each new generation of young people must learn the reasons that they should *not* use drugs. Otherwise, their natural curiosity and desires for new experiences will lead a great many of them to use drugs.

Another lesson that derives from the epidemiological data in this study is that social influences that tend to reduce the *initiation* of substance use also have the potential to deter the *continuation* of use by those who have already begun to use—particularly if they are not yet deeply involved in use. Chapter 5 of Volume I shows how increased quitting rates have contributed importantly to downturns in the use of a number of drugs at different historical periods. The lesson for prevention is that primary prevention should not be the only goal of intervention programs; early-stage users are also susceptible to being influenced when their beliefs and attitudes regarding drugs are changed.

The following facts help to put into perspective the magnitude and variety of substance use problems that presently remain among American young people:

- By the end of 8th grade, nearly a third (30%) of American young people have tried an *illicit drug* (if inhalants are included as an illicit drug), and by 12th grade, more than half (53%) have done so.
- By their late 20s, 6 in every 10 (61%) of today's American young adults have tried an *illicit drug*, and a third (35%) have tried some *illicit drug other than marijuana* (usually in addition to marijuana). (These figures do not include inhalants.)

- Today more than one in seven Americans (16% in 2004) has tried *cocaine* by the age of 30, and 8% have tried it by their senior year of high school (i.e., by age 17 or 18). More than 1 in every 26 seniors (3.9%) has tried *crack*. In the young adult sample, 1 in 13 (7.5%) has tried crack by age 29-30.
- Over 1 in every 18 high school seniors (5.6%) in 2004 smokes *marijuana daily*, and this rate has shown very little decline so far. Among young adults aged 19 to 28, the percentage is slightly less (5.0%). Among those same seniors in 2004, one in every six (18%) has been a daily marijuana smoker at some time for at least a month, and among young adults the comparable figure is one in five (21%).
- Three in ten high school seniors (29%) consumed *five or more drinks in a row* at least once in the two weeks prior to the survey, and such behavior tends to increase among young adults one to four years past high school. About half (49%) of male college students report such behavior.
- A quarter (25%) of high school seniors in 2004 were current *cigarette* smokers, and 16% already were current daily smokers. In addition, we know from studying previous cohorts that many young adults increase their rates of smoking within a year or so after they leave high school.
- Despite the substantial improvement in this country's drug situation in the 1980s and the early 1990s, and then some further improvement beginning in the late 1990s, it is still true that this nation's secondary school students and young adults show a level of involvement with illicit drugs that is among the highest in the world's industrialized nations.¹³ Even by longer-term historical standards in this country, these rates remain extremely high, though in general they are not as high as in the peak years of the epidemic in the late 1970s. Heavy drinking also remains widespread and troublesome; and certainly the continuing initiation to cigarette smoking of a large, albeit declining, proportion of young people remains a matter of the greatest public health concern. The declines in youth smoking have decelerated sharply, indicating that improvements in youth smoking may soon come to a halt.
- Finally, we note the seemingly unending capacity of pharmacological experts and amateurs to discover new substances with abuse potential that can be used to alter mood and consciousness. There is also a great capacity for our young people to discover the abuse potential of existing products, such as Roflumilast, and to "rediscover" older drugs, such as *LSD* and *heroin*. While as a society we have made significant progress on a number of fronts in the fight against drug abuse, we must remain vigilant against the

¹³A published report from an international collaborative study, modeled largely after Monitoring the Future, provides comparative data from national school surveys of 15- to 16-year-olds that was completed in 2003 in 35 European countries. It also includes MTF data from 10th graders in the United States. See Hibell, B., Andersson, B., Bjarnasson, T., Ahlström, S., Balakireva, O., Kokkevi, A., & Morgan, M. (Eds.). (2004). *The ESPAD report 2003 (The European School Survey Project on Alcohol and Other Drugs): Alcohol and other drug use among students in 35 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs, and the Council of Europe.

opening of new fronts, as well as the reemergence of trouble on older ones, particularly after a period of generational forgetting.

In fact, one of the dynamics that keeps the drug epidemic rolling is the emergence of new drugs whose hazards are little known. In 1999 we saw this happen with the drug *ecstasy* (*MDMA*). Other drugs like *Rohypnol*, *ketamine*, *GHB*, and *OxyContin* have appeared in the past decade and now must be added to the list of drugs under study. The spread of such new drugs appears to be facilitated and hastened today by young people's widespread use of chat rooms and other sites on the Internet. We predict a continuous flow of such new substances onto the scene and believe that the task of rapidly documenting their emergence, establishing their adverse consequences, and quickly demystifying them will remain important means by which policymakers, researchers, and educators deal with the continuing threats posed by such drugs.

The drug problem is not an enemy that can be vanquished, as in a war. It is more a recurring and relapsing problem that must be contained to the extent possible on a long-term, ongoing basis. Therefore, it is a problem that requires an ongoing, dynamic response from our society—one that takes into account the continuing generational replacement of our children, the generational forgetting of the dangers of drugs that can occur with that replacement, and the perpetual additional tracking of new abusable substances that will come onto the scene and threaten to lure our young people into involvement with drugs.

TABLE 2-1
Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	<u>Lifetime</u>														'03-'04 change
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	
Any Illicit Drug^a															
8th Grade	18.7	20.6	22.5	25.7	28.5	31.2	29.4	29.0	28.3	26.8	26.8	24.5	22.8	21.5	-1.4
10th Grade	30.6	29.8	32.8	37.4	40.9	45.4	47.3	44.9	46.2	45.6	45.6	44.6	41.4	39.8	-1.6
12th Grade	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	+0.1
College Students	50.4	48.8	45.9	45.5	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	-1.7
Young Adults	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	+0.3
Any Illicit Drug Other Than Marijuana^{a,b}															
8th Grade	14.3	15.6	16.8	17.5	18.8	19.2	17.7	16.9	16.3	15.8†	17.0	13.7	13.6	12.2	-1.4
10th Grade	19.1	19.2	20.9	21.7	24.3	25.5	25.0	23.6	24.0	23.1†	23.6	22.1	19.7	18.8	-0.8
12th Grade	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0†	30.7	29.5	27.7	28.7	+1.0
College Students	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	+0.5
Young Adults	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3	31.6	32.8	33.9	35.2	+1.3
Any Illicit Drug Including Inhalants^{a,c}															
8th Grade	28.5	29.6	32.3	35.1	38.1	39.4	38.1	37.8	37.2	35.1	34.5	31.6	30.3	30.2	0.0
10th Grade	36.1	36.2	38.7	42.7	45.9	49.8	50.9	49.3	49.9	49.3	48.8	47.7	44.9	43.1	-1.8
12th Grade	47.6	44.4	46.6	49.1	51.5	53.5	56.3	56.1	56.3	57.0	56.0	54.6	52.8	53.0	+0.2
College Students	52.0	50.3	49.1	47.0	47.0	49.1	50.7	55.4	54.4	54.6	53.1	52.3	54.1	52.9	-1.3
Young Adults	63.4	61.2	61.2	58.5	59.0	58.2	58.4	58.5	58.5	59.5	59.0	59.6	60.6	62.5	+1.9
Marijuana/Hashish															
8th Grade	10.2	11.2	12.6	16.7	19.9	23.1	22.6	22.2	22.0	20.3	20.4	19.2	17.5	16.3	-1.2
10th Grade	23.4	21.4	24.4	30.4	34.1	39.8	42.3	39.6	40.9	40.3	40.1	38.7	36.4	35.1	-1.3
12th Grade	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	-0.4
College Students	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8	51.2	51.0	49.5	50.7	49.1	-1.6
Young Adults	58.6	56.4	55.9	53.7	53.6	53.4	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	+0.2
Inhalants^{c,d}															
8th Grade	17.6	17.4	19.4	19.9	21.6	21.2	21.0	20.5	19.7	17.9	17.1	15.2	15.8	17.3	+1.5 s
10th Grade	15.7	16.6	17.5	18.0	19.0	19.3	18.3	18.3	17.0	16.6	15.2	13.5	12.7	12.4	-0.3
12th Grade	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	-0.3
College Students	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4	12.9	9.6	7.7	9.7	8.5	-1.2
Young Adults	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	-0.6
Nitrites^e															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.6	1.5	1.4	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	1.4	1.2	1.3	1.0	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}															
8th Grade	3.2	3.8	3.9	4.3	5.2	5.9	5.4	4.9	4.8	4.6†	5.2	4.1	4.0	3.5	-0.5
10th Grade	6.1	6.4	6.8	8.1	9.3	10.5	10.5	9.8	9.7	8.9†	8.9	7.8	6.9	6.4	-0.5
12th Grade	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0†	14.7	12.0	10.6	9.7	-0.9
College Students	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8	14.4	14.8	13.6	14.5	12.0	-2.5 s
Young Adults	15.7	15.7	15.4	15.4	16.1	16.4	16.8	17.4	18.0	18.4	18.3	19.6	19.7	19.3	-0.4

(Table continued on next page)

TABLE 2-1 (cont'd)
Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Lifetime														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
LSD															
8th Grade	2.7	3.2	3.5	3.7	4.4	5.1	4.7	4.1	4.1	3.9	3.4	2.5	2.1	1.8	-0.4
10th Grade	5.6	5.8	6.2	7.2	8.4	9.4	9.5	8.5	8.5	7.6	6.3	5.0	3.5	2.8	-0.7
12th Grade	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	-1.3 s
College Students	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7	11.8	12.2	8.6	8.7	5.6	-3.1 ss
Young Adults	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	-1.2
Hallucinogens Other Than LSD^b															
8th Grade	1.4	1.7	1.7	2.2	2.5	3.0	2.6	2.5	2.4	2.3†	3.9	3.3	3.2	3.0	-0.3
10th Grade	2.2	2.5	2.8	3.8	3.9	4.7	4.8	5.0	4.7	4.8†	6.6	6.3	5.9	5.8	-0.1
12th Grade	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9†	10.4	9.2	9.0	8.7	-0.3
College Students	6.0	5.7	5.4	4.4	6.5	6.5	7.5	8.7	8.8	8.2	10.7	11.0	12.8	10.1	-2.7 s
Young Adults	8.4	8.0	7.6	7.4	7.8	7.9	8.5	9.4	9.3	9.9	12.0	15.0	16.4	15.6	-0.9
PCP^g															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	-0.9
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	-0.3
MDMA (Ecstasy)^h															
8th Grade	—	—	—	—	—	3.4	3.2	2.7	2.7	4.3	5.2	4.3	3.2	2.8	-0.4
10th Grade	—	—	—	—	—	5.6	5.7	5.1	6.0	7.3	8.0	6.6	5.4	4.3	-1.1 s
12th Grade	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	-0.8
College Students	2.0	2.9	2.3	2.1	3.1	4.3	4.7	6.8	8.4	13.1	14.7	12.7	12.9	10.2	-2.7
Young Adults	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	+0.8
Cocaine															
8th Grade	2.3	2.9	2.9	3.6	4.2	4.5	4.4	4.6	4.7	4.5	4.3	3.6	3.6	3.4	-0.2
10th Grade	4.1	3.3	3.6	4.3	5.0	6.5	7.1	7.2	7.7	6.9	5.7	6.1	5.1	5.4	+0.3
12th Grade	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	+0.5
College Students	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4	9.1	8.6	8.2	9.2	9.5	+0.3
Young Adults	21.0	19.5	16.9	15.2	13.7	12.9	12.1	12.3	12.8	12.7	13.1	13.5	14.7	15.2	+0.4
Crackⁱ															
8th Grade	1.3	1.6	1.7	2.4	2.7	2.9	2.7	3.2	3.1	3.1	3.0	2.5	2.5	2.4	-0.2
10th Grade	1.7	1.5	1.8	2.1	2.8	3.3	3.6	3.9	4.0	3.7	3.1	3.6	2.7	2.6	-0.1
12th Grade	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	+0.3
College Students	1.5	1.7	1.3	1.0	1.8	1.2	1.4	2.2	2.4	2.5	2.0	1.9	3.1	2.0	-1.1
Young Adults	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	-0.5
Other Cocaine^j															
8th Grade	2.0	2.4	2.4	3.0	3.4	3.8	3.5	3.7	3.8	3.5	3.3	2.8	2.7	2.6	-0.1
10th Grade	3.8	3.0	3.3	3.8	4.4	5.5	6.1	6.4	6.8	6.0	5.0	5.2	4.5	4.8	+0.3
12th Grade	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	+0.6
College Students	9.0	7.6	6.3	4.6	5.2	4.6	5.0	7.4	7.8	8.1	8.3	8.6	8.5	9.3	+0.7
Young Adults	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	+0.9

(Table continued on next page)

TABLE 2-1 (cont'd)
Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Lifetime														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Heroin ^k															
8th Grade	1.2	1.4	1.4	2.0	2.3	2.4	2.1	2.3	2.3	1.9	1.7	1.6	1.6	1.6	0.0
10th Grade	1.2	1.2	1.3	1.5	1.7	2.1	2.1	2.3	2.3	2.2	1.7	1.8	1.5	1.5	0.0
12th Grade	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	-0.1
College Students	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9	1.7	1.2	1.0	1.0	0.9	-0.1
Young Adults	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	0.0
With a Needle ^l															
8th Grade	—	—	—	—	1.5	1.6	1.3	1.4	1.6	1.1	1.2	1.0	1.0	1.1	+0.2
10th Grade	—	—	—	—	1.0	1.1	1.1	1.2	1.3	1.0	0.8	1.0	0.9	0.8	-0.1
12th Grade	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.0
College Students	—	—	—	—	0.4	0.1	0.2	0.5	0.8	0.7	0.2	0.3	0.1	0.1	0.0
Young Adults	—	—	—	—	0.4	0.4	0.3	0.4	0.6	0.4	0.6	0.4	0.5	0.4	-0.1
Without a Needle ^l															
8th Grade	—	—	—	—	1.5	1.6	1.4	1.5	1.4	1.3	1.1	1.0	1.1	1.0	-0.1
10th Grade	—	—	—	—	1.1	1.7	1.7	1.7	1.6	1.7	1.3	1.3	1.0	1.1	+0.1
12th Grade	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	-0.3
College Students	—	—	—	—	0.5	1.0	1.2	2.1	1.0	2.5	1.3	1.2	1.1	1.0	-0.1
Young Adults	—	—	—	—	0.9	1.3	1.5	1.7	1.9	2.1	2.1	1.8	2.2	2.1	-0.2
Other Narcotics ^{m,n}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9†	13.5	13.2	13.5	+0.3
College Students	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7	8.9	11.0†	12.2	14.2	13.8	-0.4
Young Adults	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5†	13.9	16.8	17.6	+0.8
Amphetamines ^m															
8th Grade	10.5	10.8	11.8	12.3	13.1	13.5	12.3	11.3	10.7	9.9	10.2	8.7	8.4	7.5	-0.9
10th Grade	13.2	13.1	14.9	15.1	17.4	17.7	17.0	16.0	15.7	15.7	16.0	14.9	13.1	11.9	-1.2
12th Grade	15.4	13.9	15.1	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	+0.6
College Students	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9	12.3	12.4	11.9	12.3	12.7	+0.4
Young Adults	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	+0.7
Methamphetamine ^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	4.5	4.2	4.4	3.5	3.9	2.5	-1.4 ss
10th Grade	—	—	—	—	—	—	—	—	7.3	6.9	6.4	6.1	5.2	5.3	0.0
12th Grade	—	—	—	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	0.0
College Students	—	—	—	—	—	—	—	—	7.1	5.1	5.3	5.0	5.8	5.2	-0.6
Young Adults	—	—	—	—	—	—	—	—	8.8	9.3	9.0	9.1	8.9	9.0	0.0
Ice ^p															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	+0.2
College Students	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8	1.3	2.3	2.0	2.9	2.2	-0.7
Young Adults	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	0.0

(Table continued on next page)

TABLE 2-1 (cont'd)
Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	<u>Lifetime</u>														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Sedatives (Barbiturates)^m															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8	9.9	+1.0
College Students	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7	6.9	6.0	5.9	5.7	7.2	+1.5
Young Adults	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	+0.9
Methaqualone^{m,q}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}															
8th Grade	3.8	4.1	4.4	4.6	4.5	5.3	4.8	4.6	4.4	4.4†	5.0	4.3	4.4	4.0	-0.4
10th Grade	5.8	5.9	5.7	5.4	6.0	7.1	7.3	7.8	7.9	8.0†	9.2	8.8	7.8	7.3	-0.5
12th Grade	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9†	10.3	11.4	10.2	10.6	+0.4
College Students	6.8	6.9	6.3	4.4	5.4	5.3	6.9	7.7	8.2	8.8	9.7	10.7	11.0	10.6	-0.4
Young Adults	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5	11.9	13.4	13.8	14.9	+1.1
Rohypnol^f															
8th Grade	—	—	—	—	—	1.5	1.1	1.4	1.3	1.0	1.1	0.8	1.0	1.0	0.0
10th Grade	—	—	—	—	—	1.5	1.7	2.0	1.8	1.3	1.5	1.3	1.0	1.2	+0.2
12th Grade	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol^s															
Any Use															
8th Grade	70.1	69.3‡	55.7	55.8	54.5	55.3	53.8	52.5	52.1	51.7	50.5	47.0	45.6	43.9	-1.6
10th Grade	83.8	82.3‡	71.6	71.1	70.5	71.8	72.0	69.8	70.6	71.4	70.1	66.9	66.0	64.2	-1.8
12th Grade	88.0	87.5‡	80.0	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	+0.2
College Students	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0	86.6	86.1	86.0	86.2	84.6	-1.7
Young Adults	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	0.0
Flavored Alcoholic Beverages^{g,o}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	37.9	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	58.6	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	71.0	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	79.0	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	83.2	—
Been Drunk^p															
8th Grade	26.7	26.8	26.4	25.9	25.3	26.8	25.2	24.8	24.8	25.1	23.4	21.3	20.3	19.9	-0.3
10th Grade	50.0	47.7	47.9	47.2	46.9	48.5	49.4	46.7	48.9	49.3	48.2	44.0	42.4	42.3	-0.1
12th Grade	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	+2.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page)

TABLE 2-1 (cont'd)
**Trends in Lifetime Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)**

	<u>Lifetime</u>														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Cigarettes															
Any Use															
8th Grade	44.0	45.2	45.3	46.1	46.4	49.2	47.3	45.7	44.1	40.5	36.6	31.4	28.4	27.9	-0.5
10th Grade	55.1	53.5	56.3	56.9	57.6	61.2	60.2	57.7	57.6	55.1	52.8	47.4	43.0	40.7	-2.3 s
12th Grade	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	-0.9
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smokeless Tobacco [†]															
8th Grade	22.2	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	-0.4
10th Grade	28.2	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	-0.8
12th Grade	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	-0.4
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^P															
8th Grade	1.9	1.7	1.6	2.0	2.0	1.8	1.8	2.3	2.7	3.0	2.8	2.5	2.5	1.9	-0.6 s
10th Grade	1.8	1.7	1.7	1.8	2.0	1.8	2.0	2.0	2.7	3.5	3.5	3.5	3.0	2.4	-0.6 s
12th Grade	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	+0.1

SOURCE: The Monitoring the Future Study, the University of Michigan.

Footnotes for Table 2-1 to Table 2-4

NOTES: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001.

'—' indicates data not available.

'*' indicates less than .05 percent but greater than 0 percent.

'‡' indicates some change in the question. See relevant footnote for that drug. See relevant figure to assess the impact of the wording changes.

Any apparent inconsistency between the change estimate and the prevalence of use estimates for the two most recent classes is due to rounding error.

Approximate

Weighted Ns	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
8th Graders	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000
10th Graders	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400
12th Graders	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600
College Students	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400
Young Adults	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700

^aFor 12th graders, college students, and young adults only: Use of "any illicit drug" includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin or any use of other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders. For 8th and 10th graders only: The use of other narcotics and barbiturates has been excluded because these younger respondents appear to overreport use (perhaps because they include the use of nonprescription drugs in their answers).

^bIn 2001 the question text was changed on half of the questionnaire forms for each age group. "Other psychedelics" was changed to "other hallucinogens," and "shrooms" was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. For 8th, 10th, and 12th graders only: The 2001 data presented here are based on the changed forms only; N is one-half of N indicated. In 2002 the remaining forms were changed to the new wording. The data are based on all forms beginning in 2002. Data for "any illicit drug other than marijuana" and "hallucinogens" are also affected by these changes and have been handled in a parallel manner.

^cFor 12th graders, college students, and young adults only: Data based on five of six forms in 1991–98; N is five-sixths of N indicated. Data based on three of six forms beginning in 1999; N is one-half of N indicated.

^dInhalants are unadjusted for underreporting of amyl and butyl nitrites.

^eFor 12th graders only: Data based on one of six forms; N is one-sixth of N indicated. For college students and young adults only: Data based on two of six forms; N is two-sixths of N indicated. Questions about nitrite use were dropped from the young adult questionnaires in 1995.

^fHallucinogens are unadjusted for underreporting of PCP.

^gFor 12th graders and young adults only: Data based on one of six forms; N is one-sixth of N indicated.

^hFor 8th and 10th graders only: Data based on one of two forms in 1996; N is one-half of N indicated. Data based on one-third of N indicated in 1997–2001 due to changes in the questionnaire forms. Data based on two of four forms beginning in 2002; N is one-half of N indicated. For 12th graders only: Data based on one of six forms in 1996–2001; N is one-sixth of N indicated. Data based on two of six forms beginning in 2002; N is two-sixths of N indicated. For college students and young adults only: Data based on two of six forms in 1991–2001; N is two-sixths of N indicated. Data based on three of six forms beginning in 2002; N is one-half of N indicated.

ⁱFor college students and young adults only: Data based on five of six forms beginning in 2002; N is five-sixths of N indicated.

^jFor 12th graders only: Data based on four of six forms; N is four-sixths of N indicated. For college students and young adults only: Data based on four of six forms; N is four-sixths of N indicated.

^kIn 1995, the heroin question was changed in one of two forms for 8th and 10th graders, in three of six forms for 12th graders, and in two of six forms for college students and young adults. Separate questions were asked for use with injection and without injection. In 1996, the heroin question was changed in all remaining 8th- and 10th-grade forms. Data presented here represent the combined data from all forms.

^lFor 8th and 10th graders only: Data based on one of two forms in 1995; N is one-half of N indicated. Data based on all forms beginning in 1996. For 12th graders only: Data based on three of six forms; N is one-half of N indicated. For college students and young adults only: Data based on two of six forms; N is two-sixths of N indicated.

^mOnly drug use not under a doctor's orders is included here.

ⁿIn 2002 the question text was changed in half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced with Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one-half of N indicated. In 2003, the remaining forms were changed to the new wording. The data are based on all forms beginning in 2003.

^oFor 8th and 10th graders only: Data based on one of four forms; N is one-third of N indicated.

^pFor 12th graders, college students, and young adults only: Data based on two of six forms; N is two-sixths of N indicated.

^qFor 12th graders only: Data based on one of six forms; N is one-sixth of N indicated.

^rFor 8th and 10th graders only: Data based on one of two forms in 1996; N is one-half of N indicated. Data based on three of four forms in 1997–98; N is two-thirds of N indicated. Data based on two of four forms in 1999–2001; N is one-third of N indicated. Data based on one of four forms beginning in 2002; N is one-sixth of N indicated. For 12th graders only: Data based on one of six forms; N is one-sixth of N indicated. Data for 2001 and 2002 are not comparable due to changes in the questionnaire forms. For college students and young adults only: Data based on two of six forms; N is two-sixths of N indicated.

^sFor 8th, 10th, and 12th graders only: In 1993, the question text was changed slightly in half of the forms to indicate that a “drink” meant “more than just a few sips.” The 1993 data are based on the changed forms only; N is one-half of N indicated for these groups. In 1994 the remaining forms were changed to the new wording. The data are based on all forms beginning in 1994. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. For college students and young adults: The revision of the question text resulted in rather little change in the reported prevalence of use. The data for all forms are used to provide the most reliable estimate of change.

^tFor 8th and 10th graders only: Data based on one of two forms for 1991–96 and on two of four forms beginning in 1997; N is one-half of N indicated. For 12th graders only: Data based on one of six forms; N is one-sixth of N indicated. For college students and young adults only: Questions about smokeless tobacco use were dropped from the analyses in 1989.

^uFor 12th graders only: Data based on two of six forms in 2000; N is two-sixths of N indicated. Data based on three of six forms in 2001; N is one-half of N indicated. Data based on one of six forms beginning in 2002; N is one-sixth of N indicated. For college students and young adults only: Data based on two of six forms; N is two-sixths of N indicated.

^vFor 12th graders only: Data based on two of six forms in 2000; N is two-sixths of N indicated. Data based on three of six forms beginning in 2001; N is one-half of N indicated. For college students and young adults only: Data based on two of six forms; N is two-sixths of N indicated.

^wFor 12th graders only: The 2003 flavored alcoholic beverage data were created by adjusting the 2004 data to reflect the observed 2003 to 2004 change in a slightly different version of the flavored alcoholic beverage question. In 2004 the original question was revised to include wine coolers among the examples—a change that had very little effect on the observed prevalence of use rate in 2004.

^xDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes and smokeless tobacco, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

TABLE 2-2
Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Annual														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Any Illicit Drug^a															
8th Grade	11.3	12.9	15.1	18.5	21.4	23.6	22.1	21.0	20.5	19.5	19.5	17.7	16.1	15.2	-0.9
10th Grade	21.4	20.4	24.7	30.0	33.3	37.5	38.5	35.0	35.9	36.4	37.2	34.8	32.0	31.1	-0.9
12th Grade	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	-0.5
College Students	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	-0.3
Young Adults	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	+0.6
Any Illicit Drug Other Than Marijuana^{a,b}															
8th Grade	8.4	9.3	10.4	11.3	12.6	13.1	11.8	11.0	10.5	10.2†	10.8	8.8	8.8	7.9	-0.8
10th Grade	12.2	12.3	13.9	15.2	17.5	18.4	18.2	16.6	16.7	16.7†	17.9	15.7	13.8	13.5	-0.2
12th Grade	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4†	21.6	20.9	19.8	20.5	+0.6
College Students	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6	16.4	16.6	17.9	18.6	+0.6
Young Adults	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	+0.7
Any Illicit Drug Including Inhalants^{a,c}															
8th Grade	16.7	18.2	21.1	24.2	27.1	28.7	27.2	26.2	25.3	24.0	23.9	21.4	20.4	20.2	-0.3
10th Grade	23.9	23.5	27.4	32.5	35.6	39.6	40.3	37.1	37.7	38.0	38.7	36.1	33.5	32.9	-0.6
12th Grade	31.2	28.8	32.5	37.6	40.2	41.9	43.3	42.4	42.8	42.5	42.6	42.1	40.5	39.1	-1.4
College Students	29.8	31.1	31.7	31.9	33.7	35.1	35.5	39.1	37.4	37.0	38.2	37.7	36.0	35.9	0.0
Young Adults	27.8	29.2	28.9	29.2	30.4	30.2	30.1	30.6	30.6	31.2	33.2	32.4	32.7	34.9	+2.2
Marijuana/Hashish															
8th Grade	6.2	7.2	9.2	13.0	15.8	18.3	17.7	16.9	16.5	15.6	15.4	14.6	12.8	11.8	-0.9
10th Grade	16.5	15.2	19.2	25.2	28.7	33.6	34.8	31.1	32.1	32.2	32.7	30.3	28.2	27.5	-0.8
12th Grade	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3	-0.6
College Students	26.5	27.7	27.9	29.3	31.2	33.1	31.6	35.9	35.2	34.0	35.6	34.7	33.7	33.3	-0.4
Young Adults	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	+0.2
Inhalants^{c,d}															
8th Grade	9.0	9.5	11.0	11.7	12.8	12.2	11.8	11.1	10.3	9.4	9.1	7.7	8.7	9.6	+0.9
10th Grade	7.1	7.5	8.4	9.1	9.6	9.5	8.7	8.0	7.2	7.3	6.6	5.8	5.4	5.9	+0.5
12th Grade	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	+0.2
College Students	3.5	3.1	3.8	3.0	3.9	3.6	4.1	3.0	3.2	2.9	2.8	2.0	1.8	2.7	+0.9
Young Adults	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	+0.3
Nitrites^e															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.9	0.5	0.9	1.1	1.1	1.6	1.2	1.4	0.9	0.6	0.6	1.1	0.9	0.8	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.2	0.1	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}															
8th Grade	1.9	2.5	2.6	2.7	3.6	4.1	3.7	3.4	2.9	2.8†	3.4	2.6	2.6	2.2	-0.3
10th Grade	4.0	4.3	4.7	5.8	7.2	7.8	7.6	6.9	6.9	6.1†	6.2	4.7	4.1	4.1	0.0
12th Grade	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1†	9.1	6.6	5.9	6.2	+0.2
College Students	6.3	6.8	6.0	6.2	8.2	6.9	7.7	7.2	7.8	6.7	7.5	6.3	7.4	5.9	-1.4
Young Adults	4.5	5.0	4.5	4.8	5.6	5.6	5.9	5.2	5.4	5.4	5.4	4.7	5.2	4.7	-0.5

(Table continued on next page)

TABLE 2-2 (cont'd)
**Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)**

	<u>Annual</u>														'03-'04 change
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	
LSD															
8th Grade	1.7	2.1	2.3	2.4	3.2	3.5	3.2	2.8	2.4	2.4	2.2	1.5	1.3	1.1	-0.2
10th Grade	3.7	4.0	4.2	5.2	6.5	6.9	6.7	5.9	6.0	5.1	4.1	2.6	1.7	1.6	-0.1
12th Grade	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	+0.4
College Students	5.1	5.7	5.1	5.2	6.9	5.2	5.0	4.4	5.4	4.3	4.0	2.1	1.4	1.2	-0.1
Young Adults	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	-0.3
Hallucinogens Other Than LSD^b															
8th Grade	0.7	1.1	1.0	1.3	1.7	2.0	1.8	1.6	1.5	1.4†	2.4	2.1	2.1	1.9	-0.3
10th Grade	1.3	1.4	1.9	2.4	2.8	3.3	3.3	3.4	3.2	3.1†	4.3	4.0	3.6	3.7	+0.2
12th Grade	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4†	5.9	5.4	5.4	5.6	+0.2
College Students	3.1	2.6	2.7	2.8	4.0	4.1	4.9	4.4	4.5	4.4	5.5	5.8	7.1	5.6	-1.5
Young Adults	1.7	1.9	1.9	2.0	2.5	2.8	3.1	3.0	3.0	3.4	3.5	4.0	4.9	4.5	-0.5
PCP^g															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.4	1.4	1.6	1.8	2.6	2.3	2.1	1.8	2.3	1.8	1.1	1.3	0.7	-0.6
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	-0.2
MDMA (Ecstasy)^h															
8th Grade	—	—	—	—	—	2.3	2.3	1.8	1.7	3.1	3.5	2.9	2.1	1.7	-0.3
10th Grade	—	—	—	—	—	4.6	3.9	3.3	4.4	5.4	6.2	4.9	3.0	2.4	-0.6
12th Grade	—	—	—	—	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	-0.6
College Students	0.9	2.0	0.8	0.5	2.4	2.8	2.4	3.9	5.5	9.1	9.2	6.8	4.4	2.2	-2.2 s
Young Adults	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	-1.0 s
Cocaine															
8th Grade	1.1	1.5	1.7	2.1	2.6	3.0	2.8	3.1	2.7	2.6	2.5	2.3	2.2	2.0	-0.2
10th Grade	2.2	1.9	2.1	2.8	3.5	4.2	4.7	4.7	4.9	4.4	3.6	4.0	3.3	3.7	+0.4
12th Grade	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	+0.5
College Students	3.6	3.0	2.7	2.0	3.6	2.9	3.4	4.6	4.6	4.8	4.7	4.8	5.4	6.6	+1.1
Young Adults	6.2	5.7	4.7	4.3	4.4	4.1	4.7	4.9	5.4	5.4	5.8	5.8	6.6	7.1	+0.5
Crackⁱ															
8th Grade	0.7	0.9	1.0	1.3	1.6	1.8	1.7	2.1	1.8	1.8	1.7	1.6	1.6	1.3	-0.2
10th Grade	0.9	0.9	1.1	1.4	1.8	2.1	2.2	2.5	2.4	2.2	1.8	2.3	1.6	1.7	+0.1
12th Grade	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	+0.1
College Students	0.5	0.4	0.6	0.5	1.1	0.6	0.4	1.0	0.9	0.9	0.9	0.4	1.3	1.3	0.0
Young Adults	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	+0.3
Other Cocaine^j															
8th Grade	1.0	1.2	1.3	1.7	2.1	2.5	2.2	2.4	2.3	1.9	1.9	1.8	1.6	1.6	-0.1
10th Grade	2.1	1.7	1.8	2.4	3.0	3.5	4.1	4.0	4.4	3.8	3.0	3.4	2.8	3.3	+0.5
12th Grade	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	+0.5
College Students	3.2	2.4	2.5	1.8	3.3	2.3	3.0	4.2	4.2	4.1	4.1	5.0	5.1	6.3	+1.1
Young Adults	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	+0.3

(Table continued on next page)

TABLE 2-2 (cont'd)
Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Annual														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Heroin ^k															
8th Grade	0.7	0.7	0.7	1.2	1.4	1.6	1.3	1.3	1.4	1.1	1.0	0.9	0.9	1.0	0.0
10th Grade	0.5	0.6	0.7	0.9	1.1	1.2	1.4	1.4	1.4	1.4	0.9	1.1	0.7	0.9	+0.2
12th Grade	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	+0.1
College Students	0.1	0.1	0.1	0.1	0.3	0.4	0.3	0.6	0.2	0.5	0.4	0.1	0.2	0.4	+0.2
Young Adults	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	-0.1
With a Needle ^l															
8th Grade	—	—	—	—	0.9	1.0	0.8	0.8	0.9	0.6	0.7	0.6	0.6	0.7	+0.1
10th Grade	—	—	—	—	0.6	0.7	0.7	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.0
12th Grade	—	—	—	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.0
College Students	—	—	—	—	0.1	0.0	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.1	0.0
Young Adults	—	—	—	—	0.1	0.1	0.1	0.1	0.1	*	0.3	0.0	*	0.1	+0.1
Without a Needle ^l															
8th Grade	—	—	—	—	0.8	1.0	0.8	0.8	0.9	0.7	0.6	0.6	0.6	0.6	-0.1
10th Grade	—	—	—	—	0.8	0.9	1.1	1.0	1.1	1.1	0.7	0.8	0.5	0.7	+0.1
12th Grade	—	—	—	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	-0.1
College Students	—	—	—	—	0.0	0.8	0.4	0.9	0.3	0.8	0.6	0.2	0.1	0.6	+0.5
Young Adults	—	—	—	—	0.3	0.4	0.4	0.7	0.6	0.5	0.9	0.2	0.4	0.3	-0.1
Other Narcotics ^{m,n}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7†	9.4	9.3	9.5	+0.3
College Students	2.7	2.7	2.5	2.4	3.8	3.1	4.2	4.2	4.3	4.5	5.7†	7.4	8.7	8.2	-0.5
Young Adults	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0†	7.1	8.5	9.0	+0.5
OxyContin ^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	1.3	1.7	1.7	0.0
10th Grade	—	—	—	—	—	—	—	—	—	—	—	3.0	3.6	3.5	-0.1
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.0	4.5	5.0	+0.5
College Students	—	—	—	—	—	—	—	—	—	—	—	1.5	2.2	2.5	+0.3
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.9	2.6	3.1	+0.5
Vicodin ^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	2.5	2.8	2.5	-0.3
10th Grade	—	—	—	—	—	—	—	—	—	—	—	6.9	7.2	6.2	-1.0
12th Grade	—	—	—	—	—	—	—	—	—	—	—	9.6	10.5	9.3	-1.3
College Students	—	—	—	—	—	—	—	—	—	—	—	6.9	7.5	7.4	0.0
Young Adults	—	—	—	—	—	—	—	—	—	—	—	8.2	8.6	8.9	+0.3
Amphetamines ^m															
8th Grade	6.2	6.5	7.2	7.9	8.7	9.1	8.1	7.2	6.9	6.5	6.7	5.5	5.5	4.9	-0.6
10th Grade	8.2	8.2	9.6	10.2	11.9	12.4	12.1	10.7	10.4	11.1	11.7	10.7	9.0	8.5	-0.5
12th Grade	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	+0.1
College Students	3.9	3.6	4.2	4.2	5.4	4.2	5.7	5.1	5.8	6.6	7.2	7.0	7.1	7.0	-0.2
Young Adults	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	+0.5

(Table continued on next page)

TABLE 2-2 (cont'd)
Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Annual														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Ritalin^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	—	—	2.9	2.8	2.6	2.5	-0.1
10th Grade	—	—	—	—	—	—	—	—	—	—	4.8	4.8	4.1	3.4	-0.7
12th Grade	—	—	—	—	—	—	—	—	—	—	5.1	4.0	4.0	5.1	+1.1
College Students	—	—	—	—	—	—	—	—	—	—	—	5.7	4.7	4.7	0.0
Young Adults	—	—	—	—	—	—	—	—	—	—	—	2.9	2.9	2.7	-0.2
Methamphetamine^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	3.2	2.5	2.8	2.2	2.5	1.5	-1.0 ss
10th Grade	—	—	—	—	—	—	—	—	4.6	4.0	3.7	3.9	3.3	3.0	-0.3
12th Grade	—	—	—	—	—	—	—	—	4.7	4.3	3.9	3.6	3.2	3.4	+0.2
College Students	—	—	—	—	—	—	—	—	3.3	1.6	2.4	1.2	2.6	2.9	+0.3
Young Adults	—	—	—	—	—	—	—	—	2.8	2.5	2.8	2.5	2.7	2.8	+0.1
Ice^p															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.0	2.0	2.1	+0.1
College Students	0.1	0.2	0.7	0.8	1.1	0.3	0.8	1.0	0.5	0.5	0.6	0.8	0.9	1.1	+0.2
Young Adults	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	+0.3
Sedatives (Barbiturates)^m															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	+0.5
College Students	1.2	1.4	1.5	1.2	2.0	2.3	3.0	2.5	3.2	3.7	3.8	3.7	4.1	4.2	+0.1
Young Adults	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	+0.5
Methaqualone^{m,q}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	0.2	0.8	0.7	1.1	1.0	1.1	1.1	0.3	0.8	0.9	0.6	0.8	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}															
8th Grade	1.8	2.0	2.1	2.4	2.7	3.3	2.9	2.6	2.5	2.6†	2.8	2.6	2.7	2.5	-0.1
10th Grade	3.2	3.5	3.3	3.3	4.0	4.6	4.9	5.1	5.4	5.6†	7.3	6.3	5.3	5.1	-0.2
12th Grade	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7†	6.9	7.7	6.7	7.3	+0.6
College Students	2.4	2.9	2.4	1.8	2.9	2.8	3.8	3.9	3.8	4.2	5.1	6.7	6.9	6.7	-0.2
Young Adults	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6	5.5	7.0	6.8	7.4	+0.6
Rohypnol^f															
8th Grade	—	—	—	—	—	1.0	0.8	0.8	0.5	0.5	0.7	0.3	0.5	0.6	+0.1
10th Grade	—	—	—	—	—	1.1	1.3	1.2	1.0	0.8	1.0	0.7	0.6	0.7	+0.1
12th Grade	—	—	—	—	—	1.1	1.2	1.4	1.0	0.8	0.9†	1.6	1.3	1.6	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	0.7	0.4	0.3	-0.1
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.3	0.5	0.1	-0.4 s

(Table continued on next page)

TABLE 2-2 (cont'd)
Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	Annual														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
GHB^{o,u}															
8th Grade	—	—	—	—	—	—	—	—	—	1.2	1.1	0.8	0.9	0.7	-0.2
10th Grade	—	—	—	—	—	—	—	—	—	1.1	1.0	1.4	1.4	0.8	-0.6 s
12th Grade	—	—	—	—	—	—	—	—	—	1.9	1.6	1.5	1.4	2.0	+0.6
College Students	—	—	—	—	—	—	—	—	—	—	—	0.6	0.3	0.7	+0.4
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.8	0.6	0.5	-0.1
Ketamine^{o,v}															
8th Grade	—	—	—	—	—	—	—	—	—	1.6	1.3	1.3	1.1	0.9	-0.3
10th Grade	—	—	—	—	—	—	—	—	—	2.1	2.1	2.2	1.9	1.3	-0.6 s
12th Grade	—	—	—	—	—	—	—	—	—	2.5	2.5	2.6	2.1	1.9	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	1.3	1.0	1.5	+0.5
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.2	0.9	0.6	-0.3
Alcohol^f															
Any Use															
8th Grade	54.0	53.7†	45.4	46.8	45.3	46.5	45.5	43.7	43.5	43.1	41.9	38.7	37.2	36.7	-0.5
10th Grade	72.3	70.2†	63.4	63.9	63.5	65.0	65.2	62.7	63.7	65.3	63.5	60.0	59.3	58.2	-1.1
12th Grade	77.7	76.8†	72.7	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	+0.5
College Students	88.3	86.9	85.1	82.7	83.2	82.9	82.4	84.6	83.6	83.2	83.0	82.9	81.7	81.2	-0.5
Young Adults	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	+1.1
Flavored Alcoholic Beverages^{g,o,w}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	30.4	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	49.7	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	55.2	55.8	+0.6
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	63.2	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	62.7	—
Been Drunk^p															
8th Grade	17.5	18.3	18.2	18.2	18.4	19.8	18.4	17.9	18.5	18.5	16.6	15.0	14.5	14.5	0.0
10th Grade	40.1	37.0	37.8	38.0	38.5	40.1	40.7	38.3	40.9	41.6	39.9	35.4	34.7	35.1	+0.4
12th Grade	52.7	50.3	49.6	51.7	52.5	51.9	53.2	52.0	53.2	51.8	53.2	50.4	48.0	51.8	+3.8
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes															
Any Use															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
College Students	35.6	37.3	38.8	37.6	39.3	41.4	43.6	44.3	44.5	41.3	39.0	38.3	35.2	36.7	+1.5
Young Adults	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.11	39.1	38.6	39.0	+0.5
Bidis^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	—	3.9	2.7	2.7	2.0	1.7	-0.3
10th Grade	—	—	—	—	—	—	—	—	—	6.4	4.9	3.1	2.8	2.1	-0.7
12th Grade	—	—	—	—	—	—	—	—	—	9.2	7.0	5.9	4.0	3.6	-0.4
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page)

TABLE 2-2 (cont'd)
**Trends in Annual Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)**

	<u>Annual</u>														'03-'04 change
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	
Kretex ^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	—	—	2.6	2.6	2.0	1.9	-0.1
10th Grade	—	—	—	—	—	—	—	—	—	—	6.0	4.9	3.8	3.7	-0.2
12th Grade	—	—	—	—	—	—	—	—	—	—	10.1	8.4	6.7	6.5	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^p															
8th Grade	1.0	1.1	0.9	1.2	1.0	0.9	1.0	1.2	1.7	1.7	1.6	1.5	1.4	1.1	-0.3 s
10th Grade	1.1	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.7	2.2	2.1	2.2	1.7	1.5	-0.2
12th Grade	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	+0.4
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.4	0.5	0.5	0.0

NOTE: See Table 2-1 for relevant footnotes.

SOURCE: The Monitoring the Future Study, the University of Michigan.

TABLE 2-3
Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	30-Day														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Any Illicit Drug^a															
8th Grade	5.7	6.8	8.4	10.9	12.4	14.6	12.9	12.1	12.2	11.9	11.7	10.4	9.7	8.4	-1.3 s
10th Grade	11.6	11.0	14.0	18.5	20.2	23.2	23.0	21.5	22.1	22.5	22.7	20.8	19.5	18.3	-1.2
12th Grade	16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	-0.8
College Students	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	-0.2
Young Adults	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	-0.8
Any Illicit Drug Other Than Marijuana^{a,b}															
8th Grade	3.8	4.7	5.3	5.6	6.5	6.9	6.0	5.5	5.5	5.6†	5.5	4.7	4.7	4.1	-0.6
10th Grade	5.5	5.7	6.5	7.1	8.9	8.9	8.8	8.6	8.6	8.5†	8.7	8.1	6.9	6.9	0.0
12th Grade	7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4†	11.0	11.3	10.4	10.8	+0.4
College Students	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9	7.5	7.8	8.2	9.1	+0.9
Young Adults	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	+0.2
Any Illicit Drug Including Inhalants^{a,c}															
8th Grade	8.8	10.0	12.0	14.3	16.1	17.5	16.0	14.9	15.1	14.4	14.0	12.6	12.1	11.2	-0.8
10th Grade	13.1	12.6	15.5	20.0	21.6	24.5	24.1	22.5	23.1	23.6	23.6	21.7	20.5	19.3	-1.2
12th Grade	17.8	15.5	19.3	23.0	24.8	25.5	26.9	26.6	26.4	26.4	26.5	25.9	24.6	23.3	-1.3
College Students	15.1	16.5	15.7	16.4	19.6	18.0	19.6	21.0	21.8	22.6	21.9	21.9	21.6	21.7	+0.1
Young Adults	15.4	15.3	15.1	16.1	16.1	16.4	16.9	16.7	17.4	18.8	19.2	19.5	20.1	19.6	-0.5
Marijuana/Hashish															
8th Grade	3.2	3.7	5.1	7.8	9.1	11.3	10.2	9.7	9.7	9.1	9.2	8.3	7.5	6.4	-1.2 s
10th Grade	8.7	8.1	10.9	15.8	17.2	20.4	20.5	18.7	19.4	19.7	19.8	17.8	17.0	15.9	-1.2
12th Grade	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	-1.3
College Students	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	20.0	20.2	19.7	19.3	18.9	-0.4
Young Adults	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	-0.8
Inhalants^{c,d}															
8th Grade	4.4	4.7	5.4	5.6	6.1	5.8	5.6	4.8	5.0	4.5	4.0	3.8	4.1	4.5	+0.4
10th Grade	2.7	2.7	3.3	3.6	3.5	3.3	3.0	2.9	2.6	2.6	2.4	2.4	2.2	2.4	+0.1
12th Grade	2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	0.0
College Students	0.9	1.1	1.3	0.6	1.6	0.8	0.8	0.6	1.5	0.9	0.4	0.7	0.4	0.4	0.0
Young Adults	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	-0.1
Nitrites^e															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	*	0.1	0.2	0.1	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}															
8th Grade	0.8	1.1	1.2	1.3	1.7	1.9	1.8	1.4	1.3	1.2†	1.6	1.2	1.2	1.0	-0.2
10th Grade	1.6	1.8	1.9	2.4	3.3	2.8	3.3	3.2	2.9	2.3†	2.1	1.6	1.5	1.6	+0.1
12th Grade	2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6†	3.3	2.3	1.8	1.9	+0.1
College Students	1.2	2.3	2.5	2.1	3.3	1.9	2.1	2.1	2.0	1.4	1.8	1.2	1.8	1.3	-0.4
Young Adults	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2	1.2	0.9	1.2	0.9	-0.3

(Table continued on next page)

TABLE 2-3 (cont'd)
Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	30-Day														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
LSD															
8th Grade	0.6	0.9	1.0	1.1	1.4	1.5	1.5	1.1	1.1	1.0	1.0	0.7	0.6	0.5	-0.1
10th Grade	1.5	1.6	1.6	2.0	3.0	2.4	2.8	2.7	2.3	1.6	1.5	0.7	0.6	0.6	0.0
12th Grade	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	+0.1
College Students	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	0.9	1.0	0.2	0.2	0.2	-0.1
Young Adults	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.0
Hallucinogens Other Than LSD^b															
8th Grade	0.3	0.4	0.5	0.7	0.8	0.9	0.7	0.7	0.6	0.6†	1.1	1.0	1.0	0.8	-0.2
10th Grade	0.4	0.5	0.7	1.0	1.0	1.0	1.2	1.4	1.2	1.2†	1.4	1.4	1.2	1.4	+0.2
12th Grade	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7†	1.9	2.0	1.5	1.7	+0.2
College Students	0.6	0.7	1.1	0.8	1.6	1.2	1.2	0.7	1.2	0.8	0.8	1.1	1.7	1.2	-0.4
Young Adults	0.3	0.5	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.7	0.6	0.8	1.2	0.9	-0.3
PCP^g															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.2	0.2	0.0	0.0	0.1	0.1	0.1	0.0
MDMA (Ecstasy)^h															
8th Grade	—	—	—	—	—	1.0	1.0	0.9	0.8	1.4	1.8	1.4	0.7	0.8	+0.1
10th Grade	—	—	—	—	—	1.8	1.3	1.3	1.8	2.6	2.6	1.8	1.1	0.8	-0.4
12th Grade	—	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	0.0
College Students	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	2.5	1.5	0.7	1.0	0.7	-0.3
Young Adults	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	-0.1
Cocaine															
8th Grade	0.5	0.7	0.7	1.0	1.2	1.3	1.1	1.4	1.3	1.2	1.2	1.1	0.9	0.9	0.0
10th Grade	0.7	0.7	0.9	1.2	1.7	1.7	2.0	2.1	1.8	1.8	1.3	1.6	1.3	1.7	+0.4
12th Grade	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	+0.3
College Students	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	1.4	1.9	1.6	1.9	2.4	+0.5
Young Adults	2.0	1.8	1.4	1.3	1.5	1.2	1.6	1.7	1.9	1.7	2.2	2.2	2.4	2.2	-0.1
Crackⁱ															
8th Grade	0.3	0.5	0.4	0.7	0.7	0.8	0.7	0.9	0.8	0.8	0.8	0.8	0.7	0.6	-0.1
10th Grade	0.3	0.4	0.5	0.6	0.9	0.8	0.9	1.1	0.8	0.9	0.7	1.0	0.7	0.8	+0.1
12th Grade	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	+0.1
College Students	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.1	0.3	0.4	0.4	0.0
Young Adults	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	+0.1
Other Cocaine^j															
8th Grade	0.5	0.5	0.6	0.9	1.0	1.0	0.8	1.0	1.1	0.9	0.9	0.8	0.7	0.7	0.0
10th Grade	0.6	0.6	0.7	1.0	1.4	1.3	1.6	1.8	1.6	1.6	1.2	1.3	1.1	1.5	+0.4 s
12th Grade	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	+0.4
College Students	1.0	0.9	0.6	0.3	0.8	0.6	1.3	1.5	1.0	0.9	1.5	1.4	1.9	2.2	+0.4
Young Adults	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	0.0

(Table continued on next page)

TABLE 2-3 (cont'd)
Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	30-Day														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Heroin^k															
8th Grade	0.3	0.4	0.4	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.0
10th Grade	0.2	0.2	0.3	0.4	0.6	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.5	+0.1
12th Grade	0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	+0.1
College Students	0.1	0.0	*	0.0	0.1	*	0.2	0.1	0.1	0.2	0.1	0.0	*	0.1	+0.1
Young Adults	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.0
With a Needle^l															
8th Grade	—	—	—	—	0.4	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.0
10th Grade	—	—	—	—	0.3	0.3	0.3	0.4	0.3	0.3	0.2	0.3	0.2	0.3	+0.1
12th Grade	—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	-0.1
College Students	—	—	—	—	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.0	0.1	0.1	0.0
Young Adults	—	—	—	—	0.0	0.0	0.1	*	0.1	*	0.2	0.0	*	0.1	0.0
Without a Needle^l															
8th Grade	—	—	—	—	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.0
10th Grade	—	—	—	—	0.3	0.3	0.4	0.5	0.5	0.4	0.2	0.4	0.2	0.3	+0.1
12th Grade	—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	-0.1
College Students	—	—	—	—	0.0	0.1	0.2	0.2	0.3	0.4	0.3	0.0	0.0	0.3	+0.3
Young Adults	—	—	—	—	0.1	*	0.1	0.2	0.2	0.2	0.4	*	0.1	0.1	+0.1
Other Narcotics^{m,n}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0†	4.0	4.1	4.3	+0.2
College Students	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	1.7	1.7†	3.2	2.3	3.0	+0.7
Young Adults	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7†	2.9	2.9	3.0	+0.1
Amphetamines^m															
8th Grade	2.6	3.3	3.6	3.6	4.2	4.6	3.8	3.3	3.4	3.4	3.2	2.8	2.7	2.3	-0.5
10th Grade	3.3	3.6	4.3	4.5	5.3	5.5	5.1	5.1	5.0	5.4	5.6	5.2	4.3	4.0	-0.2
12th Grade	3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	-0.3
College Students	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	2.9	3.3	3.0	3.1	3.2	+0.1
Young Adults	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	-0.1
Methamphetamine^{o,p}															
8th Grade	—	—	—	—	—	—	—	—	1.1	0.8	1.3	1.1	1.2	0.6	-0.6 ss
10th Grade	—	—	—	—	—	—	—	—	1.8	2.0	1.5	1.8	1.4	1.3	-0.1
12th Grade	—	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	-0.3
College Students	—	—	—	—	—	—	—	—	1.2	0.2	0.5	0.2	0.6	0.2	-0.4
Young Adults	—	—	—	—	—	—	—	—	0.8	0.7	1.0	1.0	0.7	0.6	-0.1
Ice^p															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	+0.1
College Students	0.0	0.0	0.3	0.5	0.3	0.1	0.2	0.3	0.0	0.0	0.1	0.0	0.3	0.1	-0.2
Young Adults	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.4	0.4	0.0

(Table continued on next page)

TABLE 2-3 (cont'd)
Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	30-Day														'03-'04
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	change
Sedatives (Barbiturates) ^m															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9	2.9	0.0
College Students	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	1.1	1.5	1.7	1.7	1.5	-0.3
Young Adults	0.5	0.5	0.6	0.6	0.8	0.8	0.9	0.9	1.1	1.3	1.7	1.5	1.5	1.8	+0.3
Methaqualone ^{m,q}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	+0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{b,m}															
8th Grade	0.8	0.8	0.9	1.1	1.2	1.5	1.2	1.2	1.1	1.4†	1.2	1.2	1.4	1.2	-0.2
10th Grade	1.2	1.5	1.1	1.5	1.7	1.7	2.2	2.2	2.2	2.5†	2.9	2.9	2.4	2.3	-0.1
12th Grade	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6†	2.9	3.3	2.8	3.1	+0.4
College Students	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	2.0	1.5	3.0	2.8	2.7	0.0
Young Adults	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8	2.1	2.8	2.4	2.7	+0.3
Rohypnol ^f															
8th Grade	—	—	—	—	—	0.5	0.3	0.4	0.3	0.3	0.4	0.2	0.1	0.2	+0.1
10th Grade	—	—	—	—	—	0.5	0.5	0.4	0.5	0.4	0.2	0.4	0.2	0.3	0.0
12th Grade	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^s															
Any Use															
8th Grade	25.1	26.1†	24.3	25.5	24.6	26.2	24.5	23.0	24.0	22.4	21.5	19.6	19.7	18.6	-1.2
10th Grade	42.8	39.9†	38.2	39.2	38.8	40.4	40.1	38.8	40.0	41.0	39.0	35.4	35.4	35.2	-0.2
12th Grade	54.0	51.3†	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	+0.5
College Students	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	67.4	67.0	68.9	66.2	67.7	+1.5
Young Adults	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.0	68.3	67.0	68.4	+1.4
Flavored Alcoholic Beverages ^{g,o}															
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	14.6	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	25.1	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	31.1	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	34.1	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	—
Been Drunk ^p															
8th Grade	7.6	7.5	7.8	8.7	8.3	9.6	8.2	8.4	9.4	8.3	7.7	6.7	6.7	6.2	-0.5
10th Grade	20.5	18.1	19.8	20.3	20.8	21.3	22.4	21.1	22.5	23.5	21.9	18.3	18.2	18.5	+0.3
12th Grade	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	+1.6
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page)

TABLE 2-3 (cont'd)
**Trends in Thirty-Day Prevalence of Use of Various Drugs for Eighth, Tenth,
and Twelfth Graders, College Students, and Young Adults (Ages 19-28)**

	<u>30-Day</u>														'03-'04 change
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	
Cigarettes															
Any Use															
8th Grade	14.3	15.5	16.7	18.6	19.1	21.0	19.4	19.1	17.5	14.6	12.2	10.7	10.2	9.2	-1.0
10th Grade	20.8	21.5	24.7	25.4	27.9	30.4	29.8	27.6	25.7	23.9	21.3	17.7	16.7	16.0	-0.7
12th Grade	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	+0.6
College Students	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	28.2	25.7	26.7	22.5	24.3	+1.9
Young Adults	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	+0.9
Smokeless Tobacco [†]															
8th Grade	6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	0.0
10th Grade	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	-0.4
12th Grade	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ^P															
8th Grade	0.4	0.5	0.5	0.5	0.6	0.4	0.5	0.5	0.7	0.8	0.7	0.8	0.7	0.5	-0.1
10th Grade	0.6	0.6	0.5	0.6	0.6	0.5	0.7	0.6	0.9	1.0	0.9	1.0	0.8	0.8	+0.1
12th Grade	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.2	0.1	0.0	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.1	-0.2

NOTE: See Table 2-1 for relevant footnotes.

SOURCE: The Monitoring the Future Study, the University of Michigan.


TABLE 2-4
Trends in Thirty-Day Prevalence of Daily Use of Various Drugs for Eighth, Tenth, and Twelfth Graders, College Students, and Young Adults (Ages 19-28)

	<u>Daily</u>														'03-'04 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
Marijuana/Hashish, daily ^x															
8th Grade	0.2	0.2	0.4	0.7	0.8	1.5	1.1	1.1	1.4	1.3	1.3	1.2	1.0	0.8	-0.2
10th Grade	0.8	0.8	1.0	2.2	2.8	3.5	3.7	3.6	3.8	3.8	4.5	3.9	3.6	3.2	-0.5
12th Grade	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	-0.4
College Students	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	4.6	4.5	4.1	4.7	4.5	-0.2
Young Adults	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	-0.3
Alcohol ^{s,x}															
Any daily use															
8th Grade	0.5	0.6†	1.0	1.0	0.7	1.0	0.8	0.9	1.0	0.8	0.9	0.7	0.8	0.6	-0.2
10th Grade	1.3	1.2†	1.8	1.7	1.7	1.6	1.7	1.9	1.9	1.8	1.9	1.8	1.5	1.3	-0.2
12th Grade	3.6	3.4†	3.4	2.9	3.5	3.7	3.9	3.9	3.4	2.9	3.6	3.5	3.2	2.8	-0.4
College Students	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	3.6	4.7	5.0	4.3	3.7	-0.5
Young Adults	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	-0.5
Been Drunk, daily ^{p,x}															
8th Grade	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.2	0.0
10th Grade	0.2	0.3	0.4	0.4	0.6	0.4	0.6	0.6	0.7	0.5	0.6	0.5	0.5	0.4	-0.1
12th Grade	0.9	0.8	0.9	1.2	1.3	1.6	2.0	1.5	1.9	1.7	1.4	1.2	1.6	1.8	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
5+ drinks in a row in last 2 weeks															
8th Grade	12.9	13.4	13.5	14.5	14.5	15.6	14.5	13.7	15.2	14.1	13.2	12.4	11.9	11.4	-0.5
10th Grade	22.9	21.1	23.0	23.6	24.0	24.8	25.1	24.3	25.6	26.2	24.9	22.4	22.2	22.0	-0.2
12th Grade	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	+1.3
College Students	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	39.3	40.9	40.1	38.5	41.7	+3.3
Young Adults	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	+1.3
Cigarettes															
Any daily use															
8th Grade	7.2	7.0	8.3	8.8	9.3	10.4	9.0	8.8	8.1	7.4	5.5	5.1	4.5	4.4	-0.2
10th Grade	12.6	12.3	14.2	14.6	16.3	18.3	18.0	15.8	15.9	14.0	12.2	10.1	8.9	8.3	-0.6
12th Grade	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	-0.3
College Students	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	17.8	15.0	15.9	13.8	13.8	-0.1
Young Adults	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	+0.5
1/2 pack+/day															
8th Grade	3.1	2.9	3.5	3.6	3.4	4.3	3.5	3.6	3.3	2.8	2.3	2.1	1.8	1.7	-0.1
10th Grade	6.5	6.0	7.0	7.6	8.3	9.4	8.6	7.9	7.6	6.2	5.5	4.4	4.1	3.3	-0.9
12th Grade	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0	-0.3
College Students	8.0	8.9	8.9	8.0	10.2	8.4	9.1	11.3	11.0	10.1	7.8	7.9	7.6	6.8	-0.8
Young Adults	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	-0.4
Smokeless Tobacco, daily [†]															
8th Grade	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	+0.2
10th Grade	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	-0.2
12th Grade	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	+0.6
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

NOTE: See Table 2-1 for relevant footnotes.

SOURCE: The Monitoring the Future Study, the University of Michigan.

FIGURE 2-1
Trends in Annual Prevalence of an Illicit Drug Use Index
Across Five Populations



NOTES: Use of "any illicit drugs" includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin, or any use which is not under a doctor's orders of other opiates, stimulants, barbiturates, methaqualone (excluded since 1990), or tranquilizers.

Beginning in 1982, the question about stimulant use (i.e., amphetamines) was revised to get respondents to exclude the inappropriate reporting of nonprescription stimulants. The prevalence rate dropped slightly as a result of this methodological change.

Chapter 3

STUDY DESIGN AND PROCEDURES

The Monitoring the Future study's design is one that yields analytic power well beyond the sum of its component parts. Those parts include the cross-sectional study, the repeated cross-sectional study, and the panel study. As a cross-sectional study, it provides point estimates of various behaviors and conditions at any given point in time. Repeating these cross-sectional studies over time allows an assessment of change across years in the same segments of the population. The panel-study feature permits the examination of change over time in the same individuals as they enter adult roles and environments and assume adult responsibilities. These are all important research objectives. However, with a *series* of panel studies of sequential graduating class cohorts of students, in what is known as a cohort-sequential design, we are able to distinguish among, and explain, three fundamentally different types of change: period-related, age-related, and cohort-related. It is this last feature that creates the synergistic effect in terms of analytic power.

This chapter describes this complex research design, including the sampling plans and field procedures used in both the annual in-school cross-sectional surveys of 8th-, 10th-, and 12th-grade students and the follow-up surveys into early and middle adulthood—the panel studies. Related methodological issues such as response rates, population coverage, and the validity of the measures are also discussed.

We begin by describing the design that has been used consistently over the past 30 years to survey high school seniors; then we describe the more recently instituted design for 8th and 10th graders. Finally, the design for the *follow-up* surveys of former 12th graders is covered.^{14, 15}

RESEARCH DESIGN AND PROCEDURES FOR THE SURVEYS OF SENIORS

High school seniors have been surveyed in the spring of each year since 1975. Each year's data collection takes place in approximately 120 to 146 public and private high schools selected to provide an accurate representative cross section of high school seniors throughout the coterminous United States (see Figure 3-1).

¹⁴For a more detailed description of the study design, see Bachman, J. G., Johnston, L. D., & O'Malley, P. M. (2001). *The Monitoring the Future project after twenty-seven years: Design and procedures*. (Monitoring the Future Occasional Paper No. 54.) Ann Arbor, MI: Institute for Social Research.

¹⁵For a more detailed description of the full range of research objectives of Monitoring the Future, see Johnston, L. D., O'Malley, P. M., Schulenberg, J. E., & Bachman, J. G. (2001). *The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2001*. (Monitoring the Future Occasional Paper No. 52.) Ann Arbor, MI: Institute for Social Research.

The Population Under Study

We chose the senior year of high school because, for several reasons, it is an optimal point at which to monitor the drug use and related attitudes of youth. First, completion of high school represents the end of an important developmental stage in this society, demarcating both the end of universal education and, for many, the end of living full-time in the parental home. Therefore, it is a logical point at which to take stock of the cumulated influences of these two environments on American youth. Further, completion of high school represents the jumping-off point from which young people diverge into widely differing social environments and experiences. Senior year, then, represents a good time to take a “before” measure allowing calculation of changes that may be attributable to the many environmental and role transitions occurring in young adulthood. Finally, there were some important practical advantages to building the original system of data collections around samples of high school seniors. The need for systematically repeated, large-scale samples from which to make reliable estimates of change requires that considerable stress be laid on cost efficiency as well as feasibility. The last year of high school constitutes the final point at which a reasonably good national sample of an age-specific cohort can be drawn and studied economically.

The Omission of Dropouts

One limitation in the study design is the exclusion of those young men and women who drop out of high school before graduation—between 15% and 20% of each age cohort nationally, according to U.S. Census statistics. Clearly, the omission of high school dropouts introduces biases in the estimation of certain characteristics of the entire age group; however, for most purposes, the small proportion of dropouts sets outer limits on the bias. Further, since the bias from missing dropouts should remain just about constant from year to year, their omission should introduce little or no bias in *change* estimates. Indeed, we believe the changes observed over time for those who finish high school are likely to parallel the changes for dropouts in most instances. Appendix A to Volume I addresses the likely effects of the exclusion of dropouts on estimates of drug use prevalence and trends among the entire age cohort; see that appendix for a more detailed discussion of this issue.

Sampling Procedures

A multi-stage random sampling procedure is used to secure the nationwide sample of high school seniors each year. Stage 1 is the selection of particular geographic areas, stage 2 is the selection (with probability proportionate to size) of one or more high schools in each area, and stage 3 is the selection of seniors within each high school. Up to about 350 seniors in each school may be included. In schools with fewer seniors, the usual procedure is to include all of them in the data collection, though a smaller sample is sometimes taken to accommodate the needs of the school. When a subset of seniors is to be selected, it is done either by randomly sampling entire classrooms or by some other unbiased, random method. Weights are assigned to compensate for differential probabilities of selection at each stage of sampling. Final weights are normalized to average 1.0 (so that the weighted number of cases equals the unweighted number of cases overall). This three-stage sampling procedure has yielded the numbers of participating schools and students over the years shown in Table 3-1.

Questionnaire Administration

About 10 days before the questionnaire administration date, the target respondents are given flyers explaining the study. Local Institute for Social Research representatives and their assistants conduct the actual questionnaire administrations following standardized procedures detailed in a project instruction manual. The questionnaires are administered in classrooms during a normal class period whenever possible; however, circumstances in some schools require the use of larger group administrations.

Questionnaire Format

Because many questions are needed to cover all of the topic areas in the study, much of the questionnaire content intended for high school seniors is divided into six different questionnaire forms distributed to participants in an ordered sequence that ensures six virtually identical random sub-samples. (Five questionnaire forms were used between 1975 and 1988.) About one-third of each questionnaire form consists of key, or “core,” variables common to all forms. All demographic variables, and nearly all of the drug *use* variables included in this report, are contained in this core set of measures. Many of the questions dealing with attitudes, beliefs, and perceptions of relevant features of the social environment are in a single form only, and the data are thus based on one-fifth as many cases in 1975-1988 (approximately 3,300) and on one-sixth as many cases beginning in 1989 (approximately 2,600). All tables in this report list the sample sizes upon which the statistics are based, stated in terms of the weighted number of cases (which is roughly equivalent to the actual number of cases).

RESEARCH DESIGN AND PROCEDURES FOR THE SURVEYS OF LOWER GRADES

In 1991, the study expanded to include nationally representative samples of 8th- and 10th-grade students. Surveys at these two grade levels have been conducted on an annual basis since 1991.

In general, the procedures used for the annual in-school surveys of 8th- and 10th-grade students closely parallel those used for high school seniors, including the procedures for selecting schools and students, questionnaire administration, and questionnaire formats. A major exception is that only two different questionnaire forms were used from 1991 to 1996, expanding to four forms beginning in 1997, rather than the six used with seniors. Eighth and 10th graders both receive the same questionnaire forms and, for the most part, the questionnaire content is drawn from the 12th-grade questionnaires. Thus, key demographic variables and measures of drug use and related attitudes and beliefs are generally identical for all three grades. The forms used in both 8th and 10th grades have a common core (Parts B and C) that parallels the core used in 12th-grade forms. Many fewer questions about lifestyles and values are included in the 8th- and 10th-grade forms, in part because we think that many of these attitudes are likely to be more fully formed by 12th grade and, therefore, are best monitored there.

For the national survey of 8th graders each year, approximately 150 schools (mostly junior high schools and middle schools) are sampled, and approximately 17,000 students have been surveyed

annually. For the 10th graders, approximately 130 high schools are sampled and about 15,000 students surveyed annually. (See Table 3-1 for specifics.)¹⁶

Mode of Administration

From 1991 to 1993, 8th and 10th graders were followed up similarly to the 12th graders (see footnote 20). When follow-up surveys of new cohorts of 8th and 10th graders were no longer being conducted, the collection of personal identification information for follow-up purposes was no longer necessary. For confidentiality reasons, this personal information had been gathered on a tear-off sheet at the back of each questionnaire. We believed that there were potential advantages in moving toward a fully anonymous procedure for these grade levels, including the following: (a) school cooperation might be easier to obtain; (b) any suppression effect that the confidential mode of administration might have could be both eliminated and quantified; and (c) if there *were* any mode of administration effect, it would be removed from the national data—which are widely compared with results of state and local surveys (nearly all of which use anonymous questionnaires)—thus making those comparisons more valid. Therefore, in 1998 for the first time, in half of the 8th- and 10th-grade schools surveyed, the questionnaires administered were made fully anonymous. Specifically, the half-sample of schools beginning their two-year participation in Monitoring the Future in 1998 received the anonymous questionnaires, while the half-sample participating in the study for their second and final year continued to get the confidential questionnaires.

A careful examination of the 1998 results, based on the two equivalent half-samples at grade 8, and also at grade 10, revealed that there was no effect of this methodological change among 10th graders, and, at most, only a very modest effect in the self-reported substance use rates among 8th graders (with prevalence rates slightly higher in the anonymous condition). The net effect of this methodological change is a possible increase in the observed 8th-grade prevalence estimates for marijuana, alcohol, and cigarettes in 1998 from what they would have been had there been no change in questionnaire administration. For those three drugs, that means that the declines in use in 1998 may be slightly understated for the 8th graders only. In other words, the direction of the change is the same as that shown in the tables, but the actual declines may be slightly larger than those shown. For example, the annual prevalence of marijuana use among 8th graders is shown to have fallen by 0.8 percentage points between 1997 and 1998; however, the half-sample of 8th-grade schools receiving exactly the same type of questionnaire that was used in 1997 showed a slightly greater decline of 1.5 percentage points.

For cigarettes, this change in method appeared to have no effect on self-reported rates of daily use or half-pack per day use and to have had only a very small effect on 30-day prevalence.

¹⁶The research design originally called for follow-up surveys of sub-samples of the 8th and 10th graders participating in the study, carried out at two-year intervals, similar to the 12th-grade follow-up samples. From 1991 to 1994, this plan influenced the design of the cross-sectional studies of 8th and 10th graders in an important way. In order to “recapture” many of the 8th-grade participants two years later in the normal 10th-grade cross-sectional study for that year, we selected the 8th-grade schools by drawing a sample of high schools and then selecting a sample of their “feeder schools” that contained 8th graders. This extra stage in the sampling process meant that many of the 8th-grade participants in, say, the 1991 cross-sectional survey were also participants in the 1993 cross-sectional survey of 10th graders. Thus, a fair amount of panel data was generated at no additional cost. However, having followed this design from 1991 through 1993, we concluded that the saving in follow-up costs did not justify the complexities in sampling, administration, and interpretation. Therefore, since 1994, we have used a simplified design in which 8th-grade schools are drawn independently of the 10th-grade school sample. Further follow-ups (at two-year intervals) have been conducted only on panels of students drawn from the first three cohorts of students surveyed in the 8th and 10th grades—that is, those surveyed in school in 1991, 1992, and 1993. A book reporting results from these panels is now well underway.

Thus, for example, the 30-day prevalence of cigarette use among all of the 8th graders surveyed is shown to have fallen 0.3 percentage points between 1997 and 1998; while the half-sample of 8th-grade schools receiving exactly the same type of questionnaire as was used in 1997 showed a slightly greater decline of 0.6 percentage points. Finally, lifetime cigarette prevalence is shown as falling by 1.6 percentage points between 1997 and 1998, but in the half-sample of schools with a constant methodology, it fell by 2.6 percentage points.

We have examined in detail the effects of administration mode in a published journal article, in which we use multivariate controls to assess the effects of the change on the 8th-grade self-report data. It generally shows even less effect than is to be found without such controls.¹⁷

All tables and figures in Volume I use data from both half-samples of 8th graders surveyed in a given year, combined. This is also true for the 10th graders (for whom we found no methodological effect) and the 12th graders (for whom it is assumed there is no such effect, since none was found among the 10th graders). (See this chapter's later section entitled "Representativeness and Sample Accuracy, School Participation," for a further discussion of half-samples among all three grades.)

In 1999 the remaining half of the participating schools (all beginning the first of their two years of participation) received anonymous questionnaires, as well. Thus, from 1999 on, all data from 8th- and 10th-grade students are gathered using anonymous questionnaires. We continue to use confidential questionnaires with 12th graders in order to permit follow-up of the small proportion that are randomly selected into the panel studies.

Questionnaire Forms and Sample Proportions

Another consequence of not interlocking the school samples at 8th and 10th grades was that we could consider having more forms of the questionnaire.¹⁸ Beginning in 1997, the number of forms was expanded to four, but the four forms are not distributed in equal numbers. Forms 1, 2, 3, and 4 are assigned to one-third, one-third, one-sixth, and one-sixth of the students, respectively. Thus, if a question appears on only one form, it may be administered to either one-third or one-sixth of the sample. Similarly, a question in two forms may be assigned to one-third of the sample (one-sixth plus one-sixth), one-half of the sample (one-third plus one-sixth), or two-thirds of the sample (one-third plus one-third). No questions appear on three forms. Footnotes to the tables indicate what proportion of all respondents in each grade complete the question, if that proportion is other than the entire sample.

The two additional forms were introduced to allow for more questions. The new forms 1 and 2 substantially follow the content of the previous forms 1 and 2, but each was now assigned to a third of the sample instead of half. Form 3 builds on form 1, with some questions omitted to make room for more content; and form 4 builds on the content of form 2 in a similar manner.

¹⁷O'Malley, P. M., Johnston, L. D., Bachman, J. G., & Schulenberg, J. (2000). A comparison of confidential versus anonymous survey procedures: Effects on reporting of drug use and related attitudes and beliefs in a national study of students. *Journal of Drug Issues*, 30, 35-54.

¹⁸Earlier, from 1991 through 1996, two questionnaire forms were used in the surveys of 8th- and 10th-grade students, with a random half-sample of students in each grade receiving one form and the remainder receiving the other form. (By having only two forms distributed randomly at each grade, we could by chance emerge with half of the students being surveyed both times with the same form, making panel analysis possible.) With the constraint of "recapturing" students removed, we could consider having a larger number of forms.

Much of the new content was placed in both of the new forms (forms 3 and 4), each of which is administered to one-sixth of the sample, in order to assign one-third of the total sample to those new questions.

RESEARCH DESIGN AND PROCEDURES FOR THE FOLLOW-UP SURVEYS OF SENIORS

Beginning with the graduating class of 1976, some members of each senior class have been selected to be surveyed by mail after high school graduation. From the roughly 13,000 to 17,000 seniors originally participating in a given senior class, a representative sample of 2,400 individuals is chosen for follow-up. In order to ensure sufficient numbers of drug users in the follow-up surveys, seniors reporting 20 or more occasions of marijuana use in the previous 30 days (i.e., “daily users”), or *any* use of the other illicit drugs in the previous 30 days, are selected with higher probability (by a factor of 3.0) than the remaining seniors. Differential weighting is then used in all follow-up analyses to compensate for these differential sampling probabilities. Because those in the drug-using stratum receive a weight of only 0.33 in the calculation of all statistics to correct for their overrepresentation at the selection stage, there are actually more follow-up respondents than are reported in the weighted Ns given in the tables.

The 2,400 participants selected from each 12th-grade class are randomly split into two matching groups of 1,200 each—one group to be surveyed on even-numbered calendar years, and the other group to be surveyed on odd-numbered years. This two-year cycle is intended to reduce the burden on individual respondents, thus yielding a better retention rate across the years. By alternating the two half-samples, we have data from a given graduating class every year, even though any given respondent participates only every other year.

Until 2002, each respondent was followed for up to seven times; at the seventh follow-up, which would occur either 13 or 14 years after graduation, the respondents had reached modal age of 31 or 32. Beginning in 2002, the seventh follow-up was discontinued, and each respondent was followed for up to six times, corresponding to modal age of 29 or 30. Additional follow-ups still occur at modal ages 35, 40, and 45. (Age 45 follow-ups began in 2003, when the class of 1976 reached that age.) Our intention is to continue follow-ups at five-year intervals beyond age 45 to the extent that panel retention rates justify such continuation. Data like these, gathered on representative national samples over such a large part of the life span, are extremely rare and can provide needed insight into the etiology of substance use and other behaviors across the life course.

Follow-Up Procedures

Using information provided by high school senior respondents on a tear-off card (containing the respondent’s name, address, and phone number, and the name and address of someone who would always know how to reach them), mail contact is maintained with the subset of people selected for inclusion in the follow-up panels. Newsletters are sent to them each year, and name and address corrections are requested from both the U.S. Postal Service and the individual. Questionnaires are sent to each individual biennially in the spring of each year by certified mail.

A check for \$20, made payable to the respondent, is attached to the front of each questionnaire.¹⁹ Reminder letters and postcards are sent at fixed intervals thereafter; finally, those who have not responded receive a prompting phone call from the Survey Research Center's phone interviewing facility in Ann Arbor, Michigan. If requested, a second copy of the questionnaire is sent; but no questionnaire content is administered by phone. If a respondent asks not to be contacted further, that wish is honored.

Follow-Up Questionnaire Format

The questionnaires used in the young adult follow-up surveys are very much like those used in the senior year. They are optically scanned; all forms contain a common core section that includes questions on drug use, background factors, and demographic factors; and they have questions about a wide range of topics at the beginning and ending sections, many of which are unique to each questionnaire form. Many of the questions asked of seniors are retained in the corresponding follow-up questionnaires, and respondents are consistently mailed the same version (or form) of the questionnaire that they first received in senior year, so that *changes over time* in their behaviors, attitudes, experiences, and so forth can be measured. Questions specific to high school status and experiences are dropped in the follow-up, of course, and questions relevant to post-high school status and experiences are added. Thus, there are questions about college, military service, civilian employment, marriage, parenthood, and so on. Most of these are added to the core section. For the 5-year surveys that begin at age 35, the questionnaire content is streamlined (only one form is used) and directed at the major family and work issues of middle adulthood. Still, many of the questions are ones repeated from the young adult surveys.

For the early follow-up cohorts, the numbers of cases on single-form questions were one-fifth the size of the total follow-up sample because five different questionnaire forms were used. Beginning with the class of 1989, a sixth form was introduced in the senior year. That new questionnaire form was first sent to follow-up respondents in 1990; single-form data since then have Ns one-sixth the total follow-up sample size. In the follow-up studies, single-form samples from a single cohort are too small to make reliable estimates; therefore, in most cases where they are reported, the data from several adjacent cohorts are combined or concatenated.

REPRESENTATIVENESS AND SAMPLE ACCURACY

School Participation

Schools are invited to participate in the study for a two-year period. For each school that declines to participate, a similar school (in terms of size, geographic area, urbanicity, etc.) is recruited as a replacement for that "slot." In 2004, either an original school or a replacement school was obtained in 99% of the sample units, or "slots." With very few exceptions, each school participating in the first year has agreed to participate in the second year as well. Figure 3-2 provides the year-specific school participation rates and the percentage of "slots" filled since

¹⁹For the class of 1991 and all prior classes, the follow-up checks were for \$5. The rate was raised to \$10, beginning with the class of 1992, to compensate for the effects of inflation over the life of the study. An experiment was first conducted that suggested that the increased payment was justified based on the increased panel retention it achieved. Payment increased to \$20 in 2004 for much the same reason.

1977. (The data for the years prior to 1991 are for 12th grade only; beginning in 1991, the data are for 8th, 10th, and 12th grades, combined.) As shown in the figure, replacements for declining schools are obtained in the vast majority of cases.

Two questions are sometimes raised with respect to school participation rates: (a) Are participation rates so low as to compromise the representativeness of the sample? (b) Does variation in participation rates over time contribute to changes in estimates of drug use?

With respect to the first issue, the selection of replacement schools (which occurs in practically all instances of an original school refusal) almost entirely removes problems of bias in region, urbanicity, and the like that might result from certain schools refusing to participate. Other potential biases could be more subtle, however. If, for example, it turned out that most schools with “drug problems” refused to participate, the sample would be seriously biased. And if any other single factor were dominant in most refusals, that reason for refusal also might suggest a source of serious bias. In fact, however, the reasons given for a school refusing to participate tend to be varied and are often a function of happenstance specific to that particular year; only a very small proportion specifically object to the drug-related or “sensitive” nature of the content of the survey.

If it were the case that schools differed substantially in drug use, then which particular schools participated could have a greater effect on estimates of drug use. However, the great majority of variance in drug use lies within schools, not between schools. For example, for 10th graders in 1992, between-schools variance for marijuana use was 4%-6% of the total variance (depending on the specific measure); for inhalant use, 1%-2%; for LSD, 2%-4%; for crack cocaine, 1.0%-1.5%; for alcohol use, 4%-5%; and for cigarette use, 3%-4%. (Eighth- and 12th-grade values are similar.) To the extent that schools tend to be fairly similar in drug use, then which particular schools participate (within a selection framework that seeks national representation) has a small effect on estimates of drug use. The fact that the overwhelming majority of variance in drug use lies within schools implies that, at least with respect to drug use, schools are for the most part fairly similar.²⁰ Further, some, if not most, of the between-schools variance is due to differences related to region, urbanicity, etc.—factors that remain well controlled in the present sampling design because of the way in which replacement schools are selected.

With respect to the second issue, the observed data from the series make it extremely unlikely that results have been significantly affected by changes in response rates. If changes in response rates seriously affected prevalence estimates, there would be noticeable bumps up or down in concert with the changing rates. But in fact this series of surveys produces results that are very smooth and change in an orderly fashion from one year to the next. This suggests that the level of school-related error in the estimates does not vary much over time. Moreover, the fact that different substances trend in distinctly different ways further refutes any likelihood that changes in response rates are affecting prevalence estimates. We have observed, for example, marijuana

²⁰Among the schools that actually participated in the study, there is very little difference in substance use rates between the schools that were original selections, taken as a set, and the schools that were replacement schools. Averaged over the years 1991 through 2000, for grades 8, 10, and 12 combined, the difference between original schools and replacement schools averaged 0.03% in the observed prevalence rates averaged across a number of drug use measures: two indexes of annual illicit drug use, the annual prevalence of each of the major illicit drug classes, and several measures of alcohol and cigarette use. For the individual drugs and drug indexes, the differences between the original and replacement schools, averaged across grades and years, fell within $\pm 0.9\%$.

use decreasing while cocaine use was stable (in the early 1980s); alcohol use declining while cigarette use was stable (in the mid- to late 1980s); and marijuana use increasing while inhalant use was decreasing (from 1994 to 1997). All of these patterns are explainable in terms of psychological, social, and cultural factors (as described in this and previous volumes in this series) and cannot be explained by the common factor of changes in response rates.

Of course, there could be some sort of a constant bias across the years; but even in the unlikely event that there was, it seems highly improbable that it would be of much consequence for policy purposes, given that it would not affect trends and likely would have a very modest effect on prevalence rates. Thus we have a high degree of confidence that school refusal rates have not seriously biased the survey results.

Nevertheless, it is apparent that, for a host of reasons, securing high school cooperation rates has become more difficult in recent years. This is a problem common to the field, not specific to Monitoring the Future. Therefore, in the study's most recent proposal for continuation we requested funding to permit the payment of schools as a means of increasing their incentives to participate. (Several other ongoing school survey studies already use payments to schools.) Such payments were approved and were implemented in the 2003 survey.

At each grade level, schools are selected in such a way that half of each year's sample comprises schools that started their participation the previous year, and half comprises schools that began participating in the current year. (Both samples are national replicates, meaning that each is drawn to be nationally representative by itself.) This staggered half-sample design is used to check on possible errors in the year-to-year trend estimates due to school turnover. For example, separate sets of one-year trend estimates are computed based on students in the half-sample of schools that participated in both 2001 *and* 2002, then based on the students in the half-sample that participated in both 2002 *and* 2003, and so on. Thus, each one-year *matched half-sample* trend estimate derived in this way is based on a constant set of schools (about 65 in 12th grade, for example). When the trend data derived from the matched half-sample (examined separately for each class of drugs) are compared with trends based on the total sample of schools, the results are usually highly similar, indicating that the trend estimates are affected little by turnover or shifting refusal rates in the school samples. As would be expected, the *absolute* prevalence-of-use estimates for a given year are not as accurate using just the half-sample because the sample size is only half as large.

Student Participation

In 2004, completed questionnaires were obtained from 89% of all sampled students in 8th grade, 88% in 10th grade, and 82% in 12th grade. (See Table 3-1 for response rates in earlier years.) The single most important reason that students are missed is absence from class at the time of data collection; in most cases, for reasons of cost efficiency, we do not schedule special follow-up data collections for absent students. Students with fairly high rates of absenteeism also report above-average rates of drug use; therefore, some degree of bias is introduced into the prevalence estimates by missing the absentees. Much of that bias could be corrected through the use of special weighting based on the reported absentee rates of the students who *did* respond; however, we decided not to use such a weighting procedure because the bias in overall drug use

estimates was determined to be quite small *and* because the necessary weighting procedures would have introduced greater sampling variance in the estimates. Appendix A in an earlier report²¹ provides a discussion of this point, and Appendix A in Volume I illustrates the changes in trend and prevalence estimates that would result if corrections for absentees had been included. Of course, some students are not absent from class but simply refuse, when asked, to complete a questionnaire. However, the proportion of explicit refusals amounts to less than 1.5% of the target sample for each grade.

Sampling Accuracy of the Estimates

Confidence intervals (95%) are provided in Tables 4-1a through 4-1d (Volume I) for lifetime, annual, 30-day, and daily prevalence of use for 8th-, 10th-, and 12th-grade students. As can be seen in Table 4-1a, confidence intervals for lifetime prevalence for seniors average less than $\pm 1.5\%$ across a variety of drug classes. That is, if we took a large number of samples of this size from the universe of all schools containing 12th graders in the coterminous United States, 95 times out of 100 the sample would yield a result that would be less than 1.5 percentage points divergent from the result we would get from a comparable massive survey of *all* seniors in *all* schools. This is a high level of sampling accuracy, and it should permit detection of fairly small changes from one year to the next. Confidence intervals for the other prevalence periods (past 12 months, past 30 days, and current daily use) are generally smaller than those for lifetime use. In general, confidence intervals for 8th and 10th graders are very similar to those observed for 12th graders. Some drugs (smokeless tobacco, PCP, nitrites, and others, as indicated in Table 2-1 footnotes) are measured on only one or two questionnaire forms; these drugs will have somewhat larger confidence intervals due to their smaller sample sizes. Appendix C of Volume I contains information for the interested reader on how to calculate confidence intervals around other point estimates; it also provides the information needed to compare trends across time or to test the significance of differences between subgroups in any given year.

PANEL RETENTION

We discuss here the nature of the problem of panel attrition generally, the response rates we have attained in the Monitoring the Future panel surveys in recent years, and evidence relevant to assessing the impact of attrition on the study's research results.

The Problem of Panel Attrition

Virtually all longitudinal studies of drug use, including Monitoring the Future, experience attrition, which is often differential with respect to substance use.²² In addition, survey response rates in general have been declining over the past few decades,²³ highlighting an important challenge in the conduct of population-based research.

²¹Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975-1983*. DHHS (ADM) 85-1374. Washington, DC: U.S. Government Printing Office.

²²McGuigan, K. A., Ellickson, P. L., Hays, R. D., & Bell, R. M. (1997). Adjusting for attrition in school-based samples: Bias, precision, and cost trade-off of three methods. *Evaluation Review*, 21, 554-567.

²³Groves, R. M., Dillman, D. A., Eltinge, J. L., & Little, R. J. A. (Eds.). (2002). *Survey nonresponse*. New York: Wiley.

A vital feature of the Monitoring the Future panel studies is their very low cost per respondent. There are many advantages to collecting panel data through low-cost mail surveys, as we have done since the outset of the study. Indeed, given the number of panel surveys we administer each year (roughly 15,000) across the entire coterminous United States, using low-cost mail surveys is our best (and really the only) cost-effective option. One disadvantage of this mode of data collection is that attrition rates tend to be higher than those that might be obtained with much more expensive methods, for example, intensive personal tracking and interviewing. Certainly there exist a few large epidemiological/etiological surveys that have better retention rates, but their procedures are extremely expensive and not realistic for an ongoing effort like this one. Nevertheless, our retention rates compare reasonably favorably with those of most longitudinal studies (including interview studies) reported in the field.

Response Rates Attained

We begin with the college student segment in the follow-up sample. The series of survey data on American college students now goes back 24 years. We know about actual college attendance only from the follow-up questionnaire answers; however, we can use senior year questionnaire answers (i.e., college intentions and program of study) to predict college attendance with a high degree of accuracy. The study's retention of college-bound seniors remains quite good. Among those follow-up respondents who, in high school, reported plans to attend college *and* enrollment in a college-prep curriculum, the follow-up retention rates in 2001, for example, for the three most recent classes surveyed at each follow-up point were as follows: 70% in the first follow-up, one to two years past high school (based on the classes of 1998-2000); 67% in the second follow-up, three to four years past high school (based on the classes of 1996-1998); and 65% in the third follow-up, five to six years past high school (based on the classes of 1994-1996). To date we have reported in Volume II only on college students who are one to four years past high school graduation. As the average age of attendance rises, having the extended age coverage will be of growing importance. The follow-up participation rates just noted compare favorably with another major national survey of substance use among college students, the Harvard College Alcohol Study, which in both its 1997 and 1999 surveys had cross-sectional response rates of 60%.²⁴

Retention rates in the biennial follow-ups of *all* panel members ages 19-30 (corresponding to the first six follow-ups) decline with the length of the follow-up interval, of course. For the five-year period from 2000 to 2004, the response rate in the first follow-up (corresponding to 1-2 years past high school) averaged 60%; for the second through sixth follow-ups (corresponding to 3-12 years past high school) response rates averaged 54%. Among the very long-term respondents—the 35-, 40-, and 45-year-olds—the retention rates are quite good, apparently because some of the decline with age in retention rates reflects cohort differences. Among the 35-year-old respondents surveyed from 2000 to 2004 (corresponding to 17 years past high school), the average response rate was 51%. Among the 40-year-old respondents surveyed from 2000 to 2004, corresponding to a 22-year follow-up interval, the average retention rate was 58%. Among 45-year-olds surveyed in 2003 and 2004, the average retention rate was 60%.

²⁴Wechsler, H., Lee, J. E., Kuo, M., & Lee, H. (2000). College binge drinking in the 1990s: A continuing problem. Results of the Harvard School of Public Health 1999 College Alcohol Study. *Journal of American College Health*, 48, 195-198.

In sum, the response rates attained under the current design range from respectable to quite good, especially when the low-cost nature of the procedures and the substantial length of the questionnaires are taken into account. More importantly, the evidence leaves us confident that the data resulting from these follow-up panels are reasonably accurate, which brings us to our adjustments for panel attrition and the comparison of our results with those from other sources.

The Impact of Panel Attrition on Research Results

An important purpose of the Monitoring the Future follow-ups is to allow estimation of drug prevalence rates among American high school graduates at various age levels, as published annually in Volume II of this series. Thus, we have always been concerned about making the appropriate adjustments to account for panel attrition. In essence, our standard adjustment procedure is a post-stratification procedure in which we reweight the data from the individuals' obtained follow-up samples so that their reweighted senior year distribution on a given drug reproduces the original (senior year) distribution of use originally observed for that drug. This procedure is carried out (separately) for cigarettes, alcohol, marijuana, and (combined) other illicit drugs. As expected, this procedure produces estimates that are somewhat higher than those uncorrected for attrition, indicating that there is indeed some positive association between drug use and panel attrition. However, the adjustments are relatively modest, as is documented next.

One reason the adjustments are modest is that attrition rates do not differ greatly by levels of senior year substance use; they do differ, but less than one might expect. For example, among all respondents who had never used marijuana, an average of 81% of the classes of 1976-1993 participated in the first follow-up. The proportion responding is somewhat lower among those who had used marijuana once or twice in the past 12 months: 78%. This proportion decreases gradually with increasing levels of marijuana use; but even among those who used marijuana on 20-39 occasions in the past 30 days in their high school senior year, 71% participated in the first follow-up. The corresponding participation rates for the same drug-use strata at the fourth follow-up (i.e., at ages 25-26) were 68%, 65%, and 60%, respectively. Thus, even among those who in high school were quite heavy users of marijuana, response rates at the fourth follow-up were only 8 percentage points lower than among those who had never used marijuana by high school senior year. That is not to say that we assume that all types of drug users remain in the panels at comparably high rates. We believe that people who become dependent on, or addicted to, heroin or cocaine are unlikely to be retained in reasonable proportions. That is why we are careful to not quantify or characterize these special segments of the population. But we note that they constitute very low proportions of the drug-using portion of the population and even lower proportions of the entire adult population. Therefore, for a great many purposes, our samples are extremely useful.

The National Survey on Drug Use and Health (NSDUH) would seem to provide the best available data against which to validate the estimates generated for adult age groups in Monitoring the Future because it is also based on national samples but uses cross-sectional surveys that do not carry the burden of panel attrition. (Their results, of course, may be affected by their own nonresponse rates; but that will be true of any comparison survey. The overall response rates for the NSDUH were about 73% in 1997 and 1998, and 61% in 1999.)

In some earlier analyses we compared the prevalence rates on a set of drugs—cigarettes, alcohol, marijuana, and cocaine—for which there was reasonable similarity in question wording across the two studies. The comparisons that follow are for the age group 19-28 in the Monitoring the Future panel data and for 19-28 (or 19-29 for 1999 only) in the NSDUH cross-sectional data. At that time the most recent data from NSDUH that were readily available for these comparisons were for 1999, so the following comparisons are for that year. (However, similar comparisons were run for a number of prior years, and the outcomes were highly consistent.) The comparisons are not perfect; most notably, the NSDUH data contain school dropouts and, other things equal, this inclusion would lead one to expect its rates to be *higher* than those from Monitoring the Future. Nevertheless, the Monitoring the Future estimates for the 30-day prevalence of marijuana actually are higher (14.0% without post-stratification and 15.6% with it) than the NSDUH estimate (11.0%). The same is true for the 12-month cocaine prevalence estimate (4.8% without post-stratification and 5.4% with it, vs. 4.3% in the NSDUH).

The other two comparisons made were for alcohol and cigarettes. Both of these drugs show larger differences, with alcohol use consistently higher in Monitoring the Future and cigarette use consistently higher in NSDUH. We believe it likely that both are due to definitional differences in the exact question wording. In 1999, Monitoring the Future estimates of 30-day alcohol prevalence were 68.0% and 68.2% (with post-stratification) vs. 59.5% in NSDUH. For cigarettes, the 30-day Monitoring the Future prevalence estimates were 28.3% and 30.3%, respectively, vs. 37.4% in NSDUH. It is worth noting that the nature and magnitude of the differences between Monitoring the Future and NSDUH estimates tend to be quite consistent for each of the four drugs since at least 1992.

The fact that Monitoring the Future estimates for both marijuana and cocaine are higher than NSDUH estimates (especially after applying the post-stratification reweighting) suggests that attrition does not produce substantially lower estimates of drug use than would be obtained if response rates were higher. Our estimates come out as high as, and in fact somewhat higher than, the best available comparison study for estimating rates using cross-sectional data, and that despite our loss of dropouts and absentees (in high school) from the MTF samples.

It is also worth noting that even with attrition, there remain in the Monitoring the Future follow-up samples substantial proportions of recent users of the various substances. In recent years, about 15%-17% of the 19–28-year-old respondents reported marijuana use in just the prior 30 days, and about 5%-7% reported past 12-month use of cocaine. These proportions and the underlying numbers of actual cases are quite adequate for analytic purposes, particularly given that the follow-up surveys over-sample those who reported illicit drug use in the senior year surveys.

An important point worth emphasizing here is that in the present study, attrition is not necessarily as great a problem as is nonresponse in a cross-sectional study. This is because we already know a great deal about each of the follow-up nonrespondents, including their substance use, based on a lengthy questionnaire in senior year (and, for many, in subsequent years as well). Thus, adjustments can be made utilizing data that are highly informative about the lost individuals.

Effects on Relational Analyses

While differential attrition (uncorrected) may contribute to some bias in point estimates and other univariate statistics, such attrition tends to have less influence on bivariate and multivariate statistics. This was found to be true in a secondary analysis of data from seven panel studies that followed adolescents over time,²⁵ and we have found this to be true in our Monitoring the Future panel analyses²⁶ and in analyses with other panel data sets.²⁷ Thus, differential attrition may be of less concern in multivariate panel analyses focused on understanding the course, causes, and consequences of substance use. Still, as we summarized above, correcting for attrition is important, and we continue to do so.

VALIDITY OF THE MEASURES OF SELF-REPORTED DRUG USE

Are sensitive behaviors such as drug use honestly reported? Like most studies dealing with sensitive behaviors, we have no direct, totally objective validation of the present measures; however, the considerable amount of existing inferential evidence strongly suggests that the self-report questions used in Monitoring the Future produce largely valid data. A more complete discussion of the contributing evidence that leads to this conclusion may be found in other publications; here we only briefly summarize the evidence.²⁸

First, using a three-wave panel design, we established that the various measures of self-reported drug use have a high degree of reliability—a necessary condition for validity.²⁹ In essence, respondents were highly consistent in their self-reported behaviors over a three- to four-year time interval. Second, we found a high degree of consistency among logically related measures of use within the same questionnaire administration. Third, the proportion of seniors reporting some illicit drug use by senior year has reached two-thirds of all respondents in peak years and over 80% in some follow-up years, constituting *prima facie* evidence that the degree of underreporting must be very limited. Fourth, the seniors' reports of use by their unnamed friends—about whom they would presumably have less reason to distort reports of use—have been highly consistent with self-reported use in the aggregate, in terms of both prevalence *and*

²⁵Cordray, S., & Polk, K. (1983). The implication of respondent loss in panel studies of deviant behavior. *Journal of Research in Crime and Delinquency*, 20, 214-242.

²⁶Bryant, A. L., Schulenberg, J., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (2000). Understanding the links among school misbehavior, academic achievement, and cigarette use: A national panel study of adolescents. *Prevention Science*, 1(2), 71-87; Schulenberg, J., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (1994). High school educational success and subsequent substance use: A panel analysis following adolescents into young adulthood. *Journal of Health and Social Behavior*, 35, 45-62.

²⁷Bachman, J. G., O'Malley, P. M., & Johnston, J. (1978). *Youth in Transition: Vol. 6. Adolescence to adulthood: A study of change and stability in the lives of young men*. Ann Arbor, MI: Institute for Social Research; Schulenberg, J. E., Bryant, A. L., & O'Malley, P. M. (2004). Taking hold of some kind of life: How developmental tasks relate to trajectories of well-being during the transition to adulthood. *Development and Psychopathology*, 16, 1119-1140.

²⁸Johnston, L. D., & O'Malley, P. M. (1985). Issues of validity and population coverage in student surveys of drug use. In B. A. Rouse, N. J. Kozel, & L. G. Richards (Eds.), *Self-report methods of estimating drug use: Meeting current challenges to validity* (NIDA Research Monograph No. 57 (ADM) 85-1402). Washington, DC: U.S. Government Printing Office; Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975-1983*. DHHS (ADM) 85-1374. Washington, DC: U.S. Government Printing Office; Wallace, J. M., Jr., & Bachman, J. G. (1993). Validity of self-reports in student-based studies on minority populations: Issues and concerns. In M. de LaRosa (Ed.), *Drug abuse among minority youth: Advances in research and methodology*. NIDA Research Monograph. Rockville, MD: National Institute on Drug Abuse.

²⁹O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self-reports of drug use. *International Journal of the Addictions*, 18, 805-824.

trends in prevalence, as will be discussed later in this report. Fifth, we have found self-reported drug use to relate in consistent and expected ways to a number of other attitudes, behaviors, beliefs, and social situations—in other words, there is strong evidence of “construct validity.” Sixth, the missing data rates for the self-reported use questions are only very slightly higher than for the preceding non-sensitive questions, in spite of explicit instructions to respondents immediately preceding the drug section to leave blank those drug use questions they felt they could not answer honestly. Seventh, an examination of consistency in reporting of lifetime use conducted on the long-term panels of graduating seniors found quite low levels of recanting of earlier-reported use of the illegal drugs.³⁰ There was a higher level of recanting for the psychotherapeutic drugs, which we interpreted as suggesting that adolescents actually may overestimate their use of some of these drugs because of misinformation about definitions that is corrected as they get older. Finally, the great majority of respondents, when asked, say they would answer such questions honestly if they were users.³¹

This is not to argue that self-reported measures of drug use are valid in all cases. In the present study we have gone to great lengths to create a situation and set of procedures in which students feel that their confidentiality will be protected. We have also tried to present a convincing case as to why such research is needed. The evidence suggests that a high level of validity has been obtained. Nevertheless, insofar as any remaining reporting bias exists, we believe it to be in the direction of underreporting. Thus, we believe our estimates to be lower than their true values, even for the obtained samples, but not substantially so.

One additional procedure we undertake to help assure the validity of our data is worth noting. We check for logical inconsistencies in the triplets of answers about the use of each drug (i.e., about lifetime, past year, and past 30-day use), and if a respondent exceeds a minimum number of inconsistencies across the drug use questions, his or her record is deleted from the data set. Similarly, we check for improbably high rates of use of multiple drugs and delete such cases, on the assumption that the respondents are not taking the task seriously. Fortunately, relatively few cases have to be eliminated for these reasons.

Consistency and the Measurement of Trends

One further point is worth noting in a discussion of the validity of the findings. The Monitoring the Future project is designed to be sensitive to changes from one time period to another. A great strength of this study, in our opinion, is that the measures and procedures have been standardized and applied consistently across many years. To the extent that any biases remain because of limits in school and/or student participation, and to the extent that there are distortions (lack of validity) in the responses of some students, it seems very likely that such problems will exist in much the same proportions from one year to the next. In other words, biases in the survey

³⁰Johnston, L. D. & O'Malley, P. M. (1997). The recanting of earlier reported drug use by young adults. In Harrison, L. (Ed.), *The validity of self-reported drug use: Improving the accuracy of survey estimates* (pp. 59-80). (NIDA Research Monograph 167, pp 59-79). Rockville, MD: National Institute on Drug Abuse.

³¹For a discussion of reliability and validity of student self-report measures of drug use like those used in Monitoring the Future across varied cultural settings, see also Johnston, L. D., Driessen, F. M. H. M., & Kokkevi, A. (1994). *Surveying student drug misuse: A six-country pilot study*. Strasbourg, France: Council of Europe.

estimates will tend to be consistent from one year to another, which means that our measurement of *trends* should be affected very little by any such biases. The smooth and consistent nature of most trend curves reported for the various drugs provides rather compelling empirical support for this assertion.

TABLE 3-1
Sample Sizes and Response Rates

Grade:	Number of Public Schools			Number of Private Schools			Total Number of Schools				Total Number of Students				Student Response Rate		
	8th	10th	12th	8th	10th	12th	8th	10th	12th	Total	8th	10th	12th	Total	8th	10th	12th
1975	—	—	111	—	—	14	—	—	125	—	—	—	15,791	—	—	—	78%
1976	—	—	108	—	—	15	—	—	123	—	—	—	16,678	—	—	—	77
1977	—	—	108	—	—	16	—	—	124	—	—	—	18,436	—	—	—	79
1978	—	—	111	—	—	20	—	—	131	—	—	—	18,924	—	—	—	83
1979	—	—	111	—	—	20	—	—	131	—	—	—	16,662	—	—	—	82
1980	—	—	107	—	—	20	—	—	127	—	—	—	16,524	—	—	—	82
1981	—	—	109	—	—	19	—	—	128	—	—	—	18,267	—	—	—	81
1982	—	—	116	—	—	21	—	—	137	—	—	—	18,348	—	—	—	83
1983	—	—	112	—	—	22	—	—	134	—	—	—	16,947	—	—	—	84
1984	—	—	117	—	—	17	—	—	134	—	—	—	16,499	—	—	—	83
1985	—	—	115	—	—	17	—	—	132	—	—	—	16,502	—	—	—	84
1986	—	—	113	—	—	16	—	—	129	—	—	—	15,713	—	—	—	83
1987	—	—	117	—	—	18	—	—	135	—	—	—	16,843	—	—	—	84
1988	—	—	113	—	—	19	—	—	132	—	—	—	16,795	—	—	—	83
1989	—	—	111	—	—	22	—	—	133	—	—	—	17,142	—	—	—	86
1990	—	—	114	—	—	23	—	—	137	—	—	—	15,676	—	—	—	86
1991	131	107	117	31	14	19	162	121	136	419	17,844	14,996	15,483	48,323	90%	87%	83
1992	133	106	120	26	19	18	159	125	138	422	19,015	14,997	16,251	50,263	90	88	84
1993	126	111	121	30	17	18	156	128	139	423	18,820	15,516	16,763	51,099	90	86	84
1994	116	116	119	34	14	20	150	130	139	419	17,708	16,080	15,929	49,717	89	88	84
1995	118	117	120	34	22	24	152	139	144	435	17,929	17,285	15,876	51,090	89	87	84
1996	122	113	118	30	20	21	152	133	139	424	18,368	15,873	14,824	49,065	91	87	83
1997	125	113	125	27	18	21	152	131	146	429	19,066	15,778	15,963	50,807	89	86	83
1998	122	110	124	27	19	20	149	129	144	422	18,667	15,419	15,780	49,866	88	87	82
1999	120	117	124	30	23	19	150	140	143	433	17,287	13,885	14,056	45,228	87	85	83
2000	125	121	116	31	24	18	156	145	134	435	17,311	14,576	13,286	45,173	89	86	83
2001	125	117	117	28	20	17	153	137	134	424	16,756	14,286	13,304	44,346	90	88	82
2002	115	113	102	26	20	18	141	133	120	394	15,489	14,683	13,544	43,716	91	85	83
2003	117	109	103	24	20	19	141	129	122	392	17,023	16,244	15,200	48,467	89	88	83
2004	120	111	109	27	20	19	147	131	128	406	17,413	16,839	15,222	49,474	89	88	82


SOURCE: The Monitoring the Future Study, the University of Michigan.

FIGURE 3-1
Schools Included in One Year's Data Collection
Eighth, Tenth and Twelfth Grades



Note: One dot equals one school.

**FIGURE 3-2
School Response Rates**



	<u>77</u>	<u>78</u>	<u>79</u>	<u>80</u>	<u>81</u>	<u>82</u>	<u>83</u>	<u>84</u>	<u>85</u>	<u>86</u>	<u>87</u>	<u>88</u>	<u>89</u>	<u>90</u>	<u>91</u>	<u>92</u>	<u>93</u>	<u>94</u>	<u>95</u>	<u>96</u>	<u>97</u>	<u>98</u>	<u>99</u>	<u>00</u>	<u>01</u>	<u>02</u>	<u>03</u>	<u>04</u>
Original	59	63	62	63	71	71	66	72	67	66	72	71	68	70	59	55	60	53	52	53	51	51	57	62	56	49	53	62
Replacements	39	36	35	32	25	26	32	26	29	33	26	26	30	29	39	43	39	44	44	43	47	48	42	35	42	48	45	37
Total	98	99	97	95	96	97	99	98	96	99	99	98	99	99	98	98	99	97	96	96	98	99	99	97	98	97	98	99

Chapter 4

PREVALENCE OF DRUG USE IN EARLY AND MIDDLE ADULTHOOD

In this chapter we present data on several age groups of adults who are high school graduates, covering the full range of drugs that are included in the study. The respondents for the follow-up surveys come from the previous classes of high school students who participated in the study in their senior year. As described in more detail in chapter 3, the Monitoring the Future study conducts ongoing panel studies on representative samples from each graduating high school senior class, beginning with the class of 1976. Two matched subpanels, of roughly 1,200 students each, are randomly selected from each graduating class to comprise long-term follow-up panels—one panel is surveyed every even-numbered year after graduation, the other is surveyed every odd-numbered year, up through age 30. Thus, in a given year, the study encompasses one of the panels from each of the last 12 senior classes previously participating in the study.³² Because the study design calls for an end of biennial follow-up of these panels after the respondents reach approximately age 30 (i.e., six follow-ups for each half panel), in 2004 the (older) classes of 1976 through 1991 were not included in the standard, biennial follow-up surveys. Rather, representative samples of the classes of 1992 through 2003 were surveyed, using the standard young adult survey instruments. For brevity, we refer to this 19- through 30-year-old age group as “young adults” in this chapter.

Additional surveys are conducted at age 35 (that is, 17 years after high school graduation) and at 5-year intervals thereafter. In 2004, the class of 1987 received the “age-35” follow-up questionnaire, the class of 1982 received the “age-40” questionnaire, and, the class of 1977 received the “age-45” questionnaire. The findings from these special 5-year follow-up questionnaires are included here, so this chapter spans the age interval of 18 to 45, with those of modal age 18 represented by the 12th graders.

The results of these 2004 follow-up surveys should accurately characterize approximately 86% of all young adults 1 to 12 years beyond high school (modal ages 19 to 30) as well as 86% of adults 17, 22, and 27 years beyond high school (modal ages 35, 40, and 45). The remaining 14% or so—the high school dropout segment—were missing from the senior year surveys and, of course, were missing from all of the follow-up surveys as well. Thus, the results presented here are not necessarily generalizable to that small segment of the population.

Figures 4-1 through 4-21 contain the 2004 *prevalence* data by age, corresponding to those respondents 1 to 12 years beyond high school (modal ages 19 to 30), as well as 35-, 40-, and 45-year-olds. Figures provided in chapter 5 contain the *trend* data for each age group, including 12th graders and high school graduates through age 45. With the exception of the 12th graders, age groups have been paired into two-year intervals in both sets of figures in order to increase the number of cases, and thus the precision, for each point estimate. The data for ages 35, 40, and 45 are,

³²Through 2001, the follow-ups also included modal ages 31 and 32. This seventh follow-up was dropped in 2002 because we believed that the marginal costs no longer were justified by the marginal benefits of having this follow-up data, given that an age-35 survey is being conducted. Throughout the time between surveys, we send a newsletter to respondents in order to help maintain contact with them.

of necessity, based on a single age in each case. *Both* half samples from a given class cohort are included in the samples of 35-, 40-, and 45-year-olds, so in 2004 the two half samples come from the graduating classes of 1987, 1982, and 1977, respectively. Their respective weighted numbers of cases are 979, 1004, and 1041. (Actual, unweighted numbers are somewhat higher.)

It is worth noting that the pattern of age-related differences showing up in any one year can be checked in an adjacent year (i.e., the previous year's volume or the succeeding year's) for replicability, because two non-overlapping half samples of follow-up respondents in the 19 to 30 age band are surveyed on alternating years. In the case of the 35-, 40-, and 45-year-olds, two entirely different graduating classes make up the samples for any two adjacent years.

A NOTE ON ADJUSTED LIFETIME PREVALENCE ESTIMATES

In Figures 4-1 through 4-21, two different estimates of lifetime prevalence are provided. One estimate is based on the respondent's most recent statement of whether he or she ever used the drug in question (the light gray bar). The other estimate takes into account the respondent's answers regarding lifetime use gathered in *all* of the previous data collections in which he or she participated (the white bar). *To be categorized as one who has used the drug based on all past answers regarding that drug, the respondent must have reported either lifetime use in the most recent data collection and/or some use in his or her lifetime on at least two earlier occasions.* Because respondents in the age groups of 18-year-olds and 19- to 20-year-olds cannot have their responses adjusted on the basis of two earlier occasions, adjusted prevalence rates are reported only for ages 21 and older. Epidemiological studies usually present an unadjusted estimate because it must be made based on the data from just a single cross-sectional survey. An adjusted estimate of the type used here is possible only when panel data have been gathered so that a respondent can be classified as having used a drug at some time in his or her life, based on earlier answers, even though he or she no longer indicates lifetime use in the most recent survey.

The divergence of these two estimates as a function of age shows that there is more inconsistency as time passes. Obviously, there is more opportunity for inconsistency as the number of data collections increases. Our judgment is that "the truth" lies somewhere between the two estimates: the lower estimate may be depressed by tendencies to forget, forgive, or conceal earlier use, and the upper estimate may include earlier response errors or incorrect definitions of drugs that respondents appropriately corrected in later surveys as they became more knowledgeable. It should be noted that a fair proportion of those giving inconsistent answers across time had earlier reported having used the given drug only once or twice in their lifetime.

As we have reported elsewhere, cross-time stability of self-reported usage measures, which takes into account both prevalence and frequency of self-reported use, is still very high.³³ Note that the divergence between the two lifetime prevalence estimates is greatest for the psychotherapeutic drugs and for the derivative index of “use of an illicit drug other than marijuana,” which is heavily affected by the psychotherapeutic estimates. We believe this is due to respondents having greater difficulty accurately categorizing psychotherapeutic drugs (usually taken in pill form) with a high degree of certainty—especially if such a drug was used only once or twice. We expect higher inconsistency across time when the event—and in many of these cases, a single event—is reported with a relatively low degree of certainty at quite different points in time. Those who have gone beyond simple experimentation with one of these drugs would undoubtedly be able to categorize them with a higher degree of certainty. Also, those who have experimented more recently, in the past month or year, should have a higher probability of recall, as well as fresher information for accurately categorizing the drug.

We provide both estimates to make clear that a full use of respondent information provides a possible range for lifetime prevalence estimates, not a single point. However, by far the most important use of the prevalence data is to track *trends* in *current* (as opposed to lifetime) use. Thus, we are much less concerned about the nature of the variability in the lifetime estimates than we might otherwise be. The lifetime prevalence estimates are of importance primarily in showing the degree to which a drug class has penetrated the general population overall as well as particular cohorts.³⁴

The reader is reminded that the re-weighting procedures used to correct the panel data for the effects of panel attrition are described in chapter 3.

PREVALENCE OF DRUG USE AS A FUNCTION OF AGE

Figures 4-1 through 4-21 provide 2004 age comparisons of prevalence rates for each class of drugs, covering the age range 18 to 45. For virtually all drugs, available age comparisons show much higher lifetime prevalence for the older age groups, as would be expected. In fact, the figures reach impressive levels among adults in their early 30s through their mid-40s.

- The adjusted lifetime prevalence figures are most impressive for today’s 40- and 45-year-olds, who were passing through adolescence in the peak of the drug epidemic. Some 82% and 88% of them, respectively, reported trying *an illicit drug* (lifetime prevalence, adjusted), leaving only 18% and 12% who have reported not doing so. (See Figure 4-1.) Some 75% and 81% of the 40- and 45-year-olds, respectively, said they had tried marijuana, and about two-thirds (63% and 71%) said they had tried some other illicit drug, including 38% and 44%

³³O’Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self-reports of drug use. *International Journal of the Addictions*, 18, 805-824.

³⁴For a more detailed analysis and discussion, see Johnston, L. D., & O’Malley, P. M. (1997). The recanting of earlier-reported drug use by young adults. In L. Harrison, & A. Hughes (Eds.), *The validity of self-reported drug use: Improving the accuracy of survey estimates*. (NIDA Research Monograph No. 97-4147.) Washington, DC: National Institute on Drug Abuse.

who have tried cocaine specifically. Clearly, the parents of today's teenagers are themselves a very drug-experienced generation.

- In 2004 the adjusted lifetime prevalence figures among 29- to 30-year-olds reach 65% for *any illicit drug*, 61% for *marijuana*, 41% for *any illicit drug other than marijuana*, and 16% for *cocaine*. Put another way, even among young Americans who graduated from high school in 1992 and 1993—after the peak of the larger drug epidemic—only about one-third (35%) have *never* tried an illegal drug.

Their 2004 survey responses, *unadjusted* for previous answers, show somewhat lower lifetime prevalence: 59% for *any illicit drug*, 57% for *marijuana*, 32% for *any illicit drug other than marijuana*, and 14% for *cocaine*.

- Despite the higher lifetime prevalence rates among older age groups, these groups generally show levels of *annual* or *current* use that are no higher than such use among today's 12th graders. In fact, for a number of drugs, the levels reported by older respondents are lower, suggesting that the incidence of quitting more than offsets the incidence of initiating use of these drugs during the years after high school.

In analyses published elsewhere, we looked closely at patterns of change in drug use with age and identified some post-high school experiences that contribute to declining levels of annual or current use of drugs as respondents grow older. For example, the likelihood of marriage increases with age, and we have found that marriage is consistently associated with declines in *alcohol* use in general, *heavy drinking*, *marijuana* use, and *cocaine* use.³⁵

- For the use of *any illicit drug* (Figure 4-1), lifetime prevalence is 59% among 29- to 30-year-olds versus 51% among the 2004 12th graders. Annual prevalence, however, is highest among the 12th graders (39%) with progressively lower rates among the older age groups, reaching 22% among the 29- to 30-year-olds. Current (30-day) prevalence shows much the same pattern, with 12th graders having the highest rate (23%) and the rate declining gradually for each of the older age groups, reaching 11% among the 29- to 30-year-olds.
- Interestingly enough, the annual and 30-day prevalence rates found among the 35-, 40-, and 45-year-olds for *marijuana*, *any illicit drug*, and *any illicit drug other than marijuana* are all virtually identical to the rates observed among the 29- to 30-year-olds. (This is also true for many of the other specific illicit drugs.) Yet more—and sometimes substantially more—of the 35-, 40-, and 45-year-old cohorts (the classes of 1987, 1982, and 1977) have reported

³⁵Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates; and Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Schulenberg, J., O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (2000). "Spread your wings and fly": The course of well-being and substance use during the transition to young adulthood. In L. J. Crockett & R. K. Silbereisen (Eds.), *Negotiating adolescence in times of social change* (pp. 224-255). New York: Cambridge University Press. And see O'Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2004). Studying the transition from youth to adulthood: Impacts on substance use and abuse. In J. S. House, F. T. Juster, R. L. Kahn, H. Schuman, and E. Singer (Eds.), *A telescope on society: Survey research and social science at the University of Michigan and beyond* (pp. 305-329). Ann Arbor: The University of Michigan Press.

some use of marijuana and other illicit drugs in their lifetime than had the 29- to 30-year-old cohorts (the classes of 1992 and 1993). Thus, greater proportions of the older cohorts have discontinued use, but current use remains similar among the groups because a higher percentage of the 35- to 45-year-olds had used these drugs earlier in their lives.

- Among the young adults, a similar pattern exists for *marijuana*: a higher lifetime prevalence as a function of age, but considerably lower annual and 30-day prevalence rates through the late 20s (Figure 4-3). Current *daily marijuana* use shows the least variation across age (as shown in the next chapter in Figure 5-3c). Still, in 2004 it ranges from 5.6% among 12th graders to 2.0% among 29- to 30-year-olds. Daily use in 2004 ranges from 1.8% to 2.5% for 35-, 40-, and 45-year-olds.
- Statistics on the use of *any illicit drug other than marijuana* (Figure 4-2) have a similar pattern. Like marijuana and the any-illicit-drug-use index, corrected lifetime rates on this index also show an appreciable rise with age level, reaching 41% among the 29- to 30-year-old age group and 71% among the 45-year-olds. Current use shows a decline across the age bands, ranging from 11% among 12th graders to 5% among 29- to 30-year-olds. After ages 21-22, annual use is lower with increased age of the respondent through age 35 and then remains fairly constant through age 45. A number of the individual drugs that comprise this general category show lower rates of use at higher ages for annual prevalence, usually with the highest rate observed at age 18 or ages 19-20. This is particularly true for *amphetamines*, *hallucinogens*, *LSD* specifically, *inhalants*, *sedatives (barbiturates)*, *narcotics other than heroin*, and *tranquilizers*. The falloff with age is not as great nor as consistent for *cocaine*, *crack*, *other cocaine*, *ice*, *heroin*, and *MDMA (ecstasy)*, though in general, usage rates are somewhat lower among those in their 30s than among those in their early 20s. Several classes of drugs are discussed individually next.
- *Inhalants* show some very interesting differences across the age strata (see Figure 4-12). There is little difference across age in contemporaneously reported lifetime prevalence but a considerable difference in the lifetime prevalence figure *adjusted* for previous reporting of use. The adjusted pattern—an increase with age—is the one we have come to expect, and we suspect is the more accurate one. Annual prevalence rates drop off with age, while 30-day rates begin fairly low and can drop only a little. Clearly, the use of inhalants is extremely low beyond about age 20; and we know from data presented in Volume I that much of the decline in use with age already has occurred by the time young people have reached 12th grade. Questions on inhalant use are not included in the surveys administered to respondents over the age of 30.
- For *amphetamines*, lifetime prevalence is again much higher among the older age groups—reflecting the addition of new users who initiate use in their 20s but also reflecting some cohort differences (Figure 4-4). (There is also a considerable divergence between the corrected lifetime prevalence versus the contemporaneously reported lifetime prevalence, as is true for most of the psychotherapeutic drugs.) However, more recent use as reflected in the annual prevalence figure is lower among the older age groups. This has not always been true;

the present pattern is the result of a sharper decline in use among older respondents than has occurred among 12th graders. These trends are discussed in the next chapter.

- **Ritalin**, a particular amphetamine widely prescribed for the treatment of attention-deficit hyperactivity disorder (ADHD), shows a drop-off in annual prevalence of more than half after age 24, consistent with the interpretation that it is often used by college students trying to stay awake and alert for studying and completing assignments (see Table 4-3 and Figure 4-5).
- **Methamphetamine** use declines with age, with annual prevalence starting at 4.1% among the 19- to 20-year-olds, and then declining fairly steadily to 1.2% by age 29 to 30 (Table 4-3).
- Questions on the use of **crystal methamphetamine (ice)** are contained in two of the six questionnaire forms, making the estimates less reliable than those based on all six forms. (Ice use is not asked of the 35-, 40-, or 45-year-old respondents.) Among the 19- to 30-year-old respondents *combined*, 1.4% reported some use in the prior year—lower than the 2.1% reported by 12th graders (see Table 4-1 and Figure 4-17).
- **Sedatives (barbiturates)** show lifetime prevalence rates that are fairly similar across the age band 18 through 30 but are appreciably higher among 40- and 45-year-olds. Above age 24, however, annual use falls appreciably with age (Figure 4-13). At present, current usage rates are quite low in all age groups; therefore, 30-day use varies rather little by age. Because of the substantial long-term decline in sedative (barbiturate) use over the life of the study, the 45-year-olds have, by far, the highest adjusted lifetime prevalence rate (28%).³⁶
- The use of **narcotics other than heroin** (Figure 4-14) shows age differences that are similar in some ways to those seen for sedatives (barbiturates). Lifetime prevalence increases some across the early 20s, declines some across the late 20s, then is fairly constant through age 40. The 45-year-olds show the highest lifetime prevalence rate (28%, adjusted). Annual prevalence is highest in the 18 to 24 age range (at 9% to 10%), declining to 5% among 29- to 30-year-olds, then falling a bit lower in the older age bands. Thirty-day prevalence shows a similar profile across age to that for annual prevalence.³⁷
- **Tranquilizer** use shows an increase with age in lifetime prevalence—again, with the oldest cohorts having much higher levels of lifetime use—and some modest decrease with age in annual prevalence. Thirty-day prevalence is fairly flat across all age groups (Figure 4-15).

³⁶Barbiturates were the dominant form of sedatives in use when these questions were first introduced. In the intervening years, a number of non-barbiturate sedatives have entered the market and largely displaced barbiturate sedatives. Because our question did not change, we believe that a number of users of non-barbiturate sedatives are reporting them in answer to the barbiturate question, which also defines them in terms of the conditions for which they are prescribed. In recognition of this fact, we will now label them as “sedatives (barbiturates),” though to date the question specifies “barbiturates.”

³⁷In 2002 the question text for **narcotics other than heroin** was changed on three of the six questionnaire forms in order to update the list of examples of narcotics other than heroin. Talwin, laudanum, and paregoric—each of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. As a consequence of this revision, reported use rates increased in 2002 in the half of the questionnaire forms using the new question wording; however, it did not increase in those forms using the original wording, as is discussed in the next chapter. Using data from the three unchanged questionnaire forms, we derived a best guess as to the actual change in use. We added that change score to the 2001 prevalence rate that would have been observed had we not changed the measures. This adjusted value is provided in the relevant tables and figures.

- **Cocaine** generally had presented a unique case among the illicit drugs, in that lifetime, annual, and current prevalence rates have *all* tended to be higher among the older age groups. By 1994, however, 30-day cocaine use had reached such low levels that it varied rather little by age. Following the resurgence of cocaine use in the 1990s, some differences by age in annual prevalence emerged, though there are still rather few differences for current prevalence (Figure 4-6). Annual prevalence is now highest among those ages 21 through 24, who were 12th graders when the increase in cocaine use occurred. The cohort differences in lifetime cocaine use are particularly vivid, with the 40- and 45-year-olds showing 38% and 44% adjusted lifetime prevalence rates, respectively, compared to 15% among 21- to 22-year-olds in 2004.
- In 2004, lifetime prevalence of **crack** use (Figure 4-7) is fairly similar among ages 21 through 35. (Eighteen- through 20-year-olds have lower rates.) Lifetime prevalence is higher among 40- and 45-year-olds, no doubt reflecting something of a cohort effect due to the rather transient popularity of crack in the early to mid-1980s. Current prevalence is below 1% in all age groups. Annual prevalence is highest among 45-year-olds, again reflecting a lasting cohort effect; it is next highest among 18-year-olds and then falls through age 25, before leveling.

We believe that the omission of high school dropouts is likely to have a greater-than-average impact on the prevalence estimates for crack. It also seems likely that any members of the panels who are dependent on crack (or other illicit drugs like heroin) would be less likely than average to respond to the questionnaires; therefore, such extreme users are no doubt underrepresented among the panel respondents.

- **MDMA (ecstasy)** was added to two of the six forms of the follow-up surveys in 1989 to assess how widespread its use had become among young adults. It was added to a third form in 2002. Questions about its use were not asked of high school students until 1996, primarily because we were concerned that its alluring name might have the effect of stimulating interest. We were less concerned about such an effect after the name of the drug had become more widely known. (MDMA use is not asked of the 35-, 40-, or 45-year-old respondents.)

Among all 19- to 30-year-olds combined, 15% say they have ever tried **MDMA**; among 12th graders, 8% say they have used it. The age differences are quite dramatic for this drug, with lifetime prevalence now highest at ages 23-24 and generally declining with age thereafter (see Figure 4-16). This very likely reflects the fact that ecstasy use rose very rapidly between 1997 and 2001, and then declined quite sharply; therefore, recent graduating classes have less use than their predecessors, and much earlier classes had less use than their successors. Annual prevalence is fairly flat at around 4% from age 18 to age 28, which reflects an extension of the age of highest involvement from what it was in earlier years. Past-month ecstasy use, though, remains concentrated among those in their late teens, through age 26, with lower rates among the older age groups.

- In the case of **alcohol**, all prevalence rates are higher among those of post-high school age than among those in high school, and they generally increase for the first three to five years after high school, through age 23 or 24 (Figures 4-20a and b). After that, prevalence rates

vary only modestly among the different age groups. Lifetime prevalence changes very little after ages 23 to 24, due in large part to a “ceiling effect.” Current (30-day) alcohol use is considerably higher among those ages 23-24 (73%) than among 12th graders (48%); it drops some through age 30 (65%) and is at about the same level among those ages 35, 40, and 45. Current *daily drinking* (Figure 4-20b) also is slightly higher among those 21-24 years old (6%) and then is progressively lower through ages 27 to 28 (4%). The highest rate of daily drinking is among those aged 45 (9%).

- Among the various measures of alcohol consumption, *occasions of heavy drinking* in the two weeks prior to the survey show large differences among the age groups (Figure 4-20b). There is a fair difference between 18-year-olds (29%) and 21- to 22-year-olds, who have the highest prevalence of such heavy drinking (42%). Then, there is a falloff at each subsequent age level above age 24, reaching 27% by ages 29 to 30. We have interpreted this curvilinear relationship as reflecting an age effect—and not a cohort effect—because it seems to replicate across different graduating class cohorts and also because it has been linked directly to age-related events such as leaving the parental home (which increases heavy drinking) and marriage (which decreases it), both of which are, in turn, related to attending college.³⁸ Among those aged 35 to 45, about one-fifth (19% to 22%) report such heavy drinking in the prior two-week interval.
- *Cigarette smoking* also shows an unusual pattern of age-related differences (Figure 4-21). Current (30-day) smoking used to be about the same rate among those in their early 20s as among 12th graders, in part reflecting the fact that relatively few new people are recruited to smoking after high school. In 2004, however, current smoking is somewhat lower among 12th graders (25%) than among the next several age bands (peaking at 32% among 23- to 24-year-olds), almost surely due to the sharp drop in smoking that has been occurring among secondary school students—a cohort effect. On the other hand, smoking at heavier levels—such as smoking half a pack daily—is (and has been) somewhat higher among those in their 20s than among 12th graders, reflecting, at least in part, the fact that many previously light or moderate smokers move into a pattern of heavier consumption after high school.³⁹ While about a third (32%) of the current smokers in high school (seniors) smoke at the rate of a half-pack per day or more, well over one-half (57%) of the current smokers in the 29 to 30 age group do so.
- Questions about use of *steroids* were added in 1989 to one form only (and to an additional form in 1990), making it difficult to determine age-related differences with much accuracy due to the limited sample sizes. (Steroids are not asked of the 35-, 40-, or 45-year-old

³⁸O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976-1986. *American Journal of Public Health*, 78, 1315-1321 See also Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

³⁹Because age is confounded with class cohort, and because we have established that cigarette smoking shows strong cohort effects (enduring differences among cohorts), one must be careful in interpreting age-related differences in a cross-sectional sample as if they were due only to age effects—that is, changes with age consistently observable across cohorts. However, multivariate analyses conducted on panel data from multiple cohorts do show a consistent age effect of the type mentioned here (see O'Malley, Bachman, & Johnston, 1988, in previous footnote).

respondents.) Overall, 1.8% of 19- to 30-year-olds in 2004 reported having used steroids in their lifetime. Annual and 30-day use levels were very low, at 0.4% and 0.1%, respectively. (See Tables 4-2 through 4-4.) The rates among 12th graders tend to be considerably higher than the rates among older age groups, reflecting possibly both age and cohort effects. (As described in Volume I, the prevalence of steroid use among 12th graders rose sharply between 2000 and 2002. At present, the highest annual use among young adults is among 19- to 20-year-olds, at 0.8%.)

In sum, lifetime prevalence rates in some of the older age groups studied here, who passed through adolescence in the heyday of the drug epidemic, show impressively high lifetime rates of illicit drug use—particularly when lifetime prevalence is corrected for the recanting of earlier reported use. However, the *current* use of most illicit drugs is substantially lower among those in their 30s and 40s than among those in their late teens to early 20s. For the two licit drugs, alcohol and cigarettes, the picture is a more complicated one. Steroids also present a more complicated picture.

PREVALENCE COMPARISONS FOR SUBGROUPS OF YOUNG ADULTS

Subgroup differences for the group of young adults 1 to 12 years beyond high school (corresponding to modal ages 19 to 30) are presented in Tables 4-1 through 4-5. While Table 4-1 provides only gender differences, the remaining tables have prevalence rates by gender, age, region of the country, and population density. Each of these dimensions is discussed separately below.

Gender Differences

In general, most of the gender differences in drug use that pertained in high school may be found in the young adult sample as well.

- Among young adults, more males than females report using *any illicit drug* during the prior year (35% versus 29%). Males have higher annual prevalence rates for nearly all of the specific illicit drugs—with ratios greater than 2 for *steroids* and *inhalants*. For example, among the 19- to 30-year-olds, steroids were used by 0.6% of males versus 0.2% of females during the prior 12 months. (See Table 4-1.)
- All forms of *cocaine* were used in the past year by more males than females (19- to 30-year-olds). Annual *cocaine* use was reported by 8.7% of the males and 5.0% of the females, *crack* use by 1.6% of the males and 0.9% of the females, and *other cocaine* use by 7.8% of the males and 4.7% of the females.
- Other large gender differences among the 19- to 30-year-olds are found in *daily marijuana* use (6.3% for males versus 3.3% for females in 2004), *daily alcohol* use (7.6% versus 2.2%), and occasions of drinking *five or more drinks in a row* in the prior two weeks (47% versus 28%). This gender difference in occasions of heavy drinking is even greater among young adults than among 12th graders, where it is 34% for males versus 24% for females.

- **MDMA (ecstasy)** use is only slightly higher among males than among females in the young adult sample overall (annual prevalence 3.5% versus 2.8%, respectively).
- The use of **narcotics other than heroin** is fairly close, at 9.9% annual prevalence for males vs. 7.3% for females. Use of **Vicodin**, one of the most widely used drugs in the class, is also fairly close (10.8% versus 7.1%). There is a larger gender contrast for **OxyContin** (annual prevalence of 3.9% for males versus 2.0% for females).
- The use of **amphetamines**, which is now about equivalent among males and females in high school, is also fairly similar for both genders in this post-high school period (annual prevalence of 6.1% versus 5.3%, respectively).
- **Crystal methamphetamine (ice)** is used by small, and roughly similar, percentages of both genders (1.8% annual prevalence for males versus 1.2% for females).
- In the 1980s, there were few differences between males and females in rate of **cigarette** use. By the early 1990s, however, males had slightly higher rates of use. Now, there again appear few gender differences among 12th graders: past month prevalence in 2004 is 25% for males, compared to 24% for females; daily use rates are 15% for both males and females; and half-pack or more use rates are 8.0% for males and 7.5% for females. But among older respondents, the slight gender difference remains: among the 19- to 30-year-olds, males are more likely to have smoked in the past month (32% versus 25%), to have smoked daily (23% versus 18%), and to have smoked half a pack or more per day (16% versus 12%).
- **Steroid** use among young adults is much more prevalent among males than females, as is true for 12th graders. Among 12th graders, 3.3% of the males reported steroid use in the past year versus 1.7% of the females. These statistics are much lower among the 19- to 30-year-olds, but use by males remains considerably higher (0.6% for males versus 0.2% for females).

Regional Differences

Follow-up respondents are asked in what state they currently reside. States are then grouped into the same regions used in the analysis of the high school data.⁴⁰ Tables 4-2 through 4-5 present regional differences in lifetime prevalence, annual prevalence, 30-day prevalence, and current daily prevalence, for the 19- to 30-year-olds combined.

- There exist some regional differences in the use of **marijuana**, with the Northeast and the West somewhat higher than the South and North Central. They are also somewhat higher in the proportion using **any illicit drug** and **any illicit drug other than marijuana** (see Table 4-3).

⁴⁰States are grouped into regions as follows: **Northeast**—Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania; **North Central**—Ohio, Indiana, Illinois, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas; **South**—Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas; **West**—Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, and California.

- **Methamphetamine** use is clearly higher in the West (annual prevalence of 4.6%) than in the South (2.4%), North Central (2.3%), and the Northeast (0.8%) (see Table 4-3).
- The use of **crystal methamphetamine (ice)** by 19- to 30-year-olds remains considerably higher in the West, which has a 2.9% annual prevalence rate in 2004, than in the North Central (1.2%), the South (1.1%), and the Northeast (0.6%). Among 12th graders, the West also has a higher rate of use than the other three regions.
- The West and Northeast have higher rates than the other two regions for **hallucinogen** use, though the regional differences are not large.
- Regional differences in **MDMA (“ecstasy”)** use among young adults are not large at present: use is highest in the South (3.8% annual prevalence) and the West (3.2%) and slightly lower in the Northeast and the North Central (3.0% and 1.9%, respectively).
- **OxyContin** use is higher in the Northeast (3.4%) than in the other regions of the country (2.5% to 2.7%); while **Vicodin** use is highest in the West (12.4%) and lower in the North Central (9.8%), the Northeast (8.0%), and the South (5.5%).
- For the remaining illicit drugs, regional differences are not substantial (see Tables 4-3 and 4-4).
- Prevalence rates for **alcohol** are typically somewhat higher in the Northeast and North Central regions than in the South and West regions; this pattern has generally been true among 12th graders, as well. For **binge drinking**, the Northeast and North Central have prevalence rates of 39% and 40%, respectively, whereas the South and West have rates of 31% and 32%.
- As with alcohol, **cigarette smoking** among young adults is highest in the Northeast and North Central. It is lowest in the West. This difference is most pronounced at the half-pack-a-day level (Table 4-5), where the rate in the West (8.5%) is almost half the rate in the North Central (16.6%).

Population Density Differences

Population density is measured by asking respondents to select the response category that best describes the size and nature of the community where they lived during March of the year in which they were completing the follow-up questionnaire. Various categories are listed in Tables 4-2 through 4-5, and the population sizes given to the respondent to help define each level are provided in a footnote to each table. An examination of the 1987 and 1988 drug use data for the two most urban strata revealed that the modest differences in prevalence rates between the suburbs and the corresponding cities were not worth the complexity of reporting them separately; accordingly, these categories have been merged since then. See Tables 4-3 through 4-5 for the relevant tabular results that are discussed below.

- Differences in illicit drug use by population density tend to be very modest, perhaps more modest than is commonly supposed. Among the general population, use of most illicit drugs is fairly broadly distributed among all areas from rural to urban. To the extent that there are variations, almost all of the associations are positive, with rural/country areas having the lowest levels of use, and small towns having the next lowest. Medium-sized cities, large cities, and very large cities tend to be higher, with only small variations among these three categories. Positive associations with population density exist for annual prevalence of **any illicit drug, marijuana, and MDMA (ecstasy)**. The association is strongest for ecstasy, where the annual prevalence rate in the large cities (3.9%) is about twice that in the rural areas (2.0%).
- **Methamphetamine** use shows a higher rate of annual prevalence in the farm/country stratum (4.6%) than in the other strata, which range between 2.0% and 2.6% (Table 4-3). Use was highest in that stratum in 2003, as well.
- Among young adults, the lifetime, annual, and 30-day **alcohol** use measures all show a positive association with population density. **Occasions of heavy drinking** are fairly similar across all strata, although there is some positive ordinal association with urbanicity (see Table 4-5). **Daily** alcohol use falls between 3.9% and 5.0% for all community size strata, with no discernible association.
- Contrary to what we find for almost all other substances, there exists a **negative** association between population density and **daily cigarette smoking**, which is highest in the farm/country stratum and lowest in the large cities (daily prevalence rates of 25% and 17%, respectively). Smoking at the half-pack-a-day level is almost twice as high in farm/country areas (18%) as in very large cities (10%). (See Table 4-5.)

TABLE 4-1
Prevalence of Use of Various Types of Drugs by Gender, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

<i>Approx. Weighted N =</i>	<u>Males</u> 2800	<u>Females</u> 4000	<u>Total</u> 6800
Any Illicit Drug ^a			
Annual	35.4	29.3	31.8
30-Day	21.8	15.1	17.8
Any Illicit Drug ^a Other Than Marijuana			
Annual	20.1	15.9	17.6
30-Day	9.8	6.6	7.9
Marijuana			
Annual	31.5	24.0	27.0
30-Day	19.2	12.4	15.2
Daily	6.3	3.3	4.5
Inhalants ^{b,c}			
Annual	2.3	1.1	1.5
30-Day	0.3	0.3	0.3
Hallucinogens ^b			
Annual	6.1	2.9	4.2
30-Day	1.2	0.6	0.8
LSD			
Annual	1.3	0.4	0.8
30-Day	0.1	0.1	0.1
PCP ^d			
Annual	0.1	0.1	0.1
30-Day	0.1	0.1	0.1
MDMA (Ecstasy) ^c			
Annual	3.5	2.8	3.1
30-Day	0.9	0.3	0.6
Cocaine			
Annual	8.7	5.0	6.5
30-Day	2.9	1.5	2.0
Crack ^e			
Annual	1.6	0.9	1.2
30-Day	0.4	0.2	0.3
Other Cocaine ^f			
Annual	7.8	4.7	6.0
30-Day	2.7	1.5	2.0
Heroin			
Annual	0.5	0.2	0.3
30-Day	0.2	0.1	0.1
Other Narcotics ^g			
Annual	9.9	7.3	8.4
30-Day	3.3	2.5	2.8

(Table continued on next page)

TABLE 4-1 (cont.)
Prevalence of Use of Various Types of Drugs by Gender, 2004
Among Respondents of Modal Ages 19-30

(Entries are percentages)

<i>Approx. Weighted N =</i>	<u>Males</u> 2800	<u>Females</u> 4000	<u>Total</u> 6800
Amphetamines, Adjusted ^{g,h}			
Annual	6.1	5.3	5.6
30-Day	2.4	2.1	2.2
Methamphetamine ⁱ			
Annual	3.1	2.1	2.5
30-Day	0.6	0.6	0.6
Crystal Methamphetamine (Ice) ⁱ			
Annual	1.8	1.2	1.4
30-Day	0.6	0.2	0.4
Sedatives (Barbiturates) ^g			
Annual	4.7	3.5	4.0
30-Day	1.9	1.4	1.6
Tranquilizers ^g			
Annual	8.0	6.2	7.0
30-Day	3.0	2.2	2.5
Alcohol			
Annual	84.9	83.8	84.3
30-Day	72.9	64.1	67.7
Daily	7.6	2.2	4.4
5+ Drinks in a Row in the Last 2 Weeks	46.5	27.7	35.4
Flavored Alcoholic Beverages ^d			
Annual	53.9	66.7	61.5
30-Day	23.3	31.8	28.4
Cigarettes			
Annual	41.4	34.7	37.4
30-Day	32.0	25.3	28.0
Daily	22.8	18.3	20.1
Half-Pack or More per Day	15.7	11.7	13.3
Steroids ⁱ			
Annual	0.6	0.2	0.4
30-Day	0.1	0.0	0.1

Source: The Monitoring the Future Study, the University of Michigan.

*' indicates a prevalence rate of less than 0.05% but greater than true zero.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bUnadjusted for known underreporting of certain drugs. See text for details.

^cThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3400.

^dThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^eThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5600.

^fThis drug was asked about in four of the six questionnaire forms. Total N is approximately 4600.

^gOnly drug use that was not under a doctor's orders is included here.

^hBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

ⁱThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2300.

TABLE 4-2
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30

(Entries are percentages)

	<i>Approx. Weighted N</i>	Any Illicit Drug ^a	Any Illicit Drug ^a Other Than MJ	Marijuana	Inhalants ^{b,c}	Hallucinogens ^b	LSD	PCP ^d	MDMA ^c	Cocaine
Total	6800	60.3	34.6	57.4	11.8	19.3	13.7	2.6	15.1	14.9
Gender:										
Male	2800	62.2	37.1	60.5	14.7	23.4	16.4	3.1	16.8	18.9
Female	4000	58.9	32.9	55.3	9.8	16.5	11.8	2.2	14.0	12.2
Modal Age:										
19-20	1200	52.9	29.2	50.0	9.3	12.0	6.4	1.1	10.3	10.6
21-22	1200	58.5	33.1	55.9	10.5	17.1	9.7	1.2	14.4	14.3
23-24	1100	64.5	40.1	60.9	12.0	22.5	15.2	4.3	20.1	18.0
25-26	1100	63.5	37.0	60.5	12.3	22.8	18.1	4.1	19.4	16.3
27-28	1100	64.1	37.4	60.7	14.5	23.4	18.8	3.3	16.5	17.1
29-30	1100	59.1	31.8	57.4	12.8	19.2	15.1	1.9	10.4	13.8
Region:										
Northeast	1300	65.0	36.8	61.8	11.6	21.0	14.7	3.9	16.6	15.0
North Central	1900	59.6	32.9	56.9	12.3	18.2	13.3	1.4	12.3	14.0
South	2300	57.3	33.0	53.7	10.3	17.0	12.7	2.3	14.6	13.6
West	1300	61.6	38.0	59.8	13.6	23.4	15.0	3.8	18.4	18.4
Population Density:^e										
Farm/Country	800	53.7	30.3	50.2	11.6	15.4	12.0	2.7	11.4	13.2
Small Town	1800	58.3	33.7	55.2	12.5	16.9	13.0	1.9	12.3	14.2
Medium City	1600	59.7	33.2	56.6	11.1	18.6	13.5	2.5	13.0	14.0
Large City	1600	62.1	35.7	59.3	12.0	21.1	14.0	2.6	18.8	15.7
Very Large City	1000	65.7	39.4	64.0	10.6	24.4	15.5	4.2	20.4	17.3

Source: The Monitoring the Future Study, the University of Michigan.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bUnadjusted for known underreporting of certain drugs. See text for details.

^cThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3400.

^dThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.

(Table continued on next page)

TABLE 4-2 (cont.)
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Crack ^a	Heroin	Other Narcotics ^b	Amphetamines ^{b,c}	Methamphetamine ^d	Ice ^d	Sedatives (Barbiturates) ^b	Tranquilizers ^b	Alcohol	Flavored Alcoholic Beverages ^e	Cigarettes	Steroids ^d
Total	6800	4.2	1.8	16.7	15.3	9.0	4.8	9.1	14.3	89.9	83.9	NA	1.8
Gender:													
Male	2800	5.4	2.5	19.3	16.5	10.4	6.0	10.6	16.3	89.4	79.1	NA	3.7
Female	4000	3.5	1.4	14.9	14.5	8.0	4.0	8.1	13.0	90.2	87.1	NA	0.5
Modal Age:													
19-20	1200	2.8	1.1	15.7	15.1	7.5	4.8	9.2	11.5	80.4	75.1	NA	2.2
21-22	1200	4.0	1.3	17.1	14.6	7.8	4.7	9.3	14.6	90.1	80.2	NA	1.9
23-24	1100	4.8	2.1	20.2	18.3	10.8	3.2	10.4	17.1	91.0	85.9	NA	1.6
25-26	1100	4.6	3.0	17.8	16.0	8.8	5.5	10.2	16.3	92.9	88.0	NA	2.2
27-28	1100	4.8	2.1	17.2	15.7	10.2	5.5	9.2	15.4	93.8	88.1	NA	1.6
29-30	1100	4.6	1.5	12.1	12.3	9.1	5.2	6.5	11.2	92.2	87.4	NA	1.2
Region:													
Northeast	1300	3.7	2.6	18.7	14.4	4.5	1.7	10.1	14.4	92.1	88.2	NA	1.8
North Central	1900	4.0	1.3	16.8	15.5	8.9	4.7	7.7	12.5	91.9	86.5	NA	1.7
South	2300	3.9	1.9	15.0	15.1	8.6	4.1	9.6	16.4	88.5	81.5	NA	2.0
West	1300	5.9	1.6	17.4	16.3	13.8	9.1	9.2	13.4	87.2	81.2	NA	1.6
Population Density:^f													
Farm/Country	800	4.9	1.2	15.4	13.8	11.0	6.8	8.1	13.0	89.4	86.7	NA	1.3
Small Town	1800	4.6	1.7	16.2	16.6	8.4	3.9	9.0	13.3	89.3	81.5	NA	1.0
Medium City	1600	4.8	1.7	16.6	14.8	7.9	3.8	9.8	14.0	90.0	83.8	NA	2.6
Large City	1600	3.5	2.3	17.2	14.6	10.0	5.2	8.9	15.3	89.5	85.4	NA	2.6
Very Large City	1000	3.4	1.9	17.6	16.0	8.4	5.8	9.5	15.6	91.6	84.3	NA	0.8

Source: The Monitoring the Future Study, the University of Michigan.

^a'NA' indicates data not available.

^bThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5600.

^cOnly drug use that was not under a doctor's orders is included here.

^dBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^eThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2300.

^fThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^gA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents. Within each level of population density, suburban and urban respondents are combined.

TABLE 4-3
Annual Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Any Illicit Drug ^a	Any Illicit Drug ^a Other Than MJ	Marijuana	Inhalants ^{b,c}	Hallucinogens ^b	LSD	PCP ^d	MDMA ^c	Cocaine
Total	6800	31.8	17.6	27.0	1.5	4.2	0.8	0.1	3.1	6.5
Gender:										
Male	2800	35.4	20.1	31.5	2.3	6.1	1.3	0.1	3.5	8.7
Female	4000	29.3	15.9	24.0	1.1	2.9	0.4	0.1	2.8	5.0
Modal Age:										
19-20	1200	38.0	20.2	34.5	3.1	6.3	1.5	0.2	4.2	6.3
21-22	1200	36.5	21.2	32.5	2.1	6.7	1.4	0.2	2.4	8.6
23-24	1100	34.5	21.2	27.7	1.6	4.4	0.7	0.2	3.2	8.4
25-26	1100	31.6	16.3	27.6	1.2	3.2	0.5	0.0	4.0	6.7
27-28	1100	26.8	14.6	22.4	0.3	2.6	0.3	0.0	3.7	5.2
29-30	1100	22.0	11.8	16.4	0.6	1.4	0.2	0.0	0.9	3.6
Region:										
Northeast	1300	36.2	18.4	31.7	2.0	4.5	1.4	0.2	3.0	7.7
North Central	1900	31.1	16.3	26.6	1.0	3.6	0.6	0.3	1.9	6.1
South	2300	28.1	17.0	22.9	1.7	3.3	0.7	0.0	3.8	5.6
West	1300	34.9	19.9	30.2	1.5	6.0	0.5	0.0	3.2	7.4
Population Density:^e										
Farm/Country	800	24.2	16.0	18.7	0.8	3.3	0.8	0.0	2.0	5.5
Small Town	1800	31.1	17.1	26.4	1.6	4.1	0.9	0.1	2.4	6.5
Medium City	1600	31.5	16.4	27.2	1.9	4.0	0.7	0.2	2.8	5.7
Large City	1600	33.4	19.3	28.1	1.7	4.2	0.7	0.0	3.9	6.6
Very Large City	1000	36.0	19.0	31.9	1.2	4.9	0.9	0.3	3.6	8.0

Source: The Monitoring the Future Study, the University of Michigan.

^a'*' indicates a percentage of less than 0.05% but greater than true zero.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bUnadjusted for known underreporting of certain drugs. See text for details.

^cThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3400.

^dThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.

(Table continued on next page)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Crack ^a	Heroin	Other Narcotics ^b	OxyContin ^c	Vicodin ^c	Amphetamines ^{b,d}	Ritalin ^{b,e}	Methamphetamine ^c	Ice ^c
Total	6800	1.2	0.3	8.4	2.8	8.6	5.6	2.4	2.5	1.4
Gender:										
Male	2800	1.6	0.5	9.9	3.9	10.8	6.1	3.7	3.1	1.8
Female	4000	0.9	0.2	7.3	2.0	7.1	5.3	1.6	2.1	1.2
Modal Age:										
19-20	1200	1.6	0.4	10.4	3.6	9.6	8.5	3.6	4.1	2.2
21-22	1200	1.8	0.5	9.2	2.8	10.1	6.7	3.3	3.5	1.8
23-24	1100	1.2	0.3	9.5	3.8	10.5	7.1	3.5	2.6	0.6
25-26	1100	1.1	0.2	7.9	2.4	8.1	4.6	1.5	1.6	1.7
27-28	1100	0.8	0.1	7.5	2.6	6.1	3.9	1.6	1.7	1.2
29-30	1100	0.6	0.3	5.4	1.2	6.7	2.2	0.8	1.2	0.9
Region:										
Northeast	1300	1.3	0.4	9.2	3.4	8.0	4.4	2.4	0.8	0.6
North Central	1900	1.1	0.3	8.4	2.6	9.8	5.6	2.1	2.3	1.2
South	2300	0.8	0.2	7.5	2.5	5.5	6.0	2.8	2.4	1.1
West	1300	1.8	0.3	9.1	2.7	12.4	5.9	2.0	4.6	2.9
Population Density:^e										
Farm/Country	800	0.8	0.1	7.8	3.4	9.0	5.4	2.2	4.6	1.9
Small Town	1800	1.5	0.4	8.7	2.5	7.0	6.4	1.9	2.0	1.4
Medium City	1600	1.5	0.4	7.8	3.0	8.7	5.2	2.7	2.6	1.3
Large City	1600	0.9	0.2	8.7	3.2	9.2	5.5	2.6	2.2	1.0
Very Large City	1000	0.8	0.3	8.4	2.0	9.6	4.9	3.0	2.1	1.7

Source: The Monitoring the Future Study, the University of Michigan.

^a'*' indicates a percentage of less than 0.05% but greater than true zero.

^bThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5600.

^cOnly drug use that was not under a doctor's orders is included here.

^dThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2300.

^eBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^fA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.

(Table continued on next page)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Sedatives (Barbiturates) ^a	Tranquilizers ^a	Rohypnol ^b	GHB ^b	Ketamine ^b	Alcohol	Flavored Alcoholic Beverages ^c	Cigarettes	Steroids ^b
Total	6800	4.0	7.0	0.1	0.4	0.6	84.3	61.5	37.4	0.4
Gender:										
Male	2800	4.7	8.0	0.2	0.7	1.0	84.9	53.9	41.4	0.6
Female	4000	3.5	6.2	*	0.2	0.3	83.8	66.7	34.7	0.2
Modal Age:										
19-20	1200	6.0	8.0	0.3	0.1	1.3	75.2	59.0	39.7	0.8
21-22	1200	4.4	8.1	0.0	0.9	0.9	86.8	64.7	43.1	0.2
23-24	1100	5.0	8.3	0.0	0.9	0.0	87.2	65.0	41.3	0.5
25-26	1100	3.3	6.8	0.0	0.3	0.4	87.9	62.5	37.4	0.5
27-28	1100	2.8	5.6	0.4	0.4	0.6	86.1	62.6	33.1	0.3
29-30	1100	1.9	4.8	0.0	0.1	0.4	83.5	55.0	29.1	0.0
Region:										
Northeast	1300	3.8	6.8	0.1	0.5	0.9	87.9	58.7	39.2	0.2
North Central	1900	3.4	6.5	0.0	0.3	0.3	87.7	67.5	40.2	0.1
South	2300	4.5	8.2	0.2	0.5	0.8	81.1	58.9	36.0	0.5
West	1300	4.0	5.7	0.2	0.6	0.4	81.5	61.0	33.9	0.5
Population Density:^d										
Farm/Country	800	3.0	5.5	0.4	0.4	0.4	80.5	59.3	39.7	0.5
Small Town	1800	4.4	6.9	0.0	0.1	0.1	82.3	61.8	39.5	0.2
Medium City	1600	4.2	6.5	0.3	0.4	1.1	85.2	62.4	36.2	0.4
Large City	1600	3.6	8.2	0.0	0.1	0.7	85.3	61.4	35.7	0.6
Very Large City	1000	4.3	7.0	0.1	1.7	0.5	87.8	60.8	36.0	0.0

Source: The Monitoring the Future Study, the University of Michigan.

*' indicates a percentage of less than 0.05% but greater than true zero.

^aOnly drug use that was not under a doctor's orders is included here.

^bThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2300.

^cThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^dA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.

TABLE 4-4
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30

(Entries are percentages)

	<i>Approx. Weighted N</i>	Any Illicit Drug ^a	Any Illicit Drug ^a Other Than MJ	Marijuana	Inhalants ^{b,c}	Hallucinogens ^b	LSD	PCP ^d	MDMA ^c	Cocaine
Total	6800	17.8	7.9	15.2	0.3	0.8	0.1	0.1	0.6	2.0
Gender:										
Male	2800	21.8	9.8	19.2	0.3	1.2	0.1	0.1	0.9	2.9
Female	4000	15.1	6.6	12.4	0.3	0.6	0.1	0.1	0.3	1.5
Modal Age:										
19-20	1200	23.0	10.4	20.7	0.6	1.6	0.3	0.2	1.2	2.2
21-22	1200	20.9	9.8	18.3	0.5	1.5	0.2	0.2	0.7	2.7
23-24	1100	18.8	8.8	15.6	0.0	0.5	*	0.0	0.4	2.4
25-26	1100	17.7	7.2	15.1	0.2	0.6	0.0	0.0	0.7	2.6
27-28	1100	14.3	5.9	12.0	0.1	0.3	0.0	0.0	0.1	1.1
29-30	1100	11.3	4.9	8.5	0.2	0.3	0.0	0.0	0.1	1.1
Region:										
Northeast	1300	20.6	8.8	18.1	0.3	1.2	*	0.2	0.7	2.9
North Central	1900	16.9	7.3	14.4	0.1	0.8	0.1	0.1	0.4	1.9
South	2300	16.0	8.0	12.8	0.4	0.5	0.1	0.0	0.6	1.7
West	1300	19.2	7.7	16.9	0.3	0.9	*	0.0	0.3	1.8
Population Density:^e										
Farm/Country	800	13.4	6.6	10.4	0.3	0.2	0.1	0.0	0.0	1.3
Small Town	1800	17.6	8.3	14.8	0.1	0.7	0.1	0.0	0.4	2.1
Medium City	1600	18.0	7.4	15.1	0.3	0.8	0.1	0.2	0.6	2.0
Large City	1600	18.2	8.3	15.7	0.5	0.9	0.1	0.0	0.4	2.4
Very Large City	1000	20.1	8.4	18.0	0.1	1.3	0.2	0.3	1.2	2.0

Source: The Monitoring the Future Study, the University of Michigan.

* indicates a percentage of less than 0.05% but greater than true zero.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bUnadjusted for known underreporting of certain drugs. See text for details.

^cThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3400.

^dThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.

(Table continued on next page)

TABLE 4-4 (cont.)
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Crack ^a	Heroin	Other Narcotics ^b	Amphetamines ^{b,c}	Methamphetamine ^d	Ice ^d	Sedatives (Barbiturates) ^b	Tranquilizers ^b	Alcohol	Flavored Alcoholic Beverages ^e	Cigarettes	Steroids ^d
Total	6800	0.3	0.1	2.8	2.2	0.6	0.4	1.6	2.5	67.7	28.4	28.0	0.1
Gender:													
Male	2800	0.4	0.2	3.3	2.4	0.6	0.6	1.9	3.0	72.9	23.3	32.0	0.1
Female	4000	0.2	0.1	2.5	2.1	0.6	0.2	1.4	2.2	64.1	31.8	25.3	0.0
Modal Age:													
19-20	1200	0.4	0.1	4.0	3.6	0.9	0.8	2.7	3.0	56.7	33.3	27.9	0.0
21-22	1200	0.2	0.1	3.0	2.9	0.5	0.5	1.7	3.4	72.4	31.1	31.3	0.0
23-24	1100	0.6	0.1	3.2	2.4	0.7	0.1	2.1	2.6	72.8	31.1	31.5	0.1
25-26	1100	0.4	0.1	2.3	1.8	0.4	0.6	1.1	2.4	72.4	24.6	29.6	0.0
27-28	1100	*	*	2.4	1.2	0.5	0.1	1.4	1.8	68.8	26.7	25.9	0.3
29-30	1100	0.0	*	1.9	1.2	0.6	0.2	0.6	1.6	64.5	22.3	21.9	0.0
Region:													
Northeast	1300	0.3	0.2	3.0	1.9	0.1	0.3	1.4	2.1	73.1	31.4	30.2	0.0
North Central	1900	0.4	0.1	2.8	2.3	0.7	0.5	1.5	2.3	73.0	34.0	31.1	0.0
South	2300	0.2	0.1	2.8	2.4	0.3	0.1	1.9	3.3	61.6	22.6	26.6	*
West	1300	0.3	0.1	2.6	1.9	1.4	0.9	1.6	1.8	65.5	28.0	24.1	0.2
Population Density:^f													
Farm/Country	800	0.0	0.0	3.1	2.0	1.2	0.9	1.6	2.1	62.0	30.9	30.8	0.5
Small Town	1800	0.3	0.1	3.3	2.1	0.4	0.2	1.5	2.3	64.9	28.7	30.4	0.0
Medium City	1600	0.3	0.1	2.6	2.2	0.7	0.4	1.8	2.5	68.2	27.1	27.0	0.0
Large City	1600	0.4	0.1	2.3	2.2	0.5	0.2	1.6	2.9	69.2	28.0	25.4	0.0
Very Large City	1000	0.3	0.1	2.8	2.7	0.5	0.7	1.7	2.4	74.1	28.4	27.6	0.0

Source: The Monitoring the Future Study, the University of Michigan.

* indicates a percentage of less than 0.05% but greater than true zero.

^aThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5600.

^bOnly drug use that was not under a doctor's orders is included here.

^cBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^dThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2300.

^eThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1100.

^fA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents.

Within each level of population density, suburban and urban respondents are combined.


TABLE 4-5
Thirty-Day Prevalence of Daily Use of Various Types of Drugs by Subgroups, 2004
Among Respondents of Modal Ages 19-30
(Entries are percentages)

	<i>Approx. Weighted N</i>	Marijuana Daily	Alcohol Daily	Alcohol: 5+ Drinks in a Row in Past 2 Weeks	Cigarettes Daily	Cigarettes: Half-Pack or More per Day
Total	6800	4.5	4.4	35.4	20.1	13.3
Gender:						
Male	2800	6.3	7.6	46.5	22.8	15.7
Female	4000	3.3	2.2	27.7	18.3	11.7
Modal Age:						
19-20	1200	6.0	3.7	35.5	18.2	11.6
21-22	1200	5.1	5.7	41.7	21.5	12.7
23-24	1100	5.3	5.5	40.4	23.3	15.2
25-26	1100	5.5	4.3	36.4	22.7	15.6
27-28	1100	2.9	3.5	31.3	18.2	12.8
29-30	1100	2.0	3.8	26.9	16.7	12.5
Region:						
Northeast	1300	4.9	5.1	39.4	21.8	14.5
North Central	1900	4.2	4.7	40.3	23.6	16.6
South	2300	4.0	3.5	30.6	19.2	12.9
West	1300	5.6	4.6	32.3	15.1	8.5
Population Density:^a						
Farm/Country	800	3.5	4.8	31.6	25.2	18.2
Small Town	1800	4.6	4.3	34.6	23.2	15.7
Medium City	1600	4.5	4.6	35.9	18.8	12.3
Large City	1600	4.3	3.9	36.1	17.0	11.3
Very Large City	1000	5.4	5.0	38.2	17.5	10.1

Source: The Monitoring the Future Study, the University of Michigan.

^aA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000-100,000; a large city as 100,000-500,000; and a very large city as having over 500,000 residents. Within each level of population density, suburban and urban respondents are combined.

FIGURE 4-1
Any Illicit Drug: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-2


**Any Illicit Drug Other Than Marijuana: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004**

by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time.
See text for discussion.

FIGURE 4-3
Marijuana: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-4
Amphetamines: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-5
Methamphetamine: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45,* 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time.
 See text for discussion.


*This specific drug was not included in the age 35, age 40, or age 45 questionnaires.

FIGURE 4-6
Cocaine: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-7
Crack Cocaine: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-8
Other Cocaine: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-9
Hallucinogens:* Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

*Unadjusted for the possible underreporting of PCP.


FIGURE 4-10
LSD: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

*This specific drug was not included in the age 40 or age 45 questionnaires.


FIGURE 4-11
Hallucinogens Other Than LSD: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

*This specific drug was not included in the age 40 or age 45 questionnaires.


FIGURE 4-12
Inhalants:* Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

*Unadjusted for the possible underreporting of amyl and butyl nitrites. This specific drug was not included in the age 35, age 40, or age 45 questionnaires.

FIGURE 4-13
Sedatives (Barbiturates): Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time.
 See text for discussion.

FIGURE 4-14


**Narcotics Other Than Heroin: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004**

by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time.
See text for discussion.

FIGURE 4-15
Tranquilizers: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-16
MDMA: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45,* 2004
 by Age Group




NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

*This specific drug was not included in the age 35, age 40, or age 45 questionnaires.

FIGURE 4-17

**Crystal Methamphetamine ("Ice"): Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45,* 2004**


by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time.
See text for discussion.

*This specific drug was not included in the age 35, age 40, or age 45 questionnaires.


FIGURE 4-18
Steroids: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45,* 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


*This specific drug was not included in the age 35, age 40, or age 45 questionnaires.

FIGURE 4-19
Heroin: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-20a
Alcohol: Lifetime, Annual, and Thirty-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004
 by Age Group



NOTE: Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-20b
Alcohol: Two-Week Prevalence of Five or More Drinks in a Row
and Thirty-Day Prevalence of Daily Use Among High School Seniors
and Adults Through Age 45, 2004
 by Age Group





FIGURE 4-21

**Cigarettes: Annual, Thirty-Day, Daily, and Half-Pack-a-Day Prevalence
Among High School Seniors and Adults Through Age 45, 2004**

by Age Group



Chapter 5

TRENDS IN DRUG USE IN EARLY TO MIDDLE ADULTHOOD

In this chapter, we report historical trends in the use of the various licit and illicit drugs by high school graduates for particular age bands between 1 and 27 years beyond high school, spanning the modal ages 19 through 45. Use of the panel data from various senior classes for this purpose differs from the manner in which longitudinal panel data are more typically used, which is to study changes in the behaviors and attitudes of the same individuals as they age over time. Although the panel data from the many high school graduating classes encompassed in this study can be, and frequently are, used for that more typical purpose, they also can be used to track historical trends for fixed age bands across years. In other words, they can be used much as we often use the cross-sectional surveys of secondary school students.

In the early 1990s, we began to document large and important increases among secondary school students in the use of several substances, particularly marijuana and cigarettes. The increases continued among 12th graders through 1997, as discussed in Volume I. One of the important issues addressed in this chapter is whether such increases occurred only among adolescents or whether recent graduating classes have carried their higher levels of drug use with them as they have moved into young adulthood. In other words, are they exhibiting lasting cohort effects?

Figures 5-1 through 5-19 plot separate trend lines for two-year age strata (that is, 1-2 years beyond high school, 3-4 years beyond high school, etc.) We present data in two-year age strata in order to damp down the random fluctuations that would be seen with one-year strata. (Strictly speaking, these two-year strata are not age strata, because they are based on all respondents that year from two adjacent high school classes, and they do not take account of the minor differences in individual respondents' ages within each graduating class; however, they are close approximations to age strata, and we characterize them by the modal age of the respondents as ages 19 to 20, 21 to 22, and so on.) Each data point in these figures is based on approximately 1,200 weighted cases drawn from two adjacent high school classes; actual (unweighted) numbers of cases are somewhat higher. For the 2004 data, the 19- to 20-year-old stratum is composed of participating respondents from the high school graduating classes of 2003 and 2002, respectively; the 21- to 22-year-old stratum contains data from the classes of 2001 and 2000, respectively, and so on. Figures 5-1 through 5-19 also present some recent trend data from the age-35, age-40, and age-45 follow-ups. Each of these is constituted in a slightly different way, in that the two half-samples from a *single* graduating class (which through age 30 had been surveyed in alternating years) are both surveyed in the same year. In 2004, the 35-year-olds are graduates from the high school class of 1987 (weighted $n = 979$), the 40-year-olds are graduates from the high school class of 1982 (weighted $n = 1004$), and the 45-year-olds are graduates from the high school class of 1977 (weighted $n = 1014$).

Tables 5-1 through 5-5 are derived from the same data but presented in tabular form for 19- to 28-year-olds combined (i.e., those who graduated from high school 1 to 10 years earlier). Data are given for each year in which they are available for that full age band (i.e., from 1986 onward). Those aged

29 and over are omitted because their inclusion would shorten the time period over which trends can be examined. However, the full data for them are contained in Figures 5-1 through 5-19.

TRENDS IN PREVALENCE: EARLY AND MIDDLE ADULTHOOD

The trend results are as follows:

- Longer-term declines among young adults in the annual prevalence of several drugs appeared to end in 1992 or 1993 (see Table 5-2). Among the 19- to 28-year-old young adult sample this was true for the use of *any illicit drug, any illicit drug other than marijuana, marijuana, hallucinogens, narcotics other than heroin, crack, amphetamines, sedatives (barbiturates), and tranquilizers*. In 1994, annual prevalence for most drugs remained steady. *Cocaine other than crack* reached its nadir in 1994 after a period of substantial decline. In 1995 there were modest increases (a percentage point or less) in the annual prevalence of almost all of the drug classes in Table 5-2, some of which were statistically significant.

Thus, it is clear that by 1992 or 1993 the downward secular trend observable in all of these age strata (as well as among adolescents) had ended.⁴¹ (Such secular trends, in which different age groups move in parallel, are also called “period effects.”) What has happened since then, however, is quite a different form of change. Rather than being a period effect common to all age groups, it is more of a “cohort effect,” reflecting an interaction between age and period such that only adolescents showed the increase in illicit drug use initially, and then they carried those new (higher) levels of drug use with them as they entered older age bands. Figure 5-1 shows the effects due to generational replacement, as the teens of the early 1990s reached their 20s. It can be seen that, while all age groups moved fairly parallel through about 1992, the youngest age bands first showed signs of increase in their overall level of illicit drug use. The 18-year-olds shifted up first, followed by the 19- to 20-year-olds, the 21- to 22-year-olds, the 23- to 26-year-olds in 1999, the 27- to 30-year-olds in 2000, and the 35 and 40 year olds in 2004.

To repeat, in the earlier decline phase of the drug epidemic, annual prevalence of use of *any illicit drug* moved in parallel for all of the age strata, as illustrated in Figure 5-1; this pattern reflects a secular trend, because a similar change is observed simultaneously across different age levels. In what we have called the “relapse phase” of the popular drug epidemic, after 1992 a quite different pattern emerged, with the 12th graders increasing their drug use first, and rising fastest; the next oldest age group following, but with a little delay; the next oldest then following, but with a longer delay; and the oldest groups not yet showing an increase. This pattern reflects a classic cohort effect, in which different age groups are not all moving in parallel; rather, different age groups show increases when the cohorts (that is, high school classes) having heavier use at an earlier stage in development reach the relevant age level. Further, the slopes of the age bands are successively less steep in the higher age groups, suggesting that some of the cohort effect may be dissipating with maturation. But we think it

⁴¹Actually, the downturn ended at least a year earlier among the youngest adolescents—the 8th graders—who showed the beginning of an increase in 1992. (See Table 2-2.)

unlikely that only cohort effects will be occurring (in addition to the long-established age effects); also, no doubt, entering into the mix are period effects (i.e., historical effects that have an impact on all age bands).

- Use of *marijuana*, the major component of the illicit drug use index, shows an almost identical pattern to the index (Figure 5-3a). After a long and steady decline from the late 1970s to the early 1990s, use leveled for a while among young adults before beginning a gradual increase. Virtually all of this increase was attributable to the two youngest age bands (18 and 19 to 20) until 1996, when the third youngest age band (21- to 22-year-olds) began to show a rise. The older age bands then tended to show increases fairly sequentially. The 18-year-olds began a decline after 1997, and later several of the succeeding age bands through age 24 have begun to show declines in a pattern that suggests cohort differences.
- A similar pattern emerged for current *daily marijuana* use (Figure 5-3c). In the mid- to late 1990s, daily marijuana use among the 35- and 40-year-olds was as high as, or higher than use among some younger age groups, suggesting a lasting cohort effect on this behavior. However, in recent years, the 35- and 40-year-olds (and now the 45-year-olds) have been similar to those ages 27-32, who are at the lowest levels of daily use. An important finding shown in Figure 5-3c is that, although the various age groups had been moving in parallel for many years at fairly similar levels of prevalence, the trends diverged considerably in the 1990s in a staggered fashion, such that now 18- to 26-year-olds have distinctly higher levels of daily marijuana use than the older age groups, again reflecting stable cohort differences.
- The index of using *any illicit drug other than marijuana* has shown a similar transition in the pattern of change. Period effects seemed to predominate until about 1992, but a cohort-related pattern of change emerged thereafter (Figure 5-2). And, while use leveled by 1997 among 18-year-olds, it began rising during that interval among 25- to 26-year-olds and is also now rising among 27- to 28-year-olds and 29- to 30-year-olds. The primary difference from the picture for marijuana is that the increases were not as sharp in the 1990s for most of the age bands for the other illicit drugs taken as a group as they were for marijuana.
- In the 1980s and 1990s, *LSD* use also increased among those in their teens and early 20s more than among the older strata, as Figure 5-6 illustrates. Over the interval 1985 to 1996 there was a gradual but considerable increase in LSD use among those aged 18 to 24, which was sharpest among 12th graders and the 19- to 20-year-olds. (In this case the increase did not seem to radiate up the age spectrum beyond age 26.) A turnaround began among the 12th graders after 1995 and then among the older age groups in a somewhat staggered fashion, again indicative of a cohort effect. The declines in the years since have been greatest among the 18- to 24-year-olds, who had attained the highest rates of LSD use. All age bands (including 8th and 10th graders) have shown considerable decline in LSD use since 2001, suggesting that an important secular trend may have set in, quite possibly related to decreased availability of the drug.
- Several of these drug classes actually exhibited a faster decline in use among the older age groups than among 12th graders during the *earlier* period of decline. (See Figures 5-1

through 5-19.) These included *any illicit drug*, *any illicit drug other than marijuana*, *amphetamines*, *hallucinogens* (until 1987), *LSD* (through 1989), and *methaqualone*.

- In fact, there was a crossover for some drugs when 12th graders are compared to young adult graduates. In earlier years 12th graders had lower usage levels, but in recent years they have tended to have higher ones than post-high school respondents for use of *any illicit drug*, *marijuana*, *any illicit drug other than marijuana*, *hallucinogens*, *LSD* specifically, *amphetamines*, *tranquilizers*, *narcotics other than heroin*, and *crystal methamphetamine (ice)*. However, since then, as the next two age strata after high school continued to show increases on a number of these drugs, they have closed the gap with 12th graders. This has been true for *marijuana*, *hallucinogens*, *LSD*, *narcotics other than heroin*, and *tranquilizers*. (See, for example, Figure 5-3a for annual marijuana prevalence.)
- *Cocaine* (Figure 5-9) gives a quite dramatic picture of change. Unlike most of the other drugs, active use of cocaine generally has tended to rise with age after high school, usually peaking approximately three to four years past graduation. This is a classic example of an age-effect. Despite the large age differences in absolute prevalence, all age strata moved in a fairly parallel way through 1991, indicating that a secular trend was taking place. All began a sharp and sustained decline in use after 1986. The two youngest strata (12th graders and 19- to 20-year-olds) leveled by 1992, whereas use continued a decelerating decline for a few years beyond that in the older age groups, signaling the beginning of a cohort effect. From 1994 to 1999, cocaine use rose some in the five youngest strata (i.e., those younger than 27) on a somewhat staggered basis, with the three older groups still decreasing a bit more over that same period. This, to some degree, reversed the age differences that were so prominent in the 1970s and 1980s. Cohort-related change appears to have predominated in the 1990s, quite possibly as the result of “generational forgetting” of the cocaine-related casualties so evident in the early to mid-1980s. In other words, those in the older cohorts retained that learning, but those in the newer cohorts never had it. The fact that in recent years the 35- and 40-year-olds had higher lifetime prevalence levels of cocaine use than some of the younger age groups also suggests some lasting cohort-related differences established during the peak years of cocaine use.
- *Crack* use was added to the 12th graders’ questionnaires in 1986 and to the follow-up questionnaires in 1987. The subsequent decline in crack use ended in 1991 among 12th graders, and by 1994 it had ended among young adults (see Figure 5-10 and Table 5-2). Among 19- to 28-year-olds, the annual prevalence rate held at about 1%, which was down from the peak levels of just over 3% in 1986 through 1988. As was true for a number of other drugs, crack use began to rise (in this case after 1993) among 12th graders but not in the older age strata until later years, when increases were observed in a somewhat staggered pattern going up the age scale. Again, a cohort effect due to generational replacement seems to have been occurring. Since 1994, the 18-year-olds have had the highest reported rates of use, and since 1999 the 19-20 year-olds have generally had the second highest rates. Importantly, all groups now have annual prevalence rates below 2.0% except the 18-year-olds, who are at 2.3%.

- With regard to *inhalants*, the large separation of the age band lines in Figure 5-4 shows that, across many cohorts, use consistently has dropped sharply with age, particularly in the first few years after high school. In fact, of all the populations covered in this study, the 8th graders (not shown in Figure 5-4) have had the highest rate of use, indicating that the decline in use with age starts at least as early as 8th or 9th grade. Like cocaine, inhalants have shown a strong age effect, but unlike cocaine, use of inhalants *declines* with age.

Figure 5-4 also shows that there was a long-term gradual increase in annual inhalant use (unadjusted for underreporting of nitrite inhalants), one which was greatest among 12th graders, next greatest among 19- to 20-year-olds, and next greatest among 21- to 22-year-olds. Respondents more than six years past high school, who historically have had a negligible rate of use, did not exhibit the increases in use seen among the younger respondents, which began at least as early as 1977 among 12th graders and in 1983 among 19- to 20-year-olds. There was subsequently some increase among 21- to 22-year-olds and later still an increase among 23- to 24-year-olds. After 1995, this long-term trend, reflecting a cohort effect, began to reverse in the two youngest age strata and subsequently among the next two age strata. The older age strata generally have shown negligible rates of inhalant use.

- In the late 1970s, *amphetamine* use rose with age beyond high school; but after a long period of decline in use from 1981 to the early 1990s, this relationship had reversed (see Figure 5-13). The declines were greatest in the older strata and least among the 12th graders, even though use decreased substantially in all groups. As was true for many of the illicit drugs, amphetamine use began to rise among the 12th graders after 1992, and eventually among the 19- to 24-year-olds; but there has only recently been a small increase among those 25- to 30-year-old. In other words, another cohort-related pattern of change seems to have emerged in the 1990s for amphetamines, though in this case it may be dissipating quickly after respondents reach their early 20s. At present the age differences through age 45 are of considerable magnitude and mostly ordinal (with the youngest showing the highest rates of use).
- The annual prevalence for *MDMA (ecstasy)* use among the entire young adult sample (ages 19 to 28) was at about 1.5% in 1989 and 1990 (Table 5-2 and Figure 5-8). After 1991 it dropped to around 0.8% for several years before starting to rise significantly in 1995. Then ecstasy use began to rise in all of the young adult age strata but clearly rose the most among those in the younger age bands (19 through 26) through 2001. Use among 12th graders, which was not measured until 1996, was by then the highest of any of the age groups at 4.6% annual prevalence. Twelfth graders' use slipped by a full percentage point through 1998 before jumping significantly—by 2 full percentage points—in 1999. (Use by 10th graders also jumped significantly in 1999.) Thus it appears that young people from their mid-teens to mid-20s had “discovered” ecstasy, after some years of low and relatively level use. In 2000 the sharp increase in use continued among those aged 15-16 (10th graders) to 26 and also showed up among 8th graders (13- to 14-year-olds) for the first time. By 2001 the increase had slowed and even begun to reverse among those aged 18 to 26, even as the 31- to 32-year-olds showed their first appreciable increase in ecstasy use. We attributed the deceleration in 2001 to a fairly sharp increase in perceived risk that year and predicted a

turnaround in use in 2002. In 2002, and again in 2003, perceived risk increased sharply; and, as Figure 5-8 illustrates, all age bands showed a reversal, with a sharp decrease in use. Clearly, the decrease has been sharpest in the younger age bands, perhaps because a cohort effect is at work in the upper ages, helping to offset a secular downward trend.

- Since 1990, when it was first measured, the use of *crystal methamphetamine (ice)* has remained at fairly low rates in this young adult population (Figure 5-14). However, among 19- to 28-year-olds combined, annual prevalence rose from 0.4% in 1992 to 1.2% by 1995 before leveling through 2004 (Table 5-2). Use had been rising among 12th graders and 19- to 20-year-olds, specifically, between 2000 and 2002, but in 2003 and 2004 their use fell back to around the 2000 levels.
- Use of *heroin* increased appreciably in 1995 among 12th graders and young adults aged 19 to 24 but not among the older age bands (Figure 5-11). Among young adults generally, annual use had previously been quite stable at least as far back as 1986 (Table 5-2), and it stabilized again at a higher level after 1995. Heroin use among 12th graders and 19- to 20-year-olds has declined slightly since 2001, but the older age groups have maintained a fairly stable rate of low use.
- Among 19- to 28-year-olds, the use of *narcotics other than heroin* leveled after 1991, following a long period of slow, fairly steady decline (Figure 5-12). Twelfth graders showed an appreciable increase in use, beginning in 1993, which continued into 2004, while 19- to 20-year-olds showed some increase after 1994, 21- to 22-year-olds after 1996, 23- to 24-year-olds after 1997, and the older age groups after 2000. Thus, cohort-related change appears to have been occurring during the 1990s for this class of drugs, following a long period of secular trends. In 2002, the question text was changed on three of the six questionnaire forms to update the list of examples of narcotics other than heroin. Talwin, laudanum, and paregoric, each of which had negligible rates of use by 2001, were replaced by Vicodin, OxyContin, and Percocet. As a consequence of this revision, reported use rates increased in 2002. Data presented here for 2002 are from the three of six questionnaire forms with the new wording (which showed higher prevalence rates than the older question did). All six questionnaires contained the new wording in 2003, so the data presented are based on all forms. Although the older version of the question showed no significant changes occurring in 2002, there was a significant increase in narcotics use observed in 2003 (based on the new question).
- The annual prevalence rates for *Vicodin* and *OxyContin*, which were first measured in 2002 (separately from the overall question), were appreciable (8.2% and 1.9%, respectively) for all 19- to 28-year-olds. Non-statistically-significant increases were observed for these two drugs in 2003 and 2004. Among 19- to 28-year-olds (see Table 5-2), the annual prevalence of OxyContin use rose from 1.9% in 2002 to 2.6% in 2003 to 3.1% in 2004—changes that were parallel to those observed among 12th graders over the same interval (when their annual prevalence rose from 4.0% to 4.5% to 5.0%). The 2002-2004 increases in OxyContin use were significant for both 12th graders and 19- to 28-year-olds. Vicodin use rose by less, but started from a higher base, with annual prevalence increasing among 19- to 28-year-olds from 8.2% in 2002 to 8.6% in 2003 to 8.9% in 2004.

- **Sedative (barbiturate)** use (Figure 5-15) showed a long-term parallel decline in all age groups covered through the late 1970s and 1980s, leveling by about 1988. While use has remained low and quite level for most of the age bands, it began to rise by 1993 among 18-year-olds, by 1995 among 19- to 20-year-olds, by 1997 among 21- to 22-year-olds, by 1998 among 23- to 24-year-olds, and by 2001 among 25- to 28-year-olds. The same cohort-related pattern of change seen during the 1990s for many other drugs also exists for sedatives (barbiturates); like most of the other drugs, this pattern was preceded by a period of secular change.
- **Tranquilizers** (Figure 5-16) give a fairly similar picture to that just described for sedatives (barbiturates). One difference is that the 12th graders' annual prevalence rate has not always been the highest among the various age groups, as was the case for sedatives (barbiturates), although it was highest between 1994 and 2000 as a result of greater increase in tranquilizer use. In the last four years, however, as use continues to increase among those in their early 20s, the 12th graders no longer stand out as having the highest rate of tranquilizer use, though they still have considerably higher rates than those over age 26. This is another clear example of a cohort-related pattern of change.
- The use of **anabolic steroids** (Figure 5-17) is substantially lower after high school than during, and this has been true since measures of steroid use were first introduced into two of the follow-up questionnaires in 1991. The age-related differences are not consistent; the prevalence rates are all quite low and do not appear to trend in any systematic way. In general, it seems that the rise in steroid use from 1999 to 2003 among 8th and 10th graders and from 2001 to 2004 among 12th graders seems to have been specific to those age groups, at least so far.
- The **alcohol** trends for the older age groups (see Figures 5-18a-d) have been somewhat different than for the younger age groups and in some interesting ways. For **30-day prevalence** and **occasions of heavy drinking**, the declines for the two youngest age strata (12th graders and those one to two years past high school) during the 1980s were greater than for the older age groups. These differential trends were due in part to the effects of changes in minimum drinking age laws in many states, changes that would be expected to affect primarily the age groups under age 21. However, because similar (though weaker) trends were evident among 12th graders in states that maintained a constant minimum drinking age of 21, the changed laws cannot account for all the downward trends, suggesting that there was also a more general downward trend in alcohol consumption during the 1980s.⁴² By 1994, these declines in 30-day prevalence had slowed or discontinued for virtually all age groups until 1997, when they began to turn downward again for 12th graders, and 1999, when they started down among the 19- to 20-year-olds.

Those respondents three to four years past high school stand out for showing the smallest downward trend in **binge drinking** since the early 1980s (see Figure 5-18d). One important

⁴²O'Malley, P. M., & Wagenaar, A. C. (1991). Effects of minimum drinking age laws on alcohol use, related behaviors, and traffic crash involvement among American youth: 1976-1987. *Journal of Studies on Alcohol*, 52, 478-491.

segment of that age stratum is composed of college students, who showed very little downward trend (see chapter 9).

The older age groups, in general, have shown only a modest long-term decline in annual prevalence rates and no recent decline in binge drinking or in 30-day prevalence rates. Note that the binge drinking trend lines for different age groups (Figure 5-18d) are spread out on the vertical dimension, reflecting large and persisting age differentials (age effects) in this behavior. The relationship with age is curvilinear, however. In recent years the 21- to 22-year-olds consistently have shown the highest rates of binge drinking, while the two adjacent age bands have shown the next highest. Binge drinking appears to have been gradually increasing in recent years among the 23- and 24-year-olds and the 25- and 26-year-olds. This is perhaps driven in part by the fact that an increasing proportion of them are enrolled in college, where binge drinking rates tend to be high. In addition, they tend to be unmarried, which also contributes to binge drinking.⁴³

From the early 1980s through the mid 1990s, rates of *daily drinking* (Figure 5-18c) fell by considerable amounts in all age strata for which we have data, reflecting an important change in drinking patterns in the culture. Among 19- to 28-year-olds combined, daily drinking fell from 6.6% in 1987 to 3.9% in 1994, before leveling briefly and then rising as high as 5.1% in 2003 (see Table 5-4). Daily drinking rates have proven to be highest for 35-, 40-, and 45-year-olds in recent years, when data on them became available.

It is worth noting that the 35-, 40-, and 45-year-olds have had among the lowest rates of binge drinking but among the highest rates of daily drinking in recent years for which we have data available. These patterns—particularly the high rate of daily drinking—likely reflect age effects as well as perhaps some enduring cohort differences (because these cohorts had considerably higher rates of daily drinking when they were in high school).

- The prevalence rates for *cigarette smoking* show more complex trends than most other substances, due to the long-term presence of both cohort and age effects, plus slightly different patterns of such effects on different measures of smoking in the past 30 days (one or more cigarettes per month, one or more cigarettes per day, and a half-pack or more of cigarettes per day).

While in the earlier years of the study, the curves are of the same general shape for each age band (Figures 5-19a-c), each of those curves tended to be displaced to the right of the immediately preceding age group, which is two years younger. The pattern is clearest in Figure 5-19c (half-pack plus per day). This pattern is very similar to the one described in Volume I for lifetime smoking rates for various grade levels *below* senior year; it is the classic pattern exhibited by a cohort effect—that is, when cohorts (in this case, high school graduating class cohorts) differ from other cohorts in a consistent way across much or all of the life span. We interpret the cigarette data as reflecting just such a cohort effect,⁴⁴ and we

⁴³Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁴⁴O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976-1986. *American Journal of Public Health*, 78, 1315-1321.

believe that the persisting cohort differences are due to the dependence-producing characteristics of cigarette smoking.

The declining levels of cigarette smoking across cohorts at age 18, which were observed when the classes of 1978 through 1981 became 12th graders, were later observable in the early-30s age band, as those same high school graduating classes reached their early 30s (see Figures 5-19b and c). This was true at least through about 1991. After that, there was a considerable convergence of rates across age groups, largely because there were few cohort differences among the senior classes who graduated from the early to mid-1980s through the early 1990s—a period of fairly level use among succeeding senior classes.

In addition to these cohort differences, there are somewhat different age trends in which, as respondents grow older, the proportion smoking at all in the past 30 days declines some, while the proportion smoking a half-pack per day actually increases. Put another way, many of the light smokers in high school either become heavy smokers or quit smoking.⁴⁵

The picture was further complicated in the 1990s, when it appears that a new cohort effect emerged, with smoking among adolescents rising sharply (beginning after 1991 for the 8th and 10th graders and after 1992 for the 12th graders). The 19- to 20-year-olds also showed a rise at the beginning of the 1990s—responding perhaps to some of the same social forces as the adolescents (including possibly the Joe Camel advertising campaign); but the 21- to 24-year-olds did not show an increase until about 1995, and the 25- to 26-year-olds until about 1996. Those young adults over age 26 have not yet shown much increase, though they may well do so as the heavier-smoking senior class cohorts enter those age bands.

After about 1999, smoking rates among virtually all age groups leveled or declined, suggesting that general societal forces may be affecting all age groups in a similar way, giving rise to some secular trends. Large increases in price and a great deal of adverse publicity for the tobacco industry are highly plausible candidates for such forces, as are an increase in state and national anti-smoking advertising and the demise of the Joe Camel campaign and of billboard advertising.

- Apart from cigarettes, none of the other drugs included in the study showed a clear long-term pattern of enduring cohort differences in the earlier years of the study (the 1970s and 1980s), despite wide variations in their use by different cohorts at a given age. There was one exception; a modest cohort effect was observable for *daily marijuana* use during the late 1970s and early 1980s. (But as more recent classes leveled at lower rates of use, evidence for the cohort effect faded.) The emergence in the 1990s of a new epidemic of marijuana use among teens once again yielded a strong pattern of cohort effects. As can be seen in Figure 5-3c, daily use rose sharply among 12th graders and 19- to 20-year-olds after 1992, among 21- to 22-year-olds after 1993 with a sharp rise occurring in 1997, among 23- to 24-year-olds after 1998, among 25- to 26-year-olds after 2000, and among 27- to 28-year-olds in 2003. However, among those 29 and older, as of yet there has been virtually no increase in daily

⁴⁵ To illustrate, in the class of 1976, 39% were 30-day smokers in senior year, 39% by ages 19 to 20, and by age 31 to 32 only 28%—a net drop of 11 percentage points over the entire interval. By way of contrast, 19% of that class were half-pack-a-day smokers in senior year, 24% by ages 19 to 20, and 21% at ages 31 to 32—a net gain of 2% over the interval.

use. This is not unlike the pattern of change for cigarette smoking that occurred in the 1990s (Figure 5-19a). The cohort effect for daily marijuana use may be attributable, in part, to the very strong association between that behavior and regular cigarette smoking. It is noteworthy that even among the 35-, 40-, and 45-year-olds in the study, 1.8%-2.5% report that they still currently smoke marijuana on a daily basis. That amounts to 1 in every 40 to 56 adults at those ages. And we know from results published in the companion volume to this one that many more have been daily marijuana users for at least a month at some time in their life.

- In sum, except for *cigarettes* and *alcohol*, substance use prior to 1992 among 12th graders and young adults had shown *longer-term* trends that were highly parallel, indicating that general secular trends predominated in that period. Since 1992, however, there has been some considerable divergence in the trends for different age bands on a number of drugs as use among adolescents rose sharply, followed by subsequent rises among the 19- to 20-year-olds, the 21- to 22-year-olds, and so on. This divergence indicates a new cohort effect, quite possibly reflecting a “generational forgetting” of the dangers of drugs by the cohorts who reached senior year in the early to mid-1990s. The data discussed in chapter 6, “Attitudes and Beliefs About Drugs Among Young Adults,” provide additional evidence for this interpretation.

TRENDS FOR IMPORTANT SUBGROUPS OF YOUNG ADULTS

Four-year age bands have been used here to examine subgroup trends in order to yield sufficiently large numbers of cases to permit reliable estimates for the various subgroups being examined. Subgroup data for respondents of each gender and for respondents from communities of different sizes are available for 19- to 22-year-olds since 1980, 23- to 26-year-olds since 1984, and 27- to 30-year-olds since 1988. Beginning with the 1987 follow-up questionnaires, a question about state of residence was added to all follow-up questionnaires, permitting trend data to be calculated for the four regions of the country. These various subgroup data are not presented in tables or figures here because of the substantial amount of space they would require. Rather, a verbal synopsis of what they contain is presented.

Gender Differences in Trends

- Over the long term, gender differences narrowed for some drugs among young adults, primarily because of a steeper decline in use among males (who generally had higher rates of use) than among females. The overall picture, though, is one of parallel trends, with use among males remaining higher for most drugs, including the indexes of *any illicit drug* use in the prior year and use of *any illicit drug other than marijuana* (see Table 5-5, for example).
- The downward trend in *marijuana* use among 19- to 22-year-olds between 1980 and 1989 was a bit sharper among males than females, narrowing the gap between the two groups. Annual prevalence fell by 22 percentage points (to 34%) among males, compared to a drop of 14 percentage points (to 31%) among females, leaving a difference of 3 percentage points. In the late 1990s through 2004, the gap widened a bit, to 8 percentage points (38% versus 30%).

Similarly, between 1980 and 1993, *daily marijuana* use for this age group fell more steeply, from 12.9% to 2.9% among males, versus from 6.1% to 1.7% among females, narrowing the gap considerably. As use began to rise after 1993, the gap widened again. Among 23- to 26-year-olds, as daily use first began to increase in 1998 and 1999, the gap between the genders began to widen. In the oldest age group (aged 27-30), the difference had been fairly constant, with daily marijuana use among males generally being two to three times higher than among females. After 2001, however, use increased among females, while use among males remained steady.

- Males have shown slightly higher proportions using *any illicit drug other than marijuana* in all three age bands, a fact that has not changed appreciably over the years, though the differences tended to narrow some as use dropped and to widen as use increased.
- For *LSD*, males have consistently had higher rates of use than females. Among 19- to 22-year-olds, the male-female differences tended to diminish as use declined (1980–1985) and tended to increase as use increased (1986–1995). In the two older age bands there was less change in use, and differences had been relatively consistent. In the last few years, however, the pattern was accentuated: LSD use has dropped considerably since 1999 among 19-through 26-year-old males (and since 2001 among the 27- to 30-year-old males), substantially narrowing the gender differences. Males began to show these declines first, and both genders have moved to almost no use.
- *MDMA (ecstasy)* exhibited little or no gender difference in any of the three age bands before use began to grow in the late 1990s. Even since then, among the 19- to 22-year-olds there has been little gender difference, except that use among males started to decline one year ahead of use among females. But in the older age groups a gender difference did open up, with males having higher rates of use among both 23- to 26-year olds and 27- to 30-year olds. Among the 27- to 30-year-olds that gender difference disappeared in 2004 due to a decline in use among the males since 2001.
- During the period of sharp decline from the peak levels in annual *cocaine* prevalence (1986–1993), use dropped more among males than females, narrowing the gender differences. In the 19- to 22-year-old age band, annual prevalence for males declined by 16 percentage points (to 4.5%) versus 13 percentage points among females (to 2.8%) in 1993. In the 23- to 26-year-old age band, there was also a narrowing of the gender difference between 1986 and 1993, with annual prevalence down 19 percentage points (to 6.9%) among males and 13 percentage points (to 4.2%) among females. Use in the 27- to 30-year-old group also dropped faster among males (down 13.3 percentage points versus 7.1 among females) between 1988 (when data were first available) and 1997. In sum, during the period of sharp decline in overall cocaine use, the gender differences—which had been fairly large—narrowed considerably in all three of these age bands. During the more recent resurgence in cocaine use, the gap between genders expanded slightly; but males have had higher rates of use of this drug throughout the life of the study.
- A similar occurrence happened with *crack* during the earlier decline, though the proportional difference between the two genders has consistently been higher than for cocaine overall.

With crack, though, there was some convergence between 1992 and 1998 among 19- to 22-year-olds, as use among males declined slightly and use among females rose gradually. Since 1999, there has been no consistent change in differences between males and females. In the two older age bands, males have fairly consistently had somewhat higher crack usage rates.

- As *sedative (barbiturate)* use declined through the 1980s, the modest gender differences (males were higher) were virtually eliminated in all three of the age bands. Since the early 1990s, there has been a considerable increase in use by both genders among the 19- to 22-year-olds (with males being the first to rise, as is often the case), opening some gender difference that has held constant. Among the 23- to 26-year-olds, use began to rise among males since 1997, with the females catching up by 2000, but diverging again through 2004 as female use remained steady but male use continued to rise. Use by both males and females began to rise in 2001 among the 27- to 30-year-olds, and use among males has been slightly higher.
- The annual prevalence figures for *heroin* dropped among males in the 19- to 22-year-old category between 1980 and 1986 (from 0.6% to 0.2%) before leveling through 1994; thus most of the decline in use in that interval was among males, mostly eliminating the previous gender difference. Rates for both genders remained very low, between 0.1% and 0.3% throughout the period 1986 through 1994. In 1995 through 1998, use increased appreciably among both males and females in this youngest age group, but a gender difference opened up again (with males higher). After 2001, both showed some decline, followed by a gradual rise, resulting in respective annual prevalence rates of 0.6% and 0.3% in 2004. Among 23- to 26-year-olds, use also remained low (0.1% to 0.2%) over the years 1986-1994 for both genders. There was an increase from 1995 to 2001 among males, with females remaining relatively flat, and more of a gender difference emerged. However, since 2001, males have declined and females have remained stable, just about eliminating the gender gap. Among 27- to 30-year-olds there was some falloff in heroin use between 1988 (when data were first available) and 1990 in both genders, as well as a narrowing of gender differences. Use rose slightly in the early 2000s among males, and the rates among males were higher than among females for several years, though they converged in 2004 (0.3% and 0.2%, respectively).
- Among 19- to 22-year-olds, both genders showed some decline in their use of *narcotics other than heroin* between 1980 and 1991, with a near elimination of previous gender differences (males had been higher). Beginning in 1994, use by males began to rise in this age band, while use by females began to rise a year later. Some gender differences have developed as use has increased, with males at 11.7% and females at 8.5% in 2004. Among 23- to 26-year-olds, the gender difference (males higher) had been eliminated by 1988. It reemerged after 1992 as use has increased more among males. Among the 27- to 30-year-olds, there has been a smaller gender difference and the least increase in use in the 2000s. Still, use has increased in both genders since 1999 and continued to increase in 2004 among males. In fact, both genders in all three age bands (as well as among 12th graders) have shown continuing increases over a period of three to five years, helping to confirm that the increase in the use of this class of drugs is real and pervasive. Interestingly, in 2004 these upward trend lines leveled off among females in all three age bands, but continued to rise among males.

- The use of the specific drugs *OxyContin* and *Vicodin* has consistently been higher among males than among females for the three years for which we have data.
- Generally, there has been no appreciable gender difference in *amphetamine* use for some years in any of these three age bands. Between 1981 and 1991, rates of amphetamine use were similar for males and females and showed substantial and parallel downward trends for both genders. Among the 19- to 22-year-olds, use for males dropped 22 percentage points in annual prevalence (to 5.2% in 1991), and use for females dropped 21 percentage points (to 4.7% in 1991). There have been small increases in annual prevalence for both genders in the 19- to 22-year age group since 1991 and in the 23- to 26-year age group since 1995; but the genders have not diverged.
- *Crystal methamphetamine (ice)* was added to the study in 1990. In the early 1990s, use was low and very similar for both genders in all three young adult age bands. Nearly all of the increase in use that occurred in the mid-1990s in the younger two age bands occurred among males—opening a gender gap. The gap has narrowed since then, though males have tended to be slightly more likely to report use of ice.
- For *tranquilizers*, both genders showed a long, gradual decline (and very similar rates of use) from 1980 through about 1993 in all three age bands. Beginning in 1995, use increased for both genders in the 19- to 22-year-old group, followed by an increase beginning after 1997 among the 23- to 26-year-olds and after 1999 among the 27- to 30-year-olds, again reflecting generational replacement. Some gender difference has emerged in this period of increase, with males reporting somewhat higher usage rates.
- *Inhalant* use generally has been quite a bit higher among males than females in all three age groups. The 19- to 22-year-old group showed a gradual upward shift from 1980 to 1988, followed by a leveling for some years, in both genders. In 1997, female inhalant use began to decline among the 19- to 22-year-olds, followed by males in 2001; however, the gender gap did not diminish much with this decline. Among 23- to 26-year-olds there was a widening gender gap as use by males, but not females, increased between 1992 and 1999, though a decline among males since then has narrowed the gap.
- Use of three “club drugs”—*GHB*, *ketamine*, and *Rohypnol*—tends to be concentrated among males in all three age strata, with the single exception that among the 19- to 22-year-olds, GHB use has been about equivalent for both genders.
- For *alcohol*, 30-day prevalence rates have shown a long, gradual, parallel decline from 1981 through 1992 for both genders in the 19- to 22-year-old age group. Thirty-day prevalence fell from 83% to 72% among males and from 75% to 62% among females by 1992. In the two older age bands, there had also been a modest, parallel decline for both genders, from 1985 through 1992 in the case of 23- to 26-year-olds, and at least from 1988 (when data were first available) to 1991 or 1992 in the case of 27- to 30-year-olds. Since 1992, both genders in the older two age bands have shown fairly level use rates, with males somewhat higher; however, among the 19- to 22-year-olds there has been some convergence, because use by males has declined slightly while use by females has increased slightly.

There also was a general long-term decline in *daily drinking* from about 1981 or 1982 through about 1992, with daily use falling more among males, considerably reducing, but far from eliminating, what had been a large gender difference among 19- to 22-year-olds. To illustrate, in 1981, 11.8% of the males reported daily use versus 4.0% of the females; the comparable 1992 statistics were 5.3% and 2.7%. After 1995, daily drinking began to increase among the 19- to 22-year-olds for both genders but leveled a few years later. Since 2002 daily use by males has been rising, while it has been falling among females. There is still a large gender difference for daily drinking among the 19- to 22-year-old age group in 2004—8.3% for males versus 2.2% for females—but not nearly as large as it had been in 1981 (11.8% versus 4.0%). The gender differences have been similar for the older age groups (in 2004, for example, 7.6% versus 3.0% among 23- to 26-year-olds), and there has been little evidence of any convergence or divergence.

There also are long-established and large gender differences in all age groups in the prevalence of *occasional heavy drinking* or “binge drinking” (i.e., having five or more drinks in a row at least once in the past two weeks). Males in the 19- to 22-year-old band showed some longer-term decline in this statistic, from 54% in 1986 to 45% in 1995, thus narrowing the gender gap (from 24 percentage points in 1986 to 17 in 1995). After 1995 the rates for both genders have drifted up a few percentage points. Binge drinking among females in the same age band remained quite steady from 1980 through 2003, before giving some indication of an increase in 2004 (non-significant). In the two older age bands (23- to 26-year-olds and 27- to 30-year-olds), both the binge drinking rates and the sizable gender differences have been stable for the most part. However, from 1997 to 2002 among females, and from 1997 to 2004 among males, there was some increase in binge drinking in the 23- to 26-year-old group. The increases were from 22% to 29% among females and from 45% to 51% among males across the intervals stated.

- Most striking for *cigarette smoking* are the similarities between the genders in both absolute levels and in trends, though there are some differences. All three age groups showed a long-term decline in *daily smoking* rates for both males and females after data were first available for each: 19- to 22-year-olds from 1980 to 1990; 23- to 26-year-olds from 1984 to 1992; and 27- to 30-year-olds from 1988 to 1999. Male and female daily smoking rates have also been very close, particularly in the two older age groups. But among the 19- to 22-year-olds there was a crossover after 1993—up to that point females had slightly higher 30-day prevalence rates, but after that males did. Among the 23- to 26-year-olds, males reported higher 30-day smoking rates during the period 1998 to 2004. In the oldest age band, males reported higher rates in 2002 and they remain higher in 2004, but it is predictable that the difference will fade with time as the cohort effects make their way up the age bands.

There were some increases in the last decade and a half in 30-day smoking rates among the two younger groups and especially among the males. For example, from 1993 to 1999, 19- to 22-year-old males increased from 29% to 37%, while females increased from 29% to 34%. Because smoking rates in high school graduating classes after 1992 had been on the rise, and because we know that class cohorts tend to maintain their relative differences over time, we had predicted the increase in smoking among 19- to 22-year-olds and eventually in the older age bands as the heavier-smoking high school class cohorts grew older. Beginning in 1996,

smoking began to rise among the 23- to 26-year-olds, before leveling after 1998. Again, it rose more among males, opening a small gender gap. In 2004 an unusual increase is observed among males in both of the two younger strata. There was a large and statistically significant increase among males in both groups, with increases of 5.4% and 5.1% among the 19-22 and 23-26 year olds, respectively.

Regional Differences in Trends

The respondent's current state of residence was first asked in the 1987 follow-up survey; thus trend data by region exist only for the interval since then. In this case, changes have been examined for all 19- to 28-year-olds combined to increase the reliability of the estimates. Because gender and urbanicity crosscut all regions, they have less sampling error than when the sample is divided into four separate regions. (All regions are represented by between 1,000 and 2,200 cases in all years.) In general, the changes that have occurred since 1987 have been fairly consistent across regions, particularly in terms of the direction of the change.

- There were substantial drops in all four regions between 1987 (the initial measurement point) and 1991 for *any illicit drug, marijuana, any illicit drug other than marijuana, cocaine, crack, and amphetamines*. Since 1991 in most or all regions, there has been a leveling or increase in the use of these drugs (except *cocaine*, which continued to decline through the mid-1990s before beginning to inch up in the years since).
- The proportion of 19- to 28-year-olds using *any illicit drug* has been consistently lowest in the South and highest in the West and Northeast. For *marijuana* use, the South stands out as being consistently lowest, and for the most part the North Central has been second lowest. Generally, the other two regions have been fairly close to one another. For the use of *any illicit drug other than marijuana*, the West stood out as consistently highest, with the other three regions being very similar, at least until 2000; since 2001, use in the Northeast has been about as high as in the West.
- From 1991 through 1995 the West had slightly higher annual prevalence rates of *LSD* use than the other three regions among young adults (use dropped in 1995 in the West). Otherwise the usage rates have been quite similar in all four regions. All four regions have shown declines in LSD use in recent years, leaving small regional differences.
- Questions about *MDMA (ecstasy)* were added to the follow-up surveys of young adults in 1989. Through 1993, rates were highest in the West and South and lower in the Northeast and North Central regions. Subsequently, use in the Northeast began to increase (as was true among 12th graders), exceeding the levels of use found in the South and West from 1999-2001. The North Central has consistently had a much lower level of use than the other three regions. In 2000 all four regions showed a sharp and fairly parallel increase in ecstasy use; the rise decelerated in 2001 and began to decline thereafter in all regions. As we have discussed elsewhere, we believe that this decrease may be caused by a growing awareness of the hazards of ecstasy use. By 2003 and 2004, very little regional difference remained in annual prevalence, largely because the declines in use were most pronounced in the Northeast and the West.

- The declines in **cocaine** use, observed in all regions between 1987 and 1991, were greatest in the two regions that had attained the highest levels of use by the mid-1980s—the West and the Northeast. Thus, regional differences had diminished considerably by 1992. Similar to the finding for 12th graders, in 1992 these declines stalled in all regions except the Northeast. A gradual further decline then occurred in all regions through 1996 (1997 for the West) before a slight rise began to occur, likely reflecting the effects of generational replacement. Very little regional variability in cocaine use has existed since the mid-1990s.
- All four regions also exhibited an appreciable drop in **crack** use between 1987 and 1991, again with the greatest declines in the West and Northeast, where prevalence had been the highest. Use then generally leveled in all regions except the South, where it continued a gradual decline through 1997. As was true for cocaine generally, annual prevalence rates among the regions have converged; they now stand between 1.0% in the South and 1.9% in the West. (It is worth noting that lifetime use of crack stands out more in the West—and has for some years—compared to all other regions. For example, 5.6% of the 19- to 28-year-olds in the West in 2004 indicated having used crack at some time, compared to 3.5% to 4.2% in the other three regions.)
- From 1987 (when data were first available) through 1994, rates of **inhalant** use remained relatively stable, quite low, and about equal in all four regions among 19- to 28-year-olds. Annual use then rose in the Northeast in 1995 and 1996 and remained higher than in the other regions through 2000, when it dropped back to rates comparable to the other three regions. Except for that divergence, the regions have moved very much in parallel for this class of drugs.
- The regions have trended fairly similarly in their prevalence of **amphetamine** use by young adults. The only modest exception was that use declined more in the Northeast (which started out lowest) in the period 1987 to 1992, giving it a substantially lower rate than the other three regions; it remained lowest until 1998. (The West has fairly consistently had the highest rate, but not by much.) By the late 1990s, the Northeast had caught up to the North Central and South, making the regional differences very small, and there essentially have been no regional differences since 2000.
- **Methamphetamine** use in general has only been measured since 1999 (though crystal methamphetamine, discussed next, has been in the study for a longer time). It shows some divergence in rates among the regions and some differential trending, with a gradual rise in annual prevalence in the West (where use tends to be highest) and a gradual decline in the Northeast, where use is now lowest. Use in the other two regions has remained fairly flat. Lifetime prevalence has been particularly high in the West, ranging between 13% and 17% since 1999.
- The West has consistently had the highest rates for **ice (crystal methamphetamine)**, and the regional differences have been very substantial, particularly in terms of lifetime use. The Northeast generally has had the lowest rates. In fact, when data were first available on ice in 1990, the West had a lifetime prevalence of 5.1% versus a range of 1.7% to 2.3% in the other three regions. By 2004, the lifetime prevalence rate in the West increased to 8.0%, versus

4.8% in the North Central, 4.4% in the South, and 1.9% in the Northeast. This strongly suggests that ice use diffused from the West primarily to the South and North Central regions but diffused much less to the Northeast. The annual prevalence figures tell a similar story but also show that there was a bulge in use in the West in 1994 through 1996 before use there declined and then stabilized at around 2% annual prevalence.

- The use of *sedatives (barbiturates)* remained flat, and at about equivalent levels, in all four regions of the country from 1987, when regional data were first available, through 1994. Rates then rose gradually in all regions for a number of years.
- The picture for *tranquilizers* is quite similar to that for sedatives (barbiturates). The regional differences have been small, though the South tends to have a slightly higher rate than the other regions. Use generally declined in all regions from 1987 through 1993. Since then there has been some increase in all regions with the South experiencing the most, where annual prevalence stands at 8.7% in 2004 versus 5.7% to 7.2% in the other regions.
- Levels and trends in *heroin* use have been quite comparable across the four regions since 1987. All regions had low and stable rates up into the early 1990s. A gradual increase was observed from about 1993 through 2000, and there has been some decline or at least leveling since then.
- Trends in the prevalence of the use of *narcotics other than heroin* have been quite parallel for the four regions. A gradual and long-term increase has occurred since the mid 1990s.
- The annual prevalence of the specific narcotic drug, *OxyContin*, has risen some in all four regions since it was first measured in 2002. Annual prevalence of use for *Vicodin* has shown little net change in use since 2002 in three of the four regions, but shows an increase in the West, where annual prevalence is now highest at 13.0%. It should be noted that the sample sizes are more limited than usual for these two drugs, because questions about them occur on only two of the six questionnaire forms.
- The Northeast stood out in 2002, when these drugs were first measured, as having a higher rate of use of the two club drugs, *ketamine* and *GHB*; but use has dropped in the past two years in the Northeast, bringing that region's usage rates down to the same low level as the other three regions. *Rohypnol* use has remained very low in all four regions over the same interval.
- With respect to *alcohol* use, there were modest declines in 30-day prevalence in all four regions between 1987 (when the first measurement was available for 19- to 28-year-olds) and 1992. The rates for 30-day prevalence then leveled in all regions. The West and the South have consistently had lower rates of 30-day use than the Northeast and North Central, (as has generally been true among 12th graders). In 2004 the 30-day prevalence rates for each stratum are about where they were in 1992, with all regions showing a very flat pattern of prevalence over that 12-year interval.

Current *daily use* of alcohol also showed a decline from the first (1987) data collection through about 1994 or 1995 in all regions. The proportional declines were substantial—on the order of 40%-50%. (This decline corresponds to a period of appreciable decline in daily drinking among 12th graders, though we can tell from their longer-term data that their decline started in 1980; thus the decline may well have started earlier among 19- to 28-year-olds, as well.) Since the mid-1990s there has been a slight upward drift in daily prevalence in all regions; the rates are all between 3.7% and 4.9% in 2004.

- **Occasional heavy drinking** (or “binge drinking”) has remained fairly level in all regions since it was first measured in 1987. The rates have consistently been appreciably higher in the Northeast (41% in 2004) and the North Central (42%) than in the South and the West (32% and 34%, respectively). (Similar regional differences are evident among 12th graders.)
- There have been highly consistent regional differences among young adults in **cigarette smoking** since data were first available in 1987—they exist for monthly, daily, and half-pack-daily prevalence rates. The West consistently has had the lowest rates (e.g., 16% daily prevalence in 2004) and the South the next lowest (20% in 2004); the Northeast and North Central have the highest smoking rates at 23% and 24%, respectively, in 2004. After some slight decline in 30-day prevalence in all regions between 1987 and 1989, rates leveled off for about five years (roughly through 1994). There then followed a very gradual increase of a few percentage points through 1998, followed by a leveling. Daily use showed a very similar pattern. For half-pack-a-day smoking, the decline phase was longer (from 1987 through about 1992 or 1993), likely reflecting the lag between smoking initiation and regular heavy smoking. Since 1998, half-pack smoking rates have gradually declined in all four regions.

Population Density Differences in Trends

The analyses presented here for population density return to the use of four-year age groupings, which allows a longer time interval to be examined for the younger strata and for cross-age comparisons of the trends. Among the young adults, five levels of population density are distinguished based on the respondent’s answer: very large city, large city, medium-sized city, small town, and farm/country.

- The proportions of young adults using **any illicit drug** have moved in parallel among the various community size strata. In general, the farm/country stratum has tended to have lower use than all of the other strata. The other four strata have tended to differ little from one another, though the very large cities have generally ranked at the top. In 2004 the proportions of 19- to 22-year-olds reporting use of an illicit drug in the past year were 30% for the farm/country strata, 37% for small towns and medium-sized cities, 36% for large-sized cities, and 45% for very large cities.
- The use of **any illicit drug other than marijuana** tells a similar story. There was a long period of fairly parallel decline before leveling, along with some convergence of usage rates among the strata at all three age levels. In general, small, large, and very large cities all have tended to have about the same rates, and the farm/country stratum has tended to have the lowest rates, particularly prior to 1990; the differences by population density have been quite small since about 2000.

- **Marijuana** use has moved pretty much in parallel among the various strata over the time intervals for which data exist. Among 19- to 22-year-olds, the rates have been quite close among all the strata, except for the farm/country stratum. The most rural region has consistently had the lowest rate of marijuana use, and it fell less in the earlier period and rose more slowly in the subsequent increase than it did the other strata. Use also has tended to be lower in the more rural areas in the older two age bands, as well. Daily use of marijuana also has moved very much in parallel among the five population density strata within each age band.
- In general there have not been large differences in **LSD** use among young adults as a function of community size since 1983. Among the 19- to 22-year-olds (the young adult age group with by far the highest rates of LSD use), use in communities of all sizes declined appreciably in the early- to mid-1980s, particularly in the urban strata, eliminating modest prior differences by 1984. From around 1989 through 1996, there was some increase in LSD use in all strata among the 19- to 22-year-olds, with the most rural region generally continuing to have the lowest prevalence (though this has not always been true since 1998). Since about 1997, there has been a substantial decline in LSD use in all strata among the 19- to 22- year olds—the age band with by far the highest rates of use to begin with. The 23- to 26-year-old respondents had some modest increases after 1989 in all strata, though the increases had virtually ended by 1995; since about 1999, there have been declines in all strata. In the oldest age group, LSD has remained very low and for the most part quite stable, with some evidence of a decline in the last few years.
- The use of **hallucinogens other than LSD**, taken as a class, also has shown considerably higher rates in the youngest age band than in the two older ones, suggesting a sharp falloff in use with age. Use of this class of drugs fell in communities of all sizes among the young adults between 1980 and about 1988. Then there was a leveling of use for a few years, followed by an extended increase in use among all strata in the 19- to 22-year-old age band. By 2003 the rates attained by each stratum exceeded those originally observed in 1980. The 23- to 26-year-old group has shown slightly higher rates since 1998 than previously. The sharpest increase occurred in the very large cities in 1999 and 2000, possibly as a result of growing ecstasy use. Among 27- to 30-year-olds the trend lines have been very flat with only minor stratum differences, until 2001 when all strata, especially the very large cities, began to increase. In 2004, there was some decline observed in four of the five population density strata among 19- to 22-year-olds, but a more mixed pattern of change in the older groups.
- **Ecstasy (MDMA)** use was first measured in 1989 and since then has shown the largest increase among the younger adults of any of the drugs. Use in 1989 was highest among the 19- to 22-year-olds in the very large cities (5% annual prevalence); but prevalence declined in all strata between 1989 and 1994 (to 1.6% or less). By 1998, use had begun to increase in all strata within this age band, except among the farm/country stratum. The farm/country stratum moved up sharply in 1999, but then the three most urban strata jumped sharply in 2000, opening a fair gap in use as a function of population density, with the large and very large cities having rates nearly twice as high as any of the other strata in 2002. All strata showed large declines in ecstasy use since 2000 or 2001. Among the 23- to 26-year-olds use began to increase a little later, and again the most urban strata showed the most increase,

particularly in 2000; but all three urban strata showed a decline in 2001, narrowing the differences among the strata. Considerably less increase in use occurred among 27- to 30-year-olds, though there was some increase in the largest cities starting after 1996 and in the large and medium-sized cities after 1999. In the last one to four years, all strata in all three age groups have shown declines in their ecstasy use.

Ecstasy use trends in the past four years tell an interesting story. In the very large cities, where use had spiked early, use peaked in all three age bands in 2000 and then began to decline. The medium-sized cities were beginning to level or decline in all three age bands. The small town and farm/country strata peaked in 2001 in all age groups. These data support our analysis, based on the secondary schools, suggesting that the presence of this drug was still diffusing geographically—in this case from more urban to more rural areas—and, were it not for this continued diffusion, ecstasy use would actually have declined nationally a year earlier. The data from 12th graders on perceived risk provide the clue as to the most likely cause of this turnaround. They showed a large jump in the level of perceived risk associated with ecstasy use in all of the past three years. Unlike most of the other drugs that we have been discussing here, the pattern of change since the mid-1990s appears to reflect secular trends more than cohort effects.

- In the early 1980s, *cocaine* use was positively correlated with population density, with the highest use in the very large cities. The important drop in cocaine use that began after 1986 slowed considerably after 1992 or 1993 in all three of the age strata and in communities of all sizes, by which time the positive association with population density had been virtually eliminated. Among the 19- to 22-year-olds there has been a sustained increase in cocaine use among all strata since about 1993 or 1994 and among the 23- to 26-year-olds since about 1998. As just stated, usage rates among the strata tended to converge considerably during the period of decline, and this convergence remains, with the very large cities showing rates of cocaine use only slightly higher than the less densely populated areas. In the 27- to 30-year-old age group, a gradual increase in use has emerged in nearly all strata since 2000, no doubt reflecting a cohort effect. By 2004 all of these strata in the oldest age band leveled or declined, the single exception being among the farm/country stratum.
- *Crack* use among all age groups peaked in 1987 or 1988 (strongly suggesting a secular trend at work at that time) and, after declining, bottomed out in all population-density strata for several years. Among the 19- to 22-year-olds only, it may have made some comeback in the rural and small town strata in 1999 and 2000 but not in the larger cities. The crack use reported in these young adult samples at all three age levels has borne practically no systematic association with community size, and for the most part they have all trended in parallel.
- *Amphetamine* use shows virtually no differences in use associated with urbanicity in any of the three age groups, and this has been fairly consistently true since 1983.
- The use of *crystal methamphetamine (ice)*, first measured in 1990, has shown a modest increase in use since the early 1990s among the younger two young adult strata, and since the late 1990s among the oldest stratum. This has been observable in most population density

groupings. There have not been any sustained differences in use as a function of population density.

- **Methaqualone** use, which in 1981 was rather strongly associated (positively) with population density, dropped to annual prevalence rates of 0.8% or below in all size strata for all three age bands by 1989. Its use is no longer measured in the study.
- Unlike methaqualone, **sedatives (barbiturates)** have never shown much correlation with urbanicity, at least as far back as 1980. This remains true in all three age bands.
- **Tranquilizer** use among young adults also has had little or no association with population density over this time interval. However, in 2000 through 2002, tranquilizer use increased considerably in the very large cities in all three age groups, opening up a temporary differential.
- From 1980 to 1995, annual **heroin** prevalence was less than 1.0%—usually much less—in all strata for all three of the age bands. After 1994, use among 19- to 22-year-olds in all strata rose and reached 1.0% in the three urban strata by 1998. In fact, in the very large cities, it reached 1.6% in 1996 (versus 0.3% to 0.7% in the other strata). Use of heroin generally has been highest in the very large cities in this age band; but an across-the-board decrease by 2002 leaves rather little difference across the strata. Use levels are lower among the 23- to 26-year-olds and lower still among the 27- to 30-year-olds, and it is difficult to discern systematic differences among the population density strata in those age bands.
- The annual use of **narcotics other than heroin** had some positive association with degree of population density among 19- to 22-year-olds through the early 1990s; however, it has shown rather little association since then. Since 1993 in the case of 19- to 22-year-olds, since about 1996 in the case of 23- to 26-year-olds, and since about 1998 in the case of the 27- to 30-year-olds, use of narcotics other than heroin has increased very substantially in all community size strata but no systematic differentiation by community size is evident. Clearly a cohort effect is at work, but the increased use of these drugs is very widespread.
- Unfortunately, the sample sizes for two of the narcotic drugs of particular interest, **OxyContin** and **Vicodin**, are not sufficient to estimate population density differences or trends with a sufficient degree of accuracy.
- The absolute levels of **inhalant** use have remained low in these age groups, particularly above age 22. However, during the mid- to late 1980s, there was a gradual increase in use among 19- to 22-year-olds in all community-size strata. No strong or consistent association with population density has appeared, though the very large cities generally have tended to have higher rates than the other areas among 19- to 22-year-olds, particularly in the period 1998 through 2000.
- There have been few differences as a function of population density in the 30-day prevalence of drinking **alcohol** among 19- to 22-year-olds, since data were first available on them in 1980, except for the fact that the farm/country stratum has tended to have lower-than-average

use. In the two older age bands, however, there has been a fairly consistent positive correlation between urbanicity and use of alcohol in the past 30 days. But there have been no consistent differences in current ***daily drinking*** associated with urbanicity in any of the three age bands. For ***occasional heavy drinking***, all strata have been fairly close across time at all three age levels, again with the exception that the farm/country areas have fairly consistently shown the lowest rates of binge drinking at all ages.

- ***Cigarette smoking*** has been negatively associated with urbanicity in all three age strata, without much evidence of differential trends related to degree of urbanicity, with one exception. Among 19- to 22-year-olds, all smoking prevalence measures rose from 1997 through 1999 in the farm/country and small town strata, while most other strata remained level. The differences in 1999 were most striking for half-pack-a-day smoking among the 19- to 22-year-olds: farm/country (24% prevalence), small town (19%), medium-sized and large cities (both 15%), and very large cities (10%). This compares with 1985, when there was virtually no difference in half-pack-a-day smoking rates among these strata (all were at 18% or 19%). Thus, smoking among those in their early 20s has become more concentrated in the non-urban populations. Since 2000 there was a decline in 30-day prevalence in all strata (except for the farm/country) among the 19- to 22-year-olds. Continuing declines in smoking among 12th graders would lead us to expect still further declines in the young adults, as well. While smoking has been dropping among 19- to 22-year-olds, use generally has remained negatively correlated with the degree of urbanicity. In 2004, however, there was a large and significant increase in 30-day smoking prevalence in the very large cities (up 7.9 percentage points), as well as smaller and not statistically significant increases among 23- to 26-year-olds in the very large city and large city strata (up 2.3 and 3.1 percentage points, respectively). It remains to be seen if these findings indicate the emergence of a new trend in smoking among young adults.

TABLE 5-1
Trends in Lifetime Prevalence of Various Types of Drugs
Among Respondents of Modal Ages 19-28
(Entries are percentages)

	Percentage who used in lifetime																				'03-'04 change
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004		
<i>Approx. Wtd. N = (6900) (6800) (6700) (6600) (6700) (6600) (6800) (6700) (6500) (6400) (6300) (6400) (6200) (6000) (5700) (5800) (5300) (5300) (5700)</i>																					
Any Illicit Drug ^a	70.5	69.9	67.9	66.4	64.5	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	+0.3	
Any Illicit Drug ^a Other Than Marijuana	48.4	47.0	44.6	42.7	40.8	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3	31.6	32.8	33.9	35.2	+1.3	
Marijuana	66.5	66.0	63.8	62.8	60.2	58.6	56.4	55.9	53.7	53.6	53.5	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	+0.2	
Inhalants ^b	12.3	12.7	12.6	13.2	12.5	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	-0.6	
Inhalants, Adj. ^{b,c}	18.6	15.7	15.0	NA	13.5	14.1	13.9	14.5	13.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Nitrites ^d	2.6	6.9	6.2	NA	1.9	1.4	1.2	1.3	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Hallucinogens ^e	18.5	17.1	17.0	15.9	16.1	15.7	15.7	15.4	15.4	16.1	16.4	16.7	17.4	18.0	18.4	18.3	19.6	19.7	19.3	-0.4	
Hallucinogens, Adj. ^{e,f}	20.1	17.2	17.2	NA	16.5	16.0	15.9	15.5	15.5	16.2	16.5	16.7	17.5	18.2	18.5	18.4	19.6	20.0	19.6	-0.3	
LSD	14.6	13.7	13.8	12.7	13.5	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	-1.2	
PCP ^g	8.4	4.8	5.0	NA	2.5	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	-0.3	
MDMA (Ecstasy) ^h	NA	NA	NA	3.3	3.7	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	+0.8	
Cocaine	32.0	29.3	28.2	25.8	23.7	21.0	19.5	16.9	15.2	13.7	12.9	12.0	12.3	12.8	12.7	13.1	13.5	14.7	15.2	+0.4	
Crack ⁱ	NA	6.3	6.9	6.1	5.1	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	-0.5	
Other Cocaine ⁱ	NA	28.2	25.2	25.4	22.1	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	+0.9	
Heroin	1.3	1.3	1.1	1.0	0.9	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	0.0	
Other Narcotics ^{k,l}	10.7	10.6	9.8	9.6	9.4	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5	13.9	16.8	17.6	+0.8	
Amphetamines, Adj. ^{k,m}	32.3	30.8	28.8	25.3	24.4	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	+0.7	
Ice ⁿ	NA	NA	NA	NA	2.5	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	0.0	
Sedatives (Barbiturates) ^l	11.1	9.7	8.9	7.9	8.7	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	+0.9	
Sedatives, Adj. ^{k,r}	16.7	15.0	13.2	12.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Methaqualone ^k	13.1	11.6	9.7	8.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Tranquilizers ^{e,k}	17.6	16.5	15.1	13.5	12.9	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5	11.9	13.4	13.8	14.9	+1.1	
Alcohol ^o	94.8	94.9	94.8	94.5	94.3	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	0.0	
Flvd. Alcoholic Bvg. ^d	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	83.2	—	
Cigarettes	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Steroids ^p	NA	NA	NA	1.1	1.2	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	+0.1	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

'NA' indicates data not available.

See footnotes on next page.

FOOTNOTES FOR TABLES 5-1 THROUGH 5-4

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in four of the five questionnaire forms in 1986-1989, five of the six questionnaire forms in 1990-1998, and three of six questionnaire forms in 1999-2004. Total N in 2004 is approximately 2700.

^cAdjusted for underreporting of amyl and butyl nitrites. Questions about nitrite use were dropped beginning in 1995.

^dThis drug was asked about in one questionnaire form. Total N in 1994 was approximately 900. Total N in 2004 is approximately 950.

^eIn 2001 the question text was changed on half of the questionnaire forms. "Other psychedelics" was changed to "other hallucinogens," and "shrooms" was added to the list of examples. For tranquilizers, Miltown was replaced with Xanax. Beginning in 2002 the remaining forms were changed to the new wording.

^fAdjusted for underreporting of PCP.

^gThis drug was asked about in one of the five questionnaire forms in 1986-1988, and in one of the six questionnaire forms in 1990-2004. Total N in 2004 is approximately 950.

^hThis drug was asked about in two of the six questionnaire forms in 1990-2001, and in three of the six questionnaire forms in 2002-2004.

ⁱThis drug was asked about in two of the five questionnaire forms in 1987-1989, in all six questionnaire forms in 1990-2001, and in five of the six questionnaire forms in 2002-2004. Total N in 2004 is approximately 4700.

^jThis drug was asked about in one of the five questionnaire forms in 1987-1989, and in four of the six questionnaire forms in 1990-2004.

^kOnly drug use that was not under a doctor's orders is included here.

^lIn 2002 the question text was changed on half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric--all of which had negligible rates of use by 2001--were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one-half of N indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003 and beyond.

^mBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

ⁿThis drug was asked about in two of the six questionnaire forms in 1990-2004. Total N in 2004 is approximately 1900.

^oIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a "drink" meant "more than just a few sips." Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

^pThis drug was asked about in one of the five questionnaire forms in 1989, and in two of the six questionnaire forms in 1990-2004.

^qThis drug was asked about in two of the six questionnaire forms.

^r"Sedatives, adjusted" data are a combination of barbiturate and methaqualone data.

TABLE 5-2
Trends in Annual Prevalence of Various Types of Drugs
Among Respondents of Modal Ages 19-28
(Entries are percentages)

	Percentage who used in last 12 months																			'03-'04 change	
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004		
<i>Approx. Wtd. N = (6900) (6800) (6700) (6600) (6700) (6600) (6800) (6700) (6500) (6400) (6300) (6400) (6200) (6000) (5700) (5800) (5300) (5300) (5700)</i>																					
Any Illicit Drug ^a	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	+0.6	
Any Illicit Drug ^a																					
Other Than Marijuana	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	+0.7	
Marijuana	36.5	34.8	31.8	29.0	26.1	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	+0.2	
Inhalants ^b	1.9	2.1	1.8	1.9	1.9	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	+0.3	
Inhalants, Adj. ^{b,c}	3.0	2.8	2.4	NA	2.1	2.2	1.9	2.3	2.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Nitrites ^d	2.0	1.3	1.0	NA	0.4	0.2	0.1	0.4	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Hallucinogens ^e	4.5	4.0	3.9	3.6	4.1	4.5	5.0	4.5	4.8	5.6	5.6	5.8	5.2	5.4	5.4	5.4	4.7	5.2	4.7	-0.5	
Hallucinogens, Adj. ^{e,f}	4.9	4.1	3.9	NA	4.2	4.6	5.1	4.6	4.9	5.7	5.6	5.9	5.2	5.5	5.5	5.5	4.7	5.2	4.7	-0.5	
LSD	3.0	2.9	2.9	2.7	3.3	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	-0.3	
PCP ^g	0.8	0.4	0.4	NA	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	-0.2	
MDMA (Ecstasy) ^h	NA	NA	NA	1.4	1.5	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	-1.0	
Cocaine	19.7	15.7	13.8	10.8	8.6	6.2	5.7	4.7	4.3	4.4	4.1	4.6	4.9	5.4	5.4	5.8	5.8	6.6	7.1	+0.5	
Crack ⁱ	3.2	3.1	3.1	2.5	1.6	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	+0.3	
Other Cocaine ^j	NA	13.6	11.9	10.3	8.1	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	+0.3	
Heroin	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	-0.1	
Other Narcotics ^{k,l}	3.1	3.1	2.7	2.8	2.7	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0	7.1	8.5	9.0	+0.5	
OxyContin ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.9	2.6	3.1	+0.5	
Vicodin ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	8.2	8.6	8.9	+0.3	
Amphetamines, Adj. ^{k,m}	10.6	8.7	7.3	5.8	5.2	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	+0.5	
Ritalin ^{k,q}	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	2.9	2.9	2.7	-0.2	
Methamphetamine ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	2.8	2.5	2.8	2.5	2.7	+0.1	
Ice ⁿ	NA	NA	NA	NA	0.4	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	+0.3	
Sedatives (Barbiturates) ^k	2.3	2.1	1.8	1.7	1.9	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	+0.5	
Sedatives, Adj. ^{k,r}	3.0	2.5	2.1	1.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Methaqualone ^k	1.3	0.9	0.5	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Tranquilizers ^{e,k}	5.4	5.1	4.2	3.7	3.7	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6	5.5	7.0	6.8	7.4	+0.6	
Rohypnol ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.3	0.5	0.1	-0.4	
GHB ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.8	0.6	0.5	-0.1	
Ketamine ^q	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.2	0.9	0.6	-0.3	
Alcohol ^o	88.6	89.4	88.6	88.1	87.4	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	+1.1	
Flvd. Alcoholic Bvg. ^d	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	62.7	—	
Cigarettes	40.1	40.3	37.7	38.0	37.1	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.1	39.1	38.6	39.0	+0.5	
Steroids ^p	NA	NA	NA	0.5	0.3	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.5	0.5	0.5	0.0	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

*s' indicates a prevalence rate of less than 0.05% but greater than true zero.

'NA' indicates data not available.

See footnotes at end of Table 5-1.

TABLE 5-3
Trends in Thirty-Day Prevalence of Various Types of Drugs
Among Respondents of Modal Ages 19-28
(Entries are percentages)

	Percentage who used in last 30 days																			'03-'04 change
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
<i>Approx. Wtd. N = (6900) (6800) (6700) (6600) (6700) (6600) (6800) (6700) (6500) (6400) (6300) (6400) (6200) (6000) (5700) (5800) (5300) (5300) (5700)</i>																				
Any Illicit Drug ^a	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	-0.8
Any Illicit Drug ^a Other Than Marijuana	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	+0.2
Marijuana	22.0	20.7	17.9	15.5	13.9	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	-0.8
Inhalants ^b	0.4	0.6	0.6	0.5	0.6	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	-0.1
Inhalants, Adj. ^{b,c}	0.7	0.9	0.9	NA	0.7	0.6	0.7	0.7	0.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Nitrites ^d	0.5	0.5	0.4	NA	0.1	*	0.1	0.2	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Hallucinogens ^e	1.3	1.2	1.1	1.1	0.9	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2	1.2	0.9	1.2	0.9	-0.3
Hallucinogens, Adj. ^{e,f}	1.4	1.2	1.1	NA	1.0	1.2	1.6	1.2	1.4	1.7	1.3	1.5	1.5	1.3	1.2	1.2	0.9	1.2	0.9	-0.3
LSD	0.9	0.8	0.8	0.8	0.6	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.0
PCP ^g	0.2	0.1	0.3	NA	0.2	0.1	0.2	0.2	0.1	0.0	0.1	0.1	0.2	0.2	0.0	0.0	0.1	0.1	0.1	0.0
MDMA (Ecstasy) ^h	NA	NA	NA	0.4	0.2	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	-0.1
Cocaine	8.2	6.0	5.7	3.8	2.4	2.0	1.8	1.4	1.3	1.5	1.2	1.5	1.7	1.9	1.7	2.2	2.2	2.4	2.2	-0.1
Crack ⁱ	NA	1.0	1.2	0.7	0.4	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	+0.1
Other Cocaine ⁱ	NA	4.8	4.8	3.4	2.1	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	0.0
Heroin	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.0
Other Narcotics ^{k,l}	0.9	0.9	0.7	0.7	0.7	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7	2.9	2.9	3.0	+0.1
Amphetamines, Adj. ^{k,m}	4.0	3.2	2.7	2.1	1.9	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	-0.1
Ice ⁿ	NA	NA	NA	NA	0.1	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.4	0.4	0.0
Sedatives (Barbiturates) ^k	0.7	0.7	0.7	0.5	0.6	0.5	0.5	0.6	0.6	0.8	0.8	0.9	0.9	1.1	1.3	1.7	1.5	1.5	1.8	+0.3
Sedatives, Adj. ^{k,r}	0.9	0.8	0.7	0.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Methaqualone ^k	0.3	0.2	0.1	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Tranquilizers ^{s,k}	1.8	1.6	1.4	1.2	1.1	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8	2.1	2.8	2.4	2.7	+0.3
Alcohol ^o	75.1	75.4	74.0	72.4	71.2	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.2	68.3	67.0	68.4	+1.4
Flvd. Alcoholic Bvg. ^d	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	29.5	—
Cigarettes	31.1	30.9	28.9	28.6	27.7	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	+0.9
Steroids ^p	NA	NA	NA	0.2	0.1	0.2	0.1	0.0	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.2	0.1	-0.2

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

*' indicates a prevalence rate of less than 0.05% but greater than true zero.

'NA' indicates data not available.

See footnotes at end of Table 5-1.

TABLE 5-4
Trends in Thirty-Day Prevalence of Daily Use of Various Types of Drugs
Among Respondents of Modal Ages 19-28
(Entries are percentages)

	Percentage who used daily in last 30 days																			'03-'04 change
	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
<i>Approx. Wtd. N = (6900) (6800) (6700) (6600) (6700) (6600) (6800) (6700) (6500) (6400) (6300) (6400) (6200) (6000) (5700) (5800) (5300) (5300) (5700)</i>																				
Marijuana	4.1	4.2	3.3	3.2	2.5	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	-0.3
Cocaine	0.2	0.1	0.2	0.1	*	0.1	*	0.1	*	0.1	*	*	*	0.1	*	0.1	*	*	0.1	+0.1
Amphetamines, Adj. ^{k,m}	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.3	0.2	-0.1
Alcohol																				
Daily ^o	6.1	6.6	6.1	5.5	4.7	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	-0.5
5+ Drinks in a Row in Last 2 Weeks	36.1	36.2	35.2	34.8	34.3	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	+1.3
Cigarettes																				
Daily	25.2	24.8	22.7	22.4	21.3	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	+0.5
Half-Pack or More per Day	20.2	19.8	17.7	17.3	16.7	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	-0.4

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

The illicit drugs not listed here show a daily prevalence of 0.2% or less in all years.

*' indicates a prevalence rate of less than 0.05% but greater than true zero.

See footnotes at end of Table 5-1.

TABLE 5-5
Trends in Annual and Thirty-Day Prevalence of an Illicit Drug Use Index^a
Among Respondents of Modal Ages 19-28
(Entries are percentages)


	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Lifetime																				
Any Illicit Drug	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	+0.6
Males	45.3	42.6	39.5	35.7	33.6	30.0	31.4	31.1	32.3	32.1	31.6	31.9	33.6	33.9	34.4	34.9	35.6	36.0	37.0	+1.0
Females	39.0	36.5	33.6	30.5	28.3	24.5	25.8	26.1	25.3	28.1	27.3	27.1	27.1	27.6	28.2	30.1	30.2	31.0	31.4	+0.3
Any Illicit Drug																				
Other Than Marijuana	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	+0.7
Males	30.4	26.5	23.8	21.0	19.1	16.4	16.3	14.7	16.2	16.2	15.4	15.6	16.2	16.7	17.8	17.2	18.9	19.8	21.3	+1.5
Females	24.0	21.6	19.4	16.2	14.7	12.5	12.2	11.6	10.5	12.0	11.4	12.0	11.0	11.5	12.9	14.1	14.6	17.0	17.1	+0.1
Past 30 days																				
Any Illicit Drug	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	-0.8
Males	29.9	27.1	23.7	21.1	18.8	18.3	17.9	17.4	19.5	18.6	19.0	19.8	20.1	20.0	21.5	21.9	22.8	22.4	23.1	+0.7
Females	22.2	20.2	17.8	15.0	13.5	12.5	12.4	12.9	12.1	13.5	13.3	13.8	13.2	15.0	15.6	16.6	16.3	18.3	16.3	-1.9 s
Any Illicit Drug																				
Other Than Marijuana	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	+0.2
Males	15.2	12.3	10.6	9.1	6.8	6.6	6.5	5.9	7.1	6.8	5.7	6.8	7.1	7.3	7.8	8.1	8.5	9.2	10.6	+1.3
Females	11.0	9.4	8.7	6.2	5.3	4.4	4.7	4.0	3.9	4.8	4.0	4.5	4.4	5.1	5.4	6.3	7.1	7.7	7.1	-0.6
<i>Approximate Weighted N</i>																				
All Respondents	6900	6800	6700	6600	6700	6600	6800	6700	6500	6400	6300	6400	6200	6000	5700	5800	5300	5300	5700	
Males	3200	3100	3000	2900	3000	3000	3000	3000	2900	2800	2700	2800	2700	2600	2400	2400	2200	2200	2300	
Females	3700	3700	3700	3700	3700	3600	3700	3700	3600	3600	3600	3600	3500	3400	3300	3400	3100	3100	3400	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.


FIGURE 5-2
Any Illicit Drug Other than Marijuana: Trends in Annual Prevalence
Among High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	25.4	26.0	27.1	28.2	30.4	34.0	30.1	28.4	28.0	27.4	25.9	24.1	21.1	20.0	17.9	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4	21.6	20.9	19.8	20.5
19-20 Years		28.6	30.2	33.3	34.2	32.4	29.8	27.5	26.9	24.7	22.2	21.3	17.6	16.5	13.8	13.4	13.5	14.6	18.6	17.4	17.6	17.3	18.7	19.6	18.0	19.6	19.9	20.2	
21-22 Years			35.5	37.0	34.2	33.7	31.6	29.5	29.1	25.6	22.8	19.4	17.4	14.9	15.4	13.5	14.1	15.2	13.7	17.7	15.3	14.1	17.0	20.0	18.9	20.7	21.2		
23-24 Years				35.4	33.2	29.4	33.4	29.3	22.6	21.1	18.8	17.5	14.6	14.8	12.9	12.9	11.5	13.1	12.1	12.9	14.8	15.0	14.1	17.2	20.1	21.2			
25-26 Years					30.2	30.3	25.5	25.7	21.0	17.6	16.6	14.4	13.4	13.0	12.0	11.6	10.0	10.7	10.8	11.6	12.5	13.3	14.6	14.5	16.3				
27-28 Years						26.5	23.3	20.4	18.2	15.2	13.6	13.2	11.5	11.1	10.9	10.7	8.4	8.9	8.6	9.9	11.4	11.4	15.1	14.6					
29-30 Years							20.0	17.4	12.4	13.2	11.6	9.9	10.8	11.0	10.3	11.0	7.8	8.1	7.4	9.9	10.9	11.6	11.8						
31-32 Years*								13.8	13.1	10.7	9.5	11.5	8.2	10.2	10.8	9.6	8.3	7.4	9.7	NA	NA	NA							
35 Years									11.2	10.4	11.4	10.0	8.2	9.3	9.3	8.8	9.6	9.5	11.0										
40 Years										9.3	7.9	7.7	7.3	9.7	6.7	8.3													
45 Years											8.9	9.3																	


FIGURE 5-3a
Marijuana: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	44.5	47.6	50.2	50.8	48.8	46.1	44.3	42.3	40.0	40.6	38.8	36.3	33.1	29.6	27.0	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3
19-20 Years			52.8	51.0	49.7	49.0	44.9	43.0	41.4	40.3	39.1	35.8	36.2	32.2	28.4	25.4	26.9	27.9	29.3	31.8	34.2	34.8	37.2	37.9	37.0	35.4	36.4	35.9	34.5
21-22 Years				50.1	51.1	45.8	45.4	42.1	40.9	39.6	37.4	33.7	31.6	28.2	26.8	26.9	26.1	29.2	28.1	30.6	30.6	31.9	31.5	33.2	37.5	34.3	33.1	32.5	
23-24 Years					46.0	43.8	38.6	42.0	36.6	33.7	32.0	27.3	26.6	23.2	26.6	26.5	24.6	25.8	25.8	25.1	25.5	27.4	26.9	28.3	31.8	30.0	27.7		
25-26 Years						38.3	39.2	34.1	35.4	29.7	26.2	24.1	21.8	23.5	22.2	22.6	24.4	21.7	23.3	21.2	21.8	22.7	25.0	24.5	24.3	27.6			
27-28 Years							32.5	31.4	26.7	26.8	22.6	20.9	21.2	21.3	20.1	20.4	20.6	18.0	19.9	18.2	18.8	19.4	19.4	21.2	22.4				
29-30 Years								25.4	24.7	20.0	21.0	20.1	18.8	19.0	18.2	19.5	18.0	16.9	16.0	18.4	17.1	17.5	17.0	16.4					
31-32 Years*									19.8	19.9	17.7	19.9	18.6	17.2	18.6	16.7	15.8	14.8	14.5	16.7	NA	NA	NA						
35 Years																14.5	17.2	16.3	17.5	14.9	14.7	13.8	14.8	13.7	13.0	13.0			
40 Years																					17.1	13.8	13.7	12.5	14.6	13.4	13.9		
45 Years																												14.0	11.9


FIGURE 5-3c
Marijuana: Trends in Thirty-Day Prevalence of Daily Use
Among High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	8.2	9.1	10.7	10.3	9.1	7.0	6.3	5.5	5.0	4.9	4.0	3.3	2.7	2.9	2.2	2.0	1.9	2.4	3.5	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	
19-20 Years			10.5	10.9	8.1	7.9	6.6	5.2	4.7	4.6	3.5	3.4	3.5	2.8	2.3	2.1	1.4	2.3	3.1	4.7	4.9	5.4	5.2	6.2	6.0	6.1	6.0	6.5	6.0	
21-22 Years				10.9	9.4	6.4	6.2	5.3	4.5	4.1	3.9	3.5	3.1	2.5	2.4	2.6	2.3	2.9	3.4	3.2	5.3	5.2	4.6	5.5	7.0	6.0	6.0	6.0	5.1	
23-24 Years					8.1	6.7	5.5	5.8	4.9	4.3	3.1	3.0	2.7	2.1	2.3	2.7	3.1	3.3	2.3	2.6	3.1	2.6	3.1	5.1	3.8	4.7	5.5	6.6	5.3	
25-26 Years						6.0	6.1	3.6	5.0	3.4	3.3	2.7	2.5	2.6	2.5	2.7	2.3	3.1	2.5	2.4	3.1	2.5	2.4	3.1	3.4	4.6	2.7	3.5	5.5	
27-28 Years							4.8	4.6	3.0	4.1	2.4	2.6	2.5	2.3	2.2	2.5	2.5	2.7	2.3	2.8	2.0	2.3	2.8	2.0	2.3	2.5	4.0	2.9		
29-30 Years								3.2	3.2	2.2	2.6	2.9	2.7	2.4	2.5	2.2	2.3	2.4	2.5	2.2	2.6	2.8	2.1	2.6	2.9	NA	NA	NA	NA	
31-32 Years*									2.2	2.5	2.1	2.6	2.7	3.1	2.8	2.8	2.8	2.8	2.7	3.1	2.8	2.8	2.8	2.1	2.6	2.9	NA	NA	NA	
35 Years																			2.3	2.6	2.3	3.5	2.7	1.9	2.7	2.3	3.0	2.1	2.5	
40 Years																								3.2	2.1	2.6	1.8	3.0	2.4	1.8
45 Years																													2.6	2.0

FIGURE 5-5
Hallucinogens:* Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group




*Unadjusted for the possible underreporting of PCP.

**Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	9.4	8.8	9.6	9.9	9.3	9.0	8.1	7.3	6.5	6.3	6.0	6.4	5.5	5.6	5.9	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1	9.1	6.6	5.9	6.2	
19-20 Years		9.5	10.9	9.7	8.6	9.9	7.2	6.0	5.1	6.3	5.9	5.8	5.8	6.3	6.2	6.7	6.9	6.7	9.6	10.1	9.6	8.1	9.4	8.0	9.0	7.3	7.7	6.3		
21-22 Years				10.1	10.9	9.3	7.4	7.5	5.7	5.7	5.2	5.8	4.3	5.0	5.7	7.2	5.0	6.8	6.6	6.2	8.0	6.7	6.8	7.4	8.1	5.8	7.1	6.7		
23-24 Years					8.1	7.4	5.4	4.9	4.6	3.7	3.8	3.8	4.4	4.4	4.2	4.7	4.3	4.9	5.4	5.0	5.2	5.9	4.9	4.6	5.2	5.8	4.4			
25-26 Years						4.7	4.7	3.0	2.4	2.5	2.0	2.3	3.2	3.7	3.0	3.0	3.7	3.2	3.7	3.2	3.7	3.2	2.7	3.9	3.1	2.8	3.2			
27-28 Years							2.4	2.7	1.3	1.7	1.8	2.4	2.2	2.1	2.4	2.3	2.2	1.8	2.0	1.7	2.6	1.8	2.2	2.5	2.6					
29-30 Years												2.1	1.4	1.2	1.5	1.9	1.3	1.5	1.9	1.4	1.6	1.4	1.4	1.7	1.7	2.0	1.5	1.4		
31-32 Years**															1.0	1.3	1.2	0.9	1.0	1.3	0.9	1.6	0.9	1.3	0.9	1.5	NA	NA	NA	
35 Years																			0.8	0.7	0.5	1.0	0.6	0.8	0.5	0.8	0.3	0.6	1.0	
40 Years																							0.8	0.5	0.9	0.2	0.7	0.5	0.5	
45 Years																													0.6	0.3


FIGURE 5-9
Cocaine: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	6.0	7.2	9.0	12.0	12.3	12.4	11.5	11.4	11.6	13.1	12.7	10.3	7.9	6.5	5.3	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3
19-20 Years			11.8	15.0	16.3	15.9	16.9	13.8	14.6	15.4	15.9	13.4	10.6	7.6	5.6	3.8	3.7	3.2	3.2	3.9	3.7	4.5	5.3	5.7	5.8	6.0	6.5	6.3	6.3
21-22 Years				19.8	20.5	21.6	21.2	20.6	19.2	20.4	16.0	14.1	11.8	8.7	6.1	5.1	4.1	3.9	4.3	4.2	5.8	6.0	5.6	6.3	7.5	7.0	7.4	8.6	
23-24 Years						22.9	20.8	20.2	23.5	22.8	16.2	15.1	12.0	9.5	7.2	6.5	4.6	4.8	4.5	4.8	4.9	5.2	6.8	6.3	5.4	6.0	8.3	8.4	
25-26 Years								21.1	21.6	19.7	17.4	15.2	10.7	9.9	7.4	6.6	6.3	4.2	4.6	3.8	4.3	3.7	5.0	4.8	5.4	5.6	5.4	6.7	
27-28 Years										19.9	15.6	14.2	12.2	9.9	6.9	7.2	5.8	5.4	4.6	4.3	3.7	3.9	3.9	3.6	4.8	4.0	5.5	5.2	
29-30 Years												14.0	11.6	8.1	6.7	6.7	4.7	6.0	4.5	4.3	4.3	3.7	3.6	2.7	2.8	4.4	4.9	3.6	
31-32 Years*														8.9	6.8	5.7	5.1	5.5	3.8	5.0	5.1	4.4	4.1	3.0	3.5	NA	NA	NA	
35 Years																	4.7	4.3	5.1	4.1	4.4	4.6	3.9	3.5	3.6	2.7	3.3		
40 Years																						4.5	4.1	3.5	3.0	3.7	3.1	3.3	
45 Years																												3.4	3.9


FIGURE 5-10
Crack Cocaine: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years											4.1	3.9	3.1	3.1	1.9	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	
19-20 Years											2.7	2.7	1.8	1.0	0.6	1.3	1.2	1.2	1.0	1.3	1.3	1.3	1.3	2.4	1.6	1.7	1.7	1.8	1.6	
21-22 Years											4.1	2.9	3.6	1.6	1.3	1.3	1.1	1.1	1.3	1.4	1.2	1.4	1.2	1.6	1.4	1.0	1.2	1.8		
23-24 Years											3.4	4.0	3.1	2.1	1.4	1.3	1.2	0.8	1.0	1.2	1.0	1.2	1.5	1.2	1.1	1.1	1.1	1.2		
25-26 Years											2.4	2.7	1.9	2.3	1.5	1.3	1.3	1.0	1.1	0.7	0.7	0.4	0.9	1.1	1.0	0.7	0.5	1.1		
27-28 Years											3.0	3.0	2.0	1.3	1.6	1.6	1.7	1.5	0.9	0.8	0.6	0.8	0.8	0.6	1.3	0.6	0.6	0.8		
29-30 Years													3.2	2.8	1.7	1.1	0.9	1.0	1.5	1.5	1.0	1.5	0.7	1.3	0.5	0.7	0.9	0.8	0.6	
31-32 Years*															1.5	1.3	1.1	0.9	1.0	1.3	0.7	0.9	0.8	0.7	1.0	NA	NA	NA	NA	
35 Years																			1.0	1.0	1.5	0.8	1.2	1.1	0.8	0.7	1.3	0.7	0.5	
40 Years																								1.5	1.0	0.5	0.6	1.0	0.8	0.8
45 Years																													1.1	1.2


FIGURE 5-11
Heroin: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	0.8	0.8	0.8	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.6	0.5	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	
19-20 Years			0.4	0.3	0.2	0.5	0.2	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.0	0.1	0.1	0.1	0.1	0.5	0.7	0.4	1.1	0.6	0.7	1.0	0.4	0.4	0.4	
21-22 Years					0.6	0.4	0.4	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.1	0.2	0.3	0.2	0.1	0.4	0.5	0.7	0.4	0.4	0.5	0.5	0.1	0.4	0.5	
23-24 Years						0.2	0.6	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.1	0.6	0.3	0.1	0.2	0.5	0.4	0.5	0.3	0.5	0.3	
25-26 Years								0.2	0.3	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.1	0.2	
27-28 Years										0.3	0.3	0.3	0.2	0.0	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.1	0.3	0.1	0.6	0.1	
29-30 Years													0.2	0.3	0.0	0.2	0.1	0.0	0.3	0.4	0.2	0.3	0.1	0.1	0.1	0.2	0.2	0.2	0.3	
31-32 Years*															0.1	0.1	0.0	0.0	0.1	0.1	0.3	0.3	0.1	0.2	0.2	0.0	NA	NA	NA	
35 Years																				0.3	0.2	0.2	0.1	0.1	0.4	0.1	0.1	0.2	0.3	0.1
40 Years																								0.4	0.0	0.1	0.1	0.3	0.2	0.0
45 Years																													0.2	0.2

FIGURE 5-12
Narcotics Other Than Heroin: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group




*In 2002 the questions text was changed on half of the questionnaire forms for 18-30-year-olds. The list of examples of narcotics other than heroin was updated. Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2001 data presented here are based on all forms. The 2002 data presented here are based on the changed forms only. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003. Beginning in 2002 data were based on the changed question text for 35- and 40-year olds.

**Beginning in 2002 respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02*	'03*	'04			
18 Years	5.7	6.4	6.0	6.2	6.3	5.9	5.3	5.1	5.2	5.9	5.2	5.3	4.6	4.4	4.5	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7	9.4	9.3	9.5			
19-20 Years			4.7	4.7	5.6	4.9	4.4	4.2	3.9	3.4	4.2	3.7	3.1	3.0	3.9	3.2	2.2	2.5	2.7	4.7	4.7	4.3	4.3	5.5	6.2	7.0	8.3	9.9	10.4			
21-22 Years					4.9	5.0	3.5	4.0	3.3	3.8	3.8	3.6	3.6	3.4	2.7	2.4	3.4	2.9	2.9	3.1	2.6	5.0	4.3	4.2	5.0	6.8	8.9	9.6	9.2			
23-24 Years							4.4	3.3	3.5	3.8	2.7	2.4	2.3	2.4	2.7	2.4	2.2	2.0	2.6	2.5	2.9	2.7	3.5	4.1	4.2	4.3	8.2	9.7	9.5			
25-26 Years									2.7	3.4	2.0	2.5	2.5	2.4	2.3	2.4	2.6	2.0	1.8	1.8	2.1	2.2	3.0	3.0	2.9	3.7	6.0	6.4	7.9			
27-28 Years												2.7	3.0	1.6	2.9	1.5	1.8	1.7	1.4	2.1	2.4	2.0	2.0	1.9	1.8	2.1	3.1	4.3	6.7	7.5		
29-30 Years															2.2	2.1	1.5	1.8	1.9	1.3	1.7	1.9	2.1	2.6	1.5	1.8	2.5	2.9	4.2	5.1	5.4	
31-32 Years**																1.4	1.7	1.5	1.5	1.9	1.8	1.5	2.2	2.0	2.2	1.4	2.9	NA	NA	NA		
35 Years																					1.7	1.6	1.8	2.0	1.2	2.1	2.1	2.4	4.4	3.4	4.8	
40 Years																									1.7	1.4	1.7	1.6	3.4	2.3	2.9	
45 Years																															2.8	3.4


FIGURE 5-13
Amphetamines: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	15.8	16.3	17.1	18.3	20.8	26.0	20.3	17.9	17.7	15.8	13.4	12.2	10.9	10.8	9.1	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	
19-20 Years			18.2	21.5	23.8	25.5	23.9	19.7	15.8	14.5	11.0	9.1	9.2	6.9	6.6	4.9	5.6	5.4	5.4	7.2	6.5	5.9	7.5	7.9	9.3	8.7	9.1	8.6	8.5	
21-22 Years					25.5	26.7	22.4	19.9	17.4	13.0	13.0	9.9	8.1	6.8	5.5	4.9	4.3	4.8	5.3	5.7	4.9	7.3	5.0	5.0	6.0	7.9	7.1	7.5	6.7	
23-24 Years						21.8	18.3	14.0	14.1	11.4	7.9	7.6	5.1	5.3	3.8	4.0	3.8	4.5	3.0	4.1	3.8	4.3	4.5	4.8	5.2	5.8	5.8	7.1		
25-26 Years								14.9	12.5	8.6	8.3	6.4	5.5	4.0	3.4	2.7	2.9	3.9	3.5	2.5	3.2	2.9	3.4	3.6	3.6	3.9	3.1	4.6		
27-28 Years									9.1	7.9	5.0	4.3	4.3	4.0	3.5	2.6	2.9	2.7	2.5	2.0	2.3	2.6	2.7	3.2	3.3	3.6	3.9			
29-30 Years										5.5	5.0	2.7	2.9	3.3	2.4	2.6	2.5	2.6	2.7	1.8	2.4	1.4	1.9	2.1	2.6	2.2				
31-32 Years*													3.7	3.7	2.6	2.4	2.5	1.5	2.7	2.6	2.6	1.9	1.9	1.5	NA	NA	NA	NA		
35 Years																		2.4	1.9	1.9	1.7	1.7	1.9	1.8	1.9	1.2	2.2	1.5		
40 Years																								1.7	1.2	1.4	1.0	1.4	1.0	1.1
45 Years																													1.4	0.9


FIGURE 5-15
Sedatives (Barbiturates): Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	9.6	9.3	8.1	7.5	6.8	6.6	5.5	5.2	4.9	4.6	4.2	3.6	3.2	3.3	3.4	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	
19-20 Years			6.4	6.9	4.5	4.7	4.4	3.5	3.5	2.0	2.2	1.9	2.2	1.6	1.7	1.8	1.7	1.9	2.3	3.4	3.3	4.0	3.8	5.0	4.9	5.2	5.8	5.2	6.0	
21-22 Years					5.7	5.8	4.1	3.1	2.5	2.3	2.9	2.7	1.9	1.8	1.7	1.4	1.8	1.6	2.2	2.6	2.4	3.5	3.1	2.5	3.9	4.8	3.8	4.8	4.4	
23-24 Years							4.1	3.7	2.6	3.0	2.3	1.5	2.1	1.8	2.3	2.0	1.7	1.7	1.4	2.2	1.5	2.4	3.2	4.3	3.4	4.4	3.9	5.0		
25-26 Years									3.3	3.4	1.8	2.1	1.7	1.3	2.2	2.5	1.5	1.8	1.1	1.2	1.0	1.5	1.7	1.8	2.2	2.7	2.9	2.5	3.3	
27-28 Years												2.4	2.3	1.2	1.7	1.8	1.4	1.4	2.3	1.6	1.4	1.7	1.0	1.5	1.1	1.4	2.1	2.3	3.1	2.8
29-30 Years														2.1	1.4	1.6	1.6	2.0	1.1	1.4	1.7	1.6	1.8	1.0	1.2	0.9	1.3	1.7	1.5	1.9
31-32 Years*															2.2	2.2	1.7	1.5	1.2	1.1	1.4	0.8	1.0	0.9	0.9	1.1	NA	NA	NA	NA
35 Years																				1.9	1.7	1.6	1.3	1.1	1.6	0.9	1.8	1.4	1.3	1.2
40 Years																								0.9	1.5	1.6	1.4	1.0	0.7	1.0
45 Years																													1.0	1.0


FIGURE 5-16
Tranquilizers: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	10.3	10.8	9.9	9.6	8.7	8.0	7.0	6.9	6.1	6.1	5.8	5.5	4.8	3.8	3.5	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7	6.9	7.7	6.7	7.3
19-20 Years			9.4	9.8	8.8	7.4	5.6	5.1	5.4	4.4	4.2	4.0	3.5	3.4	3.0	2.7	2.2	2.1	1.9	3.7	3.5	4.7	4.2	4.1	5.5	6.1	8.8	8.0	8.0
21-22 Years					9.0	7.3	7.2	5.8	5.4	4.5	5.4	5.5	4.5	3.5	3.6	3.2	3.8	3.1	2.9	3.5	3.7	3.6	4.5	4.2	5.3	7.1	7.8	7.0	8.1
23-24 Years						8.6	6.6	5.6	6.2	5.2	4.1	4.2	3.8	3.8	4.0	3.4	3.2	3.1	3.0	3.0	2.9	3.7	4.2	5.3	5.4	6.4	7.2	8.3	
25-26 Years								6.7	7.1	5.4	5.8	4.3	2.9	5.0	3.9	4.5	3.7	3.3	3.1	2.4	1.9	3.6	3.5	3.7	5.3	7.0	6.3	6.8	
27-28 Years										6.8	6.2	4.8	4.6	3.3	3.8	3.4	3.8	3.6	3.4	2.9	2.0	2.9	2.6	3.0	3.9	4.9	5.2	5.6	
29-30 Years												4.6	4.1	3.9	4.2	3.7	2.7	3.2	3.5	3.1	4.1	2.4	2.1	2.7	4.2	5.0	4.3	4.8	
31-32 Years*														3.8	4.1	4.1	2.7	3.8	1.8	3.2	4.1	3.8	2.7	2.6	3.0	NA	NA	NA	
35 Years																		3.1	3.6	3.6	3.0	2.2	3.4	3.2	4.3	4.0	3.8	3.6	
40 Years																							3.0	2.0	3.0	3.7	4.2	2.2	3.5
45 Years																												2.9	3.1


FIGURE 5-18a
Alcohol: Trends in Annual Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	85.7	87.0	87.7	88.1	87.9	87.0	86.8	87.3	86.0	85.6	84.5	85.7	85.3	82.7	80.6	77.7	76.8	76.0	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	
19-20 Years			89.8	90.6	89.0	90.6	88.6	88.5	88.7	88.5	88.2	88.2	86.6	87.5	85.6	84.6	81.9	80.6	78.2	78.3	79.6	79.2	79.7	79.6	79.7	77.6	78.0	75.0	75.2	
21-22 Years				90.2	91.6	91.8	91.8	89.1	89.8	90.1	90.8	89.5	89.1	89.6	89.0	87.9	85.9	84.4	85.7	84.4	85.1	86.3	85.5	86.2	87.0	85.8	84.3	86.8		
23-24 Years					90.0	91.7	90.4	91.6	88.1	89.7	89.7	88.7	88.2	88.1	89.1	87.8	86.6	87.8	85.7	85.4	84.9	85.2	87.2	86.7	88.0	87.6	87.6	87.2		
25-26 Years								88.2	89.9	88.8	90.5	89.4	87.5	87.5	87.7	86.7	87.8	86.0	86.7	85.9	86.4	83.8	85.0	84.2	86.3	88.3	86.4	87.9		
27-28 Years									87.8	87.8	87.7	88.0	86.4	85.3	85.6	85.7	84.5	85.7	85.3	85.9	85.3	85.4	82.9	84.2	84.7	83.6	86.1			
29-30 Years												87.2	86.0	86.9	85.0	84.5	83.2	82.6	83.3	84.7	83.7	84.2	85.4	83.7	84.3	83.6	83.9	83.5		
31-32 Years*														84.8	83.8	85.0	83.6	83.6	81.8	82.0	83.3	83.2	85.1	82.9	84.4	NA	NA	NA		
35 Years																		82.5	82.1	83.5	82.3	82.3	81.0	81.0	82.7	85.1	82.6	86.7		
40 Years																							77.3	80.0	80.3	81.5	80.0	81.6	79.8	
45 Years																													78.9	79.2


FIGURE 5-18b
Alcohol: Trends in Thirty-Day Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04		
18 Years	68.3	71.2	72.1	71.8	72.0	70.7	69.7	69.4	67.2	65.9	65.3	66.4	63.9	60.0	57.1	54.0	51.3	51.0	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0		
19-20 Years		75.8	76.5	76.6	77.0	75.7	73.9	73.6	73.3	72.9	72.5	69.6	69.8	66.6	64.5	61.0	60.5	59.9	59.2	58.1	59.0	59.7	62.0	59.1	59.0	59.2	56.7	56.7			
21-22 Years			78.3	80.5	79.9	79.3	78.1	75.9	77.2	77.2	76.2	73.8	74.1	75.3	72.7	71.6	70.4	70.4	69.5	69.1	69.4	69.2	70.5	71.8	71.9	69.5	72.4				
23-24 Years				77.9	78.9	77.6	79.7	75.7	74.9	75.9	72.2	73.6	72.4	73.0	73.1	70.1	72.3	69.2	69.3	70.3	70.2	71.5	70.6	71.9	72.7	72.8					
25-26 Years					75.2	76.8	76.3	77.7	74.1	72.5	71.4	71.6	69.8	69.9	70.4	71.8	68.5	70.9	66.3	70.0	68.7	68.7	71.2	69.1	72.4						
27-28 Years						73.6	75.0	74.6	73.9	70.9	69.8	69.1	68.3	69.9	68.0	69.3	70.4	68.7	70.2	64.6	66.5	67.9	67.2	68.8							
29-30 Years							72.1	72.3	70.2	69.6	69.2	66.2	67.0	67.0	68.0	65.8	66.1	67.4	65.2	66.2	65.4	66.5	64.5								
31-32 Years*									68.4	68.5	67.8	66.4	67.7	67.6	65.5	65.3	65.2	66.0	66.7	67.8	NA	NA	NA								
35 Years																	65.1	66.8	64.7	65.3	62.9	64.2	64.0	63.7	67.3	63.7	70.3				
40 Years																					59.8	64.2	63.1	65.6	65.4	66.2	63.7				
45 Years																													62.2	65.7	

FIGURE 5-18c
Alcohol: Trends in Thirty-Day Prevalence of Daily Use Among
High School Seniors and Adults Through Age 45
by Age Group




*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	5.6	6.1	5.7	6.9	6.0	6.0	5.7	5.5	4.8	5.0	4.8	4.8	4.2	4.2	3.7	3.6	3.4	2.5	2.9	3.5	3.7	3.9	3.9	3.4	2.9	3.6	3.5	3.2	2.8	
19-20 Years			7.6	7.7	7.0	7.2	7.5	5.3	5.8	5.6	5.3	5.7	4.8	4.7	4.0	3.7	3.3	3.2	3.3	3.1	2.7	4.8	3.6	4.1	3.9	3.6	3.9	3.6	3.7	
21-22 Years				8.4	7.7	7.8	8.0	7.7	6.4	6.3	7.0	7.2	5.0	4.9	4.9	4.4	5.1	3.9	3.5	5.1	4.6	5.7	5.9	5.3	6.2	5.6	5.7	5.7		
23-24 Years						8.2	8.5	6.8	7.3	6.2	6.1	6.2	5.1	5.3	5.4	4.2	4.9	3.7	4.1	4.8	4.5	3.9	4.7	4.2	4.6	5.0	6.5	5.5		
25-26 Years								7.5	7.5	5.3	6.9	6.3	6.0	4.8	4.9	6.1	5.1	3.3	4.4	3.7	5.1	3.4	5.1	3.8	5.0	5.4	4.6	4.3		
27-28 Years										7.3	7.2	5.7	6.9	4.9	6.2	4.4	4.7	5.4	4.7	3.6	4.2	3.1	4.3	3.5	2.7	3.7	5.1	3.5		
29-30 Years													7.6	5.6	5.6	5.9	5.8	4.6	5.0	4.1	5.1	5.9	3.4	5.2	3.9	4.3	3.8	3.5	3.8	
31-32 Years*															6.4	5.2	6.0	5.7	4.7	4.9	5.1	5.0	4.8	4.4	4.2	4.7	NA	NA	NA	
35 Years																				7.2	5.5	7.5	4.8	6.0	5.2	5.2	5.8	4.8	3.9	6.3
40 Years																								6.9	7.5	6.5	7.5	6.6	7.8	6.6
45 Years																													7.8	9.0

FIGURE 5-18d


Alcohol: Trends in Two-Week Prevalence of Having Five or More Drinks in a Row at Least Once Among High School Seniors and Adults Through Age 45 by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	37.1	39.4	40.3	41.2	41.2	41.4	40.5	40.8	38.7	36.7	36.8	37.5	34.7	33.0	32.2	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	
19-20 Years			41.1	42.1	42.7	43.1	41.7	40.9	41.0	41.2	41.2	37.2	37.3	36.9	36.0	37.0	34.0	34.6	34.5	31.7	32.7	36.5	34.5	35.3	35.3	36.3	36.0	33.6	35.5	
21-22 Years				40.7	43.6	41.6	42.3	40.4	40.4	40.8	41.0	42.0	39.3	38.1	40.3	39.9	40.3	40.5	38.5	38.2	40.2	39.7	40.2	40.6	42.4	40.7	39.9	41.7		
23-24 Years						37.1	39.3	35.1	37.3	35.8	36.6	37.0	35.4	35.5	34.4	34.9	35.0	32.9	35.6	36.3	33.4	35.3	38.1	37.0	38.2	39.4	39.3	40.4		
25-26 Years								33.7	33.3	31.5	33.3	30.7	31.7	32.0	31.5	31.8	32.1	30.9	28.7	30.0	31.5	31.3	33.0	31.5	33.7	34.9	35.1	36.4		
27-28 Years										30.1	32.2	28.0	29.8	28.9	28.8	29.2	29.0	28.5	26.9	29.7	29.3	28.9	32.0	29.1	29.2	28.9	31.1	31.3		
29-30 Years													26.7	26.3	25.2	24.3	25.7	25.1	27.5	26.3	24.9	26.5	26.6	26.9	24.0	27.3	25.8	26.4	26.9	
31-32 Years*															25.4	25.1	23.7	24.8	24.6	24.7	24.3	24.7	22.8	24.1	24.1	24.3	NA	NA	NA	
35 Years																				21.1	20.0	21.9	22.3	20.4	21.4	22.2	20.6	22.9	22.4	21.6
40 Years																								19.7	20.5	18.3	21.3	20.8	20.7	20.2
45 Years																													20.1	19.2


FIGURE 5-19a
Cigarettes: Trends in Thirty-Day Prevalence Among
High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	38.8	38.4	36.7	34.4	30.5	29.4	30.0	30.3	29.3	30.1	29.6	29.4	28.7	28.6	29.4	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0
19-20 Years			39.3	39.3	36.0	34.9	32.1	32.5	31.5	30.9	30.0	30.1	28.4	27.7	27.2	27.6	29.5	29.0	31.3	33.4	34.0	34.0	33.9	36.1	32.2	32.8	29.8	27.0	27.9
21-22 Years				37.9	37.5	36.2	33.5	32.2	32.4	32.0	32.4	29.8	29.4	28.6	28.3	29.0	29.2	28.8	31.8	32.3	32.3	33.7	33.4	33.6	34.0	32.6	30.5	31.3	
23-24 Years					36.7	36.5	33.6	31.9	29.9	31.7	29.9	29.4	27.8	28.5	28.4	28.1	27.0	28.0	30.1	29.1	30.9	32.4	29.5	31.1	31.9	31.0	31.0	31.5	
25-26 Years							33.7	35.3	31.3	28.2	27.3	29.5	28.4	28.3	26.3	27.7	26.4	25.7	26.8	27.6	29.9	25.6	28.2	28.6	27.3	27.0	29.6		
27-28 Years								32.5	32.3	29.1	27.2	26.5	28.2	27.8	25.4	25.0	26.8	26.0	24.9	25.6	22.9	26.5	24.2	24.7	26.3	25.9			
29-30 Years									28.9	30.2	27.8	24.4	23.8	25.8	25.5	25.2	23.4	24.6	23.1	22.7	21.2	20.4	24.4	22.0	21.9				
31-32 Years*										28.3	28.1	27.5	25.3	24.9	25.0	23.8	24.3	22.5	24.0	22.7	21.2	NA	NA	NA					
35 Years																24.8	26.1	25.4	22.3	23.6	22.6	24.0	20.5	21.9	20.1				
40 Years																					24.3	23.5	23.5	22.9	18.9	21.9	20.0		
45 Years																												20.7	20.2


FIGURE 5-19b
Cigarettes: Trends in Thirty-Day Prevalence of Daily Use
Among High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	
18 Years	28.8	28.8	27.5	25.4	21.3	20.3	21.1	21.2	18.7	19.5	18.7	18.7	18.1	18.9	19.1	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	
19-20 Years			31.0	31.2	29.3	26.0	23.9	24.4	24.1	23.2	21.9	22.5	19.5	18.9	19.2	19.4	20.5	21.1	21.9	22.2	22.5	22.7	23.8	25.6	22.7	21.9	20.6	18.8	18.2	
21-22 Years					31.1	31.4	28.6	26.0	25.3	25.3	24.4	24.2	22.3	22.5	20.2	20.6	21.2	20.5	21.1	24.0	22.8	21.4	22.8	24.2	25.1	23.6	23.9	20.8	21.5	
23-24 Years							30.1	30.6	27.8	25.1	25.2	26.0	24.0	23.3	22.2	22.5	20.9	20.1	19.9	20.0	22.8	21.5	21.2	21.4	21.2	22.4	23.5	21.5	23.3	
25-26 Years									28.7	30.4	27.3	23.7	22.9	25.0	23.3	22.8	20.3	21.9	19.8	19.2	21.1	19.2	21.9	19.6	20.1	20.9	19.8	20.4	22.7	
27-28 Years											27.6	27.9	25.0	22.9	22.2	23.9	21.8	20.1	20.5	20.9	19.4	17.6	19.5	16.0	19.7	17.2	18.1	19.8	18.2	
29-30 Years													25.4	26.4	24.2	21.0	20.3	21.7	20.9	20.1	18.6	19.7	17.2	17.2	15.8	14.4	17.4	16.4	16.7	
31-32 Years*															23.9	24.9	22.8	21.4	20.9	21.2	19.8	19.1	17.9	18.9	18.1	16.1	NA	NA	NA	
35 Years																			22.5	23.0	22.1	18.3	20.4	19.7	20.1	16.5	18.2	16.3	14.8	
40 Years																								21.7	20.9	20.8	20.2	16.7	19.0	16.6
45 Years																													19.0	17.8

FIGURE 5-19c
Cigarettes: Trends in Thirty-Day Prevalence of Smoking a Half-Pack or More
Daily Among High School Seniors and Adults Through Age 45
by Age Group



*Beginning in 2002, respondents were followed through age 30 (instead of 32, as in past years), then again at ages 35 and 40. The 45-year-olds were added in 2003.

Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04
18 Years	19.2	19.4	18.8	16.5	14.3	13.5	14.2	13.8	12.3	12.5	11.4	11.4	10.6	11.2	11.3	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0
19-20 Years		23.8	24.6	21.9	19.3	18.0	17.2	17.2	16.6	16.2	15.6	13.8	13.0	14.3	12.7	14.5	14.5	15.0	15.2	14.7	15.4	16.9	16.3	14.6	13.9	12.8	11.7	11.6	
21-22 Years			25.2	25.3	23.0	19.7	21.2	20.4	19.3	19.3	17.3	16.4	15.0	14.1	15.1	14.5	15.6	18.1	15.7	14.7	16.2	16.4	17.2	15.9	14.4	13.8	12.7		
23-24 Years				24.6	25.1	22.8	20.8	21.1	21.6	18.4	18.6	17.4	17.4	15.5	15.2	15.0	15.3	16.1	16.4	14.5	14.8	14.1	15.8	15.9	15.4	15.2			
25-26 Years					24.1	24.8	22.0	19.9	18.6	20.6	19.6	18.2	15.8	17.4	15.0	14.2	15.0	13.2	15.5	15.0	14.8	15.1	14.1	14.0	12.7	12.5			
27-28 Years						23.2	23.3	20.6	19.0	18.2	19.0	17.9	16.3	15.9	16.3	14.8	12.8	14.8	12.4	14.7	12.6	13.9	14.8	12.8					
29-30 Years							22.3	22.0	20.5	16.7	17.0	17.9	16.8	16.5	15.2	15.9	12.2	13.2	12.5	11.4	14.0	12.7	12.5						
31-32 Years*								20.8	20.8	19.3	17.8	17.8	17.6	16.1	16.1	14.3	14.8	12.8	11.9	NA	NA	NA							
35 Years									19.1	19.1	18.5	15.4	16.3	17.3	15.7	13.4	13.0	12.5	10.9										
40 Years																				18.7	17.2	17.2	15.9	13.6	14.9	14.2			
45 Years																												16.8	15.4

Chapter 6

ATTITUDES AND BELIEFS ABOUT DRUGS AMONG YOUNG ADULTS

Earlier volumes in this monograph series, as well as other publications from the study, have demonstrated that shifts in certain attitudes and beliefs are important in explaining changes in actual drug-using behavior.⁴⁶ In this chapter, we review trends since 1980 in the same attitudes and beliefs among the young adult samples that were considered in Volume I for secondary school students.

PERCEIVED HARMFULNESS OF DRUGS

Table 6-1 provides trends in the percentages who perceive a great risk of harm associated with differing usage levels of various licit and illicit drugs. These questions are contained in one questionnaire form only, limiting the numbers of follow-up cases; accordingly, we use four-year age bands in order to increase the available sample size (to about 400–600 weighted cases per year for each age band) and, thus, to improve the reliability of the estimates. (The numbers of weighted cases are given at the end of Table 6-1. The actual numbers of respondents are somewhat larger.) Still, these are relatively small sample sizes compared to those available for 8th, 10th, and 12th graders, and thus the change estimates may be relatively more labile. Because of the nature of the Monitoring the Future design, trend data are available for a longer period for 19- to 22-year-olds (since 1980) than for 23- to 26-year-olds (since 1984) or for 27- to 30-year-olds (since 1988). Also displayed in this table are comparison data for 12th graders, shown here as 18-year-olds, from 1980 onward. (See also Table 8-3 in chapter 8 of Volume I for the longer-term trends in 12th graders' levels of perceived risk.) Questions about these attitudes and beliefs are not included in the questionnaires for 35-, 40-, and 45-year-old respondents.

- Table 6-1 illustrates considerable differences in the degree of risk young adults associate with various drugs. In general, the results closely parallel the distinctions made by 12th graders.
- **Marijuana** is seen as the least risky of the illicit drugs, although sharp distinctions are made between different levels of marijuana use. In 2004, experimental use of marijuana is perceived as being of “great risk” by only about one-sixth (14%-15%) of all high school

⁴⁶Bachman, J. G., Johnston, L. D., O'Malley, P. M., & Humphrey, R. H. (1988). Explaining the recent decline in marijuana use: Differentiating the effects of perceived risks, disapproval, and general lifestyle factors. *Journal of Health and Social Behavior*, 29, 92-112; Bachman, J. G., Johnston, L. D., & O'Malley, P. M. (1990). Explaining the recent decline in cocaine use among young adults: Further evidence that perceived risks and disapproval lead to reduced drug use. *Journal of Health and Social Behavior*, 31, 173-184; Bachman, J. G., Johnston, L. D., & O'Malley, P. M. (1998). Explaining recent increases in students' marijuana use: Impacts of perceived risks and disapproval, 1976 through 1996. *American Journal of Public Health*, 88, 887-892; Johnston, L. D. (1981). Characteristics of the daily marijuana user. In R. de Silva, R. L. DuPont, & G. K. Russell (Eds.), *Treating the marijuana-dependent person* (pp. 12-15). New York: The American Council on Marijuana; Johnston, L. D. (1985). The etiology and prevention of substance use: What can we learn from recent historical changes? In C. L. Jones & R. J. Battjes (Eds.), *Etiology of drug abuse: Implications for prevention* (NIDA Research Monograph No. 56, pp. 155-177). (DHHS Publication No. (ADM) 85-1335). Rockville, MD: National Institute on Drug Abuse.

graduates in the age band 19 to 30, whereas regular use is perceived to carry great risk by over half (54%-59%) of them.

It is interesting to note that in the mid-1980s and early 1990s, fewer of the older age groups attached great risk to marijuana use than did the younger age bands. Indeed, there was a quite regular negative ordinal relationship between age and perceived risk for some years after 1980, when the first comparisons were available. Although this may have first looked like an age effect, our study design allowed us to recognize it as a cohort effect: the younger cohorts initially perceived marijuana as more dangerous than the older cohorts and persisted in such beliefs as they grew older. Newer cohorts, however, have shown lower levels of perceived risk. Twelfth graders from the class of 2004 are much less likely to perceive marijuana use as dangerous as did 12th-grade cohorts in the late 1980s and early 1990s. This reflects what we have interpreted as “generational forgetting,” a phenomenon wherein younger replacement cohorts no longer carry the knowledge—and perhaps the direct or vicarious experience on which the knowledge is based—that the older cohorts had at that age.

The decline in perceived risk that began in the 1990s was greater in the younger age bands, including grades 8 and 10, and was least among the 27- to 30-year-olds. We believe that much of the decline in perceived risk that occurred in the older age bands resulted directly from generational replacement of earlier cohorts by later, less concerned ones. The credibility of this view is strengthened by the 1995–1997 reversal of the relationship between age and perceived risk of regular use. This reversal is consistent with an underlying cohort effect and could not simply reflect an association between age and a regular change in these attitudes (i.e., an “age effect”). The decline in perceived risk for regular marijuana ended in a somewhat staggered fashion—among 12th graders in 1999, among 19- to 22-year-olds in 2001, among 23- to 26-year-olds in 2002, and among 27- to 30-year-olds in 2004.

- Young adults view use of any of the other illicit drugs as distinctly more risky than use of marijuana. Even the experimental use of *amphetamines* and *sedatives (barbiturates)* is perceived as risky by about 22%-39% of young adults aged 19 to 30, and 39%-52% think trying *LSD* or *MDMA (ecstasy)* involves great risk. Trying *cocaine powder* is seen as dangerous by 46%-53%, while trying *crack* or *heroin* once or twice is seen as dangerous by 56%-68%.
- In recent years, the older age groups have been more likely than the younger age groups to see *LSD* as dangerous. Indeed, there is now a substantial age-related difference. The age distinctions for LSD use have become sharper in recent years as perceived risk has declined more in the younger age groups than in the older ones—again indicating some important cohort changes in these attitudes, quite likely as a result of the process we have labeled “generational forgetting” (see above).
- Young adults report somewhat higher perceived risk with respect to *cocaine* use than the 12th graders, who have had less experience with cocaine. Among 12th graders and the young adult age groups, the danger associated with using cocaine on a regular basis grew

considerably (by 13 and 17 percentage points, respectively) between 1980 and 1986. However, these changed beliefs did not translate into changed behavior until the perceived risk associated with *experimental* and *occasional* use began to rise sharply after 1986. When these two measures rose, a sharp decline in actual use occurred.

We hypothesized that respondents saw only these lower levels of use as relevant to them and, therefore, saw themselves as vulnerable to the dangers only of such use. (No one starts out planning to be a heavy user; further, in the early 1980s, cocaine was not believed to be addictive.) Based on this hypothesis, we included the additional question about occasional use in 1986, just in time to capture a sharp increase in perceived risk later that year. This increase occurred largely in response to the growing media frenzy about cocaine—and crack cocaine, in particular—and to the widely publicized, cocaine-related deaths of several public figures (most notably, basketball star Len Bias). After stabilizing for a few years, perceived risk began to fall off among 12th graders after about 1991 but not among the older age groups—once again suggesting that lasting cohort differences were emerging, quite possibly as a result of “generational forgetting” of the dangers of cocaine. A decline in perceived risk began among the 19- to 22-year-olds starting after 1994, among the 23- to 26-year-olds after 1997, and among the 27- to 30-year-olds after 2001, all likely reflecting generational replacement with the 12th graders who earlier had come to see cocaine as less dangerous.

- A similar situation also now exists for **crack**, for which perceived risk is highest in the oldest age band and lowest among 12th graders. Trend data (available since 1987) on the risks perceived to be associated with use of crack show increases in the 1987-1990 interval for all age groups, followed by relatively little change in the older age strata. Since 1990, the 12th graders have shown decreases in the perceived risk of experimental use of crack—perhaps reflecting the onset of “generational forgetting” of its dangers—leaving them as perceiving considerably less risk than the older groups. The young adult age groups have shown a staggered decline in this measure, with the 19- to 22-year-olds showing a decline after 1994, the 23- to 26-year-olds since 1996, and the 27- to 30-year-olds after 2001.
- Questions about perceived risk of **crystal methamphetamine (ice)** use were introduced in 1990, and the results show what may be an important reason for its lack of rapid spread. More than half of all 12th graders and young adults perceived it as a quite dangerous drug even to try, perhaps because it was likened to crack in many media accounts. (Both drugs are burned and the fumes inhaled, both are stimulants, and both can produce a strong dependence.) There was rather little age-related difference in perceived risk associated with use of ice in 1990 and 1991 (although the two youngest age groups were somewhat higher). But as perceived risk fell considerably among 12th graders (and eventually among 19- to 22-year-olds) and held steady or rose in the oldest two age groups, an age-related difference emerged. Now perceived risk has risen some among the 19- to 22-year-olds, narrowing the age-related differences that had emerged for a few years. Still, 12th graders are considerably less likely to see risk attached to trying ice (52%) than are 27- to 30-year-olds (63%), very likely due to generational forgetting.

- **MDMA (ecstasy)** questions were introduced in the follow-up surveys in 1989 but were not asked of 12th graders until 1997. At the beginning of the 1990s, all young adult age bands viewed it as a fairly dangerous drug, even for experimentation. But, again, the different age bands had diverging trends during the 1990s, with the oldest two age bands continuing to see ecstasy as quite dangerous but the 19- to 22-year-olds (and very likely the 12th graders, for whom we did not have data until 1997) coming to see it as less so. In 2000, 38% of the 12th graders saw great risk in trying ecstasy versus 49% of the 27- to 30-year-olds; in 2001, the corresponding figures were 46% and 54%. In fact, three of the four age bands showed appreciable increases in perceived risk for ecstasy in 2001, which led us to state three years ago in this chapter that “a turnaround in the use of this drug may be about to occur.” The increase in perceived risk continued in 2002 in the two youngest age strata, and their use of ecstasy did, indeed, begin to decline—and decline sharply (see chapter 5). At present there is rather limited age-related difference in perceived risk for this drug, and what difference there is derives from the 23- to 26-year-olds having the lowest level of perceived risk (which may originate in part from the fact that they have had the most experience with this drug). That age band has not shown the increase in perceived risk that the younger two age bands have shown in recent years.
- Young adults have been more cautious about **heroin** use than have 12th graders. In general, there has been relatively little change over the years in the proportions of all age groups seeing regular heroin use as dangerous; the great majority of each group (over 86%) consistently held this viewpoint. With regard to heroin experimentation, from 1975 to 1986 there had been a downward shift among the 12th graders in the proportion seeing great risk associated with trying heroin. Following this decline (although their data do not extend back as far) young adults showed an increased caution about heroin use in the latter half of the 1980s—possibly due to heroin injection’s association with the spread of HIV infection—followed by a leveling through most of the 1990s. In 1996 and 1997, young adults’ perceived risk increased some, as happened among the 12th graders (as well as among the 8th and 10th graders). These various trends may reflect, respectively, (a) the lesser attention paid to heroin by the media during the late 1970s and early 1980s; (b) the subsequent great increase in attention paid to intravenous heroin use in the latter half of the 1980s because of its important role in the spread of HIV/AIDS; (c) the emergence in the 1990s of heroin so pure that people no longer needed to use a needle to administer it, resulting in lower perceived risk; and (d) the subsequent increased attention given to heroin by the media (partly as a result of some overdose deaths by public figures and partly prompted by the emergence of “heroin chic” in the design industry), as well as through an anti-heroin campaign in the media launched by the Partnership for a Drug-Free America in June 1996. At present the young adults see heroin use as more dangerous than do the 12th graders (Table 6-1); the differences are largest for experimental and occasional use.
- A minority of young adults see **heavy drinking on weekends** as dangerous (38%-41%), which is true for 12th graders as well (44%). The belief that heavy drinking carries great risk has increased over the years in all of these age groups, rising among 12th graders from 36% in 1980 to 49% in 1992. Among 19- to 22-year-olds it rose from a low of 30% in 1981 to 42% in 1992; the increases among the older groups were smaller. The increase in this belief

may well help to explain the important decline in actual heavy drinking and may in turn be explained by the media campaigns against drunk driving and by the increase in the drinking age in a number of states.⁴⁷ After reaching peaks in the early 1990s, perceived risk for this behavior eased back some in all age strata.

- Between 1980 and 1991, a gradually increasing proportion of all four age groups viewed drinking *one or two drinks per day* as dangerous; but then they all showed a parallel decrease in perceived risk for this behavior through 2000, at least. It seems likely that the earlier increase was due to the general rising concern about the consequences of alcohol use, particularly drunk driving, and that the subsequent decline was due to increasing reports of cardiovascular health benefits of light-to-moderate daily alcohol consumption. In recent years there has been little systematic change in this belief in any of the age strata.
- About four-fifths (81%-84%) of young adults perceive regular *pack-a-day cigarette smoking* as entailing high risk, higher than the 74% of 12th graders who hold that belief and much higher than the 62% of 8th graders who do so. In recent years, the 18-year-olds have consistently shown lower perceived risk than young adults, while 10th graders have been still lower and 8th graders lowest. Clearly, there is an age effect in young people coming to understand the dangers of smoking. Unfortunately, it appears that much of the learning happens after a great deal of smoking initiation has occurred and many young people already have become addicted. These beliefs about smoking risks have strengthened very gradually in all age groups from senior year forward during the years we have monitored them. (See Table 6-1.) The parallel changes in these beliefs across the different age groups indicate a period effect, rather than a cohort effect, suggesting that all of these age groups were responding to common influences in the larger culture. These influences are discussed at some length in the chapter on attitudes and beliefs in Volume I.
- The regular use of *smokeless tobacco* is seen as dangerous by only 50%-58% of young adults and by even fewer 12th graders (45%). However, these beliefs have also gradually strengthened in all age groups over the intervals covered (Table 6-1). As with cigarettes, the change appears to reflect a secular trend or period effect because it has been occurring in parallel for all age groups.

PERSONAL DISAPPROVAL OF DRUG USE

The questions asked of 12th graders concerning the extent to which they personally disapprove of various drug-using behaviors among “people (who are 18 or older)” also are asked of follow-up respondents in one of the six questionnaire forms. Trends in the answers of young adults aged 19 to 22, 23 to 26, and 27 to 30 are contained in Table 6-2. Comparison data for 12th graders are also provided for 1980 onward. (See also Table 8-6 in chapter 8, Volume I, for the longer-term trends in 12th graders’ levels of disapproval associated with using the various drugs.)

⁴⁷See O’Malley, P. M., & Johnston, L. D. (1999). Drinking and driving among U.S. high school seniors: 1984-1997. *American Journal of Public Health, 89*, 678-684. See also O’Malley, P. M., & Johnston, L. D. (2003). Unsafe driving by high school seniors: National trends from 1976 to 2001 in tickets and accidents after use of alcohol, marijuana and other illegal drugs. *Journal of Studies on Alcohol, 64*, 305-312.

- In general, disapproval levels of adult use of the various drugs rank similarly across substances for both 12th graders and young adults. The great majority of young adults disapprove of using, or even experimenting with, all of the *illicit drugs other than marijuana*. For example, 95% or more of young adults in 2004 disapprove regular use of each of the following drugs: *LSD*, *cocaine*, *amphetamines*, *sedatives (barbiturates)*, and *heroin*. In fact, fully 80% to 96% of young adults disapprove experimentation with each of these drugs. Many of these attitudes differ rather little as a function of age, at present; when there is a difference, the younger age groups are usually the least disapproving.
- Even for *marijuana*, about half of young adults now disapprove of experimentation (from 48% to 55%). In 2004, about two-thirds (between 64% and 69%) disapproved of occasional use, and 82% to 89% disapproved of regular use.

Marijuana shows the widest fluctuations in disapproval over time—generally, fluctuations that parallel the changes in perceived risk (though sometimes with a one-year time lag, with the change in perceived risk coming first). The most fluctuation has occurred among 12th graders, nearly as much among 19- to 22-year-olds, and the least among 27- to 30-year-olds (Table 6-2). Among 12th graders, disapproval of regular marijuana use increased substantially in the 1980s, peaked in the early 1990s, declined through much of the 1990s, and then leveled around 1998. The 19- to 22-year-olds had a quite similar pattern, though the decline continued a year longer, likely due to generational replacement. Among 23- to 26-year-olds, there were some declines starting later in the 1990s, but the declines have been very modest.

- Beginning around 1990, all age groups decreased some in their disapproval of trying *LSD* (from high levels of disapproval, all at 90% or 91%). The decline was steepest among 12th graders, but there was a reversal of the decline among 12th graders in 1997, and disapproval of using LSD has increased some since then. Disapproval in the older age groups declined less, and in a staggered fashion; it has not yet shown consistent evidence of a reversal. This pattern again suggests some lasting cohort differences in these attitudes.
- Most of the disapproval statistics for *heroin* use, with regard to all three levels of use, have remained very high and stable throughout the life of the study. There was, however, a little slippage in heroin disapproval rates among 12th graders from 1991 through 1996 (from 96% to 92% for disapproval of experimental use)—a period during which heroin usage rates began to rise.
- Disapproval of regular *cocaine* use rose gradually among 19- to 22-year-olds, from 89% in 1981 to 99% in 1990, about where it has remained since. All three young adult age bands are now near the ceiling of 100%. Disapproval of experimental use of cocaine increased during the 1980s, peaking first among the 12th graders at 94% in 1991. It then peaked among 19- to 22-year-olds (at 94%) and 23- to 36-year-olds (at 92%) in 1995. Finally, it peaked among 27- to 30-year-olds at 90% in 1999. All age groups have had some falloff in disapproval of cocaine use since reaching their peaks in the 1990s. Again, the lag in inflection points between the successive age groups suggests some lasting cohort differences in these attitudes.

that show up as each cohort makes its way through the successive age bands. For the last few years, all age groups' disapproval of experimental cocaine use has hovered around 90%.

- Disapproval of experimental use of *amphetamines* and *sedatives (barbiturates)* increased significantly during the 1980s, accompanied by declining use. Trying amphetamines once or twice was disapproved of by 73%-74% of 19- to 26-year-olds in 1984, compared to 84% by 1990; the corresponding figures for trying sedatives were 84%-85% in 1984 compared to 89%-91% by 1990. Disapproval of amphetamine and sedative use slipped some among 12th graders after 1992 and among 19- to 22-year-olds after 1994, with the 23- to 26-year-olds following suit after 1996, and the 27- to 30-year-old stratum in 2004. This pattern of staggered change again suggests cohort effects, reflecting lasting cohort differences in these attitudes.
- The story for *alcohol* is quite an interesting one, in that changes in the minimum drinking age law may have led to modest changes in norms for the affected cohorts. Between 1980 and 1992, an increasing proportion of 12th graders favored total abstention, with the percent who disapproved even drinking only once or twice rising from 16% in 1980 to 33% in 1992. (This figure has fallen back some, to 26% as of 2004.) Among 19- to 22-year-olds, there was a modest increase from 15% to 22% disapproving any use between 1985 and 1989; this figure stands at 21% in 2004. For the two oldest age groups, there has been rather little change in these attitudes so far. These differing trends may reflect the fact that during the 1980s, the drinking age in a number of states was raised so that by 1987 it was 21 in all states; this change would have the greatest effect on 12th graders, who may have incorporated the legal restrictions into their normative structure and, as they entered the second age band, brought these new norms with them. Put another way, these changes could reflect a cohort effect resulting from the laws that were prevailing when the cohort passed through late adolescence. But the changes may be exhibited only when the respondents are in the age bands directly impacted by the law.

Disapproval of *daily drinking* (of one or two drinks) has not shown any such cohort effects, because all age groups have moved in parallel, at similar levels of disapproval. The three youngest age bands (which include 12th graders through 26-year-olds) showed an increase in disapproval of daily drinking up until about 1990 (there was little data yet available on the oldest age group), but disapproval has declined a fair amount in all of the age groups since then. This pattern of cross-time change closely parallels what was observed for perceived risk of light daily drinking, discussed previously; the decline in both variables may well be due to widely publicized reports that some cardiovascular benefits result from having one or two drinks per day.

There was a considerable increase in disapproval of *heavy drinking on weekends* from the early 1980s for the two youngest age groups, and this continued through 1992 for 12th graders (who then showed some drop-off) and through 1996 among 19- to 22-year-olds (who also then showed some drop-off). As Figure 5-18d illustrates, the prevalence of occasional heavy drinking declined substantially among 12th graders and 19- to 22-year-olds between 1981 and the early 1990s, as norms became more restrictive. There was little or no change in

the older age strata, either in their levels of disapproval or in their rates of occasional heavy drinking, until the early 2000s, when disapproval began to drop some in both strata.

At present, the 12th graders are most likely to disapprove of any drinking (as has been the case for some years) but are the least disapproving of heavy daily drinking. Weekend binge drinking is least disapproved by 19- to 22-year-olds, who report the most of such behavior.

- Some fluctuations in the disapproval of *cigarette smoking* have occurred over the intervals covered by the study. Twelfth graders showed some increase in disapproval of pack-a-day-or-more smoking between 1982 (69%) and 1992 (74%). Disapproval then fell through 1997 (67%) before increasing in the last several years, to 76% in 2004. The 19- to 22-year-olds showed a similar increase from 1982 (66%) to 1989 (76%). All four age strata have shown some upward drift in their level of disapproval of smoking since about 1999 or 2000, suggesting a secular change in these attitudes during this historical period.

A FURTHER COMMENTARY: COHORT DIFFERENCES AND THEIR IMPLICATIONS FOR PREVENTION AND THEORY

One theoretical point to be made—based on the strong evidence reported here for cohort effects in perceived risk and disapproval of many of the drugs under study—is that one cause for cohort effects in actual use is lasting cohort differences in these critical attitudes and beliefs. The attitudes and beliefs brought into adulthood from adolescence tend to persevere.

The second point has to do with the causes of the cohort effects in attitudes and beliefs. We noted earlier that the older respondents are more likely than the younger ones to see the use of *marijuana*, *LSD*, *heroin*, *amphetamines*, *MDMA*, *ice*, *cocaine*, *crack*, and *sedatives (barbiturates)* as dangerous. We have offered the framework for a theory of drug epidemics in which direct learning (from personal use) and vicarious learning (from observing use by others in both the immediate and mass media environments) play important roles in changing these key attitudes.⁴⁸ To the extent that the data on perceived risk represent cohort effects (enduring differences between class cohorts), these findings would be consistent with this theoretical perspective. Clearly, use of these particular drugs was greater when the older cohorts were growing up, and public attention and concern regarding the consequences of these drugs were greatest in the 1970s and early 1980s. In the early 1970s, LSD was alleged to cause brain damage and chromosomal damage, as well as bad trips, flashbacks, and behavior that could prove dangerous. Methamphetamine use was discouraged with the slogan “speed kills.” In addition, there was a serious epidemic of heroin use in the early 1970s. More recent cohorts in our study (through the mid-1990s, at least) were not exposed to those experiences. While there may have been a secular trend toward greater perceived risk for drugs in general, in the case of LSD there may also have been an operating cohort effect (with younger cohorts seeing less danger) offsetting the secular trend among 12th graders; the net effect has been a decrease in 12th graders’ perceived risk of LSD use since 1980.

⁴⁸Johnston, L. D. (1991). Toward a theory of drug epidemics. In L. Donohew, H. E. Sypher, & W. J. Bukoski (Eds.), *Persuasive communication and drug abuse prevention* (pp. 93-131). Hillsdale, NJ: Lawrence Erlbaum.

This vicarious learning process has a very practical importance for national strategy for preventing future epidemics. As future cohorts of youth grow up with less opportunity for such vicarious learning because fewer in their immediate social circles and fewer public role models are using these drugs and exhibiting the adverse consequences of use, these youth will have less opportunity to learn about the adverse consequences of these drugs in the normal course of growing up. Unless those hazards are convincingly communicated to them in *other ways*—for example, through school prevention programs, by their parents, and through the mass media, including public service advertising—they will become more susceptible to a new epidemic of use of the same or similar drugs.

Volume I, the companion volume to the present one, reported an increase in use of several drugs in 8th, 10th, and 12th grades in 1994 through 1997. This increase suggests that this form of “generational forgetting”—in which replacement cohorts lose some of the knowledge held by their predecessors and thus become more vulnerable to using drugs—may well have been taking place during those years. For the cohorts that follow such a rise in use, there is once again an increased opportunity for vicarious learning from the adverse experiences of those around them, but by that time members of affected cohorts have had to learn the hard way what consequences await those who become involved with the various drugs.

TABLE 6-1
Trends in Perceived Harmfulness of Drugs
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

Q. How much do you think people risk harming themselves (physically or in other ways), if they...

	Age Group	Percentage saying "great risk" ^a																							'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004	
Try marijuana once or twice	18	10.0	13.0	11.5	12.7	14.7	14.8	15.1	18.4	19.0	23.6	23.1	27.1	24.5	21.9	19.5	16.3	15.6	14.9	16.7	15.7	13.7	15.3	16.1	16.1	15.9	-0.2	
	19-22	8.3	7.8	9.7	9.7	12.8	11.2	13.0	12.9	16.8	16.9	17.8	19.1	19.7	19.4	18.8	13.3	16.9	14.8	13.4	12.5	14.3	11.9	13.3	17.1	15.3	-1.8	
	23-26					9.6	10.0	12.4	14.5		16.0	14.0	17.7	14.0	15.0	13.0	15.0	15.8	18.5	15.1	16.7	16.4	13.1	13.0	15.1	15.3	13.6	-1.8
	27-30										14.6	16.0	17.0	15.7	15.1	14.0	14.8	16.1	16.2	16.1	16.4	16.1	14.4	17.3	16.2	18.0	13.8	-4.2
Smoke marijuana occasionally	18	14.7	19.1	18.3	20.6	22.6	24.5	25.0	30.4	31.7	36.5	36.9	40.6	39.6	35.6	30.1	25.6	25.9	24.7	24.4	23.9	23.4	23.5	23.2	26.6	25.4	-1.3	
	19-22	13.9	14.2	16.9	16.7	21.7	20.6	22.4	23.0	28.7	29.1	30.1	30.2	29.5	30.3	31.3	25.5	25.6	22.0	22.0	19.8	25.8	18.0	21.0	24.1	23.2	-0.8	
	23-26					15.8	16.3	20.9	20.8	26.8	25.3	30.4	26.2	27.4	24.0	25.5	27.7	27.3	26.4	26.8	26.4	24.9	20.5	24.5	22.2	22.7	+0.4	
	27-30									24.2	25.7	28.7	27.4	27.5	26.8	28.1	28.3	28.1	26.0	25.8	25.3	25.8	25.0	30.2	27.9	25.1	-2.8	
Smoke marijuana regularly	18	50.4	57.6	60.4	62.8	66.9	70.4	71.3	73.5	77.0	77.5	77.8	78.6	76.5	72.5	65.0	60.8	59.9	58.1	58.5	57.4	58.3	57.4	53.0	54.9	54.6	-0.3	
	19-22	43.9	47.8	52.4	58.4	62.2	66.8	67.6	69.4	72.4	74.9	73.0	75.0	69.3	69.2	65.0	62.1	61.3	60.7	53.4	55.2	58.0	49.6	56.7	57.8	57.2	-0.6	
	23-26					52.9	57.5	59.4	65.3	68.3	72.1	71.0	70.9	67.3	64.1	63.2	64.2	62.7	64.1	62.7	60.1	60.3	55.1	53.7	56.7	54.2	-2.6	
	27-30									67.5	69.1	69.2	67.5	68.8	69.4	65.6	69.2	67.3	65.0	63.6	66.1	64.0	61.7	63.5	64.7	59.3	-5.4	
Try LSD once or twice	18	43.9	45.5	44.9	44.7	45.4	43.5	42.0	44.9	45.7	46.0	44.7	46.6	42.3	39.5	38.8	36.4	36.2	34.7	37.4	34.9	34.3	33.2	36.7	36.2	36.2	0.0	
	19-22	44.8	44.4	45.0	44.7	46.0	44.3	47.6	49.4	49.2	49.5	49.3	48.0	45.6	42.4	42.3	40.3	44.4	40.1	38.7	38.1	37.9	37.5	35.3	39.7	39.2	-0.5	
	23-26					48.3	46.9	47.9	51.5	53.7	50.7	52.0	50.1	49.7	49.0	46.8	45.8	46.1	46.6	45.7	49.3	44.9	48.5	45.7	43.8	40.7	-3.1	
	27-30									53.3	55.6	54.6	52.5	53.0	51.5	53.5	52.5	50.1	52.0	49.9	46.4	46.7	44.9	47.5	47.2	47.2	-0.3	
Take LSD regularly	18	83.0	83.5	83.5	83.2	83.8	82.9	82.6	83.8	84.2	84.3	84.5	84.3	81.8	79.4	79.1	78.1	77.8	76.6	76.5	76.1	75.9	74.1	73.9	72.3	70.2	-2.1	
	19-22	83.4	85.3	86.2	86.0	84.5	86.4	87.1	85.6	85.4	85.5	85.8	86.6	87.0	81.3	81.0	80.5	82.4	83.6	78.6	82.2	81.6	79.2	81.1	78.6	78.4	-0.2	
	23-26					89.0	86.6	88.7	90.0	89.2	89.0	88.2	89.1	87.3	85.3	87.5	86.3	84.7	85.6	82.1	85.4	84.1	86.0	85.3	84.3	83.5	-0.8	
	27-30									89.1	91.2	92.0	87.1	88.5	89.0	89.2	88.4	87.0	87.2	90.5	87.8	85.3	86.9	85.3	87.5	83.9	-3.7	
Try PCP once or twice	18								55.6	58.8	56.6	55.2	51.7	54.8	50.8	51.5	49.1	51.0	48.8	46.8	44.8	45.0	46.2	48.3	45.2	47.1	+1.9	
	19-22								63.6	63.8	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	23-26								64.8	63.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	27-30								65.9	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Try MDMA (ecstasy) once or twice	18																		33.8	34.5	35.0	37.9	45.7	52.2	56.3	57.7	+1.4	
	19-22									45.2	47.1	48.8	46.4	45.0	51.1	48.3	46.7	45.5	42.7	37.6	37.9	40.5	46.8	50.1	52.3	+2.3		
	23-26									49.5	47.2	47.4	45.5	41.9	50.6	49.3	50.4	50.5	47.7	50.0	46.7	45.7	45.6	45.9	44.9	-0.9		
	27-30									44.9	48.7	47.7	44.2	51.7	47.3	50.0	50.6	48.8	50.4	50.9	48.9		53.6	52.0	58.8	49.1	-9.7 ss	
Try MDMA (ecstasy) occasionally	18																										—	
	19-22																							72.5	77.8	81.7	78.3	-3.3
	23-26																						72.5	71.9	73.6	77.4	+3.8	
	27-30																						75.2	76.5	79.9	76.9	-3.0	
Try cocaine once or twice	18	31.3	32.1	32.8	33.0	35.7	34.0	33.5	47.9	51.2	54.9	59.4	59.4	56.8	57.6	57.2	53.7	54.2	53.6	54.6	52.1	51.1	50.7	51.2	51.0	50.7	-0.3	
	19-22	31.4	30.4	33.3	28.7	33.1	33.2	35.5	45.9	51.9	51.5	58.1	58.7	56.1	60.5	63.8	57.7	61.9	55.5	55.4	52.8	56.7	48.9	55.5	55.0	55.5	+0.5	
	23-26					31.3	31.1	35.9	48.0	47.1	51.3	51.5	50.5	53.5	54.1	56.0	58.7	57.2	63.1	60.2	62.6	63.1	62.4	61.0	55.4	52.1	-3.2	
	27-30									45.3	53.0	51.6	52.6	51.8	54.7	53.5	56.4	53.6	54.6	60.5	61.7	59.9	60.9	58.8	56.4	61.4	+5.0	

(Table continued on next page)

TABLE 6-1 (cont.)
Trends in Perceived Harmfulness of Drugs
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

Q. How much do you think people risk harming themselves (physically or in other ways), if they...

	Age Group	Percentage saying "great risk" ^a																							'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004	
Take cocaine occasionally	18							54.2	66.8	69.2	71.8	73.9	75.5	75.1	73.3	73.7	70.8	72.1	72.4	70.1	70.1	69.5	69.9	68.3	69.1	67.2	-2.0	
	19-22							53.8	61.3	67.1	72.6	74.6	72.6	74.9	75.4	78.0	73.4	76.6	76.1	71.2	68.0	72.4	70.0	69.9	70.3	70.2	-0.1	
	23-26							50.9	62.6	63.2	69.9	69.9	70.3	69.9	72.8	70.3	76.0	71.3	76.5	74.2	77.8	76.2	74.2	75.4	68.3	74.1	+5.8	
	27-30									62.6	66.6	66.6	69.1	69.9	69.1	69.9	70.0	67.8	73.8	73.2	75.4	76.5	78.1	74.3	72.6	75.3	+2.7	
Take cocaine regularly	18	69.2	71.2	73.0	74.3	78.8	79.0	82.2	88.5	89.2	90.2	91.1	90.4	90.2	90.1	89.3	87.9	88.3	87.1	86.3	85.8	86.2	84.1	84.5	83.0	82.2	-0.8	
	19-22	65.2	69.3	71.5	75.2	75.1	82.9	82.0	88.0	90.3	89.1	93.9	93.5	92.9	91.7	92.2	91.5	92.2	91.6	88.7	88.5	90.7	85.1	88.3	87.4	87.1	-0.3	
	23-26					75.6	76.9	83.0	88.9	90.9	91.2	91.2	92.7	89.9	91.9	92.6	93.3	90.6	93.2	92.9	92.7	92.9	91.1	91.5	88.5	91.5	+3.0	
	27-30									88.9	92.0	91.4	90.9	92.0	91.6	92.1	91.3	91.6	92.7	93.0	92.4	92.3	94.5	91.2	92.9	91.3	-1.6	
Try crack once or twice	18									57.0	62.1	62.9	64.3	60.6	62.4	57.6	58.4	54.6	56.0	54.0	52.2	48.2	48.4	49.4	50.8	47.3	47.8	+0.5
	19-22									59.4	67.3	68.5	69.4	66.9	65.4	63.5	70.1	61.9	65.2	62.0	59.3	56.1	52.9	54.1	54.1	55.1	56.8	+1.7
	23-26									59.1	63.5	69.8	67.3	66.9	67.1	64.2	69.3	64.8	68.6	64.7	67.3	64.6	63.2	59.8	60.9	58.5	56.4	-2.1
	27-30										66.5	64.9	68.7	66.8	64.3	68.8	65.6	66.4	66.7	68.5	66.5	65.0	62.9	69.3	67.4	66.0	62.6	-3.3
Take crack occasionally	18									70.4	73.2	75.3	80.4	76.5	76.3	73.9	73.8	72.8	71.4	70.3	68.7	67.3	65.8	65.4	65.6	64.0	64.5	+0.5
	19-22									75.0	77.3	81.8	82.3	82.7	81.9	83.6	84.3	78.8	83.5	79.1	79.1	75.5	74.9	72.3	75.3	75.3	76.0	+0.7
	23-26									70.3	74.0	79.9	81.1	83.9	84.4	81.6	83.2	81.4	85.9	80.8	84.2	81.6	84.0	80.1	82.2	77.1	76.4	-0.7
	27-30										76.4	76.7	82.6	81.8	79.1	83.6	78.6	81.1	81.3	85.3	81.7	79.8	81.6	84.4	81.5	81.9	82.1	+0.1
Take crack regularly	18									84.6	84.8	85.6	91.6	90.1	89.3	87.5	89.6	88.6	88.0	86.2	85.3	85.4	85.3	85.8	84.1	83.2	83.5	+0.3
	19-22									89.6	91.1	94.1	94.9	95.6	93.4	96.2	96.0	94.2	94.7	93.3	92.8	92.3	91.1	89.6	91.1	93.8	93.3	-0.4
	23-26									88.0	89.2	91.5	94.2	95.4	94.1	93.4	94.9	95.5	96.1	91.4	95.6	94.4	95.6	93.4	94.7	92.2	92.5	+0.3
	27-30										89.6	89.5	95.3	94.4	93.3	93.5	93.0	94.0	94.3	96.0	94.3	95.2	93.5	96.8	94.2	94.4	94.0	-0.5
Try cocaine powder once or twice	18									45.3	51.7	53.8	53.9	53.6	57.1	53.2	55.4	52.0	53.2	51.4	48.5	46.1	47.0	49.0	49.5	46.2	45.4	-0.9
	19-22									44.0	48.6	51.1	54.5	52.7	56.2	49.7	62.0	55.8	57.1	53.8	53.0	47.9	48.0	47.1	47.9	49.4	48.7	-0.8
	23-26									41.0	43.6	48.4	48.9	47.4	45.9	45.6	52.5	48.9	57.2	53.6	54.1	53.8	53.2	53.9	52.5	50.8	46.0	-4.8
	27-30										42.0	45.1	46.2	43.3	42.3	49.9	47.1	48.2	48.9	49.1	49.8	49.7	52.2	53.3	54.4	56.6	52.5	-4.1
Take cocaine powder occasionally	18									56.8	61.9	65.8	71.1	69.8	70.8	68.6	70.6	69.1	68.8	67.7	65.4	64.2	64.7	63.2	64.4	61.4	61.6	+0.2
	19-22									58.0	59.0	63.2	70.0	69.9	72.6	70.6	75.4	73.0	77.4	70.7	73.0	69.3	69.3	64.4	68.9	69.3	68.6	-0.7
	23-26									50.0	53.2	62.2	63.3	67.0	65.8	64.0	68.8	68.8	76.1	72.8	77.0	70.8	76.0	70.5	73.7	67.9	64.6	-3.3
	27-30										53.6	52.7	60.9	59.2	61.2	64.3	61.0	65.9	68.2	69.7	68.5	70.1	71.3	73.5	71.9	71.7	71.5	-0.3
Take cocaine powder regularly	18									81.4	82.9	83.9	90.2	88.9	88.4	87.0	88.6	87.8	86.8	86.0	84.1	84.6	85.5	84.4	84.2	82.3	81.7	-0.6
	19-22									86.6	87.6	91.3	92.5	93.8	92.1	94.0	94.9	93.5	93.8	92.8	91.5	92.4	90.7	89.8	91.0	92.0	91.6	-0.3
	23-26									82.9	84.1	88.5	92.4	93.8	91.3	92.4	92.8	92.1	94.8	90.8	93.7	93.6	94.2	92.2	93.4	89.1	89.4	+0.3
	27-30										85.1	86.7	92.7	91.1	91.5	92.5	90.7	92.7	91.7	93.0	92.3	93.1	91.5	94.0	93.3	94.1	93.1	-0.9

(Table continued on next page)

TABLE 6-1 (cont.)
Trends in Perceived Harmfulness of Drugs
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

Q. How much do you think people risk harming themselves (physically or in other ways), if they...

	Age Group	Percentage saying "great risk" ^a																							'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004	
Try heroin once or twice	18	52.1	52.9	51.1	50.8	49.8	47.3	45.8	53.6	54.0	53.8	55.4	55.2	50.9	50.7	52.8	50.9	52.5	56.7	57.8	56.0	54.2	55.6	56.0	58.0	56.6	-1.4	
	19-22	57.8	56.8	54.4	52.5	58.7	51.0	55.5	57.9	58.9	59.6	58.3	59.9	59.8	58.9	60.8	58.9	61.0	63.9	60.7	63.5	63.2	64.0	63.1	64.6	67.3	+2.7	
	23-26				58.2	59.2	60.8	66.6	65.4	62.3	64.1	62.4	63.7	65.0	63.3	64.1	63.5	67.3	67.3	68.0	70.7	71.9	69.8	70.6	67.5	68.4	-3.1	
	27-30								66.0	69.7	67.5	66.1	66.5	69.3	69.6	66.4	66.4	67.9	69.7	70.1	67.4	68.2	70.9	72.3	68.4	68.4	-3.9	
Take heroin occasionally	18	70.9	72.2	69.8	71.8	70.7	69.8	68.2	74.6	73.8	75.5	76.6	74.9	74.2	72.0	72.1	71.0	74.8	76.3	76.9	77.3	74.6	75.9	76.6	78.5	75.7	-2.7	
	19-22	77.5	77.8	73.6	74.5	74.9	73.6	77.2	77.6	77.5	79.8	80.8	80.2	81.6	78.8	79.0	77.9	82.1	84.7	80.4	82.5	82.0	83.6	82.2	84.9	85.1	+0.2	
	23-26				81.2	80.7	78.9	84.5	82.4	80.8	83.4	84.4	81.5	82.1	80.8	85.3	82.4	86.5	83.9	88.5	86.6	88.4	90.0	88.3	86.7	86.7	-1.5	
	27-30								86.0	86.8	85.3	84.3	84.9	86.2	86.8	83.1	83.8	85.8	86.6	87.1	86.5	86.4	87.9	87.4	88.6	88.6	+1.2	
Take heroin regularly	18	86.2	87.5	86.0	86.1	87.2	86.0	87.1	88.7	88.8	89.5	90.2	89.6	89.2	88.3	88.0	87.2	89.5	88.9	89.1	89.9	89.2	88.3	88.5	89.3	86.8	-2.5	
	19-22	87.2	89.9	87.5	88.6	86.8	90.2	90.7	90.2	89.6	90.8	91.2	91.5	92.2	89.2	91.2	89.9	94.0	93.7	92.4	92.8	94.0	91.3	92.6	93.9	94.3	+0.5	
	23-26				92.0	90.1	90.6	92.8	91.5	91.3	91.0	92.6	91.3	91.6	93.0	93.5	92.7	94.4	93.4	93.7	94.8	95.9	96.3	96.5	96.0	96.0	-0.5	
	27-30								92.7	93.5	93.0	90.7	91.3	92.6	93.8	92.4	92.1	93.8	95.0	93.7	94.2	94.5	95.9	94.9	95.0	95.0	+0.1	
Try amphetamines once or twice	18	29.7	26.4	25.3	24.7	25.4	25.2	25.1	29.1	29.6	32.8	32.2	36.3	32.6	31.3	31.4	28.8	30.8	31.0	35.3	32.2	32.6	34.7	34.4	36.8	35.7	-1.1	
	19-22	24.6	24.6	27.8	24.8	26.9	23.9	27.1	27.4	31.7	28.9	35.6	32.8	34.5	33.3	36.3	32.9	36.8	30.1	31.7	33.7	35.0	34.2	38.1	40.2	36.8	-3.3	
	23-26				29.6	29.4	29.4	34.1	33.2	32.5	35.3	31.0	32.7	32.6	32.9	34.3	34.9	37.8	40.9	41.8	39.9	41.6	38.0	38.3	33.2	33.2	-5.1	
	27-30								35.2	37.5	36.9	36.5	36.2	34.0	37.5	36.0	36.2	34.5	37.6	36.3	39.4	38.5	39.0	40.5	39.2	39.2	-1.3	
Take amphetamines regularly	18	69.1	66.1	64.7	64.8	67.1	67.2	67.3	69.4	69.8	71.2	71.2	74.1	72.4	69.9	67.0	65.9	66.8	66.0	67.7	66.4	66.3	67.1	64.8	65.6	63.9	-1.7	
	19-22	71.9	69.9	68.3	69.9	68.4	68.5	72.3	72.0	73.9	71.3	74.0	77.1	73.5	73.5	71.6	72.2	75.8	72.3	71.9	72.4	73.4	71.1	72.7	75.0	72.4	-2.5	
	23-26				75.8	77.2	75.6	78.2	77.4	76.7	77.8	79.4	76.4	76.2	73.6	80.5	78.5	79.1	77.5	78.7	79.0	77.7	77.9	80.1	75.1	75.1	-4.9	
	27-30								80.6	82.9	83.3	79.4	80.3	79.8	78.4	77.7	75.6	77.4	81.1	82.6	80.8	79.9	79.8	81.5	77.6	77.6	-3.9	
Try crystal meth (ice)	18												61.6	61.9	57.5	58.3	54.4	55.3	54.4	52.7	51.2	51.3	52.7	53.8	51.2	52.4	+1.2	
	19-22												57.8	58.6	57.7	57.5	61.4	58.9	61.1	56.4	55.8	50.6	49.2	52.5	56.5	60.0	+0.3	
	23-26												56.5	56.0	55.6	52.0	61.0	57.8	64.1	60.7	58.2	61.3	60.1	59.2	57.7	58.6	55.9	-2.7
	27-30												59.6	57.2	52.7	60.3	57.9	58.5	59.1	59.8	59.9	61.0	59.7	66.4	62.5	66.6	62.8	-3.8
Try sedatives/barbiturates^b once or twice	18	30.9	28.4	27.5	27.0	27.4	26.1	25.4	30.9	29.7	32.2	32.4	35.1	32.2	29.2	29.9	26.3	29.1	26.9	29.0	26.1	25.0	25.7	26.2	27.9	24.9	—	
	19-22	27.6	26.4	30.5	25.4	29.9	25.0	30.7	29.6	32.7	30.5	36.4	33.5	33.5	33.4	35.0	30.5	34.1	31.4	27.7	28.5	30.3	30.0	30.7	32.7	26.7	—	
	23-26				32.2	29.9	30.2	35.5	35.8	32.9	37.9	31.8	33.5	32.8	34.0	34.8	35.8	37.3	40.3	39.4	37.0	38.5	34.7	36.5	22.2	22.2	—	
	27-30								37.2	38.7	39.0	37.0	38.2	36.5	40.5	36.6	37.2	35.7	36.7	35.2	36.3	40.9	37.3	38.6	31.4	31.4	—	
Take sedatives/barbiturates^b regularly	18	72.2	69.9	67.6	67.7	68.5	68.3	67.2	69.4	69.6	70.5	70.2	70.5	70.2	66.1	63.3	61.6	60.4	56.8	56.3	54.1	52.3	50.3	49.3	49.6	54.0	—	
	19-22	74.0	73.3	72.7	71.3	71.6	71.7	74.5	73.0	74.0	71.7	75.5	75.5	73.6	71.1	69.4	66.4	70.7	69.5	65.1	64.7	64.6	61.8	64.5	63.8	60.2	—	
	23-26				77.4	77.0	74.9	79.9	79.8	76.6	80.5	77.7	76.3	75.0	74.3	77.6	77.1	75.2	73.9	75.1	73.8	73.1	73.1	72.8	63.9	63.9	—	
	27-30								81.5	83.7	84.0	79.6	78.6	80.2	78.3	77.7	74.1	77.1	79.9	80.7	75.5	78.2	75.4	79.0	70.1	70.1	—	

(Table continued on next page)

TABLE 6-1 (cont.)
Trends in Perceived Harmfulness of Drugs
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

Q. How much do you think people risk harming themselves (physically or in other ways), if they...

(Entries are percentages)
Percentage saying "great risk"^a

	Age Group	Percentage saying "great risk" ^a																				'03-'04 change					
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		2000	2001	2002	2003	2004
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	3.8	4.6	3.5	4.2	4.6	5.0	4.6	6.2	6.0	6.0	8.3	9.1	8.6	8.2	7.6	5.9	7.3	6.7	8.0	8.3	6.4	8.7	7.6	8.4	8.6	+0.2
	19-22	3.0	3.4	3.1	2.3	4.7	3.1	5.4	3.5	3.9	5.9	6.1	5.4	5.8	6.6	6.5	4.5	3.3	3.2	4.2	5.7	5.4	4.8	6.6	7.5	5.1	-2.4
	23-26					5.5	3.0	6.5	6.6	4.2	5.1	5.7	4.4	5.6	3.2	4.5	4.3	4.8	4.4	4.4	6.6	3.5	5.5	5.1	5.7	4.7	-0.9
	27-30									5.0	6.3	4.4	6.6	5.6	4.7	4.1	6.7	4.7	4.0	6.2	5.9	4.7	5.5	3.1	6.9	4.6	-2.3
Take one or two drinks nearly every day	18	20.3	21.6	21.6	21.6	23.0	24.4	25.1	26.2	27.3	28.5	31.3	32.7	30.6	28.2	27.0	24.8	25.1	24.8	24.3	21.8	21.7	23.4	21.0	20.1	23.0	+2.9 s
	19-22	22.7	22.9	23.2	23.2	25.0	26.3	27.3	26.1	26.5	28.1	30.1	29.1	30.2	28.0	27.5	24.0	23.0	24.2	22.1	23.9	22.1	19.6	22.7	19.8	21.3	+1.5
	23-26					27.8	27.4	26.9	30.2	29.1	27.8	31.1	30.4	31.6	25.9	26.2	26.1	22.0	20.2	21.0	26.0	21.7	23.5	23.4	19.1	22.9	+3.7
	27-30									27.4	31.7	32.2	31.7	30.9	28.0	27.4	27.2	24.0	24.8	20.8	25.3	22.0	22.7	21.7	21.4	21.8	+0.5
Take four or five drinks nearly every day	18	65.7	64.5	65.5	66.8	68.4	69.8	66.5	69.7	68.5	69.8	70.9	69.5	70.5	67.8	66.2	62.8	65.6	63.0	62.1	61.1	59.9	60.7	58.8	57.8	59.2	+1.5
	19-22	71.2	72.7	73.3	72.7	76.2	74.1	74.0	76.4	72.8	75.7	76.1	75.5	71.8	72.1	70.3	72.5	68.5	71.4	70.4	69.9	69.9	64.5	71.1	66.4	65.3	-1.1
	23-26					76.7	77.9	80.1	77.2	81.8	76.9	79.7	80.2	78.0	76.7	77.5	75.2	72.0	75.1	69.3	72.8	71.7	75.8	74.9	71.1	74.2	+3.1
	27-30									79.3	81.7	84.7	79.1	79.9	79.1	76.6	82.2	76.1	79.3	75.7	75.1	77.4	72.8	76.2	70.6	72.1	+1.5
Have five or more drinks once or twice each weekend	18	35.9	36.3	36.0	38.6	41.7	43.0	39.1	41.9	42.6	44.0	47.1	48.6	49.0	48.3	46.5	45.2	49.5	43.0	42.8	43.1	42.7	43.6	42.2	43.5	43.6	+0.1
	19-22	34.2	30.1	33.5	36.6	37.9	40.2	34.6	36.7	36.9	42.4	40.6	40.8	41.8	42.4	41.9	39.9	40.7	36.6	42.0	37.2	38.9	37.2	37.8	40.4	38.1	-2.3
	23-26					38.4	39.7	39.1	39.8	35.8	37.7	40.2	39.3	37.6	36.2	40.2	37.9	39.1	37.4	41.1	40.2	34.9	39.0	36.8	36.3	37.9	+1.6
	27-30									41.0	42.3	44.1	42.2	45.1	42.9	43.2	44.6	41.5	40.0	40.2	41.9	37.9	41.6	40.6	42.5	40.5	-2.0
Smoke one or more packs of cigarettes per day	18	63.7	63.3	60.5	61.2	63.8	66.5	66.0	68.6	68.0	67.2	68.2	69.4	69.2	69.5	67.6	65.6	68.2	68.7	70.8	70.8	73.1	73.3	74.2	72.1	74.0	+1.9
	19-22	66.5	61.7	64.0	62.1	69.1	71.4	70.4	70.6	71.0	73.4	72.5	77.9	72.6	76.0	71.2	71.6	73.8	76.3	77.2	75.7	77.1	76.6	80.6	77.8	81.1	+3.3
	23-26					71.1	70.1	75.7	73.6	75.5	71.4	78.5	75.3	76.3	78.4	76.4	76.0	77.6	76.5	80.9	79.7	83.9	85.1	83.6	84.1	+0.5	
	27-30									72.8	75.2	77.8	75.4	77.6	75.0	75.3	75.6	73.0	80.3	80.9	80.7	78.4	82.7	80.6	82.0	81.7	-0.3
Use smokeless tobacco regularly	18							25.8	30.0	33.2	32.9	34.2	37.4	35.5	38.9	36.6	33.2	37.4	38.6	40.9	41.1	42.2	45.4	42.6	43.3	45.0	+1.7
	19-22							29.7	34.1	31.1	37.1	33.5	38.9	40.1	43.3	37.6	42.3	40.9	46.5	47.4	47.0	52.0	48.4	53.6	50.8	49.9	-0.9
	23-26							37.0	38.5	35.8	37.9	40.1	38.9	41.6	44.6	42.9	46.6	47.2	46.2	48.4	53.1	49.8	59.8	61.4	58.9	57.8	-1.1
	27-30									42.8	42.8	43.8	44.3	44.1	47.3	46.3	44.2	43.6	50.2	52.6	53.6	49.9	53.2	56.7	58.2	55.7	-2.6
<i>Approximate Weighted N=</i>	18	3234	3604	3557	3305	3262	3250	3020	3315	3276	2796	2553	2549	2684	2759	2591	2603	2449	2579	2564	2306	2130	2173	2198	2466	2491	
	19-22	590	585	583	585	579	547	581	570	551	565	552	533	527	480	490	500	469	464	431	447	424	430	395	402	447	
	23-26					540	512	545	531	527	498	511	505	518	503	465	446	438	420	413	418	400	392	382	401	426	
	27-30									513	587	490	486	482	473	443	450	422	434	416	400	377	384	369	380	388	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aNA' indicates data not available.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

TABLE 6-2
Trends in Proportions Disapproving of Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

<i>Q. Do you disapprove of people (who are 18 or older) doing each of the following?</i>	Age Group	Percentage disapproving ^a																							'03-'04 change		
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004
Try marijuana once or twice	18	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8	68.7	69.9	63.3	57.6	56.7	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	-0.7
	19-22	38.2	36.1	37.0	42.0	44.1	46.6	51.6	52.8	55.8	62.4	59.6	60.4	57.8	60.6	63.5	57.1	55.4	56.2	55.9	54.0	55.2	49.3	48.7	54.2	48.3	-5.9
	23-26				41.2	38.6	42.6	49.1	48.7	52.5	57.5	58.8	55.0	54.6	52.3	51.9	56.3	54.5	55.3	55.7	54.8	51.2	52.4	47.8	53.4	55.4	+5.6
	27-30								49.0	50.9	53.8	54.6	51.9	56.8	55.7	57.5	54.1	59.0	55.7	52.6	58.0	54.4	56.9	54.9	55.4	55.4	+0.5
Smoke marijuana occasionally	18	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5	79.4	79.7	75.5	68.9	66.7	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	+1.2
	19-22	49.6	49.1	51.3	56.0	60.4	62.6	66.7	67.2	69.5	77.3	76.3	77.0	74.8	75.8	76.9	70.4	68.9	70.2	67.8	66.4	70.7	64.6	62.3	68.0	64.3	-3.8
	23-26				54.8	52.8	57.0	64.9	63.4	69.4	73.7	73.3	74.0	71.9	70.9	68.1	72.5	69.2	70.4	71.1	68.6	67.4	64.0	63.8	69.3	69.3	+5.5
	27-30								65.3	67.1	68.9	73.0	67.2	72.2	69.4	72.5	70.5	74.5	72.4	71.5	72.2	70.9	69.1	71.2	69.1	69.1	-2.1
Smoke marijuana regularly	18	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0	89.3	90.1	87.6	82.3	81.9	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	+2.1
	19-22	74.3	77.2	80.0	81.8	84.9	86.7	89.2	88.7	89.1	91.2	93.1	91.3	89.5	90.2	90.1	86.8	87.7	88.1	85.3	84.5	86.6	84.5	82.8	84.8	82.7	-2.1
	23-26				80.6	81.3	83.3	87.4	86.9	90.4	91.0	89.6	90.2	92.1	90.3	90.1	88.9	88.1	87.5	86.1	83.9	86.4	81.7	82.3	87.4	87.4	+5.1 s
	27-30								87.6	87.5	89.7	89.6	87.2	89.4	88.7	91.9	89.9	92.1	89.2	90.0	89.5	89.3	88.8	87.7	88.6	88.6	+0.9
Try LSD once or twice	18	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8	90.1	88.1	85.9	82.5	81.1	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	+2.4 s
	19-22	87.4	84.8	85.9	88.4	88.1	89.1	90.4	90.0	89.9	89.3	90.5	88.4	84.6	88.5	86.8	84.2	83.0	83.1	80.8	83.2	82.3	81.4	83.7	86.2	85.0	-1.2
	23-26				87.3	87.1	88.0	89.9	91.4	91.0	90.7	89.1	88.8	86.9	87.3	87.1	86.7	87.9	84.1	84.8	80.3	83.0	79.2	80.1	84.0	84.0	+3.9
	27-30								91.0	87.2	89.7	87.9	85.6	88.8	88.2	87.4	88.7	88.7	87.3	86.6	87.2	85.7	82.7	85.6	82.5	82.5	-3.1
Take LSD regularly	18	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3	96.4	95.5	95.8	94.3	92.5	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	+0.2
	19-22	98.2	97.4	97.7	97.6	97.6	98.8	98.5	98.0	98.1	97.5	99.1	97.5	97.0	97.8	97.7	96.8	97.0	97.4	96.3	97.0	96.8	96.5	96.9	98.4	97.3	-1.1
	23-26				99.2	98.0	98.5	99.0	98.0	98.4	98.3	98.4	98.3	98.1	97.7	96.7	97.7	96.1	97.6	98.0	97.0	97.1	97.9	96.9	97.1	97.1	+0.2
	27-30								98.8	97.1	98.9	98.9	97.5	98.5	98.7	98.6	98.1	97.5	97.4	97.9	98.6	98.2	98.0	98.2	98.2	98.2	0.0
Try MDMA (ecstasy) once or twice	18																	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	+3.0 s	
	19-22																					81.5	80.3	87.2	83.5	-3.8	
	23-26																					80.6	80.6	80.2	83.1	+2.9	
	27-30																					84.2	84.0	86.3	83.2	-3.1	
Try MDMA (ecstasy) occasionally	18																				NA	NA	NA	NA	—		
	19-22																					92.8	91.8	95.6	93.8	-1.8	
	23-26																					90.5	91.8	92.1	93.3	+1.2	
	27-30																					91.7	93.0	94.3	91.0	-3.4	
Try cocaine once or twice	18	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5	93.6	93.0	92.7	91.6	90.3	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	-0.8
	19-22	73.0	69.3	69.9	74.1	72.5	77.6	78.9	82.3	85.3	88.8	90.1	91.2	90.6	92.7	93.9	94.2	92.0	91.7	89.9	89.9	89.9	87.7	87.9	89.3	87.7	-1.6
	23-26				70.2	70.5	72.1	80.0	82.9	85.5	88.3	88.0	87.3	89.2	89.2	91.8	90.7	91.5	89.0	91.3	87.1	90.1	85.8	86.4	87.4	87.4	+1.0
	27-30								82.1	81.0	85.5	86.9	83.9	85.7	86.6	86.6	88.3	89.2	90.3	90.4	89.4	90.3	88.5	91.5	88.0	88.0	-3.5
Take cocaine regularly	18	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7	97.3	96.9	97.5	96.6	96.1	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	-0.4
	19-22	91.6	89.3	91.9	94.6	95.0	96.3	97.0	97.2	97.9	97.4	98.9	97.9	98.4	97.8	98.8	98.2	97.9	98.0	97.8	97.6	98.0	97.2	97.0	98.2	98.5	+0.3
	23-26				95.7	95.3	97.3	98.1	97.6	98.3	98.4	98.5	98.7	98.4	98.8	97.7	97.8	96.9	98.5	98.3	97.8	97.5	97.5	97.6	98.1	98.1	+0.5
	27-30								98.1	97.0	99.3	99.0	97.2	98.7	99.0	98.9	98.5	97.9	97.8	98.8	98.7	98.4	97.8	98.8	98.8	98.8	0.0

(Table continued on next page)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

<i>Q. Do you disapprove of people (who are 18 or older) doing each of the following?</i>	Age Group	Percentage disapproving ^a																							'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004	
Try heroin once or twice	18	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1	96.0	94.9	94.4	93.2	92.8	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	+0.1	
	19-22	96.3	95.4	95.6	95.2	95.1	96.2	96.8	96.3	97.1	96.4	98.3	95.9	95.9	96.3	96.6	95.6	95.2	95.6	95.1	95.5	94.1	94.2	95.0	96.4	95.9	-0.4	
	23-26					96.7	94.9	96.4	97.1	97.4	96.7	96.8	96.9	96.3	95.4	96.5	95.9	96.1	95.2	94.6	96.3	93.1	95.0	94.8	95.0	95.0	0.0	
	27-30									97.9	95.8	97.5	96.6	94.8	97.3	94.7	96.3	96.0	96.9	95.9	96.7	95.9	96.4	94.4	97.6	94.9	-2.6	
Take heroin occasionally	18	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7	97.3	96.8	97.0	96.2	95.7	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	+0.4	
	19-22	98.6	97.8	98.3	98.3	98.6	98.7	98.3	98.3	98.3	97.9	99.2	98.2	98.1	98.1	98.3	97.7	97.9	97.8	98.2	97.2	98.0	97.9	97.9	98.3	98.9	+0.6	
	23-26					99.2	98.2	98.8	99.1											97.4	97.5	98.5	98.2	97.8	97.5	97.2	98.5	+1.3
	27-30									99.2	97.3	99.0	98.9	97.0	98.9	98.7	98.9	98.0	98.7	97.6	98.8	98.6	98.4	98.6	98.7	98.1	-0.7	
Take heroin regularly	18	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5	97.8	97.2	97.5	97.1	96.4	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	+0.1	
	19-22	99.2	98.5	98.6	98.7	98.7	99.1	98.9	98.6	98.4	98.3	99.5	98.5	98.3	98.4	98.8	98.4	98.3	98.1	98.3	98.2	98.5	98.2	98.3	98.8	99.0	+0.2	
	23-26					99.4	98.8	99.1	99.4	98.7	98.7	98.5	99.3	99.2	98.9	98.8	98.7	98.9	97.6	98.5	98.7	98.8	98.4	98.3	98.6	98.9	+0.3	
	27-30									99.4	97.6	99.4	99.0	97.8	99.0	99.4	99.1	98.6	98.4	98.1	98.8	98.7	98.4	99.3	98.8		-0.4	
Try amphetamines once or twice	18	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3	86.5	86.9	84.2	81.3	82.2	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	-1.7	
	19-22	74.5	70.5	68.9	74.0	73.0	75.6	78.9	79.9	81.8	85.3	84.4	83.9	83.8	87.2	88.3	85.0	84.4	83.3	84.6	84.9	83.8	82.1	81.4	86.3	82.1	-4.3	
	23-26					74.2	74.2	74.6	80.3	83.5	83.3	84.1	84.8	83.4	84.8	82.7	86.0	86.4	85.7	83.5	84.5	82.4	83.9	83.5	79.9	81.6	+1.7	
	27-30									83.5	81.0	84.3	83.7	80.9	83.5	82.0	83.1	85.8	86.3	85.9	86.4	84.5	86.0	86.4	84.9	82.4	-2.5	
Take amphetamines regularly	18	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5	96.0	95.6	96.0	94.1	94.3	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	0.0	
	19-22	94.8	93.3	94.3	93.4	94.9	96.6	96.9	95.1	97.5	96.8	97.5	97.7	96.7	97.3	97.9	96.8	97.2	97.8	96.7	97.5	96.1	97.3	96.4	97.1	97.1	0.0	
	23-26					96.6	95.9	96.6	97.0	97.2	98.1	97.9	97.9	97.7	98.4	97.7	97.0	97.9	97.0	98.0	97.0	97.6	96.8	96.3	97.2	95.9	-1.3	
	27-30									98.1	96.5	98.6	97.8	96.8	97.7	99.0	98.9	98.2	98.1	97.7	98.2	98.5	97.6	97.4	98.1	98.0	-0.1	
Try sedatives/barbiturates^b once or twice	18	83.9	82.4	84.4	83.1	84.1	84.9	86.8	89.6	89.4	89.3	90.5	90.6	90.3	89.7	87.5	87.3	84.9	86.4	86.0	86.6	85.9	85.9	86.6	87.8	83.7	—	
	19-22	83.5	82.3	83.8	85.1	85.2	86.1	88.3	87.5	90.1	92.0	91.1	90.4	88.8	90.7	91.1	90.5	89.1	86.6	85.8	86.6	84.2	85.2	84.2	87.7	81.8	—	
	23-26					84	84.5	84.4	89.8	90.7	89.4	88.8	87.9	88.8	88.5	88.0	89.3	88.3	88.3	87.4	87.3	85.2	86.9	86.8	81.8	80.3	—	
	27-30									90.5	88.3	88.4	88.8	86.6	88.9	87.6	88.0	89.4	88.8	88.4	87.6	87.3	88.5	86.9	89.2	81.8	—	
Take sedatives/barbiturates^b regularly	18	95.4	94.2	94.4	95.1	95.1	95.5	94.9	96.4	95.3	95.3	96.4	97.1	96.5	97.0	96.1	95.2	94.8	95.3	94.6	94.7	95.2	94.5	94.7	94.4	94.2	—	
	19-22	96.6	95.6	97.3	96.5	96.6	98.1	98.0	97.0	97.9	97.7	98.7	98.0	97.9	98.2	98.7	97.7	97.9	97.7	97.7	97.7	97.3	97.4	96.9	97.8	98.5	96.6	—
	23-26					98.4	98.5	97.7	98.6	98.3	98.3	98.5	98.5	98.6	98.5	98.5	97.4	98.4	97.4	98.5	97.6	97.4	97.0	97.1	97.1	96.1	—	
	27-30									98.4	97.1	99.1	98.5	97.7	98.4	99.1	99.0	98.5	97.9	97.7	98.5	98.1	98.4	97.2	98.4	98.1	—	
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4	29.8	33.0	30.1	28.4	27.3	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	-1.2	
	19-22	14.8	14.5	13.9	15.5	15.3	15.4	16.9	16.0	18.4	22.4	17.6	22.2	16.9	20.8	22.2	22.0	22.0	18.3	21.5	18.3	18.4	16.3	18.3	20.1	20.7	+0.6	
	23-26					17.4	16.1	13.2	17.7	13.7	17.5	18.6	19.5	17.4	18.1	17.6	16.5	18.0	15.8	18.6	19.1	19.9	15.9	18.1	13.0	16.3	+3.2	
	27-30									19.5	19.1	18.7	18.8	17.9	19.5	18.6	18.2	16.1	17.4	15.2	15.9	14.8	15.9	18.4	15.4	18.8	+3.4	

(Table continued on next page)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30

(Entries are percentages)

<i>Q. Do you disapprove of people (who are 18 or older) doing each of the following?</i>	Age Group	Percentage disapproving ^a																							'03-'04 change		
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004
Take one or two drinks nearly every day	18	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9	76.5	75.9	77.8	73.1	73.3	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	+0.6
	19-22	67.8	69.7	71.3	73.3	74.3	71.3	77.4	75.3	76.5	80.0	79.7	77.1	76.0	75.0	78.0	74.7	73.5	73.2	70.3	67.3	66.7	68.3	63.9	66.9	68.1	+1.2
	23-26					71.4	73.7	71.6	72.7	74.6	74.4	77.6	76.9	75.5	74.2	73.3	69.7	70.6	68.4	70.2	73.4	66.3	66.5	62.7	65.0	61.7	-3.2
	27-30									76.0	73.9	73.3	76.1	69.5	73.5	72.4	71.8	71.4	71.8	69.8	67.9	65.9	68.9	70.9	63.1	66.7	+3.6
Take four or five drinks nearly every day	18	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9	90.6	90.8	90.6	89.8	88.8	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	+1.5
	19-22	95.2	93.4	94.6	94.6	94.6	94.8	94.9	95.7	94.8	96.1	95.8	96.4	95.5	95.1	96.2	95.5	94.2	93.9	92.4	92.4	92.8	94.2	92.6	92.5	92.2	-0.4
	23-26					96.2	95.0	95.5	96.9	94.3	95.9	96.9	96.1	95.7	95.7	95.2	96.5	93.8	96.1	95.1	94.3	93.5	93.7	92.6	93.1	+0.5	
	27-30									97.4	94.6	96.1	95.3	94.8	94.8	96.4	96.7	96.4	96.2	95.0	97.2	95.3	96.1	95.4	95.6	96.0	+0.5
Have five or more drinks once or twice each weekend	18	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9	67.4	70.7	70.1	65.1	66.7	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	+1.5
	19-22	57.1	56.1	58.2	61.0	59.7	59.4	60.3	61.6	64.1	66.3	67.1	62.4	65.6	63.5	68.1	66.0	69.2	66.5	63.2	63.5	65.1	58.3	57.5	61.9	59.4	-2.5
	23-26					66.2	68.3	66.5	67.5	65.2	63.2	66.9	64.6	69.6	66.8	66.9	65.3	70.9	66.6	69.5	68.1	66.2	66.0	61.2	65.5	60.9	-4.6
	27-30									73.9	71.4	73.1	72.1	68.4	73.4	73.5	73.7	72.4	73.0	71.1	73.1	73.1	73.0	70.9	71.5	73.8	+2.3
Smoke one or more packs of cigarettes per day	18	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8	71.4	73.5	70.6	69.8	68.2	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	+1.5
	19-22	68.7	68.1	66.3	71.6	69.0	70.5	71.4	72.7	73.8	75.6	73.7	73.2	72.6	72.8	75.3	69.8	72.2	74.3	72.3	70.1	73.1	73.2	73.4	73.4	74.8	+1.5
	23-26					69.9	68.7	67.5	69.7	66.4	71.1	71.5	77.2	73.6	72.9	70.3	72.2	73.0	71.7	73.9	73.8	72.7	77.3	74.8	75.7	76.2	+0.5
	27-30									72.8	69.4	73.5	71.2	70.7	73.8	72.3	73.9	72.7	74.3	71.7	71.0	78.6	75.2	78.8	76.2	77.6	+1.4
<i>Approximate Weighted N=</i>	18	3261	3610	3651	3341	3254	3265	3113	3302	3311	2799	2566	2547	2645	2723	2588	2603	2399	2601	2545	2310	2150	2144	2160	2442	2455	
	19-22	588	573	605	579	586	551	605	587	560	567	569	533	530	489	474	465	480	470	446	449	416	413	402	396	431	
	23-26					542	535	560	532	538	516	524	495	538	514	475	466	449	423	401	397	389	404	346	385	403	
	27-30									526	509	513	485	512	462	442	450	430	453	449	429	395	368	359	346	370	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

'NA' indicates data not available.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

^bIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

Chapter 7

THE SOCIAL MILIEU FOR YOUNG ADULTS

Many factors in an individual's social environment influence his or her decision about whether to use drugs. Since its inception, this study has measured three of the more important features of the social environment: (1) the amount of exposure to drug use of various kinds, (2) the norms about drug use of peer groups, and (3) the availability of the various drugs. All three factors are measured by self-reports and are, therefore, measures of the perceived environment. We believe these are all important influences on substance use, at both the individual (micro) level and the aggregate (macro) level. In Volume I, we examined these factors among secondary school students. In this chapter, we examine them for the young adult population, whose social environments are very different from what they were in high school.

Because each of these question sets is contained in only one of the six questionnaire forms, the case counts are lower than those presented in most chapters in this volume (also, in comparison to the secondary school samples covered in Volume I, follow-up samples are much smaller). Therefore, the prevalence and trend estimates are more subject to fluctuation due to relatively greater sampling error.

PEER NORMS AS PERCEIVED BY YOUNG ADULTS

Table 7-1 provides current levels and trends in perceived friends' disapproval of drug use among 12th graders, 19- to 22-year-olds, 23- to 26-year-olds, and 27- to 30-year-olds. (These are the same age groupings used in chapter 6.) Trend data are available since 1980, 1984, and 1988, respectively, for the three four-year age groupings.

The questions regarding friends' disapproval include the same answer scale (stated in terms of disapproval rates associated with different use levels of the various drugs) as the questions that ask about the respondents' own attitudes about those behaviors (discussed in chapter 6). The list of drug-using behaviors is shorter here, and the questions appear on a different questionnaire form and, therefore, have a different set of respondents. However, because the questionnaire forms are distributed randomly in senior year, there are no systematic sample differences across forms. Furthermore, the results for perceived peer norms are generally quite consistent with those for personal disapproval in the aggregate; that is, the proportion saying that they personally disapprove of a drug-using behavior tends to be similar to the proportion saying that their close friends would disapprove of that same behavior.⁴⁹ Exceptions are *trying marijuana once or twice* and *smoking one or more packs of cigarettes per day*, to which respondents have consistently reported their friends' attitudes as *more* disapproving than their own attitudes (especially in the oldest age band), and *heavy weekend drinking*, to which friends' attitudes are seen as *less* disapproving than their own.

⁴⁹The question reads, "How do you think your close friends feel (or would feel) about you... [smoking marijuana once or twice]?" The answer categories are "don't disapprove," "disapprove," and "strongly disapprove." Percentages discussed are for the last two categories combined.

Current Perceptions of Friends' Attitudes

Table 7-1 provides trends for each age band in the proportions of respondents indicating how their close friends would feel about the respondents engaging in various drug-using behaviors. For purposes of simplification, we begin by addressing results across the entire 19- to 30-year age band (tabular data for the entire age band are not presented).

- The peer norms reported by young adults 1 to 12 years past high school are quite similar to those reported by 12th graders. That is, for each of the *illicit drugs other than marijuana*, the great majority of young adults think that their close friends would disapprove of their even *trying* such drugs once or twice (86% for *LSD*, 87% for *amphetamines*, and 89% for *cocaine*).
- Well over half of the young adults (over 62%) now think their friends would disapprove of their even trying *marijuana*, while 70% think their friends would disapprove of occasional use and about 84% think they would disapprove of regular use. So, clearly the norms differ as a function of level of use of marijuana, but at all levels they tend to be restrictive for the majority of young adults.
- Fully two-thirds (67%) of young adults say their friends would disapprove if they were *daily drinkers*, and 9 out of 10 (90%) if they were *heavy daily drinkers*, defined as having four or five drinks nearly every day.
- Friends' disapproval of *heavy drinking on weekends* is distinctly lower. Only 50% to 65% of any age group thinks that their friends would disapprove of their having five or more drinks once or twice each weekend. The 19- to 22-year olds and the 23- to 26-year olds have lower levels of disapproval (52% and 50%) than the 18-year olds (59%) and the 27- to 30-year olds (65%).
- Peer disapproval of *cigarette smoking* is reasonably high in all four age bands: 79% of 12th graders say their friends would disapprove of pack-a-day smoking, as well as 82% of the 19- to 22-year-olds, 78% of the 23- to 26-year-olds, and 85% of the 27- to 30-year-olds. Since the late 1980s, anti-smoking norms have been strongest among the older age bands, but beginning in 2002, this was true only for the oldest of the four age groups, perhaps because of cohort effects working their way up the age scale.

Trends in Peer Norms

- Important changes in the social acceptability of drug-using behaviors among both 12th graders' and young adults' peers have occurred over the life of this study. (See Table 7-1 for data on trends in disapproval by friends for all of the drugs.) Among 12th graders, the proportion who said that their friends would disapprove of their trying *marijuana* rose from 41% in 1979 to 73% in 1992—a period of substantial decline in use. Friends' disapproval also grew substantially stronger in all of the young adult age bands. For example, among the 19- to 22-year-olds, the proportion thinking their friends would disapprove if they even tried marijuana rose from 41% in 1981 to 65% in 1992. A similar peaking occurred for the 23- to 26-year-olds around 1992, at 66%. In all age groups, disapproval subsequently declined,

though the declines were greatest for the younger age groups. Among the 12th graders the decline ended in 1997 and began to reverse, but it continued through 2002 among the 19- to 26-year-olds. Perceived peer disapproval of trying marijuana increased in 2004 in all age groups and for the 19- to 22-year-olds has increased significantly for occasion and regular use levels.

Among those under age 18, friends' disapproval of more frequent use of marijuana also rose until the early 1990s and then declined between 1992 and 1997. It declined through 1999 among the 19- to 22-year-olds and continues to decline among the 23- to 30-year-olds. In essence, peer norms have moved in a way consistent with the existence of some lasting cohort differences.

- There was a more gradual increase in peer disapproval levels of *amphetamine* use for all age groups through 1991, followed by definite declines evident among the 12th graders through 1996. But this weakening of norms against use occurred primarily among adolescents. As of 2004, levels among the 18- to 30-year-olds are approximately the same as they were about 10 years earlier.
- Through 1991, peer disapproval of trying *LSD* showed very little change in any of the age bands, but it fell some in the 1990s, especially among the 18-year-olds and subsequently the 19- to 22-year-olds. These declines bottomed out in a staggered fashion, beginning with the 12th graders in 1997 (who have since shown an 8-percentage-point increase in peer disapproval). Since 2000 there has been a 7-percentage-point increase among the 19- to 22-year-olds.
- Perceived peer norms regarding *cocaine* use were first measured in 1986. During the next five years, self-reported cocaine use declined substantially as peer norms in all age bands shifted considerably toward disapproval. For example, by 1994, 95% of the 19- to 22-year-olds thought their friends would disapprove of their even trying cocaine. After 1994, peer norms against use continued to strengthen a bit in the upper age bands, perhaps through generational replacement, but weakened slightly in the younger age bands, likely reflecting a new cohort effect. At present there is not much difference by age in peer norms against cocaine use.
- Peer norms regarding *occasional heavy drinking* (five or more drinks once or twice each weekend) have tended to be weakest among the 19- to 22-year-old age stratum (where such behavior is most common) and strongest among the 27- to 30-year-old stratum. Since 2002, however, the 23- to 26-year olds were also low relative to the two other age bands in their disapproval of weekend binge drinking. Among 12th graders, friends' attitudes became somewhat more restrictive between 1981 and 1992 as binge drinking declined, but they have been fairly level since then. There was a similar upward trend among the various young adult age bands that followed a staggered pattern likely reflecting a cohort effect. However, between 1997 and 2000 the 19- to 22-year-old age group became somewhat less disapproving of occasional binge drinking; this was followed by a decline in peer disapproval between 2001 and 2004 among the 23- to 26-year-olds.

- Peer norms regarding *cigarette smoking* (one or more packs per day) became somewhat more restrictive among 12th graders in the early years of this study; peer disapproval rose from 64% in 1975 to 73% in 1979. There was little further net change for 13 years (through 1992), when friends' disapproval stood at 76%. However, peer disapproval of smoking slipped some in the 1990s. Between 1992 or 1993 and 1997 or 1998, all age groups showed a decrease in perceived peer disapproval of smoking. Since then all groups, but particularly the two younger strata, have shown some strengthening of norms against smoking.

EXPOSURE TO DRUG USE BY FRIENDS AND OTHERS

Exposure to drug use is measured by two sets of questions, each appearing on a (different) single questionnaire form. The first set asks the respondent to estimate what proportion of his or her friends use each drug, while the second asks, "During the LAST TWELVE MONTHS how often were you around people who were using each of the following to get high or for 'kicks'?" The same questions are asked of 12th graders, and their results are included for comparison purposes in Tables 7-2 and 7-3. (Questions about exposure to drug use were not included in the questionnaires for 35-, 40-, and 45-year-olds, so they could not be included in Table 7-3.) We continue to deal with four-year age bands to increase the reliability of the estimates. (Ages 35, 40, and 45 are included as one-year age bands, but those years have larger numbers of cases than single years at the earlier ages. Both half samples from those two cohorts are surveyed, and only one questionnaire form is administered to those respondents.) At the bottom of each table is a summary of the weighted numbers of cases upon which each annual estimate is based. (The actual numbers of cases are somewhat higher.)

Exposure to Drug Use Among Adults

- Relatively high proportions of young adults in all of these age bands have at least *some* friends who use *some illicit drug* (Table 7-2). In recent years, the proportion declines considerably with age, although this was not always the case. The differences opened up considerably in the 1990s as use rose among the younger strata. In 2004 the proportion is highest for 12th graders (81%) and falls to 63% among 27- to 30-year-olds and to 35%-38% among 35-, 40-, and 45-year-olds. The proportions who say that *most or all* of their friends use one or more of the illicit drugs fall from 23% for 12th graders, to 4% among 27- to 30-year-olds, and between 2% and 3% for the 35-, 40-, and 45-year-olds—quite a dramatic difference, and one that is consistent with the large differences in their own self-reported current use.
- With regard to *illicit drugs other than marijuana*, taken as a whole, considerably fewer report *any* of their friends who use: 51% for 12th graders, 53% for 19- to 22-year-olds, 36% for 27- to 30-year-olds, 20% for 35-year-olds, 19% for 40-year-olds, and 25% for 45-year-olds. These age differences are considerably greater than they were throughout the 1980s. During the period of increasing drug use in the 1990s, primarily among adolescents, the 12th graders reported having the highest proportion of friends using drugs. However, as those 12th graders have aged, it is now the young adults aged 19 to 22 who are showing the highest proportion of friends using drugs, as well as the highest proportion of any of the age strata. The proportions saying that *most or all* of their friends use *illicit drugs other than*

marijuana in 2004 are 7%, 4%, and 3%, respectively, for the three youngest age bands, with fewer than 1% of respondents over the age of 26 reporting such high proportions of their friends using other illicit drugs. Thus, relatively few of these age groups appear to be deeply immersed in a drug culture that involves illicit drugs beyond marijuana.

- With respect to individual illicit drugs, exposure among all of the age groups is greatest for *marijuana*, with 79% of the 12th graders, 76% of the 19- to 22-year-olds, 60% of the 27- to 30-year-olds, and 29%-35% of the 35-, 40-, and 45-year-olds reporting that at least some of their friends use the drug. The next highest exposures are for *ecstasy* (29% among 12th graders and 28% among 19- to 22-year-olds, 31% among the 23- to 26-year-olds, declining to 21% among 27- to 30-year-olds), *amphetamines* (31% among 12th graders and 24% among 19- to 22-year-olds, declining to 14% among 27- to 30-year-olds), *hallucinogens other than LSD* (32% among 12th graders, and 27% among 19- to 22-year-olds, declining to 12% among 27- to 30-year-olds), and *LSD* (24% among 12th graders and 16% among 19- to 22-year-olds, declining to 9% among 27- to 30-year-olds). Because of the dramatic increase in its use during the 1990s and early 2000s, MDMA, or ecstasy, surpassed a number of the more traditional drugs, though its use has declined sharply in recent years.

The proportion of young adults who have some friends who use the other illicit drugs is 10% or higher in at least one of the young adult age groups for the following drugs: *cocaine* (17%-27%), *narcotics other than heroin* (12%-23%), *steroids* (10%-21%), *sedatives (barbiturates)* (12%-19%), *crack cocaine* (5%-16%), and *tranquilizers* (10%-15%).

- For most illicit drugs, the proportion of young adults having any friends who use them decreases with age, consistent with the age differentials in self-reported use. The steepest declines occur with *inhalants* (17.9% of 18-year-olds down to 2.6% of 27- to 30-year-olds). (Inhalant use is not asked of the 35-, 40-, and 45-year-olds, precisely because of this sharp decline in use with age.)
- For some years, *cocaine* was the one illicit drug that showed significantly higher rates of active use among adults than among 12th graders. That is no longer true, although there is rather little drop-off with age in early adulthood; consequently, there is not a great difference associated with age in having friends who use cocaine (17% to 29% for all four of the younger age groups). (The 35-, 40-, and 45-year-olds are asked separately about cocaine powder and crack use; far fewer, but still a fair proportion, report having friends who use cocaine powder—8% to 9% in 2004 for all three groups.)
- For *crack*, however, the story is different. Reported friends' use of crack now descends sharply with age, although this was not true in the mid-1980s, when measures of crack use were first included in the surveys. In 2004, 23% of the 12th graders report having any friends using crack, versus 3% of the 35-, 40- and 45-year-olds.
- The proportion reporting that they have any friends who use *heroin* also decreases sharply with age, from 12.7% among 12th graders to 3.0% among 27- to 30-year-olds. *Narcotics other than heroin* also follow a similar pattern. (Older respondents are not asked these questions.)

- In general, it appears that some respondents who report that their friends use illicit drugs are not directly exposed to that use themselves, judging by the differences in proportions saying they have some friends who use (Table 7-2) and the proportions who say they have not been around people who were using during the prior year (Table 7-3).
- With respect to *alcohol* use, the great majority of young adults have at least *some* friends who *get drunk at least once a week*, although this differs by age: 79% of the 12th graders, 81% of the 19- to 26-year-olds, 76% of the 23- to 26-year-olds, 74% of the 27- to 30-year-olds, 52% of the 35-year-olds, 43% of the 40-year-olds, and 42% of the 45-year-olds. Given the potential serious consequences of this behavior, these rates are impressively high across a wide age range. The proportions who say *most or all* of their friends get drunk once a week differ more substantially by age: 28% of the 12th graders and 29% of the 19- to 22-year-olds, declining sharply to 9% of the 27- to 30-year-olds and 3% of the 45-year-olds. Note in particular how high these rates are among the high school and college-aged populations. In terms of direct exposure during the past year to people who were drinking alcohol “to get high or for ‘kicks,’” having some such exposure is almost universal in the three four-year age groups of young adults: 93%, 92%, and 85%, respectively. (See Table 7-3.)
- From ages 18 through 30, nearly all respondents (84%-90%) have at least a few friends who *smoke cigarettes*, with considerable falloff by age 35. In fact, about a fifth of the 12th graders and 19- to 22-year-olds (21%) state that *most or all* of their friends smoke. Above those ages the proportions decline sharply, to 9% of the 27- to 30-year-olds, and about 6% for those 35 years of age and older. This increase in the segregation of smokers from nonsmokers may reflect the stratification of young people after high school as a function of educational attainment, which is highly correlated with cigarette smoking. Also, it can be seen in Table 7-2 that there was much less age-related difference in the late 1980s, suggesting that the sharp rise in smoking among high school students during much of the 1990s accentuated the age differentials and that those differentials remain, reflecting lasting cohort effects.

Trends in Exposure to Drug Use Among Young Adults

Tables 7-2 and 7-3 also provide *trend* data on the proportions of friends using drugs and the proportions directly exposed to drug use. (Both of these measures of exposure to use will be discussed in this section.) Once again, trends are available for the 19- to 22-year-olds since 1980, for the 23- to 26-year-olds since 1984, and for the 27- to 30-year-olds since 1988. Data for 35- and 40-year-olds are available for friends' use since 1994 and 1998, respectively. The first year for which data for the 45-year-olds were available was 2003. (Questions about frequency of being around drug users were not included in the questionnaires administered to the 35-, 40-, and 45-year-old respondents.) Data for 12th graders since 1980 also have been included in these tables for comparison purposes.

- An examination of Table 7-3 shows that exposure to illicit drug use (in the 12 months preceding the survey) declines with age for *any illicit drug, marijuana*, and *any illicit drug other than marijuana*, as well as for nearly all of the specific illicit drugs. In general, these

differences replicate across different historical periods, with the exception of cocaine, which began to show a decline in exposure with increasing age after 1996. These declines reflect age or maturational effects in both exposure to use and in personal use of most drugs.

- Until 1992, young adults' trends in exposure to use tended to parallel those observed for 12th graders. Between 1980 and 1992, that meant a decreasing number of respondents were exposed to *any illicit drug* use (Table 7-3) or reported any such use in their own friendship circle (Table 7-2). After 1992, however, an important *divergence* in trends among age groups emerged: 12th graders showed a substantial increase in both friends' use and exposure to use (and in self-reported use); the 19- to 22-year-olds showed a similar rise, but lagged by a few years; the 23- to 26-year-olds have recently shown a slight rise; while the 27- to 30-year-old age band did not show a rise until 2002. As is discussed in earlier chapters, this pattern no doubt reflects the emergence of lasting cohort differences combined with the process of generational replacement.
- *Marijuana* showed a very similar pattern of change. In addition, returning to the measures of friends' use, it is particularly noteworthy that, while 34% of the 19- to 22-year-olds in 1980 said *most or all* of their friends used marijuana, only 9% said the same in 1993. Clearly, the number of friendship groupings in which marijuana use was widespread dropped dramatically over that interval. The figure increased to 19% by 1999, where it remained for a couple of years. Friends' use has declined some in the past few years for the two youngest strata, but not yet among the older strata.
- The proportion reporting having any friends who use *any illicit drugs other than marijuana* began to decline after 1982. By 1991 or 1992 there had been a considerable drop in all four age groups. This drop appears to be due particularly to decreases in friends' use of *cocaine* and *amphetamines*, although there were decreases for *sedatives (barbiturates)* and *tranquilizers* as well. The levels then began to rise in the two youngest age bands in the early 1990s, while at the same time they continued to decline in the two oldest age bands, opening up a large age-related difference in exposure to use. Since 2001 there has been some decline in friends' use in the two youngest age strata, narrowing the age differences some.
- Between 1987 and about 1992, there was a considerable drop in the proportion of all four age groups who said they had any friends who used *crack*. (Self-reported use declined in the same period.) After that decline, the rates of friends' use increased some in the two youngest age bands and decreased some in the four oldest ones, resulting in a large age difference in the proportion of friends using crack. Of course, some of that apparent age difference could be due to a greater amount of cumulative attrition of the most drug-prone members of our panels, and crack users would certainly be among the most drug-prone. In 2004 there was a significant 3.8 percentage-point increase in the proportion of 12th graders saying they had any friends using crack (to 22.5%) and a non-significant 4.2 percentage point increase among the 19- to 22-year-olds who had friends using crack. Whether this is indicative of a new trend or simply a coincidence remains to be seen.

- It is noteworthy that there has been a substantial increase since the early 1990s in the proportion of 12th graders and 19- to 22-year-olds reporting that they have friends using *narcotics other than heroin* (though the increase was greater among the 12th graders, where it began earlier). Increases within the two older age strata began later, and have been more modest; however, unlike the case with the younger strata, they seem to be still underway.
- The proportions of all age groups for which data are available on friends' use of *ecstasy (MDMA)* increased sharply in recent years, though in a staggered fashion. Twelfth graders showed the first sharp increase beginning after 1992, 19- to 22-year-olds after 1994, 23- to 26-year-olds after 1996, and 27- to 30-year-olds after 1997. These sharp increases ended among 12th graders in 2001 and among 19- to 30-year-olds a year later. In the last year or two there have been sharp declines in all four age strata in the proportions saying that they had any friends using ecstasy, which corresponds to the sharp declines in self-reported use. In 2004, however, the decline decelerated sharply among 19- to 22-year-olds and leveled among 23- to 26-year-olds. There has not yet been a decline among the 27- to 30-year-olds, though friends use has held steady since 2002.
- For all four age groups there were modest declines between 1987 and 1992 in the proportion saying that most or all of their friends drink *alcohol*. Since 1992, there has been very little change in the lower four age bands, though a drop among 12th graders in 2002 began to open a difference before all of the older age strata showed a similar drop in 2003. The 35-, 40- and 45-year-olds report fewer friends who drink and substantially fewer who get drunk on a weekly basis than do the younger age bands; this is an age-difference that has been constant.
- Among 12th graders, the proportion who said most or all of their friends smoked *cigarettes* declined appreciably between 1975 and 1981 during the same period that self-reported use declined. Neither measure showed much change until about 1992. Thereafter, substantial increases in both measures occurred. By 1997 fully one-third (34%) of 12th graders reported that most or all of their friends smoked cigarettes, up from 21% in 1992, declining since. Among 19- to 22-year-olds a decline in friends' use occurred between 1980 (or possibly earlier) and 1985, followed by a leveling through 1994. The percentage saying most friends smoke increased from 22% in 1994 to 29% in 2000, before beginning to decline. Among 23- to 26-year-olds, a downturn was evident between at least 1984 (the first year for which data are available) and 1988, and then reported friends' use leveled. Since about 2002, some slight decline is evident. These staggered changes up until about 1998 illustrate that the cohort effects were moving up the age spectrum (as the cohorts themselves aged). Since 1998 there has been a considerable decline in the proportions saying that any of their friends smoke in both the two youngest age strata, some increase in the proportions smoking among the 23- to 26-year-old stratum (which now contain the heavier-smoking senior classes of the mid-1990s), but not much change yet in the four oldest strata.

Nearly all of these changes in exposure to drug use across the various drugs parallel changes in self-reported use by these four age groups. This pattern reinforces our trust in the validity of the self-report data, because there would presumably be less motivation to distort answers about the

proportion of an unnamed set of friends who use a drug than about one's own use of it. Also reassuring is the systematic nature of the patterns of change across age strata (whether in terms of parallel trends consistent with a secular trend or staggered ones consistent with a cohort-related trend).

PERCEIVED AVAILABILITY OF DRUGS BY YOUNG ADULTS

Young adults participating in the follow-up survey receive survey questions identical to those asked of 12th graders regarding how difficult they think it would be to get each of the various drugs if they wanted them. The questions are contained in only one of the six questionnaire forms. The data for the follow-up samples, which are grouped into the same four-year age bands, are presented in Table 7-4, along with the data for the 12th graders and the 35-, 40-, and 45-year-olds. Sample sizes are presented at the bottom.

Perceived Availability

As is true with the 12th graders, substantial proportions of the American adult population have access to various illicit drugs. (We do not ask about access to alcohol and cigarettes, because we assume access to be universal.)

- **Marijuana** is the most available illicit drug, with 86%-89% of the young adult age strata saying it would be “fairly easy” or “very easy” to get. Access generally decreases with age after age 26, but it is still 70% by age 45.
- **Ecstasy (MDMA)** is among the most widely available of all of the illicit drugs other than marijuana. Its availability is now greatest among 23- to 26-year-olds (51%). It used to be greatest for 12th graders (now 48%) and 19- to 22-year-olds (now at 46%). This reversal may reflect the aging of the heavier using cohorts, who knew how to acquire ecstasy. Availability then falls off among 27- to 30-year-olds (to 41%). (The question is not asked of those 35, 40, or 45 years of age.)
- **Amphetamines** are also among the most available of the illicit drugs (44% to 54% among young adults and 35% to 40% among 35-, 40-, and 45-year-olds). Note that there is a fair difference among the age bands, with those above age 26 reporting amphetamines to be less available.
- **Sedatives** are about as available as amphetamines to these age groups (41% to 49%), whereas **tranquilizers** are reported as available by considerably fewer (30% to 37%).
- **Cocaine** is reported as readily available by a significant proportion of young adults, with 38%-51% saying it would be fairly easy to get. **Powdered cocaine** availability does not differ much by age (38%-49%). **Crack** is available to somewhat smaller proportions than powdered cocaine—28%-37% for all three post-high school young adult age strata and 36%-40% for 35-, 40-, and 45-year-olds. Cocaine was considerably more available to the older age groups

in the 1980s (up through age 30, at least) but is now about equally available across all four lower age bands, including 12th graders.

- **Hallucinogens other than LSD** are reported as available by 47%-49% of 12th graders and 19- to 22-year-olds. Availability is lower among the 23- to 30-year-olds at 31%-44%.
- **LSD** now shows a moderate degree of availability for all ages (24%-39%).
- More than a third of young adults (36%-48%) say they can get **narcotics other than heroin** fairly easily, as do 40% of 12th graders.
- **Steroids** show some declines in perceived availability with increasing age, as has generally been the case, ranging from 43% among 12th graders down to 31% among the 27- to 30-year-olds. (The question is not asked of respondents above age 30.)
- About a quarter (22%-31%) of young adults and nearly a third of 12th graders (30%) say that they could get **heroin** fairly easily; yet a far smaller proportion of them report having used heroin.
- **Crystal methamphetamine (ice)** is perceived to be available by about a quarter to a third of each age group (26%-32%).

Trends in Perceived Availability

- **Marijuana** has been almost universally available to the adolescent and young adult age groups throughout the historical periods covered by the data (since 1975 in the case of 12th graders). There was a slight decrease through 1991 among 12th graders since the peak year of 1979 and a slightly larger decrease from 1980 through 1991 among 19- to 22-year-olds. Availability rose by a few percentage points in nearly all strata between about 1993 and 2001 or so, and since then has slipped back a few percentage points in the two younger strata. Perceived availability is now a bit higher for the younger age groups (between 86% and 89% for 12th graders through 27- to 30-year-olds versus 69% to 70% for those aged 40 and 45). In general their trends in availability have been quite parallel, though.
- **Cocaine** availability increased among all three younger age strata over the 1984-1988 interval, reaching historic highs in 1988 and 1989. (Twelfth graders showed a rise in availability in earlier years—from 1975 to 1980—followed by a leveling between 1980 and 1985. Availability was also level during the latter period among 19- to 22-year-olds.) From a policy perspective, it is worth noting that in all three age bands for which we have data, the perceived availability of cocaine *increased* in 1987—the same year that use actually dropped sharply. Between 1988 and 1989, in the two younger age strata (aged 18, and 19 to 22) the proportions reporting that they could get cocaine fairly easily were still increasing, whereas in the older age strata the proportions were beginning to decrease. In 1990 and 1991, all four groups reported decreased availability—quite parallel to the number who had friends who were users and to personal use, both of which dropped substantially in these years and then leveled in 1992. Perceived availability of cocaine dropped to between 49% and 57% for all

four age groups by 1993, with the absolute declines ranging from 4 to 7 percentage points. After that there has been some falloff in perceived availability in all age strata through age 30—particularly among those ages 23 through 30—and an increasing convergence. However, in 2004, reported availability rose significantly for 18 year olds and 23- to 26-year-olds, bringing all to about the 50% level, except the 27- to 30-year-olds (who are at 38%).

- **Crack** availability peaked in 1988-1989 for all age groups (it was first assessed in 1987) and declined through 1992, with little further change until 1995. Since 1995, crack availability has declined in all strata.
- The trends in **LSD** availability among young adults have some parallels to those for 12th graders. Among 12th graders, there was a drop of about 10 percentage points in the mid-1970s and a later drop in the interval 1980 to 1986. The latter drop, at least, was paralleled in the data from 19- to 22-year-olds. After 1986, availability increased considerably in all age bands, reaching its peak levels (the highest we have recorded since these questions were introduced) by 1995, with a considerable difference by age developing (with availability lower with increases in age); since 1995, availability has fallen considerably in the youngest two age strata and by less in the older strata, once again narrowing the differences among the age groups. Indeed, the drop-off in availability of LSD to 12th graders and 19- to 22-year-olds was quite sharp in 2002, possibly contributing to the steep decline in use that year. Availability among 35- and 40-year-olds is the lowest, decreasing slightly since data were first available.
- In the early 1980s, there was a fair decline among all age groups in the availability of **hallucinogens other than LSD**; there was little additional change through 1992. From 1992 to 1995 the three youngest age groups all showed an increase in availability, with 12th graders showing the largest increase. From 1996 to 2000, availability was fairly steady. All age groups showed substantial increases in 2001, but this was presumably due to the changed question wording in which, among other things, “shrooms” was added to the examples of hallucinogens. Since 2001 the general pattern has been one of stability, with the younger strata indicating a higher level of availability (e.g., 49% for 12th graders versus 31% for 27- to 30-year-olds).
- The availability of **MDMA (ecstasy)** rose very substantially in all of these age groups during the 1990s and early 2000s. (The questions were first introduced in 1989 and 1990.) Among the 12th graders, reported availability nearly tripled, from 22% in 1989 to 62% in 2001—the peak year for 12th graders. All four age groupings showed sharp increases in 2000 and 2001, with the oldest groups continuing to increase through 2002—their peak year. The 27- to 30-year-olds are the least likely to say that ecstasy is readily available to them; still, 41% of them say it would be fairly easy to get. In fact, the two youngest strata have shown considerable declines since 2002.
- **Heroin** availability varied within a fairly narrow range from 1980 to 1986 but then showed a fair-sized increase among both 12th graders and young adults through 1990. It then rose

further among 12th graders and 19- to 22-year-olds through 1995 before declining a bit. In the older two of the four age groups, heroin availability remained fairly flat from 1990 to 1995, and then increased some through 1999, declining thereafter. What is clear is that heroin was much more available to all of these age groups in the 1990s than it was in the 1980s. Further, all age groups have shown some decline in heroin availability over the last several years. For the past three years there is only modest variability in heroin availability across the 18 to 30 age range.

- The availability of *narcotics other than heroin* slowly rose among all age groups between 1980 and 1989, followed by considerable stability from 1989 through 1994. Since 1994, availability has increased modestly as use has been rising steadily. In general, there has not been a consistent difference by age in narcotics other than heroin availability.
- The reported availability of *amphetamines* peaked in 1982 for both 12th graders and 19- to 22-year-olds; since then it has fallen by 15 percentage points among 12th graders and 20 percentage points among the 19- to 22-year-olds. Since 1984, when data were first available, there has been a decline of 14 percentage points among the 23- to 26-year-olds, as well. For the 27- to 30-year-olds, reported availability decreased by 10 percentage points between 1988 (the first measurement point) and 2004. There have been decreases among the 35- and 40-year-olds as well. In general, the age groups above age 30 have reported somewhat lower availability than the younger strata.
- By way of contrast, *crystal methamphetamine* or “ice” exhibited an increase in availability in the 1990s, rising for all four age strata from 1991 to 1998 or 1999, before stabilizing.
- *Sedatives (barbiturates)* have exhibited a long-term decline in availability since about 1981 or 1982 through 2003 in the two younger groups—by 20 percentage points among 12th graders and 23 percentage points among 19- to 22-year-olds. All groups showed an increase in 2004, no doubt due primarily to a change in the question wording.
- *Tranquilizer* availability has declined long-term by more than half among 12th graders, from 72% in 1975 to 30% in 2004. Since 1980, when data were first available for 19- to 22-year-olds, availability declined by nearly as much over a shorter period of time (from 67% in 1980 to 30% in 2004), such that previous differences in availability between them were eliminated by 1992. The older age groups also showed a considerable decline in the availability of tranquilizers through 2003 or 2004. In general, the trend lines for the different age groups have been quite parallel, as was true for sedatives (barbiturates).
- Data on *steroid* availability were first gathered in 1990, and although there has not been much change in availability since then, availability did appear to peak in 1992 in all age strata. This was followed by a modest decline in all age groups. Twelfth graders showed some increase between 1996 and 2002, and 19- to 22-year-olds showed a very modest increase between 1998 and 2002. In 2003, decreases in the availability of steroids were observed in all age groups; however, results were more mixed in 2004.

TABLE 7-1
Trends in Proportions of Friends Who Disapprove of Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

<i>Q. How do you think your close friends feel (or would feel) about you...</i>	Age Group	Percentage saying friends disapprove ^a																						'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001		2002	2003	2004
Trying marijuana once or twice	18	42.6	46.4	50.3	52.0	54.1	54.7	56.7	58.0	62.9	63.7	70.3	69.7	73.1	66.6	62.7	58.1	55.8	53.0	53.8	55.1	58.1	57.6	54.1	58.4	59.5	+1.1
	19-22	41.0	40.6	46.9	47.1	51.6	54.5	55.2	54.7	58.7	63.0	63.6	64.7	64.7	63.4	63.7	58.5	64.3	58.4	57.0	56.5	56.0	54.2	53.4	56.5	61.0	+4.5
	23-26					47.7	47.0	49.1	53.9	58.2	62.6	61.3	64.5	65.6	65.5	63.2	63.8	61.2	59.3	66.5	62.6	64.6	55.2	53.8	51.4	57.7	+6.4
	27-30									58.6	58.7	61.4	64.6	63.5	64.4	66.3	66.1	65.8	65.0	65.4	61.8	63.9	64.9	67.1	61.9	67.2	+5.3
Smoking marijuana occasionally	18	50.6	55.9	57.4	59.9	62.9	64.2	64.4	67.0	72.1	71.1	76.4	75.8	79.2	73.8	69.1	65.4	63.1	59.9	60.4	61.6	63.9	64.3	60.3	64.2	65.0	+0.9
	19-22	50.9	49.2	54.0	57.9	59.4	64.6	64.4	65.1	69.8	71.5	74.1	73.9	74.3	73.1	73.0	66.6	71.3	65.1	65.1	64.6	61.8	61.0	62.6	63.3	70.1	+6.7 s
	23-26					54.3	56.4	57.1	63.1	68.1	73.2	71.8	72.5	75.3	73.5	72.2	70.7	70.8	68.5	73.6	70.2	70.9	63.9	64.5	61.6	63.5	+1.9
	27-30									67.8	69.4	71.9	73.7	76.0	75.1	76.4	73.8	75.6	72.4	74.9	74.5	75.0	74.2	72.9	71.4	76.9	+5.5
Smoking marijuana regularly	18	72.0	75.0	74.7	77.6	79.2	81.0	82.3	82.9	85.5	84.9	86.7	85.9	88.0	83.5	80.6	78.9	76.1	74.1	74.7	74.5	76.1	77.8	75.3	77.0	77.3	+0.3
	19-22	70.3	75.2	75.7	79.5	80.0	82.7	83.5	84.8	86.9	87.5	89.1	88.4	89.1	87.6	85.9	83.9	84.5	83.3	81.1	78.2	78.5	80.0	80.5	79.1	84.4	+5.3 s
	23-26					77.8	78.4	80.9	82.0	85.8	89.2	88.1	87.9	90.3	89.1	88.8	84.9	89.5	85.6	87.1	86.8	86.9	83.7	82.8	80.0	79.2	-0.9
	27-30									85.4	86.0	88.4	89.2	88.7	88.2	88.9	89.7	89.6	87.8	90.8	89.2	91.6	90.1	87.9	87.2	88.0	+0.9
Trying LSD once or twice	18	87.4	86.5	87.8	87.8	87.6	88.6	89.0	87.9	89.5	88.4	87.9	87.9	87.3	83.5	83.4	82.6	80.8	79.3	81.7	83.2	84.7	85.5	84.9	87.5	87.3	-0.2
	19-22	87.4	90.5	88.0	89.3	89.3	91.1	90.5	91.8	90.8	91.2	89.1	89.9	87.2	87.7	87.9	84.6	85.3	83.6	81.7	82.0	82.1	85.2	86.9	86.9	88.6	+1.7
	23-26					87.4	90.8	88.6	89.8	88.9	91.0	90.1	92.4	88.9	87.7	86.3	85.3	88.5	85.4	87.6	84.5	85.3	82.8	83.6	79.3	82.4	+3.1
	27-30									88.8	89.7	92.3	91.1	91.4	89.9	91.2	89.7	89.3	88.5	88.7	88.4	85.6	87.4	86.3	87.1	87.7	+0.7
Trying cocaine once or twice	18							79.6	83.9	88.1	88.9	90.5	91.8	92.2	91.1	91.4	91.1	89.2	87.3	88.8	88.7	90.2	89.3	89.1	91.2	87.9	-3.3 ss
	19-22							76.4	NA	84.8	87.7	89.2	92.3	91.9	92.4	94.7	91.7	91.5	91.8	90.0	91.2	89.4	89.1	91.7	90.6	90.3	-0.3
	23-26							70.8	NA	81.4	84.5	84.1	86.7	87.4	87.7	87.9	90.4	90.0	91.1	92.0	89.6	90.5	88.0	88.5	83.6	84.2	+0.7
	27-30									81.8	81.1	83.7	83.5	84.4	86.1	87.8	87.5	88.7	89.4	89.3	90.5	90.4	89.3	88.8	89.9	91.8	+2.0
Taking cocaine occasionally	18							87.3	89.7	92.1	92.1	94.2	94.7	94.4	93.7	93.9	93.8	92.5	90.8	92.2	91.8	92.8	92.2	92.2	93.0	91.0	-2.0
	19-22							84.9	NA	91.0	93.8	94.2	95.6	95.9	95.6	97.5	95.6	95.7	96.6	93.1	95.7	94.7	94.5	95.6	95.1	96.0	+0.9
	23-26							81.7	NA	88.2	91.5	92.4	94.1	93.8	93.5	94.3	94.6	95.4	95.1	95.2	95.2	96.7	94.7	93.2	91.2	90.1	-1.2
	27-30									87.7	89.5	90.0	92.2	92.3	92.8	94.6	94.1	94.6	94.2	96.1	95.4	95.9	94.2	94.0	95.1	96.3	+1.1
Trying an amphetamine once or twice	18	78.9	74.4	75.7	76.8	77.0	77.0	79.4	80.0	82.3	84.1	84.2	85.3	85.7	83.2	84.5	81.9	80.6	80.4	82.6	83.0	84.1	83.8	83.3	85.9	84.7	-1.2
	19-22	75.8	76.7	75.3	74.3	77.0	79.7	81.5	81.3	83.0	83.5	84.5	86.5	83.8	85.0	87.2	83.1	86.0	84.5	84.0	85.8	81.6	84.5	87.6	87.6	89.4	+1.7
	23-26					78.4	79.1	76.7	81.7	83.0	85.6	84.3	85.0	83.6	84.2	84.7	87.6	86.5	83.3	87.0	85.9	85.1	83.1	83.9	81.5	82.7	+1.3
	27-30									82.7	84.1	84.9	84.6	84.7	84.1	85.9	85.5	85.6	85.9	85.8	87.2	87.8	86.4	86.0	87.9	88.9	+0.9
Taking one or two drinks nearly every day	18	70.5	69.5	71.9	71.7	73.6	75.4	75.9	71.8	74.9	76.4	79.0	76.6	77.9	76.8	75.8	72.6	72.9	71.5	72.3	71.7	71.6	73.4	71.6	74.7	72.8	-1.9
	19-22	71.9	72.1	68.6	73.5	71.6	72.2	72.7	70.2	73.9	77.1	73.3	73.7	74.0	71.2	73.0	68.3	68.9	73.5	67.3	68.6	66.6	64.9	68.5	64.4	72.4	+8.0 s
	23-26					63.6	66.8	67.7	68.3	69.2	70.8	72.7	72.5	72.1	67.6	71.5	68.2	72.8	68.1	66.9	66.1	65.4	64.4	61.6	62.1	61.8	-0.3
	27-30									71.0	68.0	70.4	71.9	68.8	73.2	70.9	68.8	65.7	67.3	66.7	64.3	67.3	67.1	64.0	64.5	65.0	+0.5

(Table continued on next page)

TABLE 7-1 (cont.)
Trends in Proportions of Friends Who Disapprove of Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

<i>Q. How do you think your close friends feel (or would feel) about you...</i>	Age Group	Percentage saying friends disapprove ^a																							'03-'04 change		
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004
Taking four or five drinks nearly every day	18	87.9	86.4	86.6	86.0	86.1	88.2	87.4	85.6	87.1	87.2	88.2	86.4	87.4	87.2	85.2	84.1	82.6	82.5	82.8	82.2	82.8	84.4	80.1	83.1	82.9	-0.2
	19-22	93.7	91.7	89.9	91.9	91.7	92.5	91.5	90.8	90.4	92.5	89.9	91.7	92.6	89.6	90.1	88.8	88.1	90.0	85.9	87.9	86.6	84.6	87.7	86.8	89.8	+2.9
	23-26					90.8	90.2	92.5	92.8	93.7	92.1	92.1	92.4	91.1	93.1	92.1	92.2	92.6	90.7	93.7	89.9	92.5	91.1	88.1	89.3	87.8	-1.5
	27-30									92.8	92.0	92.9	92.7	92.7	93.9	94.0	92.9	91.9	93.8	92.1	95.3	92.4	91.2	92.7	92.6	92.5	-0.2
Having five or more drinks once or twice each weekend	18	50.6	50.3	51.2	50.6	51.3	55.9	54.9	52.4	54.0	56.4	59.0	58.1	60.8	58.5	59.1	58.0	57.8	56.4	55.5	57.6	57.7	57.8	55.6	60.3	59.4	-0.9
	19-22	53.5	51.7	51.7	53.3	50.8	53.3	47.0	49.4	50.5	56.8	53.1	51.4	53.6	51.9	54.4	55.5	52.1	56.4	52.8	51.8	45.2	47.4	50.4	47.9	52.4	+4.5
	23-26					53.8	57.3	61.0	57.2	58.8	57.5	55.1	56.8	58.4	57.6	61.4	58.9	58.4	55.6	60.0	54.5	56.6	56.9	52.9	49.5	49.5	0.0
	27-30									61.9	65.1	66.3	68.2	66.2	66.7	63.7	64.6	61.6	64.0	63.0	57.7	65.8	58.8	63.3	59.6	64.6	+5.0
Smoking one or more packs of cigarettes per day	18	74.4	73.8	70.3	72.2	73.9	73.7	76.2	74.2	76.4	74.4	75.3	74.0	76.2	71.8	72.4	69.2	69.3	68.5	69.0	71.2	72.6	74.5	75.7	79.2	78.6	-0.6
	19-22	75.6	75.1	75.4	78.5	76.2	79.7	77.7	78.6	80.2	78.4	77.5	78.3	79.0	76.0	73.8	70.9	73.9	76.5	69.2	73.9	71.1	74.3	77.3	78.3	82.1	+3.8
	23-26					73.9	77.3	80.3	80.5	79.5	80.5	78.5	83.3	82.3	77.4	80.1	78.8	78.3	75.8	76.5	78.0	79.9	77.0	75.4	78.3	77.6	-0.7
	27-30									81.2	80.9	82.9	84.5	83.1	86.8	82.5	83.4	81.9	80.5	81.9	82.6	84.0	83.6	86.1	84.0	84.6	+0.7
<i>Approx. Weighted N=</i>	18	2766	3120	3024	2722	2721	2688	2639	2815	2778	2400	2184	2160	2229	2220	2149	2177	2030	2095	2037	1945	1775	1862	1820	2133	2208	
	19-22	569	597	580	577	582	556	577	595	584	555	559	537	520	510	470	480	471	466	436	430	379	402	361	399	427	
	23-26					510	548	549	540	510	513	516	516	507	481	463	445	436	419	425	394	398	378	366	363	377	
	27-30									483	518	479	480	451	451	457	439	439	422	440	397	394	374	364	346	408	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

TABLE 7-2
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
		Take any illicit drug^a																									
% saying any friends	18	87.5	85.4	86.3	82.6	81.0	82.4	82.2	81.7	79.1	76.9	71.0	69.1	67.3	71.0	78.3	78.6	80.6	83.4	84.6	82.0	82.0	82.8	81.8	80.7	81.2	+0.5
	19-22	90.2	88.0	86.8	85.0	82.3	82.9	80.5	76.7	77.2	78.4	72.7	71.5	66.8	71.7	71.6	76.2	77.2	79.8	77.3	83.1	81.1	78.3	79.4	78.1	-1.4	
	23-26				83.6	82.7	80.3	80.9	74.4	73.8	65.8	63.0	67.3	64.6	66.7	65.3	64.6	67.0	67.6	67.9	67.8	66.9	73.4	70.8	70.8	0.0	
	27-30								74.8	72.9	69.6	67.1	61.5	60.2	57.1	58.5	59.1	60.9	58.3	59.6	55.6	57.2	61.8	58.6	63.1	+4.5	
	35														38.1	37.4	39.7	39.2	38.4	36.3	37.7	39.1	40.9	37.5	37.9	+0.4	
	40																		39.2	38.2	38.0	38.4	36.2	36.5	34.6	-1.9	
	45																							37.8	38.3	+0.6	
% saying most or all	18	32.5	29.8	26.5	23.8	20.9	22.7	21.5	18.6	15.8	15.7	11.6	11.7	12.0	15.5	20.3	21.7	23.8	23.7	25.9	25.5	24.5	25.2	23.1	23.5	23.0	-0.5
	19-22	34.9	32.8	28.1	22.4	21.9	18.2	16.2	14.0	13.5	10.9	10.5	8.8	9.0	10.4	14.9	13.1	17.3	16.2	16.8	20.6	18.9	20.3	20.2	17.3	14.7	-2.5
	23-26				19.6	15.4	16.2	11.7	9.5	9.7	9.5	7.4	6.2	6.4	8.7	7.6	8.8	10.5	9.6	8.4	9.7	10.4	10.3	10.3	11.7	+1.3	
	27-30								8.6	6.4	5.9	2.9	5.8	5.0	5.6	6.1	3.6	4.5	5.3	5.7	5.3	7.1	6.9	6.9	3.9	-3.0	
	35														2.1	1.9	2.0	3.0	3.1	2.8	3.1	3.2	2.9	3.2	2.8	-0.4	
	40																		2.3	2.0	2.0	1.6	2.2	1.6	1.6	0.0	
	45																							2.2	1.5	-0.7	
Take any illicit drug^a other than marijuana																											
% saying any friends	18	62.4	63.3	64.7	61.2	61.3	61.8	63.3	62.4	56.5	56.2	50.1	46.3	47.1	48.7	53.7	53.7	54.5	55.1	55.6	51.2	52.5	55.0	54.3	50.0	51.4	+1.3
	19-22	67.9	67.8	66.7	65.2	60.8	62.1	61.0	57.3	53.5	60.8	53.4	51.5	45.3	51.4	46.3	46.4	46.5	49.7	53.3	54.8	56.1	60.0	57.2	50.8	53.4	+2.6
	23-26				63.7	64.0	59.0	61.1	55.1	54.2	47.8	41.8	46.1	42.3	39.4	40.3	32.8	35.1	35.4	41.1	42.5	42.6	49.4	42.3	47.1	+4.9	
	27-30								55.9	55.0	49.7	47.2	37.7	38.5	33.9	37.7	36.4	33.9	34.1	35.2	31.7	33.5	36.0	34.7	35.8	+1.1	
	35														21.4	21.6	22.1	19.2	19.3	19.0	17.9	18.7	20.4	18.5	20.2	+1.8	
	40																		20.9	21.0	21.9	21.4	21.0	20.2	18.5	-1.7	
	45																						23.4	25.1	+1.7		
% saying most or all	18	11.1	11.9	10.9	11.0	10.3	10.4	10.3	9.2	6.9	7.7	5.1	4.6	5.3	7.1	7.1	7.7	8.9	7.0	8.9	7.4	7.4	7.0	6.1	6.7	7.3	+0.6
	19-22	9.8	12.9	11.8	9.8	9.3	8.6	7.6	5.0	5.3	4.0	3.2	2.6	3.3	4.0	4.4	3.5	6.2	4.1	4.3	5.1	7.7	8.0	5.7	5.1	3.5	-1.6
	23-26				10.6	6.6	8.6	5.2	3.9	4.2	3.4	1.6	1.8	2.8	2.5	1.9	1.9	2.6	2.8	2.2	3.8	3.7	2.8	3.7	3.1	-0.6	
	27-30								4.6	3.0	2.8	1.0	1.4	1.5	1.5	1.5	0.9	1.2	0.9	1.3	1.5	2.6	2.3	0.7	0.8	+0.1	
	35														0.8	0.5	0.7	0.5	0.7	0.9	1.0	0.9	0.6	0.6	0.4	-0.2	
	40																		0.4	0.8	0.7	0.5	0.3	0.3	0.2	-0.1	
	45																							0.7	0.7	0.0	

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Smoke marijuana																											
% saying any friends	18	86.4	83.0	84.4	80.3	77.7	79.5	79.2	78.4	75.3	72.5	68.3	65.8	63.1	67.4	75.6	76.1	78.0	81.4	83.2	80.7	80.5	81.2	79.4	78.9	79.5	+0.6
	19-22	88.8	86.4	85.2	83.8	81.6	81.1	78.5	75.3	75.1	73.8	67.6	68.0	63.5	67.6	67.4	68.8	74.9	74.7	77.2	73.9	81.2	78.4	77.2	76.5	75.6	-0.9
	23-26					82.0	80.8	77.7	79.4	71.6	69.8	61.8	59.6	61.3	61.2	62.6	63.2	62.6	63.5	65.0	64.4	64.8	64.5	68.8	67.7	68.4	+0.8
	27-30									71.8	68.2	65.1	62.6	58.0	57.4	52.3	55.7	55.1	58.3	55.5	57.0	51.7	56.5	59.0	55.8	60.4	+4.5
	35															36.9	36.3	36.3	35.0	34.6	33.3	34.9	35.6	37.4	32.9	34.7	+1.9
	40																			34.6	32.5	32.3	31.8	31.4	30.7	29.9	-0.8
	45																							31.1	29.4		-1.7
% saying most or all	18	31.3	27.7	23.8	21.7	18.3	19.8	18.2	15.8	13.6	13.4	10.1	10.0	10.3	13.9	18.9	20.7	22.2	22.5	23.8	24.2	23.2	24.0	21.4	21.7	21.1	-0.5
	19-22	34.1	30.6	25.6	20.6	19.4	16.0	13.3	12.5	12.2	9.0	9.2	8.3	8.2	8.5	13.0	12.5	16.3	16.2	16.4	19.4	16.6	18.5	18.6	16.0	15.0	-1.0
	23-26					17.0	14.3	13.7	10.4	7.8	8.6	8.3	6.9	5.6	5.6	7.5	6.6	8.2	9.8	9.0	8.5	8.2	9.0	8.7	9.3	9.8	+0.5
	27-30									6.8	4.4	4.0	2.8	5.1	5.2	5.0	5.6	3.5	3.9	4.8	5.5	4.9	6.3	6.2	6.7	3.5	-3.2
	35															3.0	2.5	2.9	2.9	2.8	2.6	2.8	2.6	2.7	3.1	2.7	-0.4
	40																			2.1	1.4	1.9	1.2	2.0	1.4	1.6	+0.2
	45																								1.9	0.9	-1.0
Use inhalants																											
% saying any friends	18	17.8	16.5	18.4	16.1	19.3	21.2	22.4	24.7	20.8	22.1	20.0	19.2	22.2	23.7	26.5	27.5	27.2	27.4	25.9	21.6	23.5	22.2	21.0	17.5	17.9	+0.4
	19-22	11.9	13.2	13.8	12.3	11.7	9.6	10.9	12.7	10.9	11.7	13.0	12.2	12.6	13.8	14.0	14.2	16.2	13.7	16.2	16.3	13.7	13.7	10.4	10.0	9.5	-0.4
	23-26					7.7	6.7	7.2	6.1	6.2	5.9	6.1	4.4	5.1	6.3	7.0	9.3	5.6	7.5	6.2	7.9	6.9	7.5	7.4	7.9	6.2	-1.7
	27-30									4.6	3.5	2.9	2.5	3.3	2.9	3.5	4.0	4.1	3.6	3.8	4.2	3.6	6.0	4.5	3.2	2.6	-0.6
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	1.2	0.9	1.3	1.1	1.1	1.5	2.0	1.9	1.2	1.9	1.0	0.7	1.8	1.8	2.0	2.0	2.4	1.9	2.7	1.8	1.4	1.4	1.2	1.1	1.2	+0.1
	19-22	0.5	0.4	0.7	0.3	0.5	0.6	0.7	0.7	0.7	0.4	0.6	0.2	0.8	0.7	0.7	0.6	1.1	0.7	1.3	0.8	0.6	1.2	0.4	0.6	0.0	-0.6
	23-26					0.6	0.2	0.6	0.1	0.2	0.4	0.4	0.1	0.0	0.1	0.2	0.7	0.5	0.8	0.0	0.1	0.7	0.1	0.4	0.3	0.0	-0.3
	27-30									0.3	0.0	0.2	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.0	0.0	0.0
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change		
Use nitrites																													
% saying any friends	18	19.0	17.4	17.5	14.5	15.0	15.6	18.0	18.3	13.6	13.3	10.4	8.9	9.0	10.7	10.0	10.7	11.2	11.9	12.9	10.9	11.0	11.9	11.2	8.5	9.4	+0.9		
	19-22	18.4	16.0	14.2	13.8	8.9	9.9	11.7	13.2	10.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	23-26					10.8	7.8	8.0	7.9	5.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	27-30									6.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																				NA	NA	NA	NA	NA	NA	NA	—	
	45																									NA	NA	—	
	% saying most or all	18	1.3	1.2	0.9	0.7	1.2	1.0	1.2	1.3	0.7	0.9	0.6	0.4	0.7	0.7	0.8	0.8	0.8	0.7	1.0	0.7	1.0	0.6	0.8	1.0	1.2	+0.1	
		19-22	0.3	0.4	0.9	0.6	0.6	0.6	0.4	0.4	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		23-26					0.8	0.3	0.4	0.3	0.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		27-30									0.5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		40																				NA	NA	NA	NA	NA	NA	NA	—
		45																									NA	NA	—
		Take LSD																											
% saying any friends		18	28.1	28.5	27.8	24.0	23.9	24.4	24.5	25.3	24.1	25.2	25.0	23.4	28.1	31.3	34.1	36.9	37.9	36.5	36.8	32.2	31.9	32.2	28.6	21.9	23.5	+1.6	
		19-22	30.9	25.9	26.5	22.6	21.6	18.8	18.7	18.2	19.0	20.1	20.1	22.0	22.2	28.8	23.8	26.9	28.6	24.7	29.4	28.2	27.8	28.4	24.0	15.4	15.9	+0.6	
		23-26					21.5	17.2	15.4	15.9	13.3	14.1	12.3	12.5	15.0	17.2	17.3	21.5	15.3	18.2	15.2	18.1	19.3	16.8	15.8	16.1	14.4	-1.7	
		27-30									10.4	7.7	9.1	8.6	10.9	8.7	8.1	12.0	11.6	12.3	12.6	13.4	11.8	12.5	13.1	11.4	8.9	-2.5	
		35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		40																				NA	NA	NA	NA	NA	NA	NA	—
		45																									NA	NA	—
	% saying most or all	18	1.8	2.2	2.4	1.4	2.0	1.5	1.8	1.6	1.5	2.4	1.9	1.7	2.4	3.8	4.2	4.8	5.0	3.7	4.7	3.9	3.1	2.9	1.7	1.9	1.5	-0.4	
		19-22	1.2	0.8	0.9	1.0	0.6	0.8	0.9	0.6	1.3	0.4	1.2	1.4	1.9	2.1	2.5	2.3	3.8	1.4	2.5	1.8	2.1	2.7	1.6	0.8	0.3	-0.5	
		23-26					0.8	0.5	1.0	0.2	0.6	0.5	0.6	0.2	0.4	0.7	1.1	0.7	0.7	0.6	1.0	1.5	0.9	0.3	0.4	0.8	0.4	-0.5	
		27-30									0.3	0.2	0.3	0.3	0.0	0.3	0.4	0.3	0.4	0.4	0.1	0.6	0.4	0.4	0.3	0.1	0.0	-0.1	
		35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
		40																				NA	NA	NA	NA	NA	NA	NA	—
		45																									NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Take other psychedelics/																											
hallucinogens^b																											
	18	28.2	26.3	25.6	22.1	21.3	22.0	22.3	21.7	17.8	18.1	15.9	15.1	17.0	19.3	21.4	23.8	26.4	26.3	27.4	22.5	24.0	35.4	33.6	30.1	31.9	+1.8
% saying any friends	19-22	33.4	25.5	25.1	21.0	20.2	16.6	15.8	15.0	16.1	13.9	15.3	14.2	12.0	15.0	13.8	14.9	17.2	17.2	19.1	18.9	20.9	33.6	33.5	24.8	26.8	+1.9
	23-26				20.0	16.7	13.2	13.2	11.7	9.6	8.7	8.5	9.8	9.4	10.3	11.7	10.4	13.0	11.7	9.6	11.3	18.6	22.4	20.2	24.5	+4.3	
	27-30								10.6	7.4	7.1	6.8	7.9	7.1	6.6	7.9	7.5	6.8	7.8	9.4	8.0	14.6	14.9	13.5	12.4	-1.2	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	2.2	2.1	1.9	1.6	1.9	1.4	1.3	1.2	0.9	1.4	1.0	0.8	1.0	1.7	2.2	2.2	2.3	2.6	3.1	2.4	2.4	2.9	2.3	2.4	2.6	+0.2
	19-22	1.5	0.9	1.1	1.2	0.7	1.0	0.7	0.6	0.9	0.2	0.5	0.8	0.7	0.9	1.6	1.5	1.0	1.1	1.7	0.8	2.0	2.3	2.2	1.5	1.1	-0.4
	23-26				0.8	0.3	0.5	0.3	0.2	0.3	0.8	0.1	0.4	0.7	0.6	0.8	0.1	0.8	0.7	0.8	0.3	0.6	0.7	1.0	0.8	0.8	-0.2
	27-30								0.2	0.1	0.3	0.2	0.0	0.2	0.3	0.1	0.2	0.3	0.2	0.2	0.4	0.6	1.0	0.1	0.0	0.0	-0.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
Take PCP																											
% saying any friends	18	22.2	17.2	17.3	14.2	14.2	15.9	16.1	15.5	13.5	14.7	13.0	12.0	12.7	15.6	15.5	18.3	20.3	19.7	20.2	16.8	17.5	19.1	17.2	13.6	11.8	-1.8
	19-22	24.1	15.3	15.3	12.6	9.5	8.9	10.1	9.7	10.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	23-26				11.6	6.8	7.4	6.9	5.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	27-30								6.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	1.6	0.9	0.9	1.1	1.1	1.2	1.2	1.1	0.8	1.2	0.5	0.5	0.9	1.9	1.2	1.2	1.3	1.4	1.6	1.5	1.7	1.3	1.0	1.5	1.1	-0.4
	19-22	0.5	0.3	0.3	0.5	0.7	0.7	0.2	0.1	0.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	23-26				0.6	0.0	0.4	0.0	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	27-30								0.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>		Age																							'03-'04		
	Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	change
Take MDMA (ecstasy)																											
% saying any friends	18											12.4	11.9	10.7	12.8	15.9	20.7	24.2	27.7	24.5	26.7	37.3	41.9	38.0	34.2	28.9	-5.3 ss
	19-22										16.3	14.3	12.0	12.9	13.7	11.3	17.2	20.7	21.4	26.0	30.7	42.4	43.3	43.4	31.3	27.6	-3.6
	23-26										7.6	9.0	9.5	11.0	9.8	11.4	11.2	11.3	15.1	13.7	15.2	25.9	29.4	36.8	27.0	31.2	+4.2
	27-30										5.6	6.3	5.4	4.6	6.6	5.8	6.9	10.1	7.4	8.5	12.4	13.1	17.8	20.6	19.4	20.6	+1.3
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18											2.2	1.7	2.1	1.2	1.7	2.8	3.0	2.6	2.5	2.7	4.8	5.2	3.7	2.7	3.2	+0.5
	19-22										0.4	0.7	0.2	0.7	0.7	0.5	0.5	0.8	1.7	2.0	2.9	4.9	5.8	2.7	1.9	1.9	0.0
	23-26										0.5	0.2	0.1	0.1	0.5	0.1	0.4	0.1	0.8	0.8	0.4	2.9	1.7	1.2	2.0	1.1	-0.9
	27-30										0.5	0.3	0.0	0.1	0.3	0.2	0.5	0.1	0.3	0.0	0.8	0.4	0.3	0.9	0.5	0.6	+0.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
Take cocaine																											
% saying any friends	18	41.6	40.1	40.7	37.6	38.9	43.8	45.6	43.7	37.7	37.4	31.7	26.8	26.3	24.5	26.1	24.8	28.1	28.2	31.2	27.8	27.2	27.1	26.8	23.8	29.3	+5.5 sss
	19-22	51.0	48.9	49.8	46.5	47.6	45.9	48.3	45.7	42.0	42.7	33.2	29.7	22.8	24.3	21.5	22.0	19.4	22.2	26.8	25.7	24.8	27.4	28.2	25.5	26.2	+0.6
	23-26					52.4	53.2	51.6	50.7	47.1	40.8	34.8	29.0	28.8	27.1	22.3	24.4	18.1	19.7	18.7	20.1	20.3	19.4	23.7	21.9	27.4	+5.4
	27-30									47.9	43.3	38.3	35.7	29.9	27.6	22.6	26.2	20.8	21.5	18.6	20.7	16.5	19.7	16.0	17.0	17.0	0.0
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	6.1	6.3	4.9	5.1	5.1	5.8	6.2	5.1	3.4	3.7	2.1	1.5	1.5	2.1	1.5	2.0	2.2	2.0	3.2	2.9	2.0	1.7	1.7	2.4	2.3	-0.1
	19-22	7.0	8.6	7.8	6.1	6.3	6.1	6.1	3.3	3.5	2.1	1.2	1.1	1.0	0.5	1.5	0.9	1.0	0.8	1.5	1.1	1.0	1.8	1.0	1.4	0.7	-0.7
	23-26					9.1	5.3	7.0	4.1	3.1	2.7	2.1	0.6	0.9	0.8	1.0	0.3	0.4	1.1	0.9	0.5	0.8	1.6	1.0	1.6	1.0	-0.6
	27-30									3.8	2.0	2.3	0.9	1.2	0.8	0.8	0.4	0.4	0.6	0.1	0.4	0.4	0.5	0.6	0.3	0.3	-0.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Take crack																											
% saying any friends	18								27.4	25.4	26.1	19.2	17.6	17.8	17.9	20.0	19.2	21.6	22.2	24.4	19.0	21.4	23.4	21.5	18.7	22.5	+3.8 s
	19-22								23.8	21.8	20.6	14.6	14.3	11.8	13.6	13.8	14.0	9.4	13.1	16.4	15.7	16.5	17.4	18.0	11.8	16.0	+4.2
	23-26								26.4	22.4	19.8	14.4	10.8	10.8	8.8	8.8	11.1	8.2	8.3	8.3	8.8	7.9	8.6	10.1	10.4	10.8	+0.4
	27-30									22.1	18.4	16.6	11.6	10.3	10.2	10.4	10.3	8.6	6.3	6.4	8.7	6.0	7.1	6.4	6.5	5.2	-1.3
	35															4.5	5.1	4.4	3.1	2.8	3.2	3.9	2.8	3.2	2.8	3.1	+0.3
	40																			3.8	3.0	2.9	3.5	2.6	2.7	2.6	-0.1
	45																								3.7	3.3	-0.4
% saying most or all	18								2.2	1.1	2.1	0.6	0.6	0.7	0.9	1.0	1.1	0.9	1.1	1.7	1.5	1.4	0.8	0.8	1.4	1.6	+0.2
	19-22								0.7	0.8	1.0	0.6	0.2	0.1	0.3	0.4	0.3	0.5	0.3	0.9	0.9	0.5	0.3	0.2	0.4	0.1	-0.3
	23-26								0.8	0.9	0.8	0.5	0.1	0.1	0.5	0.2	0.0	0.3	0.5	0.4	0.0	0.5	0.3	0.0	0.3	0.5	+0.3
	27-30									1.2	0.9	0.9	0.3	0.0	0.6	0.3	0.1	0.2	0.2	0.1	0.0	0.0	0.0	0.3	0.1	0.0	-0.1
	35															0.6	0.3	0.4	*	0.1	0.3	0.5	0.2	0.3	0.3	0.1	-0.2
	40																			*	0.2	0.2	0.1	*	0.0	0.0	0.0
	45																								0.4	0.3	-0.1
Take cocaine powder																											
% saying any friends	18								NA	NA	25.3	24.6	19.8	19.7	18.1	20.7	19.2	22.8	24.8	22.9	22.0	21.3	20.1	22.4	23.2	25.4	+2.2
	19-22								NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	23-26								NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	27-30									NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	35															14.2	12.9	15.4	11.1	10.4	10.0	10.3	9.4	9.4	8.2	9.2	+0.9
	40																			10.8	8.9	8.8	8.8	8.5	7.6	7.6	-0.1
	45																								8.3	8.0	-0.3
% saying most or all	18								NA	NA	2.3	2.5	1.8	2.0	1.6	1.9	1.7	1.9	2.0	1.9	1.9	1.8	1.5	1.9	1.9	3.3	+1.4 s
	19-22								NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	23-26								NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	27-30									NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	35															0.8	0.3	0.6	0.4	0.4	0.6	0.7	0.4	0.4	0.4	0.2	-0.3
	40																				0.2	0.2	*	0.2	0.1	0.1	0.0
	45																								0.5	0.5	0.0

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Take heroin																											
% saying any friends	18	13.0	12.5	13.2	12.0	13.0	14.5	15.3	13.9	12.4	14.0	11.4	11.4	13.2	13.3	14.3	14.5	15.6	15.6	16.5	12.7	14.9	13.1	12.9	10.3	12.7	+2.4 s
	19-22	11.0	8.1	9.4	7.5	7.1	6.5	8.5	8.5	7.8	6.8	6.5	6.1	4.7	7.0	8.1	10.4	6.7	7.4	9.4	9.7	7.7	8.7	8.9	5.3	7.0	+1.7
	23-26					6.1	4.4	4.3	6.5	3.6	5.2	4.2	3.6	3.8	4.5	4.9	5.8	4.0	6.2	5.8	4.8	4.7	5.0	5.2	6.1	2.9	-3.2 s
	27-30									3.8	2.8	4.5	2.7	3.1	3.6	4.2	3.6	4.4	4.2	3.5	3.8	2.8	4.3	3.9	3.4	3.0	-0.4
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	1.0	0.5	0.7	0.8	0.8	0.9	1.1	0.9	0.7	1.1	0.4	0.4	0.7	1.1	1.0	1.1	0.9	0.8	1.3	1.0	1.1	0.9	0.7	0.9	0.9	0.0
	19-22	0.3	0.5	0.1	0.2	0.4	0.6	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.4	0.4	0.4	0.2	0.5	0.1	0.3	0.6	0.0	0.3	0.0	-0.3
	23-26					0.4	0.2	0.2	0.0	0.2	0.4	0.2	0.3	0.4	0.1	0.2	0.2	0.0	0.7	0.0	0.0	0.3	0.0	0.1	0.0	0.0	0.0
	27-30									0.2	0.1	0.2	0.2	0.0	0.2	0.3	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.3	0.0	0.0	0.0
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
Take other narcotics																											
% saying any friends	18	22.4	23.1	23.9	20.8	21.4	22.8	21.8	23.2	19.2	19.2	17.2	13.7	14.9	16.1	18.5	19.5	21.8	22.2	24.8	22.9	23.1	24.0	27.5	21.6	24.6	+3.0
	19-22	22.8	20.4	21.9	17.9	17.4	16.9	14.6	15.4	14.1	15.0	12.9	14.1	10.8	13.2	10.5	15.9	13.4	13.2	15.2	19.8	23.2	23.0	21.8	21.9	22.6	+0.7
	23-26					16.0	14.9	14.0	13.0	10.6	10.8	10.5	8.5	8.4	8.7	8.0	10.5	8.9	9.9	9.4	10.4	11.2	13.5	14.6	18.4	16.8	-1.7
	27-30									12.1	8.6	9.1	9.3	7.5	8.2	8.0	7.7	9.5	7.9	8.3	7.2	8.4	11.2	11.8	11.0	12.0	+1.0
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	1.7	1.5	1.4	1.4	1.6	1.4	1.8	1.4	1.2	1.4	0.9	0.5	1.1	1.2	1.0	1.6	1.5	1.4	2.9	1.8	2.0	2.0	2.1	2.4	2.4	-0.1
	19-22	0.9	0.7	0.6	0.5	0.8	1.0	0.5	0.4	0.9	0.1	0.6	0.4	0.5	0.6	0.6	0.6	0.4	0.4	0.8	0.4	1.2	1.8	1.3	1.0	0.5	-0.5
	23-26					0.4	0.3	0.7	0.0	0.3	0.2	0.2	0.0	0.0	0.0	0.3	0.2	0.0	0.6	0.3	0.0	0.4	0.5	0.6	0.6	0.5	0.0
	27-30									0.3	0.0	0.2	0.2	0.1	0.2	0.2	0.0	0.2	0.0	0.0	0.2	0.0	0.0	0.3	0.1	0.0	-0.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Take amphetamines																											
% saying any friends	18	43.9	48.8	50.6	46.1	45.1	43.3	41.8	39.5	33.4	33.5	28.7	24.3	24.3	27.5	28.1	30.3	32.2	32.7	33.8	30.8	32.9	33.2	34.4	28.1	31.4	+3.3
	19-22	54.1	52.2	51.3	49.7	46.1	42.1	38.5	34.5	26.8	29.6	23.3	26.2	19.5	21.0	20.9	21.7	21.6	21.1	24.4	25.5	28.4	28.0	28.6	24.0	23.5	-0.5
	23-26					45.6	40.1	33.5	32.1	28.4	23.1	20.6	17.1	15.1	16.8	16.2	18.2	12.5	14.4	14.1	14.2	14.5	17.5	18.4	18.0	18.8	+0.8
	27-30									26.1	21.6	19.3	17.0	15.3	14.0	13.1	13.7	15.5	12.9	11.0	11.8	11.9	12.9	12.3	12.0	13.5	+1.5
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	4.8	6.4	5.4	5.1	4.5	3.4	3.4	2.6	1.9	2.6	1.9	1.3	1.3	2.0	1.8	2.0	2.8	2.4	3.4	2.8	3.1	2.2	2.4	2.1	2.9	+0.8
	19-22	3.8	5.7	4.6	3.8	3.3	2.9	1.3	1.9	1.4	0.7	1.0	0.6	0.9	0.2	1.1	1.2	0.7	0.7	1.2	0.7	1.7	1.6	1.3	1.2	0.5	-0.6
	23-26					1.9	1.8	1.7	1.2	0.3	0.6	0.7	0.8	0.4	1.5	0.9	0.5	0.2	0.8	0.5	0.6	0.3	0.5	0.3	0.7	0.1	-0.5
	27-30									0.6	0.4	0.5	0.5	0.1	0.5	0.5	0.3	0.3	0.1	0.3	0.6	0.1	0.5	0.9	0.1	0.0	-0.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
Take sedatives/barbiturates^c																											
% saying any friends	18	30.5	31.1	31.3	28.3	26.6	27.1	25.6	24.3	19.7	20.3	17.4	14.8	16.4	17.8	18.2	17.8	21.6	20.4	22.8	20.9	21.6	22.1	25.3	18.1	25.2	—
	19-22	33.2	27.9	27.7	23.6	22.0	17.2	18.8	15.5	14.0	14.1	11.9	12.8	10.7	11.7	9.7	13.3	11.6	12.1	14.8	16.0	15.2	18.6	17.1	14.4	18.8	—
	23-26					22.2	18.7	16.3	14.1	11.2	10.4	8.9	8.3	8.7	8.2	7.6	9.6	6.9	8.4	7.9	8.3	6.6	11.1	10.9	12.9	16.7	—
	27-30									12.0	8.5	8.8	7.1	6.6	6.7	7.4	7.2	6.7	6.5	6.1	5.7	6.4	7.9	7.4	7.3	11.5	—
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—
% saying most or all	18	2.6	2.1	1.8	1.7	1.7	1.6	1.4	1.1	1.1	1.4	0.6	0.5	0.6	1.0	1.1	1.4	1.6	1.1	2.5	1.4	1.7	1.1	1.7	1.9	2.0	—
	19-22	1.1	1.3	1.0	0.8	0.8	0.5	0.3	0.4	0.8	0.1	0.2	0.3	0.1	0.1	0.3	0.8	0.2	0.7	0.4	0.4	1.0	0.9	0.8	0.7	0.3	—
	23-26					0.4	0.3	0.3	0.3	0.1	0.2	0.2	0.1	0.1	0.3	0.2	0.0	0.0	0.8	0.0	0.0	0.4	0.4	0.0	0.2	0.4	—
	27-30									0.2	0.0	0.4	0.2	0.2	0.2	0.0	0.0	0.3	0.0	0.0	0.2	0.0	0.3	0.6	0.1	0.0	—
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	—

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

<i>Q. How many of your friends would you estimate...</i>	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change	
Take quaaludes																												
% saying any friends	18	32.5	35.0	35.5	29.7	26.1	26.0	23.5	22.0	17.1	16.6	14.3	12.0	13.1	14.2	14.2	15.5	18.1	16.1	17.4	15.5	16.2	17.8	18.0	14.2	16.6	+2.4	
	19-22	38.3	36.2	35.4	30.5	24.6	19.9	20.3	16.9	12.5	10.9	10.0	10.6	9.2	10.0	7.8	11.5	10.1	9.3	10.6	11.4	13.1	14.6	13.0	10.3	8.3	-2.0	
	23-26					25.7	21.0	17.4	15.0	12.1	10.3	8.6	5.9	6.4	7.6	7.7	9.0	6.3	6.5	6.6	6.4	4.9	7.7	8.5	8.9	6.5	-2.4	
	27-30									11.8	7.9	8.2	7.0	7.1	6.5	6.6	4.5	6.9	4.9	4.1	5.1	5.0	4.9	6.6	4.3	4.4	+0.1	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																								NA	NA	—	
% saying most or all	18	3.6	3.6	2.6	2.6	1.7	1.3	1.6	1.0	1.0	1.3	0.8	0.5	0.8	1.1	1.1	1.3	1.7	1.1	2.0	1.4	1.4	1.2	1.2	1.2	1.6	+0.3	
	19-22	1.9	2.7	1.2	1.3	1.2	0.6	0.2	0.4	0.4	0.2	0.6	0.2	0.1	0.1	0.2	0.7	0.1	0.6	0.5	0.4	0.9	0.8	0.1	0.4	0.0	-0.4	
	23-26					0.6	0.3	0.7	0.2	0.2	0.4	0.2	0.1	0.2	0.6	0.2	0.2	0.0	0.8	0.0	0.2	0.3	0.3	0.1	0.2	0.1	0.0	
	27-30									0.5	0.2	0.2	0.2	0.0	0.2	0.0	0.0	0.2	0.0	0.0	0.2	0.3	0.0	0.3	0.0	0.0	0.0	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																								NA	NA	—	
Take tranquilizers																												
% saying any friends	18	29.7	29.5	29.9	26.7	26.6	25.8	24.2	23.3	19.9	18.0	14.9	13.5	14.6	15.5	16.5	15.8	18.1	17.9	19.7	16.4	19.4	18.6	21.2	17.2	18.3	+1.1	
	19-22	37.5	33.9	28.7	22.9	22.0	19.7	20.6	18.0	16.4	14.8	13.4	13.0	11.3	11.9	9.5	13.6	10.5	11.7	13.7	16.2	16.7	21.3	18.1	14.5	12.3	-2.2	
	23-26					29.3	26.3	22.3	20.8	15.5	13.1	14.8	12.1	12.5	11.0	13.4	10.4	10.7	9.6	8.5	9.8	11.2	12.4	14.9	12.9	15.1	+2.2	
	27-30									20.1	16.6	16.9	14.9	12.0	12.5	13.9	11.9	11.0	10.8	12.6	10.4	10.6	9.6	10.6	10.4	9.9	-0.5	
	35															14.3	12.2	13.1	10.8	10.7	11.4	10.8	12.2	12.5	11.4	12.7	+1.3	
	40																			13.7	14.8	15.2	15.1	15.6	15.0	13.6	-1.4	
	45																									17.3	19.8	+2.6
% saying most or all	18	1.9	1.4	1.1	1.2	1.5	1.2	1.3	1.0	0.7	1.5	0.5	0.4	0.7	0.9	0.9	1.1	1.4	0.8	2.3	1.3	2.1	1.3	1.6	1.5	1.7	+0.2	
	19-22	0.7	0.9	0.5	0.8	0.3	0.7	0.3	0.6	0.4	0.1	0.4	0.5	0.1	0.1	0.2	0.7	0.7	0.8	0.6	0.3	0.6	0.9	0.4	0.4	0.3	0.0	
	23-26					0.4	0.3	0.5	0.0	0.3	0.4	0.2	0.3	0.1	0.4	0.2	0.0	0.0	1.1	0.1	0.0	0.5	0.8	0.1	0.0	0.5	+0.5	
	27-30									0.5	0.3	0.4	0.2	0.1	0.2	0.4	0.0	0.2	0.0	0.0	0.4	0.0	0.4	0.6	0.1	0.0	-0.1	
	35															0.5	0.3	0.3	0.1	0.2	0.6	0.6	0.2	0.1	0.2	0.3	+0.1	
	40																			0.0	0.4	0.1	0.3	0.2	*	0.2	+0.1	
	45																									0.3	0.2	-0.1

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Q. How many of your friends would you estimate...

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Drink alcoholic beverages																											
% saying any friends	18	96.1	94.7	95.7	95.5	94.6	94.6	95.6	95.4	95.7	95.1	92.0	91.2	90.5	88.9	90.1	90.9	89.6	90.7	91.2	90.2	89.8	89.2	88.0	87.9	87.8	-0.2
	19-22	96.3	96.7	96.6	97.3	96.8	95.8	96.9	95.6	97.0	97.6	96.1	95.2	93.1	95.1	92.5	94.8	93.7	94.5	94.5	92.8	95.2	93.4	94.5	92.5	90.4	-2.1
	23-26					96.8	96.8	96.2	95.9	95.3	95.4	94.7	93.9	95.1	94.4	94.0	94.1	92.7	95.4	95.5	93.3	94.5	93.1	95.3	92.8	94.9	+2.1
	27-30									96.1	96.0	95.2	94.4	95.6	93.4	93.3	93.3	93.1	95.1	93.1	94.4	92.7	91.4	92.8	90.5	94.4	+3.8 s
	35														89.6	89.9	90.3	89.5	88.1	88.7	89.6	89.3	90.1	87.4	93.4	+5.9 sss	
	40																		88.4	88.9	90.7	89.6	90.5	89.2	90.5	+1.3	
	45																							87.9	90.3	+2.4	
% saying most or all	18	68.9	67.7	69.7	69.0	66.6	66.0	68.0	71.8	68.1	67.1	60.5	58.6	56.9	57.0	59.6	56.4	56.4	60.9	61.0	58.2	57.2	59.2	53.7	53.1	53.9	+0.7
	19-22	76.6	77.6	75.2	75.1	74.9	71.9	74.2	71.3	73.4	74.1	70.0	71.4	67.4	66.5	68.7	63.9	67.0	63.8	69.4	67.8	70.1	65.4	68.8	63.9	66.4	+2.5
	23-26					73.2	74.4	69.5	74.9	68.9	69.8	67.1	69.3	68.8	68.7	70.7	67.0	68.9	66.6	67.4	63.6	70.8	65.7	73.4	66.0	71.3	+5.3
	27-30									66.7	67.8	62.0	62.7	63.3	61.3	63.2	62.6	64.1	66.6	62.9	64.4	64.8	64.9	66.3	61.5	69.0	+7.5 s
	35														43.8	45.1	49.5	46.6	47.1	46.0	49.1	48.4	52.9	51.6	53.7	+2.1	
	40																		37.7	41.4	42.5	44.7	44.8	47.2	43.3	-3.9	
	45																							38.9	41.7	+2.8	
Get drunk at least once a week																											
% saying any friends	18	83.1	81.8	83.1	83.9	81.5	82.5	84.7	85.6	84.4	82.8	79.2	79.8	79.9	79.2	81.4	78.9	78.5	82.4	81.1	81.5	79.5	79.6	78.3	77.3	79.0	+1.7
	19-22	80.9	79.9	80.0	80.4	79.8	76.7	82.0	81.1	80.6	80.4	80.1	80.8	76.5	81.1	79.6	83.2	80.9	79.2	82.3	82.8	82.2	81.9	81.5	81.5	80.5	-1.0
	23-26					73.1	72.7	73.5	73.7	72.1	73.1	72.2	74.0	73.1	74.3	72.1	73.1	74.5	71.9	74.1	71.0	76.5	74.7	81.0	76.4	75.8	-0.6
	27-30									66.3	61.8	65.4	65.2	65.5	64.5	62.7	67.1	66.7	65.4	65.5	65.9	64.3	64.7	68.9	66.5	73.8	+7.3 s
	35														44.3	43.2	44.9	42.9	46.1	44.5	46.9	47.6	48.3	47.9	52.0	+4.1	
	40																		41.6	40.6	42.2	41.3	42.6	42.9	43.2	+0.3	
	45																							41.6	42.2	+0.6	
% saying most or all	18	30.1	29.4	29.9	31.0	29.6	29.9	31.8	31.3	29.6	31.1	27.5	29.7	28.6	27.6	28.4	27.4	29.0	30.9	31.7	30.1	32.4	32.7	28.3	27.1	27.6	+0.6
	19-22	21.9	23.3	22.0	20.2	22.7	21.7	20.8	21.3	24.0	22.6	23.6	24.9	22.6	28.8	26.3	28.2	26.0	26.6	29.8	29.3	28.1	30.2	31.0	29.6	29.0	-0.6
	23-26					11.4	11.6	12.5	11.9	12.8	12.0	13.9	11.6	14.6	13.2	15.2	15.2	14.0	17.0	16.0	16.8	17.4	19.1	19.2	18.3	24.0	+5.8
	27-30									5.2	6.3	6.7	6.6	5.9	6.7	6.4	7.9	8.6	7.7	9.3	12.1	9.8	11.7	8.9	13.0	9.4	-3.6
	35														3.6	3.6	5.4	3.2	4.4	4.9	4.6	4.8	4.5	5.2	5.3	+0.1	
	40																		2.8	3.0	2.5	2.9	3.8	3.9	3.0	-0.9	
	45																							3.6	2.7	-0.9	

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Q. How many of your friends would you estimate...

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change	
Smoke cigarettes																												
% saying any friends	18	90.6	88.5	88.3	87.0	86.0	87.0	87.8	88.3	87.7	86.5	84.9	85.7	84.4	84.8	88.1	87.9	88.3	89.9	89.5	89.3	87.2	86.8	85.4	83.3	83.7	+0.5	
	19-22	94.4	94.3	93.4	93.1	91.9	91.6	91.1	90.3	89.3	90.0	86.1	86.1	86.7	86.7	86.1	88.8	89.2	91.3	92.6	91.0	90.9	90.9	89.7	86.5	89.7	+3.2	
	23-26					93.9	95.0	91.6	92.1	89.8	90.1	88.7	89.6	85.6	88.3	86.4	86.8	85.3	85.4	88.7	84.1	86.5	86.7	86.4	86.5	87.0	+0.5	
	27-30									92.6	89.8	90.7	90.4	88.0	85.8	84.8	84.9	85.4	84.1	81.1	86.3	85.1	84.9	87.0	82.8	83.5	+0.7	
	35															72.7	71.7	71.7	72.4	71.8	69.9	70.8	69.2	66.6	67.0	67.7	+0.7	
	40																			70.2	70.0	67.8	64.3	65.5	65.1	62.4	-2.7	
	45																							66.1	67.0	+1.0		
	% saying most or all	18	23.3	22.4	24.1	22.4	19.2	22.8	21.5	21.0	20.2	23.1	21.4	21.8	21.4	25.0	25.3	27.5	30.4	34.4	33.9	31.1	28.2	25.0	23.0	19.6	20.6	+1.0
		19-22	31.8	27.6	25.6	25.2	25.6	22.7	21.9	22.5	19.3	19.9	19.2	20.2	20.3	22.2	21.7	28.4	24.0	25.1	28.8	26.8	29.4	27.0	25.7	20.2	20.7	+0.5
		23-26					25.6	22.7	19.7	18.5	16.5	20.5	16.9	18.1	16.0	15.5	16.6	13.9	17.6	17.0	16.8	17.5	17.0	15.5	15.1	18.3	19.8	+1.5
		27-30									15.8	14.2	11.6	12.9	11.9	14.3	10.9	12.3	10.4	12.1	12.3	13.4	11.7	10.2	12.9	12.2	9.2	-2.9
		35															7.9	7.2	9.3	7.2	8.0	9.0	6.7	8.8	6.6	6.3	6.9	+0.5
		40																			8.1	7.4	6.8	5.7	5.8	5.9	6.0	+0.1
	45																								5.7	5.9	+0.2	
	Take steroids																											
% saying any friends	18										25.9	24.7	21.5	19.0	18.1	19.5	17.9	18.9	18.3	20.0	19.8	21.7	21.6	21.1	22.8	+1.7		
	19-22									23.4	21.5	22.2	19.7	20.7	16.8	16.6	16.1	16.8	20.0	20.6	18.9	20.0	19.3	17.1	21.4	+4.3		
	23-26									15.3	15.0	12.3	14.5	11.1	10.5	12.4	7.3	13.0	9.2	15.0	12.2	13.6	14.3	12.9	12.4	-0.5		
	27-30									9.9	10.5	7.5	8.0	8.0	8.0	8.0	10.2	9.1	7.0	11.2	9.3	10.7	6.4	11.6	10.1	-1.5		
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																							NA	NA	NA	—	
	% saying most or all	18										1.8	1.0	1.7	0.9	1.2	1.3	0.8	1.7	1.4	0.9	1.9	1.2	1.5	1.5	2.6	+1.1 s	
19-22										0.2	0.6	0.0	0.1	0.4	0.2	0.1	0.0	0.1	0.3	0.1	0.3	0.7	0.7	0.4	0.0	-0.4		
23-26										0.4	0.0	0.0	0.2	0.1	0.1	0.0	0.0	0.5	0.0	0.1	0.3	0.2	0.1	0.0	0.1	+0.1		
27-30										0.5	0.0	0.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0		
35																NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
40																				NA	NA	NA	NA	NA	NA	NA	—	
45																							NA	NA	NA	—		

(Table continued on next page)

TABLE 7-2 (cont.)
Trends in Proportions of Friends Using Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45

	Age																									
	Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
<i>Approx. Weighted N=</i>	18	2987	3307	3303	3095	2945	2971	2798	2948	2961	2587	2361	2339	2373	2410	2337	2379	2156	2292	2313	2060	1838	1923	1968	2233	2271
	19-22	576	592	564	579	543	554	579	572	562	579	556	526	510	468	435	470	469	467	437	426	402	402	375	388	443
	23-26					527	534	546	528	528	506	510	507	516	495	449	456	416	419	394	414	387	403	358	362	411
	27-30									516	507	499	476	478	461	419	450	464	454	428	424	363	359	348	369	396
	35															1200	1187	1187	1209	1067	1071	1033	1005	918	968	985
	40																			1098	1156	1144	1119	1083	945	1004
	45																								976	1074

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

'NA' indicates data not available.

'*' indicates a prevalence rate of less than 0.05% but greater than true zero.

^aThese estimates were derived from responses to the questions listed above. For the young adult sample, "any illicit drug" includes all of the drugs listed except cigarettes and alcohol. For the 35-, 40-, and 45-year-olds, "any illicit drug" includes marijuana, tranquilizers, crack, cocaine powder, and "other illicit drugs."

^bIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^cIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

TABLE 7-3
Trends in Exposure to Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Any illicit drug^a																											
% saying any exposure																											
	18	84.3	82.7	81.4	79.4	77.9	77.7	75.5	73.9	71.3	68.6	67.6	64.2	61.3	66.1	70.8	75.3	78.0	78.8	77.2	77.9	76.0	76.5	76.5	73.6	74.3	+0.8
	19-22	80.6	81.0	81.5	76.5	76.3	77.4	74.6	72.7	69.5	61.5	60.8	58.9	58.6	58.4	60.7	66.4	67.2	65.3	69.1	65.8	64.7	69.7	65.7	68.0	67.6	-0.5
	23-26					68.9	70.2	68.0	62.4	62.7	58.3	54.6	52.1	48.2	49.9	47.1	54.2	50.3	55.4	50.6	50.5	55.1	56.4	56.5	57.0	53.5	-3.5
	27-30									52.4	50.2	47.0	39.6	41.7	38.9	45.6	42.4	44.9	41.6	37.5	41.1	40.8	42.2	47.0	46.7	43.3	-3.4
% saying often exposed																											
	18	36.3	36.1	31.4	29.8	28.3	27.2	26.3	23.3	20.8	22.0	20.7	18.2	18.0	24.0	29.3	32.3	33.8	34.7	33.2	35.6	32.6	33.6	32.6	31.8	30.3	-1.5
	19-22	34.6	34.0	32.1	24.4	24.4	23.7	21.1	18.9	19.9	16.2	16.4	17.6	21.4	16.1	18.1	23.7	20.4	25.3	24.2	24.0	21.3	26.1	25.2	26.5	26.8	+0.2
	23-26					20.7	23.3	18.5	17.4	18.2	13.8	13.7	13.3	12.2	11.1	11.1	12.5	12.8	14.3	14.2	15.0	15.9	16.4	15.9	17.8	15.1	-2.7
	27-30									13.7	12.0	10.8	8.2	10.5	9.0	12.5	8.5	10.1	10.3	8.5	9.6	9.4	10.4	13.8	13.9	10.3	-3.6
Any illicit drug^a other than marijuana																											
% saying any exposure																											
	18	58.5	62.6	62.5	59.4	59.8	59.3	55.3	51.7	47.8	47.1	45.4	40.0	41.6	42.6	45.3	47.2	49.7	47.9	47.3	46.5	47.2	49.9	49.3	46.3	48.3	+2.0
	19-22	56.9	58.4	61.6	54.9	57.1	53.3	53.4	48.5	46.4	36.5	39.4	33.8	37.1	29.4	33.9	36.8	36.5	39.4	40.0	36.4	38.1	39.2	38.0	40.2	40.9	+0.6
	23-26					51.5	51.9	51.5	43.6	42.9	36.8	34.0	30.0	27.3	27.8	24.9	26.8	23.2	25.6	27.1	28.0	31.0	31.4	31.5	32.2	32.6	+0.4
	27-30									35.8	33.7	31.5	25.8	26.6	24.2	25.8	21.1	21.8	21.4	15.4	19.5	17.2	22.2	23.1	26.1	23.2	-2.9
% saying often exposed																											
	18	14.1	17.1	16.6	14.2	14.6	12.9	12.1	10.2	9.6	10.7	9.2	7.9	7.5	9.6	9.4	11.1	12.1	11.7	9.9	11.7	10.5	11.9	12.6	10.8	11.4	+0.7
	19-22	11.8	15.6	13.5	11.1	10.7	10.2	8.2	8.1	7.5	6.7	4.5	4.4	5.5	4.1	5.1	7.7	3.9	7.6	7.0	4.8	6.4	7.8	8.6	5.2	7.9	+2.7
	23-26					9.0	10.4	9.3	8.5	6.7	5.0	5.1	3.5	2.6	3.0	2.2	3.5	3.4	3.1	3.1	4.3	3.5	3.4	5.0	5.4	5.4	0.0
	27-30									6.0	4.7	4.1	3.2	3.7	2.4	3.4	2.9	3.4	3.2	1.0	2.5	1.6	3.7	4.7	4.9	2.4	-2.5
Marijuana																											
% saying any exposure																											
	18	82.0	80.2	77.9	76.2	74.4	73.5	72.0	70.4	67.0	64.8	63.4	59.6	56.8	61.0	67.2	72.7	75.6	76.8	75.5	75.8	73.8	74.9	74.2	71.4	72.2	+0.8
	19-22	79.8	79.8	78.7	72.7	74.1	75.5	72.4	70.5	66.3	59.3	57.5	55.0	56.4	55.4	56.8	64.0	64.8	63.4	67.1	63.5	63.9	68.0	64.6	64.8	65.1	+0.3
	23-26					65.3	66.0	64.1	59.0	57.6	55.0	50.6	47.9	44.6	45.9	44.4	51.0	47.8	53.1	48.8	48.1	51.8	54.2	53.5	54.4	50.6	-3.8
	27-30									49.1	47.4	42.1	36.0	38.2	35.3	41.9	38.3	41.8	39.1	35.7	38.7	38.8	37.0	44.6	44.1	40.4	-3.7
% saying often exposed																											
	18	33.8	33.1	28.0	26.1	24.8	24.2	24.0	20.6	17.9	19.5	17.8	16.0	15.6	20.9	27.6	30.7	31.8	32.9	31.4	34.4	30.3	30.8	30.7	30.4	28.0	-2.4
	19-22	32.6	30.5	30.3	21.1	21.9	20.3	18.6	16.4	18.3	14.2	14.7	15.9	19.9	14.7	17.0	22.1	20.3	23.7	22.8	23.0	20.4	24.5	24.8	24.2	24.5	+0.3
	23-26					17.5	20.6	14.6	14.8	15.6	11.6	11.2	11.6	10.9	10.4	10.4	11.1	11.5	12.9	13.6	13.2	15.2	15.6	14.9	16.2	13.7	-2.4
	27-30									10.9	9.8	8.5	6.7	8.9	7.6	10.7	7.4	9.1	8.9	8.1	8.8	8.6	8.4	11.7	11.7	9.6	-2.1

(Table continued on next page)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
LSD																											
% saying any exposure	18	17.2	17.4	16.1	13.8	12.5	13.2	13.1	12.9	13.4	15.0	14.9	15.7	17.8	21.0	24.2	26.1	27.6	25.9	23.1	23.6	22.0	21.6	17.2	14.2	12.4	-1.8
	19-22	17.4	15.8	16.0	13.5	12.8	12.7	10.8	10.9	12.0	12.1	13.1	19.3	13.4	16.5	18.6	20.7	22.3	21.0	20.1	15.9	15.2	13.6	10.0	8.5	-1.5	
	23-26					8.3	9.3	8.8	7.3	6.3	6.7	8.4	8.6	8.8	7.8	8.4	9.9	8.6	7.6	9.8	9.4	9.8	11.1	9.3	5.5	4.4	-1.2
	27-30									3.6	3.2	3.3	3.6	3.9	4.9	5.3	5.5	4.3	3.9	3.2	3.7	3.2	4.3	4.8	3.0	4.7	+1.8
% saying often exposed	18	1.4	2.0	1.9	1.4	1.5	1.3	1.6	1.8	1.6	2.2	2.6	2.9	3.0	3.9	4.2	6.1	4.7	5.1	3.2	4.1	3.3	2.8	2.6	1.8	1.6	-0.3
	19-22	1.4	1.5	1.4	0.6	0.8	0.7	0.5	1.2	0.6	1.1	1.2	1.0	2.0	1.1	0.4	3.6	1.4	1.8	2.0	1.7	1.4	2.4	0.9	0.2	0.1	-0.1
	23-26					0.3	0.4	0.4	0.7	0.6	0.3	0.5	0.2	0.8	0.3	0.5	0.5	0.4	0.2	0.1	0.3	0.2	0.0	0.3	0.3	0.0	-0.3
	27-30									0.3	0.2	0.5	0.2	0.2	0.5	0.5	0.2	0.2	0.0	0.0	0.1	0.0	0.0	0.0	0.3	0.3	0.0
Other psychedelics/ hallucinogens^b																											
% saying any exposure	18	20.4	17.6	16.8	13.1	12.7	12.5	11.8	10.0	9.0	8.8	9.4	9.4	9.7	12.1	14.0	15.8	16.6	17.8	15.9	17.7	16.3	28.1	26.4	25.8	24.8	-1.0
	19-22	18.3	16.3	16.3	12.5	10.5	11.0	9.2	9.1	7.7	8.4	8.3	8.9	10.6	6.7	8.3	12.8	13.1	15.0	15.0	12.4	11.8	22.8	23.4	18.9	18.7	-0.2
	23-26					8.4	8.9	9.1	6.0	5.1	4.8	5.7	5.5	5.1	5.7	5.2	5.5	6.9	5.6	8.7	5.8	8.9	14.8	14.7	11.9	10.1	-1.7
	27-30									5.0	3.4	3.4	3.4	2.1	3.7	3.4	4.2	3.2	2.9	2.6	3.0	3.0	6.4	7.7	6.3	7.9	+1.7
% saying often exposed	18	2.2	2.0	2.6	1.1	1.7	1.4	1.5	1.2	1.1	1.3	1.2	1.3	1.1	1.9	2.3	2.5	2.7	2.8	1.7	2.7	2.1	3.6	4.5	3.2	3.2	0.0
	19-22	1.1	0.9	0.9	0.7	0.8	0.8	0.2	0.8	0.3	0.4	0.4	0.5	0.7	0.4	0.2	1.6	0.7	0.7	0.5	0.6	0.8	2.6	2.4	0.4	0.7	+0.3
	23-26					0.1	0.3	0.5	0.6	0.8	0.1	0.4	0.0	0.2	0.4	0.3	0.3	0.2	0.0	0.0	0.4	0.2	0.4	0.0	0.0	0.0	0.0
	27-30									0.2	0.4	0.5	0.3	0.1	0.5	0.2	0.3	0.2	0.5	0.0	0.1	0.0	0.4	0.0	0.0	0.3	+0.3
Cocaine																											
% saying any exposure	18	37.7	36.3	34.9	33.3	35.6	38.3	37.4	34.9	30.2	30.2	27.7	21.3	19.8	19.2	18.8	21.6	25.0	25.6	26.6	25.8	24.2	24.5	24.9	24.8	24.4	-0.5
	19-22	37.6	42.3	43.6	36.6	38.9	39.4	41.5	37.0	36.2	26.6	24.0	18.5	19.8	13.5	14.7	14.1	19.3	18.8	21.6	18.5	19.1	20.6	22.5	18.4	23.6	+5.3
	23-26					38.5	40.6	42.0	34.5	35.9	28.0	24.0	19.9	16.7	14.6	14.3	14.1	12.5	14.0	16.0	18.2	16.4	16.9	18.3	17.4	18.7	+1.3
	27-30									28.9	28.3	24.2	18.6	19.4	16.6	14.3	11.4	12.1	11.4	8.6	11.6	10.2	11.6	12.2	12.6	13.0	+0.4
% saying often exposed	18	5.9	6.6	6.6	5.2	6.7	7.1	7.8	5.9	5.1	5.4	4.7	3.4	2.7	2.9	2.5	3.2	4.0	4.2	3.7	4.6	4.6	4.5	5.3	5.0	4.7	-0.3
	19-22	5.8	7.6	6.5	4.3	6.5	7.0	5.4	5.2	4.8	4.3	2.2	1.6	1.7	1.7	1.8	1.7	1.2	2.4	3.2	1.4	3.8	3.0	4.1	1.6	2.6	+0.9
	23-26					5.3	8.5	7.0	6.0	5.4	3.5	2.5	1.7	1.4	1.7	1.0	1.7	1.3	1.8	1.5	2.2	1.8	1.0	2.5	1.9	2.9	+1.0
	27-30									4.4	3.9	2.9	2.2	2.0	1.2	1.5	1.4	1.9	1.6	0.8	1.5	0.3	1.6	2.4	1.7	0.7	-1.0

(Table continued on next page)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Heroin																											
% saying any exposure	18	7.4	6.6	7.1	5.1	6.0	5.5	6.0	5.8	5.7	6.5	5.4	5.1	5.4	5.7	7.3	7.9	8.6	9.1	8.7	8.1	9.1	8.7	8.3	7.3	6.6	-0.7
	19-22	4.4	3.3	4.1	2.9	3.1	4.8	2.9	2.9	2.9	2.9	2.5	3.0	2.7	2.0	3.7	3.8	3.6	3.7	6.4	3.2	5.2	3.2	5.3	3.4	3.0	-0.3
	23-26					2.3	3.3	3.2	2.9	1.7	2.3	2.3	1.8	1.7	1.5	1.9	2.8	2.9	2.7	3.1	2.9	2.6	2.4	3.8	2.0	3.1	+1.1
	27-30									2.1	1.4	1.5	0.9	1.0	2.0	2.0	1.7	1.5	1.3	1.4	1.9	1.9	2.3	2.7	1.3	3.2	+1.8
% saying often exposed	18	0.4	0.6	1.0	0.7	1.1	0.5	1.0	0.9	0.8	1.0	0.5	0.9	0.7	1.1	0.7	1.2	1.6	1.2	0.9	1.3	1.5	0.7	1.3	1.2	1.2	0.0
	19-22	0.2	0.3	0.3	0.1	0.2	0.5	0.2	0.1	0.2	0.1	0.2	0.4	0.6	0.4	0.6	1.2	0.2	0.4	0.7	0.8	0.7	0.8	0.6	0.2	0.0	-0.2
	23-26					0.0	0.7	0.3	0.6	0.4	0.3	0.6	0.3	0.0	0.0	0.0	0.2	0.2	0.3	0.5	1.0	0.0	0.0	0.8	0.5	0.5	0.0
	27-30									0.3	0.3	0.5	0.2	0.2	0.9	0.3	0.6	0.6	0.0	0.0	0.2	0.0	0.0	0.7	0.3	0.0	-0.3
Other narcotics																											
% saying any exposure	18	19.6	17.5	18.5	17.3	18.0	18.4	15.6	14.4	14.8	13.8	14.2	11.3	11.1	12.4	14.9	15.5	18.5	20.4	20.7	21.9	21.1	21.6	22.5	21.8	20.3	-1.5
	19-22	14.4	14.4	15.2	10.9	12.4	13.7	9.8	12.2	11.2	9.0	9.4	9.2	8.5	6.8	10.1	12.1	11.5	14.5	15.3	13.9	17.0	18.3	18.7	13.6	14.5	+1.0
	23-26					9.0	12.3	9.2	9.7	7.4	8.0	5.9	8.3	7.0	4.6	6.9	7.8	7.4	6.5	8.1	9.4	10.9	12.2	12.0	12.6	12.6	0.0
	27-30									6.5	6.5	5.8	5.5	3.7	5.6	5.9	5.7	4.7	4.9	3.6	5.2	6.5	9.0	7.9	9.5	8.8	-0.7
% saying often exposed	18	1.7	1.7	2.4	2.2	2.0	1.8	2.1	1.7	1.7	1.7	1.6	1.4	1.3	1.7	1.7	2.1	3.4	2.5	2.8	3.9	2.9	3.0	3.8	3.0	3.3	+0.3
	19-22	0.7	0.5	0.5	0.9	0.7	1.0	0.5	0.4	0.9	0.3	0.2	1.0	0.9	0.6	0.8	1.4	0.7	1.5	1.7	1.1	2.4	1.6	3.0	1.2	0.8	-0.3
	23-26					0.4	0.5	1.3	0.8	0.8	0.5	1.6	0.7	0.1	0.3	0.1	0.1	0.3	0.7	0.5	1.1	0.7	1.0	0.9	1.6	1.4	-0.3
	27-30									0.7	0.5	1.0	0.3	0.8	1.2	0.8	0.8	0.7	0.5	0.0	0.2	1.1	1.0	0.7	1.2	0.1	-1.1
Amphetamines																											
% saying any exposure	18	40.8	49.5	50.2	46.1	45.0	41.0	36.5	31.7	27.9	27.4	28.3	23.6	24.5	24.7	28.2	28.1	31.5	31.0	29.9	30.1	29.5	31.5	30.6	27.4	27.2	-0.2
	19-22	42.3	48.6	48.4	39.7	41.3	35.9	31.3	26.7	21.2	18.5	19.5	17.4	21.3	15.1	20.3	21.0	22.3	24.6	24.8	21.2	24.8	23.3	25.5	21.6	23.7	+2.0
	23-26					32.3	30.5	29.1	20.9	18.8	14.0	16.8	14.6	11.8	13.2	11.2	13.0	11.1	11.7	14.6	12.3	18.5	18.2	17.9	15.4	18.8	+3.4
	27-30									15.6	14.3	13.5	10.7	11.4	11.3	11.0	10.6	7.6	9.1	6.6	10.4	7.4	11.1	11.5	12.2	11.4	-0.8
% saying often exposed	18	8.3	12.1	12.3	10.1	9.0	6.5	5.8	4.5	4.1	4.7	4.1	3.1	3.0	3.9	4.1	4.5	5.6	5.2	4.7	6.3	4.4	6.0	6.4	4.9	5.3	+0.4
	19-22	7.4	9.9	7.7	6.9	5.4	4.4	3.1	3.3	2.2	1.5	1.1	1.9	2.6	1.5	3.3	5.0	1.3	4.1	2.9	2.2	2.4	2.6	5.6	1.7	4.1	+2.4
	23-26					3.9	3.2	2.2	3.3	1.9	0.7	2.0	1.3	0.2	0.8	0.9	1.6	1.3	1.4	2.2	1.7	1.4	2.2	0.7	1.3	1.7	+0.4
	27-30									2.0	2.0	1.2	0.8	0.8	1.3	0.7	1.6	1.8	1.0	0.2	1.1	0.4	0.6	1.5	1.0	1.2	+0.2

(Table continued on next page)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
High School Seniors (Age 18) and Young Adults in Modal Age Groups of 19-22, 23-26, and 27-30
(Entries are percentages)

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change	
Sedatives/Barbiturates^c																												
% saying any exposure	18	25.2	25.9	25.7	22.5	21.2	18.9	15.8	13.1	12.4	11.8	13.3	10.0	10.2	11.9	13.0	14.5	15.5	16.1	16.1	17.1	16.3	17.1	17.7	14.8	21.5	—	
	19-22	25.6	23.1	21.8	18.3	15.7	14.7	12.8	12.0	8.2	8.3	6.5	7.9	7.3	7.2	7.4	10.1	8.8	11.7	13.4	11.6	13.1	13.1	16.0	11.9	17.2	—	
	23-26					16.1	13.1	11.0	7.1	7.1	6.6	6.9	5.9	6.5	3.8	4.2	5.7	6.6	4.9	8.5	7.1	9.3	9.0	9.8	7.9	15.9	—	
	27-30									8.0	6.8	5.9	5.4	5.2	5.7	4.5	5.2	3.5	3.8	2.7	4.1	2.9	5.3	6.0	6.1	9.2	—	
% saying often exposed	18	3.4	4.0	4.3	3.0	2.7	1.7	2.1	1.5	1.4	1.7	1.7	1.2	1.1	1.6	1.7	2.0	2.9	2.5	2.7	3.8	2.7	2.7	4.6	2.8	4.1	—	
	19-22	2.5	2.8	1.1	1.4	0.7	1.3	0.5	0.7	0.7	0.3	0.7	0.4	0.7	0.7	1.3	1.3	0.4	0.9	1.4	0.9	1.6	1.2	1.8	0.8	1.7	—	
	23-26					0.7	0.9	1.7	0.8	0.6	0.3	1.1	0.3	0.3	0.0	0.2	0.3	0.8	0.5	0.9	0.7	0.2	0.3	0.4	0.7	—		
	27-30									0.7	0.4	0.6	0.2	0.4	1.2	0.2	0.6	0.5	0.2	0.0	0.6	0.2	0.9	0.4	0.6	0.4	—	
Tranquilizers^b																												
% saying any exposure	18	29.1	29.0	26.6	23.5	23.1	23.4	19.6	18.4	18.2	15.1	16.3	14.2	12.7	13.8	16.5	15.7	17.9	18.9	17.3	18.2	17.7	23.8	22.7	21.0	22.1	+1.1	
	19-22	29.6	26.9	28.5	19.5	21.2	19.5	16.4	18.5	13.8	12.0	12.7	12.6	11.0	10.0	12.0	11.8	10.7	15.6	16.9	14.3	18.5	21.3	23.6	20.0	21.9	+1.9	
	23-26					23.1	21.0	16.9	15.9	13.4	12.9	12.0	10.4	9.7	10.9	9.8	10.3	10.1	9.4	10.9	10.8	12.3	16.4	20.1	18.7	19.9	+1.2	
	27-30									15.0	11.6	11.1	9.7	10.3	10.4	9.0	11.2	9.6	9.6	6.1	8.8	7.6	12.6	13.6	15.3	14.6	-0.7	
% saying often exposed	18	3.2	4.2	3.5	2.9	2.9	2.2	2.5	2.6	2.2	2.1	1.9	1.4	1.9	1.7	1.8	2.3	3.5	3.2	2.8	3.7	3.5	4.9	5.8	4.2	4.1	-0.1	
	19-22	3.2	2.6	1.8	2.1	1.5	1.7	0.9	1.1	1.8	1.0	1.1	1.1	1.5	1.1	1.3	1.5	0.5	1.3	1.6	1.5	1.7	3.1	3.6	2.3	2.7	+0.4	
	23-26					2.0	1.6	2.6	1.8	1.2	0.8	0.5	1.0	0.6	0.7	0.1	1.1	1.5	0.7	1.1	1.5	1.7	1.3	2.1	1.6	2.0	+0.4	
	27-30									1.4	0.3	1.7	0.8	1.3	1.3	1.0	1.1	0.8	1.2	0.2	0.9	0.4	1.6	1.6	1.9	0.8	-1.1	
Alcoholic beverages																												
% saying any exposure	18	94.7	94.0	94.0	94.0	94.0	94.0	94.1	93.9	93.1	92.3	93.6	91.7	90.6	91.8	90.0	91.2	91.5	91.4	92.2	91.8	90.7	90.8	89.5	88.3	87.6	-0.6	
	19-22	94.3	93.8	94.5	93.4	94.2	92.7	93.6	94.4	92.5	91.8	92.4	94.0	93.3	92.9	93.7	93.1	93.7	93.1	91.8	91.0	93.3	94.3	93.7	93.6	92.5	-1.1	
	23-26					90.3	92.7	91.4	90.6	91.1	92.9	91.3	91.0	91.4	90.3	89.5	91.9	89.6	89.2	86.4	88.4	88.7	89.8	91.2	89.0	90.0	85.3	-4.8
	27-30									87.1	88.4	86.2	87.7	87.3	86.6	86.2	89.3	89.2	86.4	88.4	88.7	89.8	91.2	89.0	90.0	85.3	-4.8	
% saying often exposed	18	60.2	61.0	59.3	60.2	58.7	59.5	58.0	58.7	56.4	55.5	56.1	54.5	53.1	51.9	54.0	54.0	54.5	53.9	54.5	53.5	50.2	52.7	50.8	49.0	48.2	-0.8	
	19-22	59.6	61.2	62.5	56.6	59.3	61.8	59.9	61.4	55.4	53.8	56.0	53.9	56.1	56.8	57.0	56.3	52.3	54.2	57.9	54.7	54.3	53.4	54.9	55.7	54.3	-1.4	
	23-26					52.1	54.8	51.4	53.0	48.1	50.9	49.7	48.4	45.4	45.4	43.3	47.5	44.8	49.8	44.6	45.7	49.6	48.8	46.3	50.5	48.3	-2.1	
	27-30									39.9	39.5	38.7	38.0	39.9	38.1	39.3	38.0	34.7	37.1	36.6	38.3	34.4	40.0	39.6	40.6	36.8	-3.9	
<i>Approx. Weighted N=</i>	18	3259	3608	3645	3334	3238	3252	3078	3296	3300	2795	2556	2525	2630	2730	2581	2608	2407	2595	2541	2312	2153	2147	2162	2454	2456		
	19-22	582	574	601	569	578	549	591	582	556	567	567	532	528	489	460	464	485	471	445	450	415	412	403	396	432		
	23-26					533	532	557	529	531	514	523	494	532	513	471	467	447	424	400	398	389	406	345	385	404		
	27-30									522	507	506	478	502	457	425	452	432	455	449	430	395	369	359	347	370		

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change and prevalence estimates for the two most recent years is due to rounding.

^aThese estimates were derived from responses to the questions listed above. For the young adult sample, "any illicit drug" includes all of the drugs listed except cigarettes and alcohol.

^bIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. For tranquilizers, Xanax was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^cIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

TABLE 7-4
Trends in Reported Availability of Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Percentage saying "fairly easy" or "very easy" to get^a

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a																								'03-'04 change		
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		2004	
Marijuana	18	89.0	89.2	88.5	86.2	84.6	85.5	85.2	84.8	85.0	84.3	84.4	83.3	82.7	83.0	85.5	88.5	88.7	89.6	90.4	88.9	88.5	88.5	87.2	87.1	85.8	-1.3	
	19-22	95.6	91.1	92.4	89.7	88.3	89.5	87.2	85.9	87.1	87.1	86.2	86.0	87.8	85.6	87.2	87.9	89.3	90.6	89.9	87.4	89.6	91.7	88.1	87.7	87.3	-0.4	
	23-26					92.5	88.8	88.8	90.3	86.9	88.7	83.3	82.5	83.8	84.6	87.1	86.2	85.3	84.4	87.5	85.9	88.4	87.0	89.1	87.2	88.8	+1.6	
	27-30									89.3	86.0	83.1	83.8	80.7	82.8	80.3	83.3	82.6	84.5	82.1	83.0	81.5	84.8	83.6	81.8	86.0	+4.1	
	35															75.7	75.6	73.0	77.1	76.0	74.9	77.1	75.3	76.5	75.1	75.6	+0.5	
	40																			73.4	71.7	73.1	70.4	72.1	72.3	68.9	-3.4	
	45																								68.5	69.9	+1.4	
Amyl & butyl nitrites	18	NA	NA	NA	NA	NA	NA	NA	23.9	25.9	26.8	24.4	22.7	25.9	25.9	26.7	26.0	23.9	23.8	25.1	21.4	23.3	22.5	22.3	19.7	20.0	+0.4	
	19-22	NA	NA	NA	NA	NA	NA	NA	22.8	26.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	23-26					NA	NA	NA	23.1	28.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	27-30									26.7	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	35																NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																				NA	NA	NA	NA	NA	NA	—	
LSD	18	35.3	35.0	34.2	30.9	30.6	30.5	28.5	31.4	33.3	38.3	40.7	39.5	44.5	49.2	50.8	53.8	51.3	50.7	48.8	44.7	46.9	44.7	39.6	33.6	33.1	-0.6	
	19-22	39.6	38.4	35.1	31.8	32.7	29.6	30.5	29.9	33.9	36.4	36.6	37.8	42.5	44.9	43.7	50.5	50.8	47.7	51.1	43.8	47.1	42.5	37.9	34.1	30.3	-3.8	
	23-26					32.7	29.1	30.0	27.5	32.7	32.6	30.2	32.8	33.5	33.4	40.1	41.0	43.6	39.2	40.4	41.2	40.4	38.3	37.2	34.1	38.5	+4.4	
	27-30									29.4	29.9	32.3	27.0	30.9	30.5	27.2	35.6	33.6	35.2	32.9	35.7	35.6	38.3	32.3	33.5	30.0	-3.5	
	35																33.8	32.4	28.4	32.9	31.2	27.7	32.2	28.7	29.1	29.8	25.6	-4.2
	40																				31.1	31.0	28.5	25.7	27.4	25.0	24.4	-0.6
	45																								24.2	27.0	+2.8	
Some psychedelic/hallucinogen other than LSD^b	18	35.0	32.7	30.6	26.6	26.6	26.1	24.9	25.0	26.2	28.2	28.3	28.0	29.9	33.5	33.8	35.8	33.9	33.9	35.1	29.5	34.5	48.5	47.7	47.2	49.4	+2.1	
	19-22	42.1	37.7	33.5	31.0	28.9	28.7	26.3	27.5	28.7	28.1	28.9	26.6	28.3	29.5	28.6	31.5	31.5	33.4	34.1	31.1	33.4	45.9	48.8	45.1	46.9	+1.8	
	23-26					31.8	29.6	26.4	25.6	29.6	28.7	27.0	25.7	27.7	25.3	28.3	29.2	32.6	31.0	32.4	31.5	28.5	38.3	39.7	39.2	44.4	+5.1	
	27-30									28.6	29.6	30.8	24.9	24.8	25.4	24.7	29.3	25.9	28.0	25.2	30.3	25.0	38.6	33.3	35.6	31.2	-4.4	
	35																NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																				NA	NA	NA	NA	NA	NA	—	
	45																								NA	NA	—	

(Table continued on next page)

TABLE 7-4 (cont.)
Trends in Reported Availability of Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Percentage saying "fairly easy" or "very easy" to get^a

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

	Age Group	Year																							'03-'04 change			
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002		2003	2004	
PCP	18	NA	NA	NA	NA	NA	NA	NA	22.8	24.9	28.9	27.7	27.6	31.7	31.7	31.4	31.0	30.5	30.0	30.7	26.7	28.8	27.2	25.8	21.9	24.2	+2.3	
	19-22	NA	NA	NA	NA	NA	NA	NA	21.7	24.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	23-26					NA	NA	NA	21.2	27.6	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	27-30									24.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	NA	—
MDMA (ecstasy)	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	21.7	22.0	22.1	24.2	28.1	31.2	34.2	36.9	38.8	38.2	40.1	51.4	61.5	59.1	57.5	47.9	-9.6 sss	
	19-22	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	26.6	24.9	27.1	23.9	27.0	29.3	33.4	35.6	39.4	43.2	49.9	55.5	59.7	52.1	45.8	-6.2	
	23-26					NA	NA	NA	NA	NA	NA	21.4	23.1	26.4	24.0	26.0	27.8	28.7	31.1	30.1	34.9	41.8	51.5	52.9	49.3	51.3	+2.0	
	27-30									NA	NA	27.1	20.8	22.2	22.8	21.9	27.1	29.3	24.3	26.4	30.0	35.5	40.6	41.2	41.0	41.1	+0.1	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	NA	—
Cocaine	18	47.9	47.5	47.4	43.1	45.0	48.9	51.5	54.2	55.0	58.7	54.5	51.0	52.7	48.5	46.6	47.7	48.1	48.5	51.3	47.6	47.8	46.2	44.6	43.3	47.8	+4.5 s	
	19-22	55.7	56.2	57.1	55.2	56.2	56.9	60.4	65.0	64.9	66.8	61.7	54.3	54.5	49.2	49.9	49.4	44.4	49.7	47.7	52.6	52.1	49.6	47.6	46.7	47.0	+0.3	
	23-26					63.7	67.2	65.8	69.0	71.7	70.0	65.6	58.0	61.1	53.8	54.4	54.7	50.2	46.9	51.8	45.7	45.0	44.6	47.8	40.8	50.7	+9.9 ss	
	27-30									68.6	68.2	64.0	60.0	63.1	56.8	53.1	57.0	53.0	50.4	46.9	50.0	44.6	45.5	46.3	42.9	38.0	-5.0	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	NA	—
	45																								NA	NA	NA	—
Crack	18	NA	NA	NA	NA	NA	NA	NA	41.1	42.1	47.0	42.4	39.9	43.5	43.6	40.5	41.9	40.7	40.6	43.8	41.1	42.6	40.2	38.5	35.3	39.2	+3.9 s	
	19-22	NA	NA	NA	NA	NA	NA	NA	41.9	47.3	47.2	46.9	42.1	42.1	38.4	41.6	40.7	32.9	39.9	40.0	40.8	40.2	37.3	35.7	37.5	33.7	-3.8	
	23-26					NA	NA	NA	44.5	53.0	49.9	46.9	42.0	42.6	42.5	42.4	42.3	37.9	37.2	38.4	35.0	31.9	37.1	33.9	32.8	36.5	+3.7	
	27-30									46.5	46.8	46.8	43.1	45.2	45.8	41.1	44.7	39.9	36.5	33.3	38.8	35.9	36.9	33.4	33.7	28.0	-5.7	
	35															49.6	48.2	43.1	44.3	45.0	41.6	45.0	41.2	38.9	40.5	36.1	-4.3	
	40																			43.3	44.3	42.0	38.7	39.5	39.0	35.8	-3.3	
	45																								37.0	40.0	+3.0	

(Table continued on next page)

TABLE 7-4 (cont.)
Trends in Reported Availability of Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Percentage saying "fairly easy" or "very easy" to get^a

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

	Age Group	Year																				'03-'04 change						
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		2000	2001	2002	2003	2004	
Cocaine powder	18	NA	NA	NA	NA	NA	NA	NA	52.9	50.3	53.7	49.0	46.0	48.0	45.4	43.7	43.8	44.4	43.3	45.7	43.7	44.6	40.7	40.2	37.4	41.7	+4.3	
	19-22	NA	NA	NA	NA	NA	NA	NA	58.7	60.2	61.7	56.5	52.5	48.9	45.7	47.8	45.5	41.3	46.0	47.1	45.2	45.2	43.3	43.9	45.5	43.2	-2.3	
	23-26					NA	NA	NA	64.9	69.1	60.1	58.6	53.2	56.4	50.5	49.7	49.6	45.9	43.6	44.4	44.3	41.8	44.4	40.7	43.4	48.5	+5.1	
	27-30									63.5	62.8	57.9	55.8	56.8	55.0	48.9	52.9	48.4	45.1	43.9	46.5	43.9	42.7	42.4	39.7	37.9	-1.8	
	35															53.9	52.1	46.7	48.3	47.0	43.4	47.9	43.1	41.7	42.0	39.6	-2.4	
	40																			46.0	46.7	44.7	41.5	41.5	40.7	38.5	-2.2	
	45																								39.0	40.2	+1.3	
Heroin	18	21.2	19.2	20.8	19.3	19.9	21.0	22.0	23.7	28.0	31.4	31.9	30.6	34.9	33.7	34.1	35.1	32.2	33.8	35.6	32.1	33.5	32.3	29.0	27.9	29.6	+1.7	
	19-22	18.9	19.4	19.3	16.4	17.2	20.8	21.2	24.4	28.5	31.6	30.7	25.3	30.2	30.0	33.2	35.2	29.1	31.4	32.1	32.7	29.4	30.2	26.4	26.9	22.6	-4.3	
	23-26					18.6	18.1	21.0	22.3	28.4	31.2	28.1	25.6	25.7	25.7	29.2	29.3	32.3	30.5	35.1	31.9	25.7	26.6	27.2	25.5	30.9	+5.4	
	27-30									23.6	27.4	29.5	22.1	25.6	28.5	24.4	30.7	29.5	30.0	28.3	33.0	29.3	29.9	27.0	27.5	22.0	-5.5	
	35																NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																			NA	NA	NA	NA	NA	NA	NA	NA	—
	45																									NA	NA	—
Some other narcotic	18	29.4	29.6	30.4	30.0	32.1	33.1	32.2	33.0	35.8	38.3	38.1	34.6	37.1	37.5	38.0	39.8	40.0	38.9	42.8	40.8	43.9	40.5	44.0	39.3	40.2	+0.8	
	19-22	32.7	32.4	30.8	31.0	28.7	34.3	32.6	33.8	37.9	37.9	35.6	35.4	35.2	33.5	35.1	38.7	37.3	38.3	38.9	39.5	41.1	44.1	40.4	40.6	39.4	-1.3	
	23-26					32.8	32.1	33.6	32.2	35.9	36.4	34.7	33.2	33.9	33.1	35.8	32.6	36.7	35.7	39.9	38.2	38.1	35.8	40.0	40.3	47.7	+7.3	
	27-30									31.6	36.2	36.1	29.0	31.8	33.0	34.8	36.9	37.2	35.2	32.2	36.9	32.4	39.4	38.5	38.9	35.8	-3.1	
	35																NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
	40																				NA	NA	NA	NA	NA	NA	NA	—
	45																									NA	NA	—
Amphetamines	18	61.3	69.5	70.8	68.5	68.2	66.4	64.3	64.5	63.9	64.3	59.7	57.3	58.8	61.5	62.0	62.8	59.4	59.8	60.8	58.1	57.1	57.1	57.4	55.0	55.4	+0.4	
	19-22	71.7	72.6	73.5	69.7	69.1	69.1	63.1	61.8	61.3	62.2	57.7	58.3	56.3	56.0	56.6	60.3	56.9	55.5	56.3	57.6	60.2	56.5	53.7	55.1	53.9	-1.2	
	23-26					65.8	66.0	64.5	65.3	62.2	60.1	55.8	54.8	54.5	52.6	52.9	56.0	52.8	51.2	53.2	49.1	51.1	49.4	48.2	50.3	51.8	+1.6	
	27-30									54.3	58.6	55.3	54.4	50.4	52.9	48.3	53.7	51.7	48.1	41.4	48.2	47.6	49.3	45.6	48.7	43.9	-4.8	
	35															45.6	43.5	39.1	40.9								-3.8	
	40																				41.0	41.9	39.4	37.5	39.4	38.7	37.9	-0.7
	45																									35.8	39.8	+4.0

(Table continued on next page)

TABLE 7-4 (cont.)
Trends in Reported Availability of Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)
Percentage saying "fairly easy" or "very easy" to get^a

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?

	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a																								'03-'04 change		
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		2004	
Crystal meth. (ice)	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	24.0	24.3	26.0	26.6	25.6	27.0	26.9	27.6	29.8	27.6	27.8	28.3	28.3	26.1	26.7	+0.6	
	19-22	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	24.0	21.8	22.5	20.9	24.7	25.5	25.4	29.3	31.0	31.8	27.4	28.4	31.2	26.5	27.1	+0.6	
	23-26					NA	NA	NA	NA	NA	NA	22.3	20.0	21.3	22.9	24.5	24.7	24.7	25.8	30.2	28.5	25.8	26.4	25.1	26.4	32.3	+5.8	
	27-30									NA	NA	27.3	19.7	22.0	21.2	21.7	25.8	26.1	25.1	22.6	29.1	25.3	27.6	29.5	30.9	25.5	-5.4	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																				NA	NA	NA	NA	NA	NA	—	
Sedatives/Barbiturates^c	18	49.1	54.9	55.2	52.5	51.9	51.3	48.3	48.2	47.8	48.4	45.9	42.4	44.0	44.5	43.3	42.3	41.4	40.0	40.7	37.9	37.4	35.7	36.6	35.3	46.3	—	
	19-22	59.5	61.1	56.8	54.2	48.1	52.7	46.8	44.6	45.5	47.7	44.2	41.7	43.4	41.9	40.6	42.9	41.1	39.8	39.2	42.3	40.6	39.3	40.8	38.4	43.8	—	
	23-26					52.7	47.7	46.4	45.9	47.4	44.8	41.6	39.6	42.0	38.8	40.3	42.1	40.6	39.1	42.6	39.7	37.6	36.1	36.4	37.8	49.4	—	
	27-30									43.2	44.5	44.2	38.5	37.8	39.7	37.4	39.9	41.2	39.1	33.9	38.4	36.1	38.1	34.8	35.6	40.5	—	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																				NA	NA	NA	NA	NA	NA	—	
Tranquilizers	18	59.1	60.8	58.9	55.3	54.5	54.7	51.2	48.6	49.1	45.3	44.7	40.8	40.9	41.1	39.2	37.8	36.0	35.4	36.2	32.7	33.8	33.1	32.9	29.8	30.1	+0.3	
	19-22	67.4	62.8	62.0	62.3	52.5	55.6	52.9	50.3	50.0	49.4	45.4	44.8	40.7	40.9	41.0	40.2	37.6	37.8	36.8	37.1	36.5	34.9	34.6	34.2	29.7	-4.5	
	23-26					60.2	54.3	54.1	56.3	52.8	51.4	47.8	45.1	48.1	43.2	45.9	44.3	42.3	36.4	39.4	38.3	37.6	38.7	33.7	32.5	36.6	+4.1	
	27-30									55.3	54.4	54.9	47.5	47.8	47.4	44.4	44.8	46.2	41.9	39.9	41.5	36.7	42.9	38.1	35.9	30.6	-5.3	
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																				NA	NA	NA	NA	NA	NA	—	
Steroids	18	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	46.7	46.8	44.8	42.9	45.5	40.3	41.7	44.5	44.6	44.8	44.4	45.5	40.7	42.6	+1.8	
	19-22	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	44.1	44.8	46.3	41.7	40.9	41.8	40.8	39.2	39.2	40.5	40.3	38.1	41.4	39.4	37.8	-1.6
	23-26					NA	NA	NA	NA	NA	NA	NA	37.6	35.8	39.3	35.8	37.0	37.4	33.9	35.5	34.9	37.1	34.0	34.7	33.1	31.1	34.7	+3.6
	27-30									NA	NA	NA	36.4	30.6	35.0	31.6	30.5	33.1	35.6	32.5	30.5	34.5	36.2	34.6	33.0	32.6	30.6	-2.1
	35															NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
	40																			NA	NA	NA	NA	NA	NA	NA	—	
	45																				NA	NA	NA	NA	NA	NA	—	

(Table continued on next page)

TABLE 7-4 (cont.)
Trends in Reported Availability of Drugs
High School Seniors (Age 18) and Adults in Modal Age Groups of 19-22, 23-26, 27-30, 35, 40, and 45
(Entries are percentages)

Age																										
Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	
<i>Approximate</i>	18	3240	3578	3602	3385	3269	3274	3077	3271	3231	2806	2549	2476	2586	2670	2526	2552	2340	2517	2520	2215	2095	1850	2138	2391	2169
<i>Weighted N=</i>	19-22	582	601	582	588	559	571	592	581	568	572	571	534	512	480	459	470	467	463	433	425	400	398	375	386	441
	23-26				540	541	548	539	526	514	532	511	523	500	463	449	418	419	395	415	388	401	362	356	411	
	27-30								519	513	510	487	475	473	437	446	468	459	425	424	365	357	349	368	393	
	35														1142	1141	1146	1150	1032	1022	981	977	890	934	963	
	40																		1029	1093	1096	1065	1037	898	967	
	45																							911	1026	

Source: The Monitoring the Future Study, the University of Michigan

NOTES: Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

'NA' indicates data not available

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, and (5) Very easy.

^bIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^cIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

Chapter 8

PREVALENCE OF DRUG USE AMONG COLLEGE STUDENTS

The Monitoring the Future study has been able to generate an excellent (and unparalleled) national sample of college students every year since 1980 by following representative samples of high school classes after they graduate. The graduating class of 1976 was the first such class followed in this way, and by 1980 the survey included college students one to four years post-high school. The 2004 survey is thus the 25th such survey covering this important segment of the general population.

We believe that one of the more important functions of the study over the past quarter of a century has been to track various forms of substance use among the nation's college students, in part because they often have been the harbingers of social and political changes that spread eventually to all segments of the population up and down the age spectrum. This was certainly the case for the epidemic of illicit drug use that emerged in the American population in the late 1960s, and it continues today.

The absence of dropouts in the original high school senior samples should have practically no effect on the representativeness of these college samples because very few dropouts go on to college. Perhaps the major limitation of the present design for the purpose of characterizing college students is that it limits the age range of the college sample. For trend estimation purposes, we decided to limit the age band to the most typical one for college attendance, that is, one to four years past high school, which corresponds to the modal ages of 19 to 22. According to the latest statistics available from the United States Bureau of the Census,⁵⁰ this age band should encompass about 73% of all undergraduate college students enrolled full-time in 2003, down some from the 79% covered in 1989. Although expanding the age band to be covered by an additional two years would cover 84% of all enrolled college students, it would also reduce by two years the interval over which we could report trend data. Some special analyses conducted in 1985 indicated that the differences in *prevalence-of-use* estimates under the two definitions were extremely small. The annual prevalence of all drugs except cocaine shifted only about one- or two-tenths of a percent. Cocaine, which has the greatest amount of age-related change, would have had an annual prevalence rate only 0.8 percentage points higher if the six-year age span were included rather than the four-year age span. A replication of these analyses in 1997 yielded virtually the same results. Thus, for purposes of estimating all prevalence rates except lifetime prevalence, the four-year and six-year intervals are nearly interchangeable.

On the positive side, controlling the age band may allow for trend estimation by controlling for changes in the age composition of college students over the years. Otherwise, college students characterized in one year might represent a noncomparable segment of the larger population when compared to college students surveyed in another year.

⁵⁰U.S. Bureau of the Census, October 2003. Available at <http://www.census.gov>.

Definition of college students. *College students are defined here as those follow-up respondents one to four years past high school who say they were registered as full-time students in a two- or four-year undergraduate college at the beginning of March in the year in question.* Note that students at two-year colleges, such as community colleges, are included. The definition excludes those who are currently enrolled in college part-time and those who previously may have been college students or may have completed college.

Prevalence-of-use rates for college students, as well as their same-age peers who are also high school graduates, are provided in Tables 8-1 to 8-5. Having statistics for both groups, a unique feature of the Monitoring the Future panels, makes it possible to see whether college students' substance use rates are higher or lower than those of their age peers (one to four years past high school, i.e., of modal ages 19 to 22). The college-enrolled sample now constitutes over half (59%) of the entire follow-up sample one to four years past high school. The differences reported here pertain to differences between those who are full-time in college versus those who are not, *among high school graduates*. If data from the missing high school dropout segment were available for inclusion as part of the noncollege segment, any difference between the two groups likely would be enlarged; therefore, any differences observed here are only an indication of the direction and relative size of differences between the college and the *entire* noncollege population, not an absolute estimate of them.

PREVALENCE OF DRUG USE: COLLEGE STUDENTS VERSUS THOSE NOT ENROLLED IN COLLEGE

In the year 2004, lifetime prevalence of use among college students is lower for all illicit drugs than among their age peers, but the degree of difference varies considerably by drug, as Table 8-1 shows. However, there is much less difference between the two groups on annual or 30-day prevalence-of-use rates. (See Tables 8-2 and 8-3.) Annual use rates for *alcohol*, *Rohypnol*, *GHB*, *ketamine*, *Ritalin*, and possibly *inhalants* stand apart from the others as being higher among college students than among those not enrolled in college.

- In 2004, annual prevalence for the use of *any illicit drug* among college students stands at 36%, compared to 39% among those high school graduates not in college—a rather modest difference. A larger proportional difference exists for the annual prevalence of *any illicit drug other than marijuana* (19% versus 24%).
- Annual *marijuana* use is very similar among college students and high school graduates of the same age (33% versus 34%). However, the rate of current *daily marijuana* use is lower among college students (4.5% versus 7.2%). (See Table 8-4 for the prevalence of current daily use.)
- Among those illicit drugs for which annual prevalence is *lower* among the college students, *Vicodin*, *narcotics other than heroin*, and *tranquilizers* show the largest absolute difference in annual prevalence. For Vicodin it is 7.4% for college students versus 12.9% for those not in college; for narcotics other than heroin generally, 8.2% versus 12.3%; and for tranquilizers, 6.7% versus 9.9%.

- Smaller absolute differences occur for MDMA (“ecstasy”), with 2.2% of the college students versus 4.7% of their noncollege age peers reporting use in the past year; *sedatives (barbiturates)*, at 4.2% versus 6.7%; *crystal methamphetamine (ice)*, at 1.1% versus 3.3%; *methamphetamine*, at 2.9% versus 5.1%; *cocaine*, at 6.6% versus 8.5%; *OxyContin*, at 2.5% versus 4.3%; *amphetamines*, at 7.0% versus 8.5%; *hallucinogens*, at 5.9% versus 7.3%; and *crack*, at 1.2% versus 2.3%.
- Annual use of *LSD* and *heroin* is also less prevalent (though not significantly) among college students than among their noncollege age peers, at 1.2% versus 1.7% for LSD and 0.4% versus 0.5% for heroin.
- It should be noted that, while the absolute differences are not great between the two groups on many of the low prevalence drugs—including *crack*, *MDMA* (“ecstasy”), *OxyContin*, *methamphetamine*, and *ice*—the ratio of the differences tends to be high (on the order of two or three to one in several cases). Further, these differences are diminished by the absence of the high school dropouts from the noncollege group. So, it is clear that use of a number of the illicit drugs other than marijuana tends to be concentrated among those not in college.
- *Ritalin*, a drug in the amphetamine class and newly added to the MTF questionnaires in 2002, shows quite a different pattern, with use considerably higher among college students (4.7% annual prevalence) than among those not in college (1.6%). This considerable difference may well be explained by college students using Ritalin to stay awake late at night to finish assignments or to study for tests.
- In 2004, college students were not significantly different in prevalence of lifetime or annual use of *alcohol* but were significantly higher than their age peers in monthly use (68% versus 59%).
- College students also had a significantly higher prevalence of *occasions of heavy drinking* (five or more drinks in a row in the past two weeks)—42% versus 34% among their age peers—but they reported significantly lower rates of *daily drinking* than their age peers (3.7% versus 5.8%). It is noteworthy that in high school, college-bound students, especially in earlier grades, were far less likely to drink alcohol at any level compared to their noncollege-bound peers; thus the relative and absolute increases in alcohol use in the first few years following high school are striking for college students.
- Questions on the use of *flavored alcoholic beverages* were introduced in 2004. College students showed a level of use (63%) similar to their noncollege counterparts (61%).
- Among all substances studied, both licit and illicit, the largest absolute difference between the two groups occurs for *cigarette smoking*. For example, the college student prevalence of daily smoking is “only” 14% versus 28% for high school graduates who are of the same age and currently not full-time college students. Smoking at the rate of a half-pack per day stands at 7% versus 19% for these two groups, respectively. Recall that the 12th-grade data show the college-bound to have much lower smoking rates in high school than the noncollege-

bound; thus, in contrast to what was true for alcohol use, these substantial differences observed at college age actually preceded college attendance.⁵¹ The smoking differences would be even greater if dropouts were included in the noncollege group because they have an exceptionally high rate of smoking.

GENDER DIFFERENCES IN PREVALENCE OF USE AMONG COLLEGE STUDENTS

Tabular data are provided separately in Tables 8-1 to 8-5 for male and female college students and their same-age peers.

- Most of the gender differences among college students replicate those discussed earlier for all young adults 1 to 12 years past high school, and they in turn replicate gender differences among secondary school students for the most part. That means that among college students, males have higher annual prevalence rates for most of the illicit drugs. The rates for use of *any illicit drug* are 41% versus 33%; for *any illicit drug other than marijuana*, 22% versus 17%; for *marijuana*, 40% versus 30%; for *hallucinogens*, 9.2% versus 4.0%; and for *LSD* specifically, 2.0% versus 0.8%.
- *Daily marijuana* use is higher among male college students (6.8%) than among females (3.1%).
- Both male and female college students have higher rates of *occasional heavy drinking* than their counterparts not in college (38% versus 27% for females and 49% versus 42% for males).
- *Flavored alcoholic beverages* are favored by significantly more females regardless of whether they attend college or not (69% of females versus 54% of males for college students and 68% versus 50%, respectively, for the noncollege group).
- *Cigarette smoking* among both college students and their noncollege age peers is more prevalent among males. The college segment has rates of daily smoking at 16% among males and 12% among females. Within the noncollege segment, 29% of men were daily smokers compared to 27% of women. Smoking a half-pack or more per day is higher among noncollege men than noncollege women (20% versus 18%); similarly the rates for college men were higher than for college women (8.9% versus 5.6%). As Figure 9-15b in the next chapter shows, there has generally not been a consistent gender difference among college students in their smoking rates for the past several years.
- For a number of drugs in which college students have lower annual prevalence overall, those differences are caused largely or exclusively by the differences between college and noncollege males. (Put another way, the females from these two groups are not nearly as

⁵¹See also Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

different in their use of these drugs as are the males. See Table 8-2.) These drugs include *marijuana*, *LSD*, *Rohypnol*, *GHB*, and *ketamine*.

- On the other hand, it is the noncollege females who account for a disproportionately large part of the overall college versus noncollege differences in the use of *inhalants*, *hallucinogens*, *cocaine*, *narcotics other than heroin*, *Vicodin*, *amphetamines*, *methamphetamine*, and *tranquilizers*.

In sum, the noncollege segment is generally more drug-experienced than the college student segment. This pattern is a continuation of the high school scenario in which those without college plans are more likely to use drugs. The only drugs for which college students are more likely to be users are *alcohol* (including *binge drinking*), *Ritalin*, and the very low prevalence club drugs *Rohypnol*, *GHB*, and *ketamine*. The gender differences observed among college students generally parallel those observed among high school students.

TABLE 8-1
Lifetime Prevalence of Use for Various Types of Drugs, 2004:
Full-Time College Students vs. Others
Among Respondents 1-4 Years Beyond High School
(Entries are percentages)

	Total		Males		Females	
	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>
Any Illicit Drug ^a	52.2	60.0	54.9	59.4	50.6	60.6
Any Illicit Drug ^a Other Than Marijuana	28.0	35.5	31.1	35.6	26.2	35.3
Marijuana	49.1	57.8	53.7	58.4	46.5	57.3
Inhalants ^{b,c}	8.5	11.4	12.3	13.1	6.1	10.1
Hallucinogens ^c	12.0	17.9	16.1	20.5	9.6	15.8
LSD	5.6	11.4	6.9	12.3	4.9	10.7
Cocaine	9.5	16.5	12.5	17.5	7.8	15.6
Crack ^d	2.0	5.5	2.4	5.8	1.7	5.2
MDMA (Ecstasy) ^b	10.2	15.4	9.2	16.1	10.9	14.9
Heroin	0.9	1.7	1.2	1.6	0.7	1.8
Other Narcotics ^e	13.8	20.2	18.2	20.8	11.2	19.6
Amphetamines, Adjusted ^{e,f}	12.7	18.0	13.8	16.7	12.1	19.0
Ice ^g	2.2	8.3	3.5	9.8	1.5	7.1
Sedatives (Barbiturates) ^e	7.2	12.2	8.4	12.4	6.5	12.0
Tranquilizers ^e	10.6	16.6	13.5	17.4	8.8	16.1
Alcohol	84.6	85.8	83.2	85.3	85.3	86.2
Flavored Alcoholic Bvg. ^h	79.0	76.3	73.9	74.3	82.3	77.6
Cigarettes	NA	NA	NA	NA	NA	NA
<i>Approximate Weighted N =</i>	<i>1400</i>	<i>990</i>	<i>520</i>	<i>450</i>	<i>880</i>	<i>540</i>

Source: The Monitoring the Future Study, the University of Michigan.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N in 2004 for college students is approximately 700.

^cUnadjusted for known underreporting of certain drugs. See text for details.

^dThis drug was asked about in five of the six questionnaire forms. Total N in 2004 for college students is approximately 1160.

^eOnly drug use that was not under a doctor's orders is included here.

^fBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^gThis drug was asked about in two of the six questionnaire forms. Total N in 2004 for college students is approximately 470.

^hThis drug was asked about in one of the six questionnaire forms. Total N in 2004 for college students is approximately 230.

TABLE 8-2
Annual Prevalence of Use for Various Types of Drugs, 2004:
Full-Time College Students vs. Others
Among Respondents 1-4 Years Beyond High School
(Entries are percentages)

	Total		Males		Females	
	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>
Any Illicit Drug ^a	36.2	38.8	40.9	40.1	33.4	37.8
Any Illicit Drug ^a Other Than Marijuana	18.6	23.7	22.1	24.3	16.5	23.1
Marijuana	33.3	33.7	39.6	35.8	29.6	32.0
Inhalants ^{b,c}	2.7	2.4	3.5	3.6	2.2	1.6
Hallucinogens ^c	5.9	7.3	9.2	9.3	4.0	5.6
LSD	1.2	1.7	2.0	2.6	0.8	1.0
Cocaine	6.6	8.5	9.3	9.1	4.9	8.1
Crack ^d	1.3	2.3	1.2	2.4	1.3	2.2
MDMA (Ecstasy) ^b	2.2	4.7	2.1	4.0	2.2	5.2
Heroin	0.4	0.5	0.6	0.7	0.3	0.3
Other Narcotics ^e	8.2	12.3	10.6	13.2	6.8	11.5
OxyContin ^f	2.5	4.3	3.5	6.2	1.9	2.7
Vicodin ^f	7.4	12.9	11.1	13.0	5.2	12.9
Amphetamines, Adjusted ^{c,g}	7.0	8.5	8.2	8.2	6.3	8.8
Ritalin ^f	4.7	1.6	6.8	2.8	3.5	0.7
Methamphetamine ^f	2.9	5.1	4.4	5.3	2.0	4.9
Ice ^f	1.1	3.3	1.9	4.2	0.6	2.6
Sedatives (Barbiturates) ^e	4.2	6.7	4.6	6.8	3.9	6.6
Tranquilizers ^c	6.7	9.9	8.2	10.3	5.8	9.6
Rohypnol ^f	0.3	0.0	0.8	0.0	0.0	0.0
GHB ^f	0.7	0.2	1.3	0.0	0.3	0.4
Ketamine ^f	1.5	0.5	2.8	0.5	0.8	0.6
Alcohol	81.2	80.0	79.2	80.2	82.4	79.9
Flavored Alcoholic Bvg. ^h	63.2	60.6	53.6	50.3	69.3	67.6
Cigarettes	36.7	47.6	42.6	49.1	33.2	46.4
<i>Approximate Weighted N =</i>	<i>1400</i>	<i>990</i>	<i>520</i>	<i>450</i>	<i>880</i>	<i>540</i>

Source: The Monitoring the Future Study, the University of Michigan.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N in 2004 for college students is approximately 700.

^cUnadjusted for known underreporting of certain drugs. See text for details.

^dThis drug was asked about in five of the six questionnaire forms. Total N in 2004 for college students is approximately 1160.

^eOnly drug use that was not under a doctor's orders is included here.

^fThis drug was asked about in two of the six questionnaire forms. Total N in 2004 for college students is approximately 470.

^gBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^hThis drug was asked about in one of the six questionnaire forms. Total N in 2004 for college students is approximately 230.

TABLE 8-3
Thirty-Day Prevalence of Use for Various Types of Drugs, 2004:
Full-Time College Students vs. Others
Among Respondents 1-4 Years Beyond High School
(Entries are percentages)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Any Illicit Drug ^a	21.2	23.2	26.1	25.3	18.4	21.5
Any Illicit Drug ^a Other Than Marijuana	9.1	11.6	11.3	13.8	7.8	9.7
Marijuana	18.9	20.3	24.6	22.1	15.6	18.9
Inhalants ^{b,c}	0.4	0.7	0.5	0.4	0.4	0.9
Hallucinogens ^c	1.3	1.8	1.9	2.7	1.0	1.2
LSD	0.2	0.4	0.2	0.6	0.2	0.1
Cocaine	2.4	2.4	3.6	3.1	1.8	1.9
Crack ^d	0.4	0.3	0.5	0.2	0.4	0.3
MDMA (Ecstasy) ^b	0.7	1.2	1.0	1.5	0.5	1.0
Heroin	0.1	0.1	0.3	0.0	0.1	0.2
Other Narcotics ^e	3.0	4.3	3.8	5.0	2.5	3.7
Amphetamines, Adjusted ^{e,f}	3.2	3.4	3.6	3.6	3.1	3.2
Ice ^g	0.1	1.4	0.3	2.1	0.0	0.8
Sedatives (Barbiturates) ^e	1.5	3.2	1.5	3.7	1.5	2.8
Tranquilizers ^e	2.7	4.0	3.2	4.5	2.4	3.5
Alcohol	67.7	59.3	66.7	62.8	68.2	56.3
Flavored Alcoholic Bvg. ^h	34.0	30.0	27.4	22.3	38.3	35.2
Cigarettes	24.3	36.5	29.8	39.0	21.1	34.4
<i>Approximate Weighted N =</i>	<i>1400</i>	<i>990</i>	<i>520</i>	<i>450</i>	<i>880</i>	<i>540</i>

Source: The Monitoring the Future Study, the University of Michigan.

*' indicates a percentage of less than 0.05% but greater than true zero.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N in 2004 for college students is approximately 700.

^cUnadjusted for known underreporting of certain drugs. See text for details.

^dThis drug was asked about in five of the six questionnaire forms. Total N in 2004 for college students is approximately 1160.

^eOnly drug use that was not under a doctor's orders is included here.

^fBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^gThis drug was asked about in two of the six questionnaire forms. Total N in 2004 for college students is approximately 470.

^hThis drug was asked about in one of the six questionnaire forms. Total N in 2004 for college students is approximately 230.

TABLE 8-4
Thirty-Day Prevalence of Daily Use for Various Types of Drugs, 2004:
Full-Time College Students vs. Others
Among Respondents 1-4 Years Beyond High School
(Entries are percentages)

	Total		Males		Females	
	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>
Marijuana	4.5	7.2	6.8	8.9	3.1	5.8
Cocaine	0.0	0.2	0.0	0.2	0.0	0.2
Amphetamines, Adjusted ^{a,b}	0.2	0.4	0.0	0.2	0.2	0.5
Alcohol						
Daily	3.7	5.8	7.2	9.0	1.8	3.1
5+ Drinks in a Row in Past 2 Weeks	41.7	33.7	48.9	41.7	37.7	27.0
Cigarettes						
Daily	13.8	28.0	16.2	29.4	12.3	26.8
Half-Pack or More per Day	6.8	19.2	8.9	20.1	5.6	18.4
<i>Approximate Weighted N =</i>	<i>1400</i>	<i>990</i>	<i>520</i>	<i>450</i>	<i>880</i>	<i>540</i>

Source: The Monitoring the Future Study, the University of Michigan.

*' indicates a prevalence rate of less than 0.05% but greater than true zero.

^aOnly drug use that was not under a doctor's orders is included here.

^bBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

TABLE 8-5**Lifetime, Annual, and Thirty-Day Prevalence of an Illicit Drug Use Index,^a 2004:
Full-Time College Students vs. Others****Among Respondents 1-4 Years Beyond High School**

(Entries are percentages)

	Total		Males		Females	
	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>	<u>Full-Time College</u>	<u>Others</u>
Lifetime						
Any Illicit Drug	52.2	60.0	54.9	59.4	50.6	60.6
Any Illicit Drug Other Than Marijuana	28.0	35.5	31.1	35.6	26.2	35.3
Past 12 Months						
Any Illicit Drug	36.2	38.8	40.9	40.1	33.4	37.8
Any Illicit Drug Other Than Marijuana	18.6	23.7	22.1	24.3	16.5	23.1
Past 30 Days						
Any Illicit Drug	21.2	23.2	26.1	25.3	18.4	21.5
Any Illicit Drug Other Than Marijuana	9.1	11.6	11.3	13.8	7.8	9.7
<i>Approximate Weighted N =</i>	<i>1400</i>	<i>990</i>	<i>520</i>	<i>450</i>	<i>880</i>	<i>540</i>

Source: The Monitoring the Future Study, the University of Michigan.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

Chapter 9

TRENDS IN DRUG USE AMONG COLLEGE STUDENTS

College students are often the harbingers of social change in society, and such was the case in the emergence of the illicit drug epidemic of the last forty years. Illicit drug use increased dramatically among American college students in the mid-1960s, then spread quickly to their noncollege age peers and eventually down the age spectrum to high school students and even to middle school students. College students were thus the leading edge of that social change in illicit drug use. As we shall show in this chapter, the diffusion process seems to have reversed during the more recent “relapse” of the epidemic in the 1990s, when use first increased among those in early adolescence and then radiated *up* the age spectrum as those cohorts grew older.

The definition of college students is the same here as described in chapter 8: *high school graduates one to four years past high school who are enrolled full-time in a two-year or four-year college at the beginning of March in the year in question*. For comparison purposes, trend data are provided on the remaining follow-up respondents in this age band, who are also one to four years past high school. (See Figures 9-1 through 9-15.) Because the rate of college enrollment declines steadily with number of years beyond high school, this comparison group is slightly older on average than the college-enrolled group. It is also worth noting that the proportion of young adult high school graduates one to four years beyond high school who are enrolled full-time in college has increased considerably over the past 24 surveys. In 2004, about 59% of the weighted number of follow-up respondents one to four years past high school met our definition of college students, compared with only 38% in the 1980 survey. That 21-percentage-point increase represents a substantial rise over the past two and one-half decades in the proportion of high school graduates attending college.

The reader is reminded that the difference between the enrolled group and the other group provides an estimate of the degree to which college students are above or below average for other high school *graduates* in this age band. Were we able to include the high school dropout segment in the calculation for the noncollege group, many of the differences with the college-enrolled likely would be accentuated.

For each year given, there are approximately 1,100-1,500 weighted respondents constituting the college student sample (see Table 9-5 for Ns per year) and roughly 900-1,700 respondents constituting the “other” group one to four years past high school. Comparisons of the trends for these two groups are provided in this chapter. Because it was not until 1980 that enough follow-up years had accrued to characterize young people one to four years past high school, the comparisons begin with that year.

TRENDS IN PREVALENCE 1980-2004: COLLEGE STUDENTS VERSUS THOSE NOT ENROLLED IN COLLEGE

- The proportion of college students using *any illicit drug* in the 12 months prior to the survey (i.e., the annual prevalence rate) dropped fairly steadily between 1980 and 1991 (from 56% to 29%). (See Table 9-2.) In other words, illicit drug use fell by nearly half over the 11-year period 1980–1991. After 1991, annual (and also 30-day) prevalence held fairly steady for a couple of years before beginning to rise, reaching 38% in 1998—still well below the peak of 56% in 1980. There has been little change since then (36% in 2004). The noncollege group moved similarly from 1980 to 1998. Twelfth graders also showed a similar trajectory in the decline phase through 1991, but their rise in use after 1992 was distinctly sharper, as Figure 9-1 illustrates. All three groups showed a leveling after 1998. However, in 2000, the noncollege group exhibited a 4-percentage-point increase that was due largely to their sharper increases in marijuana, amphetamine, and tranquilizer use in that year, and their level remains above the college student sector.
- Use of *any illicit drugs other than marijuana* declined fairly steadily among college students between 1980 and 1994, with annual prevalence dropping by nearly two-thirds from 32% to 12% (Table 9-2). This generally paralleled the trend for the noncollege group, as well as for 12th graders. All three groups showed some increase in use during the 1990s: the 12th graders after 1992, the noncollege group after 1993, and the college students after 1994. However, the rise in use of illicit drugs other than marijuana was not as sharp among college students as it was in either of the two other groups (Figure 9-2). Since 1999 the college students and noncollege segment have shown some further increase (particularly the noncollege group), whereas use among the 12th graders held steady and has even begun to decline. In fact, the college students continue to show an increase in their annual prevalence rate into 2004. This divergence most likely reflects some cohort effects working their way up the age spectrum.

In general, among those enrolled in college, the trends during the 1980s for most individual classes of illicit drugs tended to parallel those for the noncollege group and those observed among 12th graders. During the 1990s, however, there was more divergence in the trends, with the college students usually showing less increase than the 12th graders and, for some drugs, less increase than their age-mates not in college.

- The annual prevalence of *marijuana* use among college students decreased steadily from 1981 through 1991, dropping by nearly half from 51% to 27% (Figure 9-3a). Their noncollege peers showed a comparable decline over the same time interval. Use among 12th graders rose sharply after 1992, while use among college students and their age peers rose more gradually. From 1991 through 1998, annual prevalence rose by nearly 10 percentage points among college students, by 7 percentage points among their age peers not in college, but by 14 percentage points among 12th graders. The 12th graders were the first to show a leveling off in marijuana use (in 1998), followed by the college students in 1999 and the noncollege group in 2002. All three groups have very similar rates of use in 2004 after some gradual decline in use.

- **Daily marijuana** use among college students (Figure 9-3b) fell appreciably between 1980 and 1986, from 7.2% to 2.1%, as it did for those not in college and among 12th graders. (The latter two groups were able to show sharper declines because they started higher than the college students in 1980.) After 1986, the decline decelerated, and by 1991 the rate stood at 1.8%. In sum, the proportion of American college students who actively smoked marijuana on a daily basis dropped by about three-fourths between 1980 and 1991. Daily use then leveled until 1994 and began increasing thereafter, reaching 4.6% in 2000. Since then, daily use for college students has held level (4.5% in 2004). The other two groups showed considerably larger increases after 1993 than did college students, with 12th graders' daily use rates leveling after 2000. The noncollege segment showed further increase in 2001, reaching 9.4%, and then falling to 7.2% in 2004, still well above college students.
- A very appreciable decline occurred for **amphetamine** use between 1981 and 1991 (Figure 9-11). Annual prevalence among college students dropped by more than eight-tenths, from 22% in 1981 to 4% in 1991. Proportionately, this was a larger drop than among 12th graders but fairly parallel to the overall change among their age peers not in college. Amphetamine use among college students and their noncollege age peers leveled for a year before beginning to increase in both groups after 1992 and 1993, respectively, through 2001, with a leveling through 2004. Over the years, those not in college consistently have reported a higher rate of amphetamine use than the college students, and since the mid-1980s 12th graders have reported higher rates still.
- During the early 1980s, one of the largest proportional declines observed among college students was for **LSD** (see Figure 9-6). Annual prevalence fell from 6.3% in 1982 to 2.2% in 1985. After 1985, their use began to increase, reaching 6.9% by 1995. Since 1995 use has fallen among college students, their age-mates, and 12th graders; in 2002 there was a particularly sharp decrease in all groups. All three groups now have annual prevalence rates between 1.2% and 2.2%. College students have remained at lower levels of use than the other two groups since the mid-1990s.
- After 1997 there was a sharp increase in the use of **ecstasy (MDMA)** by American college students (Figure 9-8). Their annual prevalence rose three- to fourfold in just three years, from 2.4% in 1997 to 9.2% in 2001, before it began to decrease, reaching 2.2% in 2004. The trends among college students have run fairly parallel to those for the noncollege segment and 12th graders through 2004. Since 2000, the noncollege segment has exhibited the highest rate of ecstasy use—reaching 14% in 2001, when use among the college students and 12th graders was at 9%. All three groups showed sharp declines in the following three years.
- When the college data were first available in 1980, **sedative (barbiturate)** use already was quite low among college students (at 2.9% annual prevalence), but it still fell by more than half to 1.3% by 1985. This proportional decline was, once again, sharper than among 12th graders and less sharp than among the young adults not in college, both of whom started at a higher level of use. Annual prevalence remained essentially unchanged between 1985 and 1993 among all three groups (see Figure 9-12). The groups then showed a gradual increase in use between 1993 (or 1994 in the case of the college students) and 2001, with 12th graders

showing a significant increase in 2002 and use in the other two groups leveling off. The college students have shown a fairly steady increase over the ten-year period 1994-2004, with pauses in 1998 and 2002; but the other two groups remained at higher levels than the college students throughout this period.

- Figure 9-13 shows that the annual prevalence of *tranquilizer* use among college students dropped by half in the period 1980-1984, from 6.9% to 3.5%, and again fell by half between 1984 and 1994, to 1.8%.⁵² After this long period of gradual decline, tranquilizer use then began to increase gradually, returning to 6.9% by 2003. Use by the noncollege segment and by 12th graders dropped more sharply, eliminating the differences among the three groups by 1992. Use rose after 1992 for all but the college students, again opening some differences, and then rose in all three groups after 1994; in 2002, tranquilizer use was at or near its recent high in all three groups. From 1999 through 2002 the increase in use was particularly sharp among the noncollege segment, making them the highest-using group. In 2003, however, they and the 12th graders showed their first declines in recent years, thus narrowing the differences among the three groups. No further decline was observed in any group in 2004.
- The overall trends in the use of *narcotics other than heroin*⁵³ have been quite parallel to those for sedatives (barbiturates) and tranquilizers. By 1994 the use of narcotics other than heroin (Figure 9-10a) by college students was about half what it was in 1980 (2.4% in 1994 versus 5.1% in 1980) as a result of a gradual decline over the interval. This trend closely parallels use among noncollege young adults and 12th graders. As with a number of other drugs, use among 12th graders began to rise after 1992, but use among college students did not begin to increase until after 1994, likely due to a cohort effect. In 2003, annual prevalence among college students reached an historic high point of 8.7% while use among 12th graders finally leveled off after reaching an historic high in 2002. The noncollege group reached an all-time high of 12% in 2004, and—unlike the other two groups—does not yet show signs of leveling.
- Although data were not collected until 2002, it is clear that *OxyContin* and *Vicodin* (Figures 9-10b and 9-10c) help to explain the difference between the college and noncollege segments in their use of narcotics other than heroin. The noncollege group has had prevalence rates up to twice that for the college students in the use of both drugs (see Table 8-2). Annual prevalence of OxyContin use rose continuously among college students, from 1.5% in 2002 to 2.5% in 2004, as it did among 12th graders (from 4.0% to 5.0%). Use among the noncollege segment also rose from 2002-2003, from 3.3% to 4.5%; but then held constant at 4.3% in 2004. Vicodin use showed a somewhat different pattern over the two years: annual prevalence among the college students is higher in 2004 (7.4%) than in 2002 (6.9%), but all of that increase occurred by 2003, followed by a leveling. The other two groups have rates in 2004 about what they were in 2002.

⁵²The use of sedatives (barbiturates) and tranquilizers very likely dropped during the latter half of the 1970s, as well, judging by the trends among 12th graders.

⁵³As discussed in the text in chapters 4 and 5, because the questions about narcotics other than heroin were changed in 2002, the prevalence figures are adjusted estimates. See the earlier discussion for details.

- Like the 12th graders, college students showed a relatively stable pattern of *cocaine* use between 1980 and 1986, when their usage levels (and those of their age peers) were considerably higher than those observed among 12th graders. (See Figure 9-9.) This level period was followed by a dramatic drop of nearly nine-tenths in annual prevalence among college students, from 17.1% in 1986 to 2.0% in 1994. Their noncollege counterparts also showed a large, but somewhat less dramatic decline, from 18.9% in 1986 to 5.1% in 1994. Because use among college students also dropped more sharply than among 12th graders, there was little or no difference between the two groups in annual prevalence rates for cocaine between 1990 and 1995. After 1995, cocaine use rose the least among the college students, creating a reversal of the previous gap with the 12th graders having higher levels of use than the college students. Between 1994 and 1998 annual cocaine prevalence for college students increased significantly, from a 14-year low of 2.0% in 1994 to 4.5% by 1998, roughly where it stayed through 2002. Use has shown a gradual rise since 2002, however, with annual prevalence rising from 4.8% to 6.6%. Twelfth graders and noncollege students also exhibited an increase in annual prevalence of cocaine use after 1992 and 1993, respectively. Use has been level among 12th graders since 2000 but continued to increase among those not in college through 2001, considerably widening the gap between the noncollege segment and the other two groups.
- College students have shown some shifts in *alcohol* use that are different from those observed either among their age peers not in college or among 12th graders. As can be seen in Figure 9-14d, both the noncollege segment and the 12th graders showed fairly substantial declines from 1981 through 1990 in the prevalence of having *five or more drinks in a row* during the prior two weeks. (The 12th graders then showed further decline for three more years.) In contrast, the college students showed no decline in binge drinking from 1981 to 1986 and then only a modest decline of 5 percentage points from 1986 through 1993. In the 11-year period between 1981 (when all three populations were very close in use) and 1992, this measure of heavy drinking dropped by 14 percentage points among 12th graders, by 11 percentage points among the noncollege 19- to 22-year-olds, but by only 2 percentage points among college students. After 1992, binge drinking began to rise among 12th graders while it still was declining some among college students—likely reflecting a cohort effect emerging during this period, similar to that observed for a number of illicit drugs—narrowing the gap somewhat. Binge drinking also then began to increase among the noncollege segment after 1995 and by less among college students after 1996—increases that continued into 2001, after which noncollege students showed some decline. Meanwhile, among 12th graders, binge drinking started a gradual decline after 1998, enlarging the difference between them and the other two groups among whom this behavior was still rising. Once again there is evidence of cohort effects at work here since the beginning of the 1990s, with the inflection points being later for the older strata. Despite the different patterns of trends, perhaps the most noteworthy facts are that college students have exhibited the highest level of, and greatest constancy of rate in, binge drinking throughout the entire 24-year interval that college students have been covered by the study.

It is interesting to conjecture why college students did not show much decline in heavy drinking for a decade (1981–1991) while their noncollege peers and 12th graders did. One possibility is that campuses provided some insulation to the effects of changes in the

drinking age laws. Similarly, entrenched in many college campuses is a “culture of binge drinking” which has proven impervious to many societal trends (and intervention attempts) regarding excessive alcohol use.⁵⁴ Also, individuals who are under the legal drinking age in college are mixed in with peers who are of legal age to purchase alcohol in a way that is no longer true in high schools and less true, perhaps, for those aged 19 to 22 who are not in college. Finally, much alcohol advertising was directed specifically at the college student population.

It is worth noting that in 2004 there was a 3.3 percentage point increase in binge drinking reported by college students, reaching the highest level observed in recent years (but note that this large one-year increase in binge drinking, the largest seen in the study’s history, did not reach statistical significance). However, there is a possible methodological artifact present, in that the first question in the series of questions about alcohol use was slightly modified in 2004, in half of the six questionnaire forms. The observed increase of 3.3 percentage points was entirely due to higher reported binge drinking by college students in the three changed forms. Responses to the other alcohol use questions (lifetime, annual, 30-day prevalence) by college students, and responses to all the alcohol use questions, including binge drinking, by other high school graduates, as well as secondary school students, appeared not to be affected; thus, it is possible that the reported increase was simply a transitory quirk. Another year of data may clarify the trends.

College students generally have had slightly lower rates of *daily drinking* than their age group taken as a whole, though by the early 1990s such differences nearly disappeared (Figure 9-14c). Daily drinking among the young adults (1-4 years past high school) not enrolled in college declined from 8.7% in 1981 to 6.5% in 1984, remained essentially unchanged through 1988, declined further (to 3.2%) by 1994, then increased to 5.8% by 2004. College students’ daily drinking estimates—which appear a little less stable, perhaps due to smaller sample sizes in the 1980s—showed little or no decline between 1980 (6.5%) and 1984 (6.6%) but a considerable decline from 1984 through 1995 (to 3.0%), followed by a period of some increase, reaching 5.0% in 2002. Since 2002 their daily drinking has dropped, to 3.7% in 2004. Twelfth graders showed a similar pattern of daily drinking with a long period of decline, followed by a somewhat earlier reversal, beginning in 1994. After 1998 their daily drinking rate actually declined a little, increased a bit in 2001, and then declined again up through 2004. While daily drinking has been dropping over the past two years among 12th graders and college students, it has been rising among the 19- to 22-year-olds not in college, leaving them with the highest current rate of daily drinking, as they have had over most of the past 24 years.

- **Cigarette smoking** among American college students (Figure 9-15a) declined modestly in the first half of the 1980s. Thirty-day prevalence fell from 26% to 22% between 1980 and 1985, remained fairly stable through 1990 (22%), then increased gradually but substantially, reaching 31% by 1999. It was not until 2000 that the first evidence of a decline in smoking among college students began to appear, two years after smoking had begun to decline

⁵⁴Schulenberg, J. E., & Maggs, J. L. (2002). A developmental perspective on alcohol use and heavy drinking during adolescence and the transition to young adulthood. *Journal of Studies on Alcohol, Supplement 14*, 54-70.

among 12th graders. This lag no doubt reflects a cohort effect operating through generational replacement. The noncollege group showed the first evidence of a decline in their smoking rate in 2002—considerably later than the college students; further, their smoking rate leveled thereafter. College students as well as 12th graders showed a slight increase in 30-day smoking prevalence in 2004 (neither change was statistically significant)—a potentially noteworthy development in that it may signal the end to the important decline in smoking that had been taking place in both of these populations.

The *daily smoking* rate for college students (Figure 9-15b) fell from 18% in 1980 to 13% in 1986, as the cohorts who had lower initiation rates by senior year replaced the earlier, heavier-smoking cohorts. It remained fairly level through 1990 (12%) but by 1999 rose to 19%, the highest level of daily smoking we have recorded among American college students since we began tracking them in 1980. (The 1999 30-day prevalence rate was also the highest we have recorded.) After 1999, both statistics declined among college students until 2004, when daily smoking leveled and 30-day prevalence rose.

While smoking rates have consistently been lower among college students than among those who were of the same age and were not in college, the trends for these two groups converged some after 1984, as smoking rates more or less stabilized among college students but continued to decline among young adults not in college (see Figure 9-15a). In fact, between 1989 and 1991, use began to rise among college students while continuing to decline among their peers. Both groups showed fairly parallel increases in smoking between about 1991 and 1999, after which use continued to increase among the noncollege segment but began to decline among college students. (Twelfth graders exhibited an increase from 1992 to 1997, and their use has declined significantly since.) The popularity of Camel cigarettes among the college-bound, which we have reported elsewhere, may help to explain some of the narrowing of the gap between college students and their age peers.⁵⁵ The Joe Camel advertising and promotion campaign, commenced in the late 1980s and ended in the late 1990s, may have succeeded in initiating more college students (particularly male college students) smoking than had been the case previously or since.

- For many drugs—*amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—differences between college students and their noncollege age peers narrowed over the years. Much of this is due to general overall declines in usage rates during the 1980s, but some may also reflect the increasing proportion of the age group going to college.

The overall drug use trends among college students also are parallel, for the most part, to the trends among 12th graders; still declines in many drugs over the decade of 1980 to 1990 were proportionately larger among college students, and for that matter, among all young adults of college age, than among 12th graders. Despite parallel trends in the early 1990s, the 12th graders have shown a larger, and often earlier, increase in the use of a number of drugs in the years since; as indicated in Volume I, the 8th and 10th graders in secondary school

⁵⁵Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (1999). *Cigarette brand preferences among adolescents*. (Monitoring the Future Occasional Paper No. 45.) Ann Arbor, MI: Institute for Social Research.

showed increases a year earlier than the 12th graders. It is clear that this most recent upsurge, or what we have called a “relapse phase” in the illicit drug epidemic, did not originate on the nation’s campuses, as did the original epidemic. It originated among secondary school students—and the younger ones at that—and has been carried up the age spectrum through generational replacement, at least in part. Put in more general terms, there is clear evidence of some important cohort effects at work here.

GENDER DIFFERENCES IN TRENDS AMONG COLLEGE STUDENTS

One trend that is not obvious from the figures included here is the slow rise in the proportion of college students who are female. Females constituted 50% of our 1980 sample of college students compared to 63% of our 2004 sample. Given that substantial gender differences exist in the use of some drugs, we have been concerned all along that apparent long-term trends in the levels of drug use among college students might actually be attributable to changes in the gender composition of that population. For that reason, in particular, we have consistently presented separate trend lines for the male and female segments of the college student population. Differences in the trends observed for the two genders, illustrated in the lower panels of Figures 9-1 through 9-15, are discussed next.

In general, trends in the use of the various drugs and in the overall drug use indexes have been highly parallel for male and female college students, as an examination of the relevant figures will show. The most noteworthy exceptions are mentioned below.

- Certain drug use measures showed a convergence of usage levels between the genders as they dropped to very low levels. *Marijuana* use was one example, with some convergence in the rates between 1980 and 1991 as overall use declined, and then some divergence between 1991 and 1999 as usage rates rose. Since 2001, however, the two genders have been moving in opposite directions, with use among males increasing and use among females decreasing (see Figure 9-3a). *Daily marijuana* use presents a clearer example, with the decline among males between 1980 and 1986 narrowing the gap between the genders. Between 1986 and 1993 there was no further narrowing; but as use began to rise in the mid-1990s, a greater increase among males widened the gap again. Use among males held fairly steady in the late 1990s while use among females rose, once again narrowing the gender gap somewhat. In 2004 the rates for the two genders seemed to be moving in opposite directions—6.8% versus 3.1% for male and female college students, respectively. (See Figure 9-3b.)
- *LSD* use dropped more steeply among males over the four-year interval 1999-2003 than among females, bringing the genders closer together at quite low prevalence rates (Figure 9-6). Prior to that period of convergence, use had quite consistently been higher among males.
- After 1986, *cocaine* use dropped more steeply for males than for females in general and among male college students in particular, considerably narrowing the sizable gap between the genders (see Figure 9-9). Since 1991 both genders moved closely parallel, with males reporting somewhat higher usage rates (6.2% versus 5.0% for females in 2003). In 2004 males showed an increase in prevalence (not significant) not observed among females.

Whether this reflects a real shift or simply a brief perturbation in the data will be clearer with another year's data.

- Like a number of other drugs, *methaqualone* also showed a convergence in use through 1989, with use among males declining more than among females (no figure given).
- *Amphetamine* use (Figure 9-11) also showed some convergence in the early 1980s due to a greater decline among males. In fact, male and female college student use has been essentially equal since 1989.
- The annual prevalence of *alcohol* use has been virtually identical for the two genders throughout the duration of the study (Figure 9-14a) and the 30-day prevalence rate has been quite close, with males slightly higher; but college males have consistently had higher rates of *daily drinking* and *binge drinking* (Figures 9-14c and 9-14d). From 1988 through 1994, binge drinking among college females decreased some (from 37% to 31%); but binge drinking among college males declined more, from a high point in 1986 of 58% to a low of 47% in 1995 (see Figure 9-14d). Since 1998, there has been some closing of the gender gap in binge drinking, as the rate among college females has risen (from 31% in 1998 to 38% in 2004, including a not quite significant increase of 4.5 percentage points in 2004), while it actually declined a bit from 52% to 49% among college males.
- Between 1980 and 1992, the 30-day prevalence of *cigarette smoking* was consistently higher among college females than males (Figure 9-15a). However, the gap in 30-day prevalence narrowed because use by female college students declined considerably between 1980 and 1989, while use by male college students did not decline. After 1989, the gap remained quite small and the genders reversed position, with males catching up to, and passing, females in their rate of smoking by 1994 and then remaining higher through at least 2000. (A similar reversal had occurred among 12th graders a few years earlier.) Both genders exhibited a considerable decrease in 30-day smoking between 1999 and 2003, leaving very little difference between them (22% for males, 23% for females). In 2004, the decline continued for females, while males exhibited a significant increase.

While the rise in smoking among college students was longer-term and more gradual than in the other two groups, it nevertheless was substantial, rising by nearly half between 1989 (21%) and 1999 (31%). The increase in smoking after 1988 was sharper among college males than among college females, consistent with the notion that Camel cigarettes' promotion and advertising—which ended in the late 1990s as a part of the tobacco settlement—may have played a role in the overall increase. Camels proved considerably more popular among males, especially among those college-bound and from more educated families.⁵⁶

⁵⁶Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (1999). *Cigarette brand preferences among adolescents* (Monitoring the Future Occasional Paper No. 45.) Ann Arbor, MI: Institute for Social Research; see pp. 6-7.

TABLE 9-1
Trends in Lifetime Prevalence of Various Types of Drugs
Among College Students 1-4 Years Beyond High School

Percentage who used in lifetime

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
<i>Approx. Wtd. N =</i>	1040	1130	1150	1170	1110	1080	1190	1220	1310	1300	1400	1410	1490	1490	1410	1450	1450	1480	1440	1440	1350	1340	1260	1270	1400	
Any Illicit Drug ^a	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	-1.7
Any Illicit Drug ^a																										
Other Than Marijuana	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	+0.5
Marijuana	65.0	63.3	60.5	63.1	59.0	60.6	57.9	55.8	54.3	51.3	49.1	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8	51.2	51.0	49.5	50.7	49.1	-1.6
Inhalants ^{b,c}	10.2	8.8	10.6	11.0	10.4	10.6	11.0	13.2	12.6	15.0	13.9	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4	12.9	9.6	7.7	9.7	8.5	-1.2
Hallucinogens ^{c,d}	15.0	12.0	15.0	12.2	12.9	11.4	11.2	10.9	10.2	10.7	11.2	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8	14.4	14.8	13.6	14.5	12.0	-2.5 s
LSD	10.3	8.5	11.5	8.8	9.4	7.4	7.7	8.0	7.5	7.8	9.1	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7	11.8	12.2	8.6	8.7	5.6	-3.1 ss
MDMA (Ecstasy) ^e	NA	NA	NA	NA	NA	NA	NA	NA	NA	3.8	3.9	2.0	2.9	2.3	2.1	3.1	4.3	4.6	6.8	8.4	13.1	14.7	12.7	12.9	10.2	-2.7
Cocaine	22.0	21.5	22.4	23.1	21.7	22.9	23.3	20.6	15.8	14.6	11.4	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4	9.1	8.6	8.2	9.2	9.5	+0.3
Crack ^f	NA	NA	NA	NA	NA	NA	NA	3.3	3.4	2.4	1.4	1.5	1.7	1.3	1.0	1.8	1.2	1.4	2.2	2.4	2.5	2.0	1.9	3.1	2.0	-1.1
Heroin	0.9	0.6	0.5	0.3	0.5	0.4	0.4	0.6	0.3	0.7	0.3	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9	1.7	1.2	1.0	1.0	0.9	-0.1
Other Narcotics ^{g,h}	8.9	8.3	8.1	8.4	8.9	6.3	8.8	7.6	6.3	7.6	6.8	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7	8.9	11.0	12.2	14.2	13.8	-0.4
Amphetamines ^g	29.5	29.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Amphetamines, Adj. ^{g,i}	NA	NA	30.1	27.8	27.8	25.4	22.3	19.8	17.7	14.6	13.2	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9	12.3	12.4	11.9	12.3	12.7	+0.4
Crystal Meth. (Ice) ^j	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.0	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8	1.3	2.3	2.0	2.9	2.2	-0.7
Sedatives (Barbiturates) ^g	8.1	7.8	8.2	6.6	6.4	4.9	5.4	3.5	3.6	3.2	3.8	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7	6.9	6.0	5.9	5.7	7.2	+1.5
Sedatives, Adj. ^{g,k}	13.7	14.2	14.1	12.2	10.8	9.3	8.0	6.1	4.7	4.1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Methaqualone ^g	10.3	10.4	11.1	9.2	9.0	7.2	5.8	4.1	2.2	2.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Tranquilizers ^{d,g}	15.2	11.4	11.7	10.8	10.8	9.8	10.7	8.7	8.0	8.0	7.1	6.8	6.9	6.3	4.4	5.4	5.4	6.9	7.7	8.2	8.8	9.7	10.7	11.0	10.6	-0.4
Alcohol ^l	94.3	95.2	95.2	95.0	94.2	95.3	94.9	94.1	94.9	93.7	93.1	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0	86.6	86.1	86.0	86.2	84.6	-1.7
Flvd. Alcoholic Bvg. ^m	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	79.0	—
Cigarettes	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. 'NA' indicates data not available.

See footnotes on next page.

FOOTNOTES FOR TABLES 9-1 THROUGH 9-5

^a"Any illicit drug" includes use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in four of the five questionnaire forms in 1980-1989, in five of the six forms in 1990-1998, and in three of the six forms in 1999-2004. Total N in 2004 (for college students) is 700.

^cUnadjusted for known underreporting of certain drugs. See text for details.

^dIn 2001 the question text was changed on half the questionnaire forms. "Other psychedelics" was changed to "other hallucinogens," and "shrooms" was added to the list of examples. For tranquilizers, "Miltown" was replaced with "Xanax" in the list of examples. Beginning in 2002 the remaining forms were changed to the new wording.

^eThis drug was asked about in two of the five questionnaire forms in 1989, in two of the six questionnaire forms in 1990-2001, and in three of the six questionnaire forms in 2002-2004. Total N in 2004 (for college students) is 700.

^fThis drug was asked about in two of the five questionnaire forms in 1987-1989, in all six questionnaire forms in 1990-2001, and in five of the six questionnaire forms in 2002-2004. Total N in 2004 (for college students) is 1160.

^gOnly drug use that was not under a doctor's orders is included here.

^hIn 2002 the question text was changed on half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric--all of which had negligible rates of use by 2001--were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one-half of N indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003-2004.

ⁱBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^jThis drug was asked about in two of the six questionnaire forms. Total N in 2004 (for college students) is 470.

^k"Sedatives, adjusted" data are a combination of barbiturate and methaqualone data.

^lIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a "drink" meant "more than just a few sips." Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

^mThis drug was asked about in one of the six questionnaire forms. Total N in 2004 (for college students) is 230.

ⁿRevised questions about amphetamine use were introduced in 1982 to exclude more completely inappropriate reporting of nonprescription amphetamines. The data in italics are therefore not strictly comparable to the other data.

TABLE 9-2
Trends in Annual Prevalence of Various Types of Drugs
Among College Students 1-4 Years Beyond High School

Percentage who used in past year

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change	
<i>Approx. Wtd. N =</i>	1040	1130	1150	1170	1110	1080	1190	1220	1310	1300	1400	1410	1490	1490	1410	1450	1450	1480	1440	1440	1350	1340	1260	1270	1400		
Any Illicit Drug ^a	56.2	55.0	49.5	49.8	45.1	46.3	45.0	40.1	37.4	36.7	33.3	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	-0.3	
Any Illicit Drug ^a																											
Other Than Marijuana	32.3	31.7	29.9	29.9	27.2	26.7	25.0	21.3	19.2	16.4	15.2	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6	16.4	16.6	17.9	18.6	+0.6	
Marijuana	51.2	51.3	44.7	45.2	40.7	41.7	40.9	37.0	34.6	33.6	29.4	26.5	27.7	27.9	29.3	31.2	33.1	31.6	35.9	35.2	34.0	35.6	34.7	33.7	33.3	-0.4	
Inhalants ^{b,c}	3.0	2.5	2.5	2.8	2.4	3.1	3.9	3.7	4.1	3.7	3.9	3.5	3.1	3.8	3.0	3.9	3.6	4.1	3.0	3.2	2.9	2.8	2.0	1.8	2.7	+0.9	
Hallucinogens ^{c,d}	8.5	7.0	8.7	6.5	6.2	5.0	6.0	5.9	5.3	5.1	5.4	6.3	6.8	6.0	6.2	8.2	6.9	7.7	7.2	7.8	6.7	7.5	6.3	7.4	5.9	-1.4	
LSD	6.0	4.6	6.3	4.3	3.7	2.2	3.9	4.0	3.6	3.4	4.3	5.1	5.7	5.1	5.2	6.9	5.2	5.0	4.4	5.4	4.3	4.0	2.1	1.4	1.2	-0.1	
MDMA (Ecstasy) ^e	NA	NA	NA	NA	NA	NA	NA	NA	NA	2.3	2.3	0.9	2.0	0.8	0.5	2.4	2.8	2.4	3.9	5.5	9.1	9.2	6.8	4.4	2.2	-2.2 s	
Cocaine	16.8	16.0	17.2	17.3	16.3	17.3	17.1	13.7	10.0	8.2	5.6	3.6	3.0	2.7	2.0	3.6	2.9	3.4	4.6	4.6	4.8	4.7	4.8	5.4	6.6	+1.1	
Crack ^f	NA	NA	NA	NA	NA	NA	1.3	2.0	1.4	1.5	0.6	0.5	0.4	0.6	0.5	1.1	0.6	0.4	1.0	0.9	0.9	0.9	0.4	1.3	1.3	0.0	
Heroin	0.4	0.2	0.1	*	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.4	0.3	0.6	0.2	0.5	0.4	0.1	0.2	0.4	+0.2	
Other Narcotics ^{g,h}	5.1	4.3	3.8	3.8	3.8	2.4	4.0	3.1	3.1	3.2	2.9	2.7	2.7	2.5	2.4	3.8	3.1	4.2	4.2	4.3	4.5	5.7	7.4	8.7	8.2	-0.5	
OxyContin ⁱ	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.5	2.2	2.5	+0.3	
Vicodin ^j	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	6.9	7.5	7.4	0.0	
Amphetamines ^g	22.4	22.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Amphetamines, Adj. ^{g,i}	NA	NA	21.1	17.3	15.7	11.9	10.3	7.2	6.2	4.6	4.5	3.9	3.6	4.2	4.2	5.4	4.2	5.7	5.1	5.8	6.6	7.2	7.0	7.1	7.0	-0.2	
Ritalin ^j	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	5.7	4.7	4.7	0.0	
Methamphetamine ^l	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	3.3	1.6	2.4	1.2	2.6	2.9	+0.3	
Crystal Meth. (Ice) ^j	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.1	0.1	0.2	0.7	0.8	1.1	0.4	0.8	1.0	0.5	0.5	0.6	0.8	0.9	1.1	+0.2	
Sedatives (Barbiturates) ^g	2.9	2.8	3.2	2.2	1.9	1.3	2.0	1.2	1.1	1.0	1.4	1.2	1.4	1.5	1.2	2.0	2.3	3.0	2.5	3.2	3.7	3.8	3.7	4.1	4.2	+0.1	
Sedatives, Adj. ^{g,k}	8.3	8.0	8.0	4.5	3.5	2.5	2.6	1.7	1.5	1.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Methaqualone ^g	7.2	6.5	6.6	3.1	2.5	1.4	1.2	0.8	0.5	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Tranquilizers ^{d,g}	6.9	4.8	4.7	4.6	3.5	3.6	4.4	3.8	3.1	2.6	3.0	2.4	2.9	2.4	1.8	2.9	2.8	3.8	3.9	3.8	4.2	5.1	6.7	6.9	6.7	-0.2	
Rohypno ^l	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.7	0.4	0.3	-0.1	
GHB ^l	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.6	0.3	0.7	+0.4	
Ketamine ^l	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	1.3	1.0	1.5	+0.5	
Alcohol ^l	90.5	92.5	92.2	91.6	90.0	92.0	91.5	90.9	89.6	89.6	89.0	88.3	86.9	85.1	82.7	83.2	83.0	82.4	84.6	83.6	83.2	83.0	82.9	81.7	81.2	-0.5	
Flvd. Alcoholic Bvg. ^m	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	63.2	—	
Cigarettes	36.2	37.6	34.3	36.1	33.2	35.0	35.3	38.0	36.6	34.2	35.5	35.6	37.3	38.8	37.6	39.3	41.4	43.6	44.3	44.5	41.3	39.0	38.3	35.2	36.7	+1.5	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '*' indicates a percentage of less than 0.05% but greater than true zero. 'NA' indicates data not available.

See footnotes at end of Table 9-1.

TABLE 9-3
Trends in Thirty-Day Prevalence of Various Types of Drugs
Among College Students 1-4 Years Beyond High School

Percentage who used in last 30 days

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change	
<i>Approx. Wtd. N =</i>	1040	1130	1150	1170	1110	1080	1190	1220	1310	1300	1400	1410	1490	1490	1410	1450	1450	1480	1440	1440	1350	1340	1260	1270	1400		
Any Illicit Drug ^a	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	-0.2	
Any Illicit Drug ^a																											
Other Than Marijuana	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9	7.5	7.8	8.2	9.1	+0.9	
Marijuana	34.0	33.2	26.8	26.2	23.0	23.6	22.3	20.3	16.8	16.3	14.0	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	20.0	20.2	19.7	19.3	18.9	-0.4	
Inhalants ^{b,c}	1.5	0.9	0.8	0.7	0.7	1.0	1.1	0.9	1.3	0.8	1.0	0.9	1.1	1.3	0.6	1.6	0.8	0.7	0.6	1.5	0.9	0.4	0.7	0.4	0.4	0.0	
Hallucinogens ^{c,d}	2.7	2.3	2.6	1.8	1.8	1.3	2.2	2.0	1.7	2.3	1.4	1.2	2.3	2.5	2.1	3.3	1.9	2.1	2.1	2.0	1.4	1.8	1.2	1.8	1.3	-0.4	
LSD	1.4	1.4	1.7	0.9	0.8	0.7	1.4	1.4	1.1	1.4	1.1	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	0.9	1.0	0.2	0.2	0.2	-0.1	
MDMA (Ecstasy) ^e	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.3	0.6	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	2.5	1.5	0.7	1.0	0.7	-0.3	
Cocaine	6.9	7.3	7.9	6.5	7.6	6.9	7.0	4.6	4.2	2.8	1.2	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	1.4	1.9	1.6	1.9	2.4	+0.5	
Crack ^f	NA	NA	NA	NA	NA	NA	NA	0.4	0.5	0.2	0.1	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.1	0.3	0.4	0.4	0.0	
Heroin	0.3	0.0	0.0	0.0	*	*	0.0	0.1	0.1	0.1	0.0	0.1	0.0	*	0.0	0.1	*	0.2	0.1	0.1	0.2	0.1	0.0	*	0.1	+0.1	
Other Narcotics ^{g,h}	1.8	1.1	0.9	1.1	1.4	0.7	0.6	0.8	0.8	0.7	0.5	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	1.7	1.7	3.2	2.3	3.0	+0.7	
Amphetamines ^g	13.4	12.3	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Amphetamines, Adj. ^{g,i}	NA	NA	9.9	7.0	5.5	4.2	3.7	2.3	1.8	1.3	1.4	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	2.9	3.3	3.0	3.1	3.2	+0.1	
Crystal Meth. (Ice) ^j	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	0.0	0.0	0.0	0.3	0.5	0.3	0.1	0.2	0.3	0.0	0.0	0.1	0.0	0.3	0.1	-0.2	
Sedatives (Barbiturates) ^g	0.9	0.8	1.0	0.5	0.7	0.4	0.6	0.5	0.5	0.2	0.2	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	1.1	1.5	1.7	1.7	1.5	-0.3	
Sedatives, Adj. ^{g,k}	3.8	3.4	2.5	1.1	1.0	0.7	0.6	0.6	0.6	0.2	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Methaqualone ^g	3.1	3.0	1.9	0.7	0.5	0.3	0.1	0.2	0.1	0.0	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—	
Tranquilizers ^{l,g}	2.0	1.4	1.4	1.2	1.1	1.4	1.9	1.0	1.1	0.8	0.5	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	2.0	1.5	3.0	2.8	2.7	0.0	
Alcohol ^l	81.8	81.9	82.8	80.3	79.1	80.3	79.7	78.4	77.0	76.2	74.5	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	67.4	67.0	68.9	66.2	67.7	+1.5	
Flvd. Alcoholic Bvg. ^m	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	34.0	—	
Cigarettes	25.8	25.9	24.4	24.7	21.5	22.4	22.4	24.0	22.6	21.1	21.5	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	28.2	25.7	26.7	22.5	24.3	+1.9	

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '*' indicates a percentage of less than 0.05% but greater than true zero. 'NA' indicates data not available.

See footnotes at end of Table 9-1.

TABLE 9-4
Trends in Thirty-Day Prevalence of Daily Use of Various Types of Drugs
Among College Students 1-4 Years Beyond High School

	Percentage who used daily in last 30 days																									'03-'04
	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	change
<i>Approx. Wtd. N =</i>	1040	1130	1150	1170	1110	1080	1190	1220	1310	1300	1400	1410	1490	1490	1410	1450	1450	1480	1440	1440	1350	1340	1260	1270	1400	
Marijuana	7.2	5.6	4.2	3.8	3.6	3.1	2.1	2.3	1.8	2.6	1.7	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	4.6	4.5	4.1	4.7	4.5	-0.2
Cocaine	0.2	0.0	0.3	0.1	0.4	0.1	0.1	0.1	0.1	*	0.0	*	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	0.0	0.0
Amphetamines ^g	0.5	0.4	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	—
Amphetamines, Adj. ^{g,i}	NA	NA	0.3	0.2	0.2	*	0.1	0.1	*	*	0.0	0.1	0.0	0.1	0.1	0.1	*	0.2	0.1	0.1	0.1	0.2	0.1	0.3	0.2	-0.2
Alcohol ^k																										
Daily	6.5	5.5	6.1	6.1	6.6	5.0	4.6	6.0	4.9	4.0	3.8	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	3.6	4.7	5.0	4.3	3.7	-0.5
5+ Drinks in a Row in Last 2 Weeks	43.9	43.6	44.0	43.1	45.4	44.6	45.0	42.8	43.2	41.7	41.0	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	39.3	40.9	40.1	38.5	41.7	+3.3
Cigarettes																										
Daily	18.3	17.1	16.2	15.3	14.7	14.2	12.7	13.9	12.4	12.2	12.1	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	17.8	15.0	15.9	13.8	13.8	-0.1
Half-Pack or More per Day	12.7	11.9	10.5	9.6	10.2	9.4	8.3	8.2	7.3	6.7	8.2	8.0	8.9	8.9	8.0	10.2	8.5	9.1	11.3	11.0	10.1	7.8	7.9	7.6	6.8	-0.8

Source: The Monitoring the Future Study, the University of Michigan.

NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '*' indicates a percentage of less than 0.05% but greater than true zero. 'NA' indicates data not available.

See footnotes at end of Table 9-1.

TABLE 9-5
Trends in Lifetime, Annual, and Thirty-Day Prevalence of an Illicit Drug Use Index^a
Among College Students 1-4 Years Beyond High School, by Gender


	<i>1980ⁿ</i>	<i>1981ⁿ</i>	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	'03-'04 change
Lifetime																										
Any Illicit Drug	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	-1.7	
Males	71.0	67.5	68.1	71.3	66.4	69.8	64.7	63.5	56.0	56.5	52.5	51.3	50.8	45.7	49.5	47.3	50.3	52.1	54.4	58.4	54.4	53.9	54.3	54.1	54.9	+0.8
Females	67.5	66.3	61.5	63.0	59.2	61.6	59.4	57.4	60.2	54.9	55.1	49.7	47.1	46.0	42.6	44.3	45.6	46.7	52.0	49.6	53.2	53.5	50.2	53.7	50.6	-3.1
Any Illicit Drug																										
Other Than Marijuana	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	+0.5
Males	42.8	39.8	45.1	44.6	40.9	42.1	38.2	37.2	31.8	30.6	26.2	27.6	26.3	24.3	24.6	26.6	25.0	27.3	27.3	29.4	28.9	27.0	30.4	27.6	31.1	+3.5
Females	41.6	42.6	34.7	39.2	36.4	38.3	37.0	34.6	34.6	30.4	30.1	24.3	26.1	24.3	20.1	22.9	21.2	22.2	23.3	22.8	23.5	25.9	24.6	27.5	26.2	-1.3
Past 12 months																										
Any Illicit Drug	56.2	55.0	49.5	49.8	45.1	46.3	45.0	40.1	37.4	36.7	33.3	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	-0.3
Males	58.9	56.2	54.6	53.4	48.4	50.9	49.8	43.3	37.0	38.2	34.2	30.2	32.8	32.6	33.9	36.1	36.6	38.3	40.1	42.5	38.0	38.8	39.5	39.2	40.9	+1.7
Females	53.3	54.0	44.9	46.7	41.9	42.7	41.1	37.7	37.6	35.4	32.5	28.4	28.7	29.1	29.5	31.7	32.7	31.1	36.4	33.2	34.7	37.3	35.4	34.8	33.4	-1.3
Any Illicit Drug																										
Other Than Marijuana	32.3	31.7	29.9	29.9	27.2	26.7	25.0	21.3	19.2	16.4	15.2	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6	16.4	16.6	17.9	18.6	+0.6
Males	33.7	32.8	33.4	33.5	29.2	29.7	28.6	23.5	19.4	18.7	15.7	14.4	13.8	15.0	14.9	19.5	15.1	18.1	17.0	19.0	18.6	17.2	19.2	19.3	22.1	+2.8
Females	31.1	30.8	26.9	26.8	25.2	24.4	22.1	19.6	19.0	14.6	14.8	12.1	12.6	10.5	10.2	13.3	11.3	14.1	12.1	12.8	13.5	15.8	15.0	17.1	16.5	-0.5
Past 30 days																										
Any Illicit Drug	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	-0.2
Males	42.9	40.6	37.7	33.8	30.4	29.9	31.0	24.0	18.8	20.0	18.2	16.0	18.0	16.0	20.5	23.7	20.6	23.4	23.1	26.7	24.0	25.0	25.1	22.8	26.1	+3.3
Females	34.0	34.8	25.6	25.5	23.7	23.2	21.7	21.1	18.3	16.7	12.7	14.6	14.5	14.5	12.7	15.7	15.8	16.2	17.6	18.1	19.6	19.8	19.3	20.5	18.4	-2.2
Any Illicit Drug																										
Other Than Marijuana	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9	7.5	7.8	8.2	9.1	+0.9
Males	22.8	18.6	20.2	16.0	16.1	12.6	14.4	9.0	8.2	8.0	4.9	4.8	5.1	7.3	6.2	8.8	6.1	7.8	8.6	7.5	8.2	9.0	8.4	8.1	11.3	+3.2
Females	18.7	18.5	14.2	12.1	11.5	11.2	9.3	8.5	8.8	6.0	4.0	3.9	4.2	3.8	3.4	4.5	3.4	6.1	4.6	5.6	6.0	6.4	7.4	8.3	7.8	-0.5
Approximate Weighted N																										
All Respondents	1040	1130	1150	1170	1110	1080	1190	1220	1310	1300	1400	1410	1490	1410	1450	1450	1480	1440	1440	1350	1340	1260	1270	1400		
Males	520	530	550	550	540	490	540	520	560	580	620	640	680	660	590	610	560	630	570	590	560	540	490	480	520	
Females	520	600	610	620	570	600	650	700	750	720	780	770	810	830	820	840	890	860	880	850	790	800	770	790	880	

Source: The Monitoring the Future Study, the University of Michigan.


NOTES: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change and prevalence estimates for the two most recent years is due to rounding.

See footnotes at end of Table 9-1.

FIGURE 9-1
Any Illicit Drug: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School




Any Illicit Drug: Trends in Annual Prevalence
Among Male vs. Female College Students



NOTE: "Others" refers to high school graduates 1-4 years beyond high school not currently enrolled full-time in college.

FIGURE 9-2

**Any Illicit Drug Other Than Marijuana: Trends in Annual Prevalence
Among College Students vs. Others
1-4 Years Beyond High School**



**Any Illicit Drug Other Than Marijuana: Trends in Annual Prevalence
Among Male vs. Female College Students**





FIGURE 9-3a
Marijuana: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Marijuana: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-3b
Marijuana: Trends in Thirty-Day Prevalence of Daily Use
Among College Students vs. Others
 1-4 Years Beyond High School



Marijuana: Trends in Thirty-Day Prevalence of Daily Use
Among Male vs. Female College Students






FIGURE 9-4
Inhalants:* Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School




Inhalants:* Trends in Annual Prevalence
Among Male vs. Female College Students




*Unadjusted for the possible underreporting of amyl and butyl nitrites.

FIGURE 9-5
Hallucinogens:* Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School




Hallucinogens:* Trends in Annual Prevalence
Among Male vs. Female College Students



*Unadjusted for the possible underreporting of PCP.

FIGURE 9-6
LSD: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



LSD: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-7
Hallucinogens Other Than LSD: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Hallucinogens Other Than LSD: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-8
MDMA (Ecstasy): Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



MDMA (Ecstasy): Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-9
Cocaine: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Cocaine: Trends in Annual Prevalence
Among Male vs. Female College Students






FIGURE 9-10a
Narcotics Other Than Heroin:* Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Narcotics Other Than Heroin:* Trends in Annual Prevalence
Among Male vs. Female College Students



*In 2002 the question text was changed on half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only. In 2003 the remaining forms were changed to the new wording.

FIGURE 9-10b
Vicodin: Trends in Annual Prevalence
Among College Students vs. Others
1-4 Years Beyond High School




FIGURE 9-10c
OxyContin: Trends in Annual Prevalence
Among College Students vs. Others
1-4 Years Beyond High School





FIGURE 9-11
Amphetamines: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Amphetamines: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-12
Sedatives (Barbiturates): Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Sedatives (Barbiturates): Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-13
Tranquilizers: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Tranquilizers: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-14a
Alcohol: Trends in Annual Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Alcohol: Trends in Annual Prevalence
Among Male vs. Female College Students





FIGURE 9-14b
Alcohol: Trends in Thirty-Day Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Alcohol: Trends in Thirty-Day Prevalence
Among Male vs. Female College Students





FIGURE 9-14c
Alcohol: Trends in Thirty-Day Prevalence of Daily Use
Among College Students vs. Others
 1-4 Years Beyond High School



Alcohol: Trends in Thirty-Day Prevalence of Daily Use
Among Male vs. Female College Students





FIGURE 9-14d

**Alcohol: Trends in Two-Week Prevalence of Five or More Drinks in a Row
Among College Students vs. Others
1-4 Years Beyond High School**



**Alcohol: Trends in Two-Week Prevalence of Five or More Drinks in a Row
Among Male vs. Female College Students**





FIGURE 9-15a
Cigarettes: Trends in Thirty-Day Prevalence
Among College Students vs. Others
 1-4 Years Beyond High School



Cigarettes: Trends in Thirty-Day Prevalence
Among Male vs. Female College Students





FIGURE 9-15b
Cigarettes: Trends in Thirty-Day Prevalence of Daily Use
Among College Students vs. Others
 1-4 Years Beyond High School



Cigarettes: Trends in Thirty-Day Prevalence of Daily Use
Among Male vs. Female College Students






FIGURE 9-15c

**Cigarettes: Trends in Thirty-Day Prevalence of Smoking a Half-Pack or More per Day Among College Students vs. Others
1-4 Years Beyond High School**



Cigarettes: Trends in Thirty-Day Prevalence of Smoking a Half-Pack or More per Day Among Male vs. Female College Students



INDEX

A

alcohol

daily use

- college enrollment, 22, 23, 227, 240, table 8-4, fig. 9-14c
- disapproval, 179, 180, table 6-2
- friends' attitudes, perceived, 190, table 7-1
- gender, 22, 83, 132, 243, table 4-1, table 4-5, fig. 9-14c
- harmfulness, perceived, 177, table 6-1
- population density (urbanicity), 86, 140, table 4-5
- prevalence, 30, 82, table 4-1, table 4-5, fig. 4-20b
- region, 136, table 4-5
- trends, 21, 22, 23, 126, table 2-4, table 5-4, table 9-4, fig. 5-18c, fig. 9-14c

disapproval, 179, 180, table 6-2

drunk, getting

- friends' use, 194, table 7-2
- prevalence, 29
- race/ethnicity, 28
- trends, tables 2-1-2-4

exposure to use, table 7-3 (*see also* friends' use)

flavored alcoholic beverages

- college enrollment, 227, tables 8-1-8-3, tables 9-1-9-3
- gender, 228
- prevalence, 227, tables 4-1-4-4
- trends, tables 2-1-2-3, tables 5-1-5-3, tables 9-1-9-3

friends' use, 196, table 7-2

harmfulness, perceived, table 6-1

heavy drinking

- college enrollment, 22, 227, 229, 239, 240, table 8-4, fig. 9-14d
- disapproval, 179, 180, 189, table 6-2
- friends' attitudes, perceived, 189, 190, 191, table 7-1
- gender, 23, 83, 132, 228, 243, table 4-1, table 4-5, fig. 9-14d
- harmfulness, perceived, 176, 177, table 6-1
- population density (urbanicity), 86, 140, table 4-5
- prevalence, 21, 30, 33, 78, 82, table 4-1, table 4-5, fig. 4-20b
- race/ethnicity, 27, 28
- region, 85, 136, table 4-5
- trends, 21, 125, 126, table 2-4, fig. 5-18d, fig. 9-14d

- prevalence, 21, 29, 30, 78, 81, 82, tables 4-1-4-4, fig. 4-20a
- college enrollment, 226, 227, 229, tables 8-1-8-3
- gender, tables 4-1-4-4
- population density (urbanicity), 86, tables 4-2-4-4
- race/ethnicity, 28
- region, 85, tables 4-2-4-4
- trends, 21, 125, tables 2-1-2-3, tables 5-1-5-3, fig. 5-18a-5-18b
- college enrollment, 239, tables 9-1-9-3, fig. 9-14a-9-14b
- gender, 131, 243, fig. 9-14a-9-14b
- population density (urbanicity), 139, 140
- region, 135

amphetamines

- availability, perceived, 197, 200, table 7-4
- daily use, table 5-4, table 8-4, table 9-4
- disapproval, 12, 178, 179, table 6-2
- exposure to use, table 7-3 (*see also* friends' use)
- friends' attitudes, perceived, 190, 191, table 7-1
- friends' use, 193, 195, table 7-2
- harmfulness, perceived, 12, 174, 180, table 6-1
- prevalence, 27, 29, 30, 79, 80, tables 4-1-4-4, fig. 4-4
- college enrollment, 19, 227, tables 8-1-8-4
- gender, 21, 84, 229, tables 4-1-4-4
- population density (urbanicity), tables 4-2-4-4
- race/ethnicity, 28
- region, tables 4-2-4-4
- trends, 11, 12, 17, 18, 120, 122, 123, tables 2-1-2-3, tables 5-1-5-3, fig. 5-13
- college enrollment, 20, 237, 241, tables 9-1-9-4, fig. 9-11
- gender, 131, 243, fig. 9-11
- population density (urbanicity), 138
- race/ethnicity, 28
- region, 133, 134

androstenedione, 18, 19

any illicit drug

- exposure to use, 194, 195, table 7-3 (*see also* friends' use)
- friends' use, 195, table 7-2
- prevalence, 6, 32, 77, 78, tables 4-1-4-4, fig. 4-1
- college enrollment, 19, 226, tables 8-1-8-3, table 8-5
- gender, 83, 228, tables 4-1-4-4
- population density (urbanicity), 86, tables 4-2-4-4

- any illicit drug, prevalence (*continued*)
region, 84, tables 4-2-4-4
trends, 7, 120, 122, tables 2-1-2-3, tables 5-1-5-3, table 5-5, fig. 2-1, fig. 5-1
college enrollment, 236, tables 9-1-9-3, table 9-5, fig. 9-1
gender, 128, fig. 9-1
population density (urbanicity), 136
region, 133
- any illicit drug other than marijuana
disapproval, 178
exposure to use, 194, table 7-3 (*see also* friends' use)
friends' attitudes, perceived, 190
friends' use, 192, 193, 195, table 7-2
prevalence, 6, 30, 32, 77, 78, 79, tables 4-1-4-4, fig. 4-2
college enrollment, 19, 226, tables 8-1-8-3, table 8-5
gender, 21, 228, tables 4-1-4-4
population density (urbanicity), tables 4-2-4-4
region, 84, tables 4-2-4-4
trends, 6, 7, 9, 31, 120, 121, 122, tables 2-1-2-3, tables 5-1-5-3, table 5-5, fig. 5-2
college enrollment, 20, 236, tables 9-1-9-3, table 9-5, fig. 9-2
gender, 128, 129, fig. 9-2
population density (urbanicity), 136
region, 133
- availability, perceived, table 7-4
- B**
barbiturates. *See* sedatives
- C**
cigarettes
bidis, table 2-2
daily use
college enrollment, 24, 26, 241, table 8-4, fig. 9-15b-9-15c
disapproval, 180, table 6-2
friends' attitudes, perceived, 189, 190, 192, table 7-1
gender, 132, 227, 228, table 4-1, table 4-5, fig. 9-15b-9-15c
harmfulness, perceived, 24, 29, 177, table 6-1
population density (urbanicity), 86, table 4-5
prevalence, 31, 33, table 4-1, table 4-5
region, table 4-5
trends, 24, 25, table 2-4, table 5-4, table 9-4, fig. 5-19b-5-19c
disapproval, 24, 180, 189
friends' use, 194, 196, table 7-2
- harmfulness, perceived, 24, 29
kreteks, table 2-2
prevalence, 29, 31, 33, 82, tables 4-1-4-4, fig. 4-21
college enrollment, 227, 228, tables 8-1-8-3
gender, 84, 228, tables 4-1-4-4
population density (urbanicity), tables 4-2-4-4
race/ethnicity, 28
region, 85, tables 4-2-4-4
trends, 7, 8, 23, 24, 25, 31, 126, 127, tables 2-1-2-4, tables 5-1-5-3, fig. 5-19a
college enrollment, 26, 240, 241, tables 9-1-9-3, fig. 9-15a
gender, 26, 27, 132, 133, 243, fig. 9-15a
population density (urbanicity), 140
race/ethnicity, 27, 28
region, 136
- club drugs. *See* GHB; ketamine; Rohypnol
- cocaine
availability, perceived, 14, 197, 198, 199, table 7-4
college enrollment, table 8-4
daily use, table 5-4, table 9-4
disapproval, 14, 178, 179, table 6-2
exposure to use, table 7-3 (*see also* friends' use)
friends' attitudes, perceived, 190, 191, table 7-1
friends' use, 193, 195, table 7-2
harmfulness, perceived, 14, 174, 175, 180, table 6-1
prevalence, 30, 33, 77, 78, 79, 81, tables 4-1-4-4, fig. 4-6
college enrollment, 19, 227, tables 8-1-8-4
gender, 21, 83, 229, tables 4-1-4-4
population density (urbanicity), tables 4-2-4-4
region, tables 4-2-4-4
trends, 13, 14, 17, 18, 31, 122, tables 2-1-2-3, tables 5-1-5-3, fig. 5-9
college enrollment, 20, 239, tables 9-1-9-3, fig. 9-9
gender, 129, 242, fig. 9-9
population density (urbanicity), 138
race/ethnicity, 28
region, 133, 134
See also cocaine powder; crack cocaine
- cocaine, other. *See* cocaine powder
- cocaine powder
availability, perceived, 14, 197, table 7-4
disapproval, 14
friends' use, 193, table 7-2
harmfulness, perceived, 14, 174, table 6-1
prevalence, 29, 30, 79, table 4-1, fig. 4-8
gender, 83, table 4-1
race/ethnicity, 27
trends, 31, 120, tables 2-1-2-3, tables 5-1-5-3

- codeine, 16
- crack cocaine
- availability, perceived, 197, 199, table 7-4
 - disapproval, 13
 - friends' use, 193, 195, table 7-2
 - harmfulness, perceived, 13, 174, 175, 180, table 6-1
 - prevalence, 29, 30, 33, 79, 81, tables 4-1-4-4, fig. 4-7
 - college enrollment, 19, 227, tables 8-1-8-3
 - gender, 21, 83, tables 4-1-4-4
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28
 - region, tables 4-2-4-4
 - trends, 12, 13, 31, 120, 122, tables 2-1-2-3, tables 5-1-5-3, fig. 5-10
 - college enrollment, tables 9-1-9-3
 - gender, 129, 130
 - population density (urbanicity), 138
 - region, 133, 134
- See also* cocaine
- creatine, 19
- crystal methamphetamine (ice)
- availability, perceived, 198, 200, table 7-4
 - harmfulness, perceived, 175, 180, table 6-1
 - prevalence, 79, 80, tables 4-1-4-4, fig. 4-17
 - college enrollment, 19, 227, tables 8-1-8-3
 - gender, 84, tables 4-1-4-4
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28
 - region, 85, tables 4-2-4-4
 - trends, 12, 122, 124, tables 2-1-2-3, tables 5-1-5-3, fig. 5-14
 - college enrollment, tables 9-1-9-3
 - gender, 131
 - population density (urbanicity), 138
 - region, 134, 135
- D**
- diet pills, 19
- disapproval of use, table 6-2
- E**
- ecstasy. *See* MDMA
- exposure to drug use, table 7-3. *See also* friends' use
- F**
- friends' attitudes, perceived, table 7-1
- friends' use, table 7-2
- G**
- GHB
- prevalence, 18, table 4-3
 - college enrollment, 19, 226, 229, table 8-2
 - gender, 21, 229, table 4-3, table 8-2
 - population density (urbanicity), table 4-3
 - region, table 4-3
 - trends, 18, 34, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 131
 - region, 135
- H**
- hallucinogens
- prevalence, 30, 79, tables 4-1-4-4, fig. 4-9
 - college enrollment, 19, 227, tables 8-1-8-3
 - gender, 228, 229, tables 4-1-4-4
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28
 - region, 85, tables 4-2-4-4
 - trends, 120, 122, tables 2-1-2-3, tables 5-1-5-3, fig. 5-5
 - college enrollment, tables 9-1-9-3, fig. 9-5
 - gender, fig. 9-5
- hallucinogens other than LSD
- availability, perceived, 198, 199, table 7-4
 - exposure to use, table 7-3 (*see also* friends' use)
 - friends' use, 193, table 7-2
 - prevalence, 29, 30, fig. 4-11
 - race/ethnicity, 27, 28
 - trends, 31, tables 2-1-2-3, fig. 5-7
 - college enrollment, fig. 9-7
 - gender, fig. 9-7
 - population density (urbanicity), 137
- harmfulness, perceived, table 6-1
- hashish. *See* marijuana
- heroin (including with and without a needle)
- availability, perceived, 198, 199, 200, table 7-4
 - disapproval, 178, table 6-2
 - exposure to use, table 7-3 (*see also* friends' use)
 - friends' use, 193, table 7-2
 - harmfulness, perceived, 15, 16, 174, 176, 180, table 6-1
 - prevalence, 29, 30, 79, tables 4-1-4-4, fig. 4-19
 - college enrollment, 19, 227, tables 8-1-8-3
 - gender, 21, tables 4-1-4-4
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28
 - region, tables 4-2-4-4
 - trends, 15, 33, 124, tables 2-1-2-3, tables 5-1-5-3, fig. 5-11
 - college enrollment, tables 9-1-9-3
 - gender, 130
 - population density (urbanicity), 139
 - region, 135
- I**
- ice. *See* crystal methamphetamine

inhalants

- friends' use, 193, table 7-2
- prevalence, 29, 79, tables 4-1-4-4, fig. 4-12
 - college enrollment, 19, 226, tables 8-1-8-3
 - gender, 21, 83, 229, tables 4-1-4-4
 - population density (urbanicity), tables 4-2-4-4
- race/ethnicity, 27, 28
- region, tables 4-2-4-4
- trends, 12, 17, 18, 31, 123, tables 2-1-2-3, tables 5-1-5-3, fig. 5-4
 - college enrollment, tables 9-1-9-3, fig. 9-4
 - gender, 131, fig. 9-4
 - population density (urbanicity), 139
 - region, 134

K

ketamine (Special K)

- prevalence, 18, table 4-3
 - college enrollment, 19, 226, 229, table 8-2
 - gender, 229, table 4-3
 - population density (urbanicity), table 4-3
 - region, table 4-3
- trends, 18, 34, table 2-2, table 5-2
 - college enrollment, table 9-2
 - gender, 131
 - region, 135

L

look-alike drugs, 19

LSD

- availability, perceived, 198, 199, table 7-4
- disapproval, 10, 178, table 6-2
- exposure to use, table 7-3 (*see also* friends' use)
- friends' attitudes, perceived, 190, 191, table 7-1
- friends' use, 193, table 7-2
- harmfulness, perceived, 10, 174, 178, 180, table 6-1
- prevalence, 29, 30, 79, tables 4-1-4-4, fig. 4-10
 - college enrollment, 19, 227, 228, tables 8-1-8-3
 - gender, 21, 228, 229, tables 4-1-4-4, tables 8-1-8-3
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 27, 28
 - region, tables 4-2-4-4
- trends, 9, 10, 17, 18, 31, 33, 121, 122, tables 2-1-2-3, tables 5-1-5-3, fig. 5-6
 - college enrollment, 237, tables 9-1-9-3, fig. 9-6
 - gender, 129, 242, fig. 9-6
 - population density (urbanicity), 137
 - region, 133

M

marijuana (hashish)

- availability, perceived, 197, 198, table 7-4
 - daily use
 - college enrollment, 226, 237, table 8-4, fig. 9-3b
 - gender, 20, 83, 129, 228, 242, table 4-5, fig. 9-3b
 - population density (urbanicity), table 4-5
 - prevalence, 30, 33, 79, table 4-5
 - region, table 4-5
 - trends, 8, 121, 127, table 2-4, table 5-4, table 9-4, fig. 5-3c
 - disapproval, 9, 178, 189, table 6-2
 - exposure to use, 194, table 7-3 (*see also* friends' use)
 - friends' attitudes, perceived, 189, 190, 191, table 7-1
 - friends' use, 193, 194, 195, table 7-2
 - harmfulness, perceived, 8, 9, 173, 174, 180, table 6-1
 - prevalence, 29, 30, 77, 78, 79, tables 4-1-4-4, fig. 4-3
 - college enrollment, 19, 226, tables 8-1-8-4
 - gender, 21, 228, 229, tables 4-1-4-4, tables 8-1-8-4
 - population density (urbanicity), 86, tables 4-2-4-4
 - race/ethnicity, 28
 - region, 84, tables 4-2-4-4
 - trends, 7, 8, 17, 18, 31, 120, 121, 122, tables 2-1-2-3, tables 5-1-5-3, fig. 5-3a-5-3b
 - college enrollment, 20, 236, tables 9-1-9-3, fig. 9-3a
 - gender, 128, 242, fig. 9-3a
 - population density (urbanicity), 137
 - region, 133
- MDMA (ecstasy)
- availability, perceived, 11, 197, 199, table 7-4
 - disapproval, table 6-2
 - friends' use, 193, 196, table 7-2
 - harmfulness, perceived, 174, 175, 176, 180, table 6-1
 - prevalence, 79, 81, tables 4-1-4-4, fig. 4-16
 - college enrollment, 19, 227, tables 8-1-8-3
 - gender, 21, 84, tables 4-1-4-4
 - population density (urbanicity), 86, tables 4-2-4-4
 - race/ethnicity, 27, 28
 - region, 85, tables 4-2-4-4
 - trends, 10, 11, 18, 34, 123, 124, tables 2-1-2-3, tables 5-1-5-3, fig. 5-8
 - college enrollment, 237, tables 9-1-9-3, fig. 9-8
 - gender, 129, fig. 9-8
 - population density (urbanicity), 137, 138

region, 133

methamphetamine

- prevalence, 29, 80, tables 4-2-4-4, fig. 4-5
- college enrollment, 19, 227, table 8-2
- gender, 229, tables 4-2-4-4, table 8-2
- population density (urbanicity), 86, tables 4-2-4-4
- region, 85, tables 4-2-4-4
- trends, 12, tables 2-1-2-3, table 5-2
- college enrollment, table 9-2
- region, 134

methaqualone

- college enrollment, tables 9-1-9-3
- gender, 243
- trends, 17, 122, 139, tables 2-1-2-3, tables 5-1-5-3

N

narcotics other than heroin

- availability, perceived, 198, 200, table 7-4
- exposure to use, table 7-3 (*see also* friends' use)
- friends' use, 193, 196, table 7-2
- prevalence, 30, 79, 80, tables 4-1-4-4, fig. 4-14
- college enrollment, 19, 226, tables 8-1-8-3
- gender, 84, 229, tables 4-1-4-4, tables 8-1-8-3
- population density (urbanicity), tables 4-2-4-4
- race/ethnicity, 28
- region, tables 4-2-4-4
- trends, 16, 17, 18, 120, 122, 124, tables 2-1-2-3, tables 5-1-5-3, fig. 5-12
- college enrollment, 20, 238, tables 9-1-9-3, fig. 9-10a
- gender, 130, fig. 9-10a
- population density (urbanicity), 139
- region, 135

nitrites

- availability, perceived, table 7-4
- friends' use, table 7-2
- trends, 12, tables 2-1-2-3, tables 5-1-5-3
- See also* inhalants

O

OxyContin

- prevalence, table 4-3
- college enrollment, 19, 227, table 8-2
- gender, 84, table 4-3, table 8-2
- population density (urbanicity), table 4-3
- region, 85, table 4-3
- trends, 16, 18, 34, 124, table 2-2, table 5-2
- college enrollment, 238, table 9-2, fig. 9-10c
- gender, 131
- population density (urbanicity), 139
- region, 135

P

PCP

- availability, perceived, table 7-4
- friends' use, table 7-2
- harmfulness, perceived, table 6-1
- prevalence, tables 4-1-4-4
- trends, 14, tables 2-1-2-3, tables 5-1-5-3
- perceived harmfulness. *See* harmfulness, perceived
- perceived risk. *See* harmfulness, perceived
- Percocet, 16
- powder cocaine. *See* cocaine powder
- psychedelics. *See* hallucinogens

Q

qualaludes, table 7-2

R

risk, perceived. *See* harmfulness, perceived

Ritalin

- prevalence, 80, table 4-3
- college enrollment, 19, 226, 227, 229, table 8-2
- gender, 21, table 4-3
- population density (urbanicity), table 4-3
- region, table 4-3
- trends, 12, table 2-2, table 5-2, table 9-2

Rohypnol

- prevalence, 18, 29, table 4-3
- college enrollment, 19, 226, 229, table 8-2
- gender, 21, 229, table 4-3
- population density, table 4-3
- region, table 4-3
- trends, 34, tables 2-1-2-3, table 5-2
- college enrollment, table 9-2
- gender, 131
- region, 135

S

sedatives (barbiturates)

- availability, perceived, 197, 200, table 7-4
- disapproval, 178, 179, table 6-2
- exposure to use, table 7-3 (*see also* friends' use)
- friends' use, 193, 195, table 7-2
- harmfulness, perceived, 174, 180, table 6-1
- prevalence, 29, 30, 79, 80, tables 4-1-4-4, fig. 4-13
- college enrollment, 19, 227, tables 8-1-8-3
- gender, tables 4-1-4-4, tables 8-1-8-3
- population density (urbanicity), tables 4-2-4-4
- race/ethnicity, 28
- region, tables 4-2-4-4
- trends, 17, 120, 125, tables 2-1-2-3, tables 5-1-5-3, fig. 5-15
- college enrollment, 20, 237, 238, 241, tables 9-1-9-3, fig. 9-12

sedatives (barbiturates), trends (*continued*)
gender, 130, fig. 9-12
population density (urbanicity), 139
race/ethnicity, 28
region, 135

smokeless tobacco
harmfulness, perceived, 177, table 6-1
prevalence
gender, 29
race/ethnicity, 28
trends, table 2-1, tables 2-3–2-4

stay-awake pills, 19

steroids
availability, perceived, 198, 200, table 7-4
friends' use, 193, table 7-2
prevalence, 29, 82, 83, tables 4-1–4-4, fig. 4-18
gender, 83, 84, tables 4-1–4-4
population density (urbanicity), tables 4-2–4-4
region, tables 4-2–4-4
trends, 18, 19, 125, tables 2-1–2-3, tables 5-1–5-3, fig. 5-17

stimulants, nonprescription, 19

T

tobacco. *See* cigarettes; smokeless tobacco

tranquilizers
availability, perceived, 197, 200, table 7-4
exposure to use, table 7-3 (*see also* friends' use)
friends' use, 193, 195, table 7-2
prevalence, 29, 30, 79, 80, tables 4-1–4-4, fig. 4-15

college enrollment, 19, 226, tables 8-1–8-3
gender, 21, 229, tables 4-1–4-4
population density (urbanicity), tables 4-2–4-4
race/ethnicity, 28
region, tables 4-2–4-4
trends, 16, 17, 18, 120, 122, 125, tables 2-1–2-3, tables 5-1–5-3, fig. 5-16
college enrollment, 20, 238, 241, tables 9-1–9-3, fig. 9-13
gender, 131, fig. 9-13
population density (urbanicity), 139
race/ethnicity, 27
region, 135

V

Valium, 17

Vicodin

prevalence, table 4-3
college enrollment, 19, 226, table 8-2
gender, 84, 229, table 4-3
population density (urbanicity), table 4-3
region, 85, table 4-3
trends, 16, 18, 124, table 2-2, table 5-2
college enrollment, 238, table 9-2, fig. 9-10b
gender, 131
population density (urbanicity), 139
region, 135

X

Xanax, 16, 17

Study Web site address:
<http://www.monitoringthefuture.org>

NIDA NATIONAL INSTITUTE
ON DRUG ABUSE

NIH Publication No. 05-5728
Printed October 2005