


Climate model suggests extreme El Niño tipping point could be reached if global warming continues

July 15 2024, by Bob Yirka



El Niño-Southern Oscillation teleconnection in PR, here represented as the median of composites of PR in December, January, and February in (a) during El Niño in COLD, in (b) during La Niña in COLD, in (c, d) same as (a, b), but here in CONTROL; in (e, f) same as (a, b) but here in WARM; stippling indicates significant values on a 99% confidence level. Credit: *Geophysical Research Letters* (2024). DOI: 10.1029/2023GL107848

A trio of physicists and oceanologists, two with the University of Cologne's Institute of Geophysics and Meteorology and the third with the GEOMAR Helmholtz Center for Ocean Research Kiel, all in Germany, has found via the CESM1 climate model that an extreme El Niño tipping point could be reached in the coming decades under current emissions.

The [study](#) by Tobias Bayr, Stephanie Fiedler and Joke Lübbecke is published in *Geophysical Research Letters*.

The El Niño-Southern Oscillation (ENSO) is a climate phenomenon in which heat released in parts of the ocean into the atmosphere results in more rainfall in places like the western coast of North and South America and droughts in places like Canada and Africa. Over the past several years, weather watchers have noticed that ENSO events have become more extreme.

Prior research has shown that such extreme events used to occur approximately eight or nine times per century. Some in the field have suggested that rising global temperatures could make them happen more often.

To find out if that might be the case, the researchers gathered historical data, as well as data from other research efforts describing possible weather impacts due to rising [global temperatures](#), and fed it to the CESM1 model then ran it under increased-temperature scenarios. Current estimates suggest that an increase of 2.9°C will occur by 2100 if [greenhouse gas emissions](#) are not stopped.

In their work, the model showed that if temperatures keep rising past 2100 to as much as 3.7°C, a tipping point could be reached where virtually all ENSO events will be extreme. Such a tipping point, they note, suggests that even if mankind was somehow able to stop [climate](#)

[change](#), it would take centuries for the weather to return to what they describe as "normal."

The model also showed that extreme ENSO events would happen more often, as well, perhaps as often as every four years. It also showed changes, such as the Gulf Stream dropping farther south, leading to far less rain in Canada and northern parts of the U.S., and more rain in southern parts of the U.S.

More information: Tobias Bayr et al, Is El Niño-Southern Oscillation a Tipping Element in the Climate System?, *Geophysical Research Letters* (2024). [DOI: 10.1029/2023GL107848](https://doi.org/10.1029/2023GL107848)

© 2024 Science X Network

Citation: Climate model suggests extreme El Niño tipping point could be reached if global warming continues (2024, July 15) retrieved 21 August 2024 from <https://phys.org/news/2024-07-climate-extreme-el-nio-global.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.