

**Research-Based Guidelines and  
Practices for School-Based Suicide  
Prevention**

**Deborah Kimoqueo, EdD  
Teachers College-Columbia University  
2006**

# **Research-Based Guidelines and Practices for School-Based Suicide Prevention**

## **Table of Contents:**

<b>Section I.</b> Background Information and Resources.....	<b>3</b>
<b>Section II.</b> Planning of School-Based Primary Suicide Prevention Program.....	<b>15</b>
<b>Section III.</b> School-Based Secondary Suicide Prevention Program.....	<b>22</b>
<b>Section IV.</b> School-Based Tertiary Suicide Prevention Program.....	<b>23</b>
<b>Section V.</b> School-Based Suicide Prevention Program Evaluation Plan.....	<b>28</b>
<b>Section VI.</b> Useful Sample Forms, Checklists, Job Descriptions & Guidelines.....	<b>29</b>
<b>Section VII. Bibliography.....</b>	<b>70</b>

## **Section I. Background Information and Resources.**

### **A. Statistical research synthesis about youth suicide prevention.**

1. Incidence of suicide among adolescents and young adults nearly tripled between 1952 and 1995 (CDC, 2002).
2. In 1999, more teenagers and young adults died from suicide than from cancer, heart disease, AIDS, birth defects, stroke, and chronic lung disease combined (CDC, 2002).
3. In 2000, suicide was the third leading cause of death among 10- to 14-year-olds and 15- to 19-year-olds in the United States (Anderson, 2002).
4. Teen suicide rates demonstrated a drop of 26 percent between 1993 and 2000 in the year 2000 (Child Trends, 2002), but still ranked as the third leading cause of death for young persons 10-24 years of age.
5. 1.5 per 100,000 children in the 10-14 year age group took their own lives in 2000 (CDC, 2003).
6. Suicide rate of children 10-14 years of age is growing faster than any other demographic, nearly doubling in the past few decades (National Institute of Mental Health, 2000).
7. Teenagers in the 15-19-age bracket took their own lives at the rate of 8.2 per 100,000 in 2000, for a total of 3,994 deaths (Child Trends, 2002).
8. Hispanic students (12.1%) were significantly more likely than black or white students, (8.8% and 7.9%, respectively), to have attempted suicide (Grunbaum, et al., 2002).
9. Female students (11.2%) were significantly more likely to have attempted to kill themselves than male students (6.2%) (Grunbaum, et al., 2002).
9. Nearly five times more 15-19 year old boys commit suicide than females (MacKay et al., 2000).
10. American Indians and Alaska Native adolescents have the highest rates of suicide, being more than twice as likely to commit suicide as other racial/ethnic group (CDC Wonder, 1999; IHS, 1999).
11. In California, 1999-2000, young people, ages 20 to 24 years made 1,722 suicide attempts and 206 suicide completions (Suicide Prevention Resource Center (SRPC), 2004).
13. In California, 1999-2000, young people, ages 15-19 years made 2,421 suicide attempts and 116 suicide completions (SRPC, 2004).
14. In California, 1999-2000, young people, ages 0 to 14 years made 649 suicide attempts and 23 suicide completions (SRPC, 2004).

## B. Major Federal Initiatives Addressing Youth Suicide Prevention: 1985-2005

Date	Initiative Sponsoring Agency	Initiative
1985	U.S. House of Representatives	Youth Suicide Prevention Act of 1985
1990	U.S. Department of Health and Human Services	Health People Initiative 2000
1998	United States Senate & House of Representatives	Senate Resolution #84 and House Resolution #212
1999	U.S. Public Health Service	Surgeon General's Call to Action to Prevent Suicide
2000	U.S. Department of Health and Human Services	Healthy People Initiative 2010
2000	U.S. Public Health Service	Surgeon General's Conference of Children's Mental Health
2001	U.S. Public Health Service	National Strategy for Suicide Prevention: Goals and Objectives for Action
2002	President's New Freedom Commission on Mental Health	Achieving the Promise: Transforming Mental Health Care in America Report
2004	U.S. Senate & House of Representatives	Garrett Smith Memorial Act Senate Bill# 2634 & House Resolution # 4799

## C. Federal Support for School-Based Suicide Prevention

1. *The Surgeon General's Call to Action to Prevent Suicide* (U. S. Public Health Service (USPHS), (1999). <http://www.surgeongeneral.gov/library>  
This federal government document has published specific, serious concerns over youth suicide and its prevention. In its call to action it states four aims, which include:
  - a. Prevent premature deaths due to suicide across the life span.
  - b. Reduce the rates of other suicidal behaviors.
  - c. Reduce the harmful after-effects associated with suicidal behaviors and the traumatic impact of suicide on family and friends.
  - d. Promote opportunities and settings to enhance resiliency, resourcefulness, respect, and interconnectedness for individuals, families, and communities.
  - e. This report calls for schools to be access and referral points for children's mental and physical health services.

2. *National Strategy for Suicide Prevention (NSSP)*  
<http://www.mentalhealth.org/suicideprevention>
  - a. This is the companion volume to *The Surgeon General's Call to Action to Prevent Suicide* (U. S. Public Health Service (USPHS) (1999). It has eleven stated goals, accompanied by 68 objectives. The eleven goals of the NSSP include (USPHS, 2001):
    - 1) Promote awareness that suicide is a public health problem that is preventable.
    - 2) Develop broad-based support for suicide prevention.
    - 3) Develop and implement strategies to reduce stigma associated with being a consumer of mental health, substance abuse and suicide prevention services.
    - 4) Develop and implement community-based suicide prevention programs.
    - 5) Promote efforts to reduce access to lethal means and methods of self-harm.
    - 6) Implement training for recognition of at-risk behavior and delivery of effective treatment.
    - 7) Develop and promote effective clinical and professional practices.
    - 8) Increase access to and community linkages with mental health and substance abuse services.
    - 9) Improve reporting and portrayals of suicidal behavior, mental illness, and substance abuse in the entertainment and news media.
    - 10) Promote and support research on suicide and suicide prevention.
    - 11) Improve and expand surveillance systems.
  - b. Within the National Strategy schools are elevated as important venues for suicide prevention activities and are challenged to develop quality, safe, and effective school-based suicide prevention programs.
3. *The President's New Freedom Commission on Mental Health (NFCMH, 2003)*. <http://www.mentalhealthcommission.gov>.  
 In July 2003, the President released the report, *Achieving the Promise: Transforming Mental Health Care in America* (NFCMH, 2003). The report was created to address the problems in the current mental health system and underlined that mental illnesses rank first among illnesses that cause disability in the United States, Canada, and Western Europe.
  - a. The six goals in the President's report with recommendations under each are:
    - (1) Americans understand that mental health is essential to overall health.
    - (2) Mental health care is consumer and family driven.
    - (3) Disparities in mental health services are eliminated.
    - (4) Early mental health screening, assessment, and referral to services are common practice.
    - (5) Excellent mental health care is delivered and research is accelerated.
    - (6) Technology is used to access mental health care and information.
  - b. The accompanying recommendations include the call to:

- (1) Swiftly advance and implement a national campaign to reduce the stigma of seeking care.
  - (2) Promote the national strategy for suicide prevention.
  - (3) Create public-private partnerships to address suicide prevention in communities, including school personnel, local leaders, and representatives of the faith community.
  - (4) Create integrated systems of care for children with serious emotional disturbances and their families, a population known to be at high risk for youth suicide.
  - (5) Promotion of the mental health of young children.
  - (6) Provide early mental health screening, assessment, and referral to service.
  - (7) Strengthen school mental health services, as evidenced by an improvement and expansion of education and training, prevention, early identification, early intervention, and treatment services within the school.
- c. The report emphasizes the growing evidence that school mental health programs improve educational outcomes by decreasing absences and discipline referrals, as well as improving test scores (Jennings, Pearson, & Harris, 2000).

#### D. Federal Agency Resources for Suicide Prevention

1. Centers for Disease Control and Prevention (CDC) National Youth Violence Prevention Resource Center. <http://safeyouth.org/home.htm>.  
Following a cluster of suicides in school aged children, in 1992, this agency created a resource guide that includes eight strategies for youth suicide prevention programs, most of which can be used to establish specific suicide prevention programs in schools. Those strategies include formation and implementation of:
  - a. School gatekeeper training.
  - b. Community gatekeeper training.
  - c. General suicide education.
  - d. Screening programs.
  - e. Peer support programs.
  - f. Crisis centers and hotlines.
  - g. Means restriction.
  - h. Intervention after a suicide.
2. *Human Resources Services Administration (HRSA)*.  
<http://www.mchb.hrsa.gov/>.  
This agency gives out Maternal Child Health block grants that can be used in many ways, including youth suicide prevention projects.
3. *Indian Health Service (IHS)*.  
<http://www.his.gov>.  
This agency seeks to raise the physical and mental health status of American Indians and Alaska Natives.
4. *National Council for Suicide Prevention Training*.  
<http://www.ncspt.org>.

This is a national on-line training program to educate people about suicide prevention.

5. *National Institutes of Health/Mental Health (NIH/NIMH).*

<http://www.nimh.nih.gov>

NIH is the principal biomedical and behavioral research agency of the United States Government. NIMH is a part of NIH, with a mission to reduce the burden of mental illness and behavioral disorders through research on mind, brain, and behavior.

6. *Substance Abuse and Mental Health Services (SAMSA).*

<http://www.samhsa.gov>.

This agency funds the Suicide Prevention Resource Center (SPRC), a good resource for suicide prevention information.

### **E. California Endeavors in School-Based Suicide Prevention.**

1. *California Education Code: [www.schoolcounselor-ca.org/downloads/California Education Code 49600.doc](http://www.schoolcounselor-ca.org/downloads/California%20Education%20Code%2049600.doc).*

The legal references to school-based suicide prevention related issues:

- a. *49602 Confidentiality of student information.*
- b. *9604 Suicide prevention training for school counselors.*

2. *California Department of Education Website.*

<http://www.cde.ca.gov/>.

This site includes current resource information and references to public and private sources of information, research, curriculum, screening tools, gatekeeper training, foundations, survivor support, prevention, intervention and postvention program materials, all related to youth suicide prevention.

3. *California Department of Education (CDE) – Counseling and Student Support Division. <http://www.newcde.ca.gov/ls/cg>.*

In 1987, the CDE Instructional Support Division, School Climate Unit published the *Suicide Prevention Program for California Public Schools (SPPCPS)*, (ISBN#:0-8011-0682-6). The SPPCPS program materials include four guides: the Implementation and Resource Guide; the Lesson Guide; the Guide for Staff Awareness In-Service Training; and Guide for Parent Awareness Meeting. The SPPCPS, along with many other suicide prevention resources published in the 1980s, is now referred to in the program evaluation literature as a “first generation program”, because it is recognized that program development and evaluation occur in an iterative cycle (Kalafat & Lazarus, 2002). While this program still contains useful information, there has been much research on youth suicide prevention in the past seventeen years since its publication. Additional information has been learned about youth suicide behavior and school-based suicide prevention. New recommendations for current school-based suicide prevention program development are listed later in this document. The CDE Counseling and Support Office has recently updated its school-based suicide prevention resource website.

4. *California School Boards Association (CSBA). <http://www.csba.org>.*

- a. CSBA has created an optional board policy and administrative regulation for districts that wish to adopt policy providing for student, staff, and parent/guardian training in suicide prevention, which are:

(1) *BP 5141.52*

(2) *AR 5141.52*

- b. The courts have ruled that a district may be liable to parents/guardians of a student who committed suicide while under supervision of school staff.
  - c. Before adopting policy on this topic, the Board should consider staff and capabilities to be sure that the policy will be fully enforced. Once a district has adopted a policy, it has a duty to enforce it.
5. California Strategy for Suicide Prevention.  
On May 6-7, 2004, a collaborative group of individuals from many different organizations and vantage points (suicide prevention advocacy groups, government officials, education and business professionals, survivors of suicide, and public health professionals) met to devise a draft of the California Strategy for Suicide Prevention, modeled after the National Strategy for Suicide Prevention. This document will be in the formation stage for a period and then published. The document called for schools to be active participants in screening, prevention, intervention, and training.
6. *California Welfare and Institutions Code*.  
[www.leginfo.ca.gov/calaw.html](http://www.leginfo.ca.gov/calaw.html).  
The legal references to school-based suicide prevention related issues:  
a. *5698 Emotionally disturbed youth; legislative intent*
7. Center for Mental Health in Schools (CMHS) – UCLA.  
<http://smhp.psych.ucla.edu>.  
The CMHS has created a comprehensive technical assistance sampler on school interventions to prevent suicide. It gives guidance about suicide, its prevention, suicide assessment, intervention planning and training, and aftermath (postvention) assistance and the prevention of contagion.

#### **F. Websites with Suicide Prevention Research**

1. *International Academy for Suicide Research*.  
<http://www.uni-wuerzburg.de/IASR>.
2. *National Library of Medicine*.  
[http://mentalhealth.org/suicide\\_prevention](http://mentalhealth.org/suicide_prevention).
3. *National Institute of Mental Health Suicide Research Consortium*.  
<http://www.nimh.nih.gov/research/suicide.cfm>.
4. *Oxford University Centre for Suicide Research*.  
<http://cebmh.warne.ox.ac.us/csr>.
5. *Suicide Information and Education Centre (SIEC)*.  
<http://www.siec.ca>.
6. *Unit for Suicide Research (Belgium)*.  
<http://allserv.rug.ac.be/~cvheerin>.
7. *Web-based Injury Statistics Query and Reporting System (WISQARS)*.  
<http://www.cdc.gov/ncipc/wisqars/default.htm>.

#### **G. Websites with Suicide Related Statistics**

1. *Centers for Disease Control and Prevention*.  
<http://www.cdc.gov/scientific.htm>.
2. *National Injury Data Technical Assistance Center*.



<http://injurypreventionweb.org/info/data/htm>.

3. *National Violent Injury Statistics System*.  
<http://www.nviss.org>.
4. *World Health Organization Statistical Information System (WHOIS)*.  
<http://www3.who.int/whois/menu.cfm>.

## **H. Professional Organizations Related to Suicide Prevention.**

### **1. National Council for Suicide Prevention.**

<http://www.ncsp.org/>.

Coalition of organizations committed to suicide prevention including:

- a. American Association of Suicidology (AAS).  
<http://www.suicidology.org>.
- b. American Foundation for Suicide Prevention (AFSP).  
<http://www.afsp.org>.
- c. The Jason Foundation.  
<http://www.jasonfoundation.com>.
- d. The Kristin Brooks Hope Center.  
<http://www.hopeline.com>.
- e. The Link's National Resource for Suicide Prevention.  
<http://www.thelink.org>.
- f. National Organization for People of Color Against Suicide (NOPCAS).  
<http://www.nopcas.com>.
- g. Organization for Attempters and Survivors of Suicide in Interfaith Service (OASSIS).  
<http://www.oassis.org>.
- h. Samaritans USA.  
<http://www.samaritans.org>.
- i. Suicide Awareness\Voices of Education (SAVE)  
<http://www.save.org>.
- j. Suicide Prevention Action Network USA, Inc. (SPAN)  
<http://www.spanusa.org>.
- k. Yellow Ribbon Suicide Prevention Program  
<http://www.yellowribbon.org>.

## **I. Ecological Theory of Suicide Prevention.**

1. Ecological theory useful to youth suicide prevention with intentional focus on specific risk factors most closely associated with completed youth suicide. Scientific criteria for selecting targeted risk factors are (Breton, Boyer, Bilodeau, Raymond, 2002):
  - a. Modifiability - ability to change in structure or function due to internal, external or hereditary influences.
  - b. Measurability – ability to be measured
  - c. Degree of association with the health problem.
  - d. Degrees of proximity – processes that involve patterns of progressively more complex reciprocal interaction with persons, objects, and symbols in the immediate environment and to be effective, occurring on a fairly regular basis over extended periods of time.

2. Risk factors most strongly associated with completed youth suicide are, in decreasing order (Bell & Clark, 1998; Brent, 1995, Moscicki, 1997):
  - a. Prior suicide attempts.
  - b. Depression.
  - c. Substance abuse (including alcohol).
  - d. Conduct disorder.
3. Vast majority of experts agree that suicide is a multi-dimensional problem and ecological model addresses multiple systems involved in emergence, development, and maintenance of suicidal behavior (Henry, Stephenson, Hanson, & Hargett, 1993; Shagle & Barber, 1995; White, 1998; Motes, Melton, & Simmons, 1999) which include:
  - a. Personal.
  - b. Family.
  - c. School.
  - d. Work.
  - e. Community.
4. An effective ecological approach to school-based suicide prevention serves to reduce suicide rates and includes:
  - a. Implementation with fidelity.
  - b. Multiple levels of school and community
  - c. Dissemination to enough sites to obtain large population samples for epidemiological impact assessment.
  - d. Institutionalization long enough to detect epidemiological trends (Kalafat & Lazarus, 2002).
5. Ecological approach to youth suicide purports suicide emerges from adolescents' interactions and interdependencies within hierarchically organized multiple level ecological contexts (Henry, et al. 1993; Bronfenbrenner, 1979), therefore guidelines, programs and procedures for school-based suicide prevention, in this report will serve to:
  - a. Target and reduce risk factors most strongly associated with completed youth suicide (Burns & Patton, 2000; Borowsky et al., 2001).
  - b. Identify and enhancing protective factors related to youth suicide prevention and resilience (Burns & Patton, 2000; Borowsky et al., 2001).
  - c. View suicide from a psychiatric illness model, rather than a stress model as many studies verify nearly 90% of those completing suicide suffer from a diagnosable mental illness that needs to be identified and treated (Beautrais, 2001; Brent et al., 1999; Shaffer et al., 1996).
  - d. Emphasis identification, diagnosis, and treatment of the underlying psychiatric disorder are the most critical pieces of suicide prevention (Beautrais, 2001; Brent et al., 1999; Shaffer et al., 1996).
6. Risk Factors for Youth Suicide from an Ecological Approach.
  - a. Personal Characteristics:
    - (1) Psychopathology (particularly mood disorders, schizophrenia, anxiety, disorders, certain personality disorders, alcohol & other substance abuse disorders). (Beautrais, 2001; DHHS, 2001).
    - (2) Prior suicide attempts (Shaffer et al., 1996).

- (3) Cognitive and personality factors (i.e. skill development delays, emotional difficulties; apathy, hopelessness, school problem, poor interpersonal problem-solving ability) (Russell & Joyner, 2001).
- (4) Sexual orientation (homosexual, trans-gender, bisexual) (Blake et al., 2001, McDaniel et. al., 2001).
- (5) Biological factors (i.e. major physical illness, abnormalities of serotonin function) (DHHS, 2001; Arango, 2001).
- b. Family Characteristics:
  - (1) Family history of suicidal behavior (Gould et al., 1996; Agerbo et al, 2002).
  - (2) Parental psychopathology (Gould et al., 1996).
  - (3) Involvement in sports & physical activity and high academic achievement are protective (King et al., 2002; Resnick, Harris, & Blum, 1993; Jessor et al., 1995).
- 7. Protective Factors for Youth Suicide from an Ecological Approach.
  - a. Emotional Well-being:
    - (1) Emotional wellness especially protective for adolescent girls (Borowsky, Resnick, Ireland & Blum, 1999).
  - b. Gender:
    - (1) Statistically, being female is more protective than being male (Kaslow, McClure & Connell, 2002; Maris, 2002).
    - (2).Females make more suicide attempts, but males complete more suicides (Lacourse, Claes, & Villeneuve, 2001).
  - c. Ethnicity/Race:
    - (1) Overall, the rate of suicide is lower in ethnic minorities as a whole than whites (Maris, 2002), though there are some individual minority group exceptions (i.e., Native Americans).
    - (2) Mexican-Americans were 1.8 times more likely to die by suicide than European-Americans.
    - (3) Suicide attempts are 19.3% higher among Latina girls than in many other ethnic-gender groups (Rew et al., 2001).
  - d. Religiosity:
    - (1) Religiosity is a protective factor (Fernquist, 2000; Forman & Kalafat, 1998; Foster, 2001; Maris, 2002).
    - (2) Greater religiosity has been touted as the reason for the historically lower suicide rate among African-Americans (Glowinski et al., 2001).

## **J. Definitions of Suicide Related Terminology.**

1. ***Assisted Suicide:***  
Completed suicide in which a person intended to die, but relied on the assistance of another individual (who becomes an agent of the suicide) to complete the act (Capuzzi, 2004).
2. ***Completed Suicide:***  
Self-inflicted lethal act resulting from intentional life-threatening action. The Centers for Disease Control and Prevention provided an Operational Criteria for the Classification of Suicides, listing the three essential elements of a completed suicide as: death; self-infliction; and intentionality (O'Carroll et al., 1998).
3. ***Durkheim's four basic types of suicide*** (Durkheim, 1897/1951):
  - a. ***Egotistical:*** Not selfish, but rather lonely and isolationist, lack of integration.
  - b. ***Altruistic:*** Enmeshed relationship to society, lack of individuation.
  - c. ***Anomic:*** Situational crises and chaos.
  - d. ***Fatalistic:*** Suppression of individual by society leading to pessimistic life outlook.
4. ***Gatekeepers:***  
Individuals who has face-to-face contact with large numbers of community members as part of their usual routine.
5. ***Gatekeeper Training in Suicide Prevention:***  
Training for gatekeepers to learn how to identify persons at risk of suicide and refer them to treatment or support services as appropriate.
6. ***Means Restriction:***  
Techniques, policies, and procedures designed to reduce access or availability to means of and methods of deliberate self-harm.
7. ***Protective Factors:***  
Factors that make it less likely an individual will develop a disorder. Protective factors may encompass biological, psychological, or social factors in the individual, family, and environment.
8. ***Resilience:***  
Capacities within a person that promote positive outcomes, such as mental health and well-being, and provide protection from factors that might otherwise place that person at risk for negative health outcomes.
9. ***Risk Factors:***  
Factors that make it more likely an individual will develop a disorder: risk factors may encompass biological, psychological, or social factors in the individual, family, and environment.
10. ***Screening:***  
Administration of an assessment tool to identify persons in need of more in-depth evaluation or treatment.

11. ***Self-harm/self injury:***  
Various methods individuals use to injure themselves, such as self-laceration, self-battering, taking overdoses, or exhibiting deliberate recklessness.
12. ***Stigma:***  
Object, idea, or label associated with disgrace or reproach.
13. ***Suicide Prevention Levels (Primary, Secondary, Tertiary):***
  - (a) ***Primary Suicide Prevention (also referred to as Prevention):***  
Primary suicide prevention involves activities and programs focused on preventing suicide from occurring, before it is considered or attempted.
  - (b) ***Secondary Suicide Prevention (also referred to as Intervention):***  
Secondary suicide prevention involves activities and programs to be implemented when a person is actively considering or has attempted suicide.
  - (c) ***Tertiary Suicide Prevention (also referred to as Postvention):***  
Tertiary prevention is the intervention conducted with survivors once a suicide has occurred.
14. ***Tri-Levels of Suicide Intervention (Universal, Selected, Indicated):***
  - (a) ***Universal Interventions:***  
Universal interventions are activities or programs directed at an entire population, rather than selected individuals, (i.e. a whole student body) to decrease suicide risk factors and enhance protective factors (Greenberg et al., 2001).
  - (b) ***Selected Interventions:***  
Selective interventions are activities or programs delineated for individuals or sub-groups believed to be high-risk based on biological or social risk factors that place them at significantly higher than average risk for committing suicide, yet they have not yet exhibited symptoms of a disorder that places them at high risk for suicide, such as depression, substance abuse, or deliberate self-harm (Greenberg et al., 2001).
  - (c) ***Indicated Interventions:***  
Indicated interventions are activities or programs designed for individuals already exhibiting symptoms of a disorder that places them at high risk for suicide, such as depression, substance abuse, or deliberate self-harm (Burns & Patton, 2000).
15. ***Suicide Attempt or Parasuicide (With or Without Injury):***  
Suicidal act that has a nonfatal outcome in which the person has some level of intention to die (Capuzzi, 2004).
16. ***Suicide Gesture:***  
Gesture meant to portray an appearance of suicide and self-harm, but is without any real intention to kill oneself to accomplish an ulterior motive. (Capuzzi, 2004).

17. ***Suicide Threat:***

Any behavior in form of some type of interpersonal communication, verbal or non-verbal, without physical harm, implying the individual has intention on some level to act in a suicidal manner (Capuzzi, 2004).

18. ***Suicidal Ideation:***

Thinking about future or fantasized suicidal action (Capuzzi, 2004).

*Definitions from:*

Capuzzi, D. (2004). From Seneca to suicidology: A history of suicide. In D. Capuzzi (Ed). *Suicide Across the Life Span: Implications for Counseling* (3-38). Alexandria, VA: American Counseling Association.

U. S. Public Health Service (2001). *National Strategy for Suicide Prevention: Goals and Objectives for Action (NSSP)*. (196-204). (DHHS Publication No. 02NLM: HV 6548.A1 2001). Rockville, MD: Author.

**Section II. Planning of School-Based Primary Suicide Prevention Program (also called Prevention), involves planning activities and programs focused on preventing suicide from before it is considered or attempted.**

**A. School District Level – Superintendent & School District Board of Education.**

1. Formation and maintenance of policies and procedures.
  - a. Imperative to create a school district board policy regarding suicide prevention to clarify and underline the district's commitment to suicide prevention (Cellota, Jacobs, Keys, & Cannon, 1989; Fetro, 1998). (See Section 6, Attachment 1 example from CA School Board Association).
  - b. Include suicide prevention policies and procedures in the school district's mandated Comprehensive School Safety Plan.
  - c. Essential to develop and implement policies and procedures in the event of a suicide attempt or completion by a staff member or student, including those occurring on and off campus. (Rowling & Holland, 2000).
    - (1) If two or more school mental health professionals assess a student as being at risk for suicide:
      - (a) Student's parents/guardians should be immediately contacted.
      - (b) Referral should be made to a mental healthy agency or hospital emergency for immediate services.
      - (c) If parent/guardian does not cooperate and agree to get the student outside help, the school must report this to local law enforcement or child protective services official (Lieberman & Davis, 2002; McEvoy & McEvoy, 1994).
    - (2) If two or more school mental health professionals deem a student to be at risk for suicide and the student's parent or guardian (or emergency contact) cannot be reached, the local law enforcement should be called to escort the student to the nearest hospital emergency room for evaluation (Lieberman & Davis, 2002; Capuzzi, 1994; Capuzzi & Gross, 2004; King, 2001; McEvoy & McEvoy, 1994).
    - (3) If two or more school mental health professionals assess a staff member as being at risk for suicide:
      - (a) Staff member should be immediately referred to a mental healthy agency or hospital emergency for services.
      - (b) Superintendent or designee must be notified and the staff member may be placed on administrative leave until cleared by a mental health
    - (4) If two or more school mental health professionals assess a staff member as being suicidal and the staff member refuses to seek treatment:
      - (a) Local law enforcement should be called to escort the staff member to the nearest hospital emergency room for evaluation.
      - (b) The superintendent or designee must be notified and the staff member may be placed on administrative leave until cleared by a mental health official approved or designated by the school district.

2. Formation and maintenance of community partnerships & mutual aid agreements in youth suicide prevention ( See Section 6, Attachment 11) to facilitate collaborative work on youth suicide prevention throughout the school and its community (McKee, Jones, & Barbe, 1993), including:
  - a. County Office of Education or Regional Education Agency.
  - b. Neighboring School Districts' Crisis Team Representatives.
  - c. Law enforcement and coroner.
  - d. City and county government officials.
  - e. Hospital emergency departments and emergency medical personnel.
  - f. Community fire and emergency services.
  - g. Youth health services.
  - h. Psychiatric facilities.
  - i. Crisis hotline and suicide prevention center representatives.
  - j. Religious/clergy associations.
  - k. Survivor groups.
  - l. Community social service and mental health agencies.
  - m. After-school care providers.
3. Make arrangements for school district representation in multi-disciplinary reviews of primary, secondary, and tertiary prevention efforts affecting decedents who are students or employees of the school district with the County Child Suicide Death Review Team.
4. Initiate school communication between hospitals and outpatient mental health and substance abuse centers to encourage a good professional working relationship, helping ensure continuity of care of district students who receive those services and return or continue in school (See Section 6, Attachment 11).
5. Formation and maintenance of a District Crisis Team as a back up support system to all schools and sites in the district for issues pertaining to suicide, crisis, death, and grief. (See Section 6, Attachment 4 for details on membership selection and roles) (Kalafat & Lazarus, 2002)
  - a. Appoint/hire a Crisis Guidance Manager (see Section 6, Attachment 3)
  - b. Have both a primary and secondary school District Crisis Teams (Section 6, Attachment 4)
  - c. Minimum two-year commitment.
6. wide suicide prevention training for school district employees and parents and guardians of district students  
must create a sound infrastructure of staff trained in assessment, and gatekeeper skills early in school year in English (and Spanish if needed). (See Section 6, Attachment 7).
  - a. Designate District Crisis Guidance Manager to recruit school or community mental health professionals knowledgeable about suicide and its prevention to conduct suicide prevention training with all district employees (See Section 6, Attachment 8 for recommended programs).
  - b. Commit to and budget for district mental health professionals involved in training and suicide risk assessment to receive continuing education in suicide prevention methodology and research.



- c. Prepare the infrastructure and community partnerships to support a systematic district suicide screening for secondary students.
  - (1) Screening via individual interviews and self-report, used to identify students at risk for suicidal behavior shown to be efficacious in literature (Joiner, Pfaff, & Acres, 2002; Shaffer & Craft, 1999; Thompson & Eggert, 1999; Zametkin, Alter, & Yemini, 2002). (See Section 6, Attachment 8)
  - (2) Must have lined up ample community referral sources to immediately direct students for follow up treatment if they have screened positive for being at risk for suicide (Gould et al., 2003).
  - (3) Participate in a screening program that has strong empirical data behind it (Aseltine, 2003; Aseltine & DeMartino, 2004; Shaffer & Greenberg, 2002).
  - (4) Anticipate resistance from professionals who prefer curriculum-based approaches (Miller et al., 2003) because screening approach is more complicated, yet defend via literature because it has been shown to be more effective.
- d. Continue to prepare the infrastructure for school-based suicide prevention by creating a plan to implement gatekeeper training for all employees district-wide on a two-year year cycle (Adelman & Taylor, 2000; King, 2001; Kalafat & Lazarus, 2002).
  - (1) Training to include (see Section 6, Attachment 7).
    - (a) Clarification of school's role, policies and procedures relative to suicide and its prevention.
    - (b) Role of mental illness in suicide.
    - (c) Identification of suicide warning signs.
    - (d) Identification of suicide contagion factors.
    - (e) Identification of and promotion of protective factors.
    - (f) Identification of referral resources and effective referral procedures.
    - (g) How to respond to depressed and suicidal people.
    - (h) Role of school connectedness, student aggression and bullying prevention, and substance abuse prevention in the prevention of suicide. (Gould et al., 2003; Kalafat & Lazarus, 2002; King, 2001; Speaker & Petersen, 2000).
    - (i) Role of media in suicide and its prevention. (See Section 6, Attachment 5).
    - (j) School mental health and nursing professionals should receive additional advanced suicide prevention training, in screening, risk assessment, and information on legal cases pertaining to student suicide on a two-year cycle.
- e. Institute a parent and guardian education program on youth suicide prevention (in English and Spanish if needed) that covers the same topics as the employee training, with an increased emphasis on:
  - (1) Means restriction strategies (Kalafat & Lazarus, 2002).

- (2) Introduction of suicide prevention strategies children in the school district will be taught.
- (3) Availability of community resources for referring their children or themselves if help is needed (Capuzzi & Gross, 2004; Kalafat & Lazarus, 2002).
- (4) Invite nursery and pre-school teachers, after-school program personnel, and known childcare providers in the district, also.
- g. Formation and maintenance media relationships and agreements. (See Section 6, Attachment 5).
  - (1) Assign a district media spokesperson.
  - (2) Designated school mental health person(s) should develop and provide press an updated information kit each August for district media spokesperson and local media outlets to include information about reporting on suicide and the sensitive issues included in media coverage of suicide, as well as contact information for district's designated media spokesperson (USPHS, 2001).

**B. School/Site Level – Building Administrators, School Mental Health and Medical Professionals**

- 1. Formation and maintenance of school/site policies and procedures (see
- 2. Formation and maintenance of a school/site crisis team (see Attachment 6).
  - a. Obtain school district level and Board of Education administrative support (Capuzzi & Gross, 2004; Huberman & Miles, 1984).
  - b. Establish and train a building/site crisis response team with a diverse group of professionals, (principal, counselor, teacher, school nurse, school psychologist) (Brock et al., 1994; Brock, 2000, 2002).
  - c. Designate a team leader (typically the principal) and alternate to ensure leadership presence in each school building at all times.
  - d. Crisis response team leader should schedule and arrange for annual suicide intervention training and suicide intervention rehearsals for entire staff (Siehl, 1990).
- 3. Formation and maintenance of student suicide prevention program plan (see Section 6, Attachment 8)
  - a. Be aware of tension in the literature over suicide awareness curriculum & and make informed decisions (see Attachment #2) (Aseltine, 2003; Capuzzi, 1998; 1994; Capuzzi & Golden, 1988; Capuzzi & Gross, 2000; Curran, 1987; Kalafat & Elias, 1994; Ross, 1980; Sudak, Ford, & Rushforth, 1984; Zenere & Lazarus, 1997).
  - b. Some advocates support suicide prevention education and discussion curriculum in a forum that allows for accurate information, asking questions, and learning how to obtain help for self and friends in the context of a school wide prevention effort (Capuzzi & Gross, 2004).
  - c. Some advocates have found suicide prevention curriculum to be without benefit or even detrimental for school students of all ages (Shaffer,

Garland & Whittle, 1988; Shaffer et al., 1991; Vieland et al. 1991; Mazza, 1997; Garland, 1989; Ciffone, 1993).

- d. Some researchers believe there remains insufficient evidence to definitively either support or not support curriculum-based suicide awareness programs in school (Guo & Harstall, 2002), making alternative school-based strategies such as identification of suicidal warning indicators, monitoring of students predisposed to suicide, and crisis intervention viable options (Helsel, 2001; Range, 1993, Capuzzi, 1994; Freeman, 1998).
4. The Suicide Prevention Resources Center (SPRC), with help from the American Foundation for Suicide Prevention (AFSP), developed a National Registry of Effective Programs, (NREP) This registry is was cancelled after January 2005 and is to be subsumed by a new registry, the National Registry of Evidence-Based Programs and Practices, (NREPP) anticipated to be in place sometime in the Fall of 2005.
5. Programs that were registered on the NREP are described more fully in District Crisis Guidance Manager – Attachment and they include:
  - a. ***C-Care/CAST*** (Thompson, Eggert, Randell, & Pike, 2001):
    - (1) SPRC Classification: Effective
    - (2) Target Age: 14-18
    - (3) Gender: Male & Female
    - (4) Ethnicity/Race: Multiple
    - (5) Level of Intervention: Selective, Indicated
  - b. ***SOS: Signs of Suicide*** (Asestine & DeMartino, 2004)
    - (1) SPRC Classification: Promising
    - (2) Target Age: 14-18
    - (3) Gender: Male & Female
    - (4) Ethnicity/Race: Multiple
    - (5) Level of Intervention: Universal
  - c. ***Columbia TeenScreen*** (Shaffer, Scott, Wilcox, Maslow, Hicks, Lucas, et al., 2004):
    - (1) SPRC Classification: Promising
    - (2) Target Age: 11-18
    - (3) Gender: Male & Female
    - (4) Ethnicity/Race: Varied
    - (5) Level of Intervention: Universal
  - d. ***Lifelines*** (Kalafat & Elias, 1994)
    - (1) SPRC Classification: Promising
    - (2) Target Age: 12-17
    - (3) Gender: Male & Female
    - (4) Ethnicity/Race: Multiple
    - (5) Level of Intervention: Universal
  - e. ***Reconnecting Youth Class*** (Eggert & Nicholas, 2004)
    - (1) SPRC Classification: Promising
    - (2) Target Age: 14-18
    - (3) Gender: Male & Female

- (4) Ethnicity/Race: Multiple
- (5) Level of Intervention: Selective, Indicated
- f. **Zuni Life Skill Development** (LaFromboise, 1995)
  - (1) SPRC Classification: Promising
  - (2) Target Age: 14-18
  - (3) Gender: Male & Female
  - (4) Ethnicity/Race: American Indian
  - (5) Level of Intervention: Selective
- g. The NREP noted two programs under consideration, but in need of more study and not registered:
  - (1) **ASIST**
    - (a) SPRC Classification: Unrated at this time
    - (b) Target Age: 18+
    - (c) Gender: Male & Female
    - (d) Ethnicity/Race: Multiple
    - (e) Level of Intervention: Selective, Universal
  - (2) **Yellow Ribbon**
    - (a) SPRC Classification: Unrated at this time
    - (b) Target Age: All Ages
    - (c) Gender: Male & Female
    - (d) Ethnicity/Race: Multiple
    - (e) Level of Intervention: Selective, Universal
- 6. Since January 2005, the NREP Registry has closed.
  - a. Programs recognized by the former program, the National Registry of Effective Programs (NREP), will need to be reviewed under the new criteria of the National Registry of Effective Programs and Practices (NREPP) and will be administered under the Substance Abuse and Mental Health Services Administration (SAMHSA) with new standards and criteria for inclusion as a model program.
  - b. SAMSHA anticipates the new NREPP website will be on-line in late 2005 at [www.nationalregistry.samhsa.gov](http://www.nationalregistry.samhsa.gov) with newly reviewed model programs.
  - c. The NREPP goal is to provide the public with contemporary and reliable information about the scientific basis and practicality of interventions to prevent suicide and mental illnesses.
- 7. Include skill instruction in: bullying prevention, conflict resolution, coping communication, decision-making, assertiveness, self-awareness, help-seeking, and cognition to help reduce suicide risk factors, (such as depression, hopelessness, and drug abuse) Gould et al., 2003; Zenere & Lazarus, 1997; Eggert, Thompson, Herting et al., 1995; Randell et al., 2001; Thompson et al., 2000, 2001).
- 8. Implement school-wide direct screening of students (see Attachment #3) (Joiner, Pfaff & Acres, 2002; Shaffer & Craft, 1999; Zametkin, Alter, & Yemini, 2002).
  - a. Screening done via individual interviews and self-report, used to identify students at risk for suicidal behavior, and has shown to be efficacious in

the recent literature (Joiner, Pfaff & Acres, 2002; Shaffer & Craft, 1999; Thompson & Eggert, 1999; Zametkin, Alter, & Yemini, 2002).

- b. Some barriers to overcome in school-wide screening suicide screening procedures include:
  - (1) Need for multiple screenings to minimize “false –negatives” on screenings (Berman & Jobes, 1995).
  - (2) Resistance by some school psychologists and high school principals who prefer curriculum based approaches (Miller et al., 2003; Miller et al., 1999).
  - (3) Need for effective, numerous referral resources for student screening positive for suicide risk (Gould et al, 2003).
9. Schedule specific yearly collaboration/in-service time among teachers, nurses, and school mental health professionals to coordinate and implement suicide prevention programs (King, 2001; CDC, 1992; Ollendick, Greene, Werst, & Oswald, 1990).
10. Establish protocol that when any staff member recognizes a student or staff member at risk for suicide, the individual must immediately convey that information to school nurse and mental health professional (counselors, psychologists, and social workers) (King, 2001) and proceed with district policy.
11. Implement school activities aimed at increasing school connectedness, as National Longitudinal Study on Adolescent Health (Resnick et al., 1993) found adolescents perceive school connectedness as a leading protective factor against youth suicide.
12. Provide and encourage opportunities for participation in after-school clubs and activities, and a comfortable physical environment because they are known protective factors against suicide (King, 2001).
13. For suicide attempts or completions on campus, principal or designee should provide information to local public mental health agencies and local child death suicide review teams to help improve and expand surveillance (USPHS, 2001).
14. Implement the parent/guardian suicide prevention education program developed by the district (see Section 6, Attachment 7).

**Section III. School-Based Secondary Suicide Prevention Program, (also called Intervention), refers to steps school personnel on scene should take when a student or staff member threatens or attempts suicide (King, 2001).**

**A. School/ Site Level**

1. Goals to maintain in extreme risk situation (McKee et al., 1993):
  - a. Prevent suicide completion.
  - b. Move others to safety.
2. Have risk person escorted by an adult employee to non-threatening environment with access to phone, **never** leave risk person alone, even to go to restroom.
3. Alert building/site crisis team (and police if warranted).
4. Do not make promises of confidentiality—no deals.
5. Question specificity of plan.
6. Trained school mental health professional, with the witness and help of another mental health or medical colleague, should conduct a suicide risk assessment when a student or staff member threatens or attempts (Capuzzi & Gross, 2004; Davis & Brock, 2002).
  - a. Extreme risk: Has specific plan, has dangerous instrument or means to carry out plan.
  - b. Severe risk: Has specific plan, but no current means to accomplish it.
  - c. Moderate risk: Suicidal ideation verbalized without plan or means Notify parents or legal guardians if student threatens or attempts suicide.
7. Failure to notify parents/guardians of suicide threats or attempts has resulted in school districts being sued. (Poland & McCormick, 1999).
  - a. Mental health professionals should document all that is done in an intervention on behalf of a suicidal student or staff member, keep a copy, and turn a copy into the building administrator (Capuzzi & Gross, 2004).
  - b. Follow up and support suicidal student and family or suicidal staff member, and document those efforts:
8. Counselor and school psychologist should offer support for child to parents/guardian, needed educational modification, and assessment as appropriate.
9. School mental health professionals should offer community agency or social support program referrals to parents/guardians in writing and keep a copy (Capuzzi & Gross, 1994; Lieberman & Davis, 2002).
10. Encourage continuity of care by facilitating communication between school, hospitals, and outpatient mental health and substance abuse providers by encouraging parents/guardians of suicidal student or the suicidal adult staff member to sign appropriate forms consenting to exchange of information between parties.
11. After an intervention with a student or staff member:
  - a. Notify and debrief school staff on a need-to-know basis, (nurse, teachers of student, supervisors of staff member, rest of the crisis team, district office contact, Crisis Guidance Manager).
  - b. Debrief and evaluate strategies employed and maintain or modify them based on their effectiveness (King, 2001; Callahan, 1996).

**Section IV. School-Based Tertiary Suicide Prevention Program, (also called Postvention), refers to the provision of crisis intervention, support, and assistance for those affected by a completed suicide (American Association of Suicidology, 1999; Brock, 2002; Leenaars & Wenckstern, 1998).**

**A. School District Level Responsibilities (with some Building/Site level overlap)**

1. Principal, or designee, verifies facts with the police, hospital, or family of the victim (Brock, 2002) and reports to the superintendent or designee, District Crisis Guidance Manager, Media Coordinator, District Crisis Team, and site/building crisis team.
2. Death should never be referred to as a suicide unless there has been verification by the coroner (Wenckstern & Leenaars, 1993; Poland & McCormick, 1999).
3. Principal or designee will notify appropriate involved school personnel, (ideally within the hour of death verification) (Siehl, 1990).
  - a. Notify all personnel in all district schools and neighboring school districts who will be affected by the death.
  - b. Notify all after-school care providers who will be affected by death.
  - c. Notify all personnel who will participate in postvention responses quickly.
  - d. If possible, notify all school employees impacted by the suicide before the school day begins if death happened outside school hours (American Association of Suicidology, 1999).
  - e. Keep Media Liaison updated about facts as they are received.

**B. Building/Site Level Responsibilities:**

1. Principal or designee mobilizes District Crisis Guidance Manager and building/site crisis team whose members generally have specific team members filling specific roles in postvention efforts (Brock, 2002). Specific roles and duties commonly included are below (see Attachment #8):
  - a. **Team Leader/Crisis Response Coordinator** (usually, school principal or designee):
    - (1) Verifies death.
    - (2) Mobilizes and coordinates crisis response team and its actions.
    - (3) Assesses impact of suicide on the school.
    - (4) Determines if postvention response is warranted.
    - (5) If response is warranted, assess what level of response needed (e.g., few individuals, entire school), being careful to estimate accurately (Brock, Sandoval, & Lewis, 2001).
    - (6) Keeps superintendent or designee informed of the events and responses.
    - (7) Contact family of the suicide victim within 24 hours of the death, in person if possible (Poland & McCormick, 1999; Siehl, 1990).
      - (a) Offer postvention assistance.
      - (b) Ask if they can help identify any of victim's friends who may need intervention (Davis & Sandoval, 1991; American Association of Suicidology, 1999).

- (c) Verify with parents what details about the death can be shared with outsiders (Brock, 2002; Roberts et al., 1998), without agreeing to keep it a secret that the death was a suicide.
- (8) With crisis team, determine how to share about the death.
  - (a) Create a basic written message regarding the death to inform students, parents, and the media.
  - (b) Inform school personnel first, if possible in pre-school meeting.
  - (c) If pre-school meeting not possible, attempt basic message delivery to each classroom simultaneously, with instructions for teacher or designee to share information with students, and place other staff members there to assist as needed.
  - (d) Make arrangements for individuals or small groups of students who are particularly vulnerable or were close to the student who committed suicide to receive counseling with school mental health professionals (American Association of Suicidology, 1999; Berman & Jobes, 1991).
  - (e) Principal generally informs parents/guardians via a letter sent home on the day of student notification of suicide and for students most affected by the death; a personal call would be in good form.
  - (f) Schedule a parent/guardian meeting to address concerns of the parent and offer helping resource information for their children (Brock, 2002) if deemed appropriate and of interest to parents/guardians.
- (9) With crisis team, initiate comprehensive crisis intervention service plans, ideally initiated in the first 24 hours following the suicide and will include (Leenaars, Wenckstern, 1998; Davis & Sandoval, 1991; Davidson, 1989):
  - (a) Individual meetings.
  - (b) Group crisis intervention.
  - (c) Classroom activities and/or presentations.
  - (d) Staff meetings.
  - (e) Parent meetings.
  - (f) Referrals to community agencies
- (10) Conduct a faculty planning session.
  - (a) Include all classified and certificated employees.
  - (b) Provide updated information about the death, background information on the suicide postvention process.
  - (c) Clarify school's postvention role.
  - (d) Inform about suicide contagion, suicide risk factors, and crisis prevention and intervention services that are planned (American Association of Suicidology, 1999; Berman & Jobes, 1991; Poland & McCormick, 1999; Thompson, 1990).
  - (e) Give teachers specific charges to perform, including (Berman & Jobes, 1991).
    - (i) Replacing rumors with facts.



- (ii) Encouraging discussions of feelings and the normality of grief and stress reactions; discouraging any romanticizing of suicide.
  - (iii) Necessity of identifying any at-risk students.
  - (iv) Make appropriate referrals
- (f) Stress the importance of keeping school open, following normal bell schedule, and continuity (Berman & Jobes, 1991; Davis & Sandoval, 1991).
- (g) Allow opportunity for staff to process their own feelings and issues (Davis & Sandoval, 1991; Brock, 2002).
- (11) Crisis team should meet at least once a day throughout the postvention period to review progress and make additional plans (Berman & Jobes, 1991)
- b. **Community Liaison:**
  - (1) Networks with community agencies involved.
  - (2) Coordinates community involvement with crisis intervention coordinator.
  - (3) Keeps records of all community agency involvement at site.
- c. **Parent Liaison:**
  - (1) Coordinates with principal reproduction and distribution of all written communication to be sent home to families of students following suicide.
  - (2) Organizes and facilitates comfortable parent gathering place on campus if needed.
  - (3) Helps prepare, advertise, and organize parent information meeting if appropriate.
- d. **Medical Liaison** (typically a school nurse):
  - (1) Work to ensure student and staff health following the suicide.
  - (2) Act as a liaison with outside medical personnel involved.
- e. **Crisis Intervention Coordinator** (usually a school-based mental health professional such as head school counselor, school psychologist, or school social worker).
  - (1) Identify students and staff most affected by the suicide and initiate postvention assistance and referral procedures.
  - (2) Create and maintain a list identifying students most affected.
  - (3) Begin this process within the first three hours following death verification, with physical and/or emotional proximity to the suicide being key identification variables (Brent et al., 1992; Callahan, 2000; Garfinkel et al., 1988b).
  - (4) Special attention must be given to students believed to be at high-risk themselves for suicide, to include (Davidson, 1989; O'Carroll et al., 1998):
    - (a) Siblings and friends of the suicide victim.
    - (b) Previous suicide attempters.
    - (c) Clinically depressed students.
    - (d) Students known to be with limited social resources

- (5) Arrange for trained counselors to screen all high-risk persons, as well as any other student who wants to talk about the death (O'Carroll et al., 1998; Davis & Sandoval, 1991).
  - (6) With community liaison, secure mutual aid community mental health resource assistance if necessary (Brock, 2002).
  - (7) Due to correlation with the presence of support and less distress among suicide survivors (Callahan, 2000), drop-in counseling centers should maintain operations for several days following the suicide.
  - (8) Crisis team member should follow the suicide victim's class schedule, help teacher discuss the death with children, and reduce the stigma of the "empty chair" (Poland & McCormick, 1999).
  - (9) Individuals who were emotionally or physically proximal to the suicide should be met with separately.
  - (10) No student should be allowed to leave campus without parent consent (Poland & McCormick, 1999).
  - (11) Notify parents in the event any student expresses suicidal ideation.
  - (12) Facilitate distancing the students from identification with the victim in all crisis interventions and ensure students do not glorify the suicidal behavior (Davis & Sandoval, 1988).
- f. **Media Liaison** (usually superintendent, school principal, or designee).
- (1) Avoid identification of the death as a suicide until it is verified by coroner (Garfinkel et al., 1988a).
  - (2) Once the death ruled a suicide by coroner, this information should be acknowledged with discretion in tandem with warnings about suicide contagion, and crisis intervention referrals (American Association of Suicidology, 1999).
  - (3) Needs to work with the press to downplay the incident and not romanticize the death, which has been associated with increases in the suicide rate (Brock, 2002; Davis & Sandoval, 1991; Gould, 2001).
  - (4) Shares essential facts, without excessive details, to quell rumors, which create greater anxiety and can stir factors related to contagion (Davidson, 1989; American Association of Suicidology, 1999).
  - (5) Acts as the single spokesperson for the district, constructing an accurate, singular account of the death for the public as appropriate, while cooperating as much as possible with wishes of decedent's family, without agreeing to keep it a secret that the death was a suicide (Davidson, 1989; Berman & Jobes, 1991).
- g. **Security Liaison** (typically a school vice-principal or designee):
- (1) Ensuring student safety after the crisis.
  - (2) Acts as a liaison with law enforcement officials
  - (3) Responsible for crowd control and monitoring common areas, finding students who are out of their classrooms who might need help (Berman & Jobes, 1991).
- h. **Staff Liaison:**
- (1) Keep staff informed of crisis action plans in progress.
  - (2) Coordinate classroom activities dealing with the trauma of the suicide.

- (3) Coordinate staff referral of highly traumatized students or staff members.
- (4) Conduct staff debriefing as necessary in between the times meetings of the whole are conducted.

### **C. Memorials.**

1. Funerals (Poland & McCormick, 1999; Ruof & Harris, 1988).
  - a. Attendance by students is a parental decision.
  - b. School should not be canceled if the service is during school time.
  - c. School should not be funeral or memorial site.
  - d. For some students, the suicide death may be first encounter with death, and they might appreciate some information about what occurs at a funeral (Berman & Jobes, 1991).
2. School should advise the family of the suicide victim (and the clergy person, if one is involved), to avoid glorification of the act and the need for survivors to distance themselves from the person who committed the act (Ruof & Harris, 1988).
3. Choice of memorial should also not glamorize or glorify the suicide act.
4. Avoid memorial plaques, yearbook dedications, and other song or event tributes in memory of the suicide victim (Brock & Sandoval, 1997; Garfinkel et al., 1988b; Ruof & Harris, 1988).
5. Consider developing a living memorial, such as a contribution to a suicide prevention or student assistance, which will serve to help others cope with problems (Brock & Sandoval, 1997).

### **D. Postvention De-Briefing.**

1. Provide a debriefing for all who helped in the crisis intervention (Ward, 1995).
2. Review and evaluate crisis intervention activities and plan for follow up actions (Brock, 2002).
3. Provide opportunity for mutual support for crisis interveners to help them cope with the crisis as they have been directing their energy to caring for others (Davidson, 1989).

## **Section V. School-Based Suicide Prevention Program Evaluation Plan**

### **A. Formation and maintenance of program evaluation plan.**

- (1) The District Crisis Guidance Manager should coordinate the program evaluation of the school district's suicide prevention program including:
  - a. Develop an assessment tool to evaluate gatekeeper training every even
  - b. Develop an assessment tool to evaluate level of functioning of the District Crisis Team every even year by its participants and building administrators.
  - c. Develop an assessment tool to evaluate the parent/guardian suicide prevention training every even year by the trainers and trainees.
  - d. Develop an assessment tool to help building/site administrators evaluate and document that selected suicide prevention programs are empirically-based and are being implemented with fidelity.
  - e. Each August update the district's suicide prevention program, ensuring the resource and referral names and contact numbers are accurate and current.

## **Section VI. Useful Sample Forms, Checklists, Job Descriptions & Guidelines**

### **A. Twelve Attachments**

1. Sample Policy from CA School Boards Relating to School-Based Suicide Prevention
2. School District Crisis Response (Including Suicide): District Office Responsibilities
3. District Crisis Guidance Manager Responsibilities & Responsibilities & Job Description
4. District Crisis Support Team: Membership & Roles
5. District Media Guidelines
6. Establishing School Crisis Team: School Site Responsibilities
7. School-Based Suicide Prevention Gatekeeper Training Checklist
8. School-Based Suicide Prevention Program Recommendations
9. School Crisis Immediate Response Checklist
10. School Crisis Team Quick Checklist Following a Suicide
11. Suicide Prevention Community Coordination.
12. Sample Form Letters
  - a. **Parent/Guardian Letters**
    - (1) Sample parent/guardian letter inviting to Gatekeeper Training
    - (2) Sample parent/guardian letter in death by accident
    - (3) Sample parent/guardian letter in death by suicide.
    - (4) Sample parent/guardian letter in death by murder
    - (5) Sample parent/guardian letter after a natural disaster
    - (6) Sample parent/guardian informational evening guidelines
  - b. **Staff Letters/Announcements/Documentation/ Forms**
    - (1) Sample staff letter advising of meeting to introduce Crisis and Suicide Prevention Handbook.
    - (2) Sample staff letter advising of schedule of mandatory gatekeeper meetings.
    - (3) Sample staff script to inform students of death of another student or staff member.
    - (4) Sample staff announcement of death of a staff member or his/her

## Attachment 1

### SAMPLE DOCUMENT FOR SCHOOL BOARDS RELATING TO SCHOOL-BASED SUICIDE PREVENTION PARTS A & B

The California School Boards Association (CSBA) provides the following sample school board policy and administrative regulations pertaining to suicide prevention for use by California public school districts.

#### *Part A: Sample School Board Policy for School-Based Suicide Prevention*

Students BP 5141.52(a)

#### SUICIDE PREVENTION

**Note: The following optional policy and regulation are offered for districts that wish to adopt policy providing student, staff and parent/guardian training in suicide prevention. The courts have ruled that a district may be liable to the parents/guardians of a student who committed suicide while under supervision of school staff. Before adopting policy on this topic, the Board should consider district staff and capabilities to ensure the policy will be fully enforced. Once a district has adopted a policy, it has a duty to enforce it.**

The Governing Board recognizes that suicide is a major cause of death among youth and that all suicide threats must be taken seriously. The Superintendent or designee shall establish procedures to be followed when a suicide attempt, threat or disclosure is reported. The district shall also provide students, parents/guardians and staff with education that helps them recognize the warning signs of severe emotional distress and take preventive measures to help potentially suicidal students.

**Note: The following sentence may be expanded to designate the grade levels and subject areas, such as health or physical education, into which this instruction will be integrated.**

**The Superintendent or designee shall incorporate suicide prevention instruction into the curriculum. The Superintendent or designee shall also offer parent education or information, which describes the severity of the youth suicide problem and the district's suicide prevention curriculum. This information shall be designed to help parents/guardians recognize warning signs of suicide, learn basic steps for helping suicidal youth and identify community resources that can help youth in crisis.**

Suicide prevention training for certificated and classified staff shall be designed to help staff recognize sudden changes in students' appearance, personality or behavior which may indicate suicidal intentions, help students of all ages develop a positive self-image and a realistic attitude towards potential accomplishments, identify helpful community resources, and follow procedures established by the Superintendent or designee for intervening when a student attempts, threatens or discloses the desire to commit suicide.

The training shall be offered under the direction of a trained district counselor/psychologist or in cooperation with one or more community mental health agencies.

*(cf. 1020 – Youth Services)*

*(cf. 4131 – Staff Development)*

*(cf. 4231 – Staff Development)*

*(cf. 4331 – Staff Development)*

*(cf. 6164.2 – Guidance/Counseling Services)*

## **SUICIDE PREVENTION**

**BP 5141.52(b)**

**Note: Education Code 49602, which requires confidentiality of information disclosed to a school counselor by students 12 years of age or older, specifically allows such information to be reported to the principal; or parents/guardians of the student when the counselor has reasonable cause to believe that disclosure is necessary to avert a clear and present danger to the student’s health or safety.**

**Staff shall promptly report suicidal threats or statements to the principal or mental health counselor, who shall promptly report the threats or statements to the student’s parents/guardians. These statements shall otherwise be kept confidential. *(cf. 5141 – Health Care and Emergencies)***

The Board endorses the use of peer counselors who can provide an effective support system for students who may be uncomfortable communicating with adults. Peer counselors shall first complete the suicide curriculum and demonstrate that they are able to identify the warning signs of suicidal behavior and rapidly refer a suicidal student to appropriate adults.

***Legal Reference:***

### **EDUCATION CODE**

***49602 Confidentiality of student information***

***49604 Suicide prevention training for school counselors***

### **WELFARE AND INSTITUTIONS CODE**

***5698 Emotionally disturbed youth; legislative intent***

***Management Resources:***

### **CDE PUBLICATIONS**

***Suicide Prevention Program for California Schools, 1987***

***Health Framework for California Public Schools, 1994***

## ***Part B: Sample School Board Administrative Regulation for Suicide Prevention***

**Students**

**AR 5141.52(a)**

## **SUICIDE PREVENTION**

### **Curriculum**

**The district's suicide prevention instruction shall be designed to help students:**

- 1. Understand how feelings of depression and despair can lead to suicide**
- 2. Identify alternatives to suicide and develop new coping skills.**
- 3. Recognize the warning signs of suicidal intention in others.**
- 4. Learn to listen, be honest, share feelings and get help when communication with friends who show signs of suicidal intent.**
- 5. Identify community intervention resources where youth can get help.**

### **Crisis Intervention Procedures**

**District procedures to be followed when a suicide attempt, threat or disclosure is reported shall:**

- 1. Ensure the student's short-term physical safety by one of the following, as appropriate:**
  - a. Securing immediate medical treatment if a suicide attempt has occurred.**
  - b. Securing police and/or other emergency assistance if a suicidal act is being actively threatened.**
  - c. When a suicidal act is less actively threatened but is a serious possibility, keeping the student under continuous adult supervision until the parent/guardian can be contacted and has the opportunity to intervene. If an unsuccessful suicide attempt has been reported or threatened, monitoring the student's actions until the parent guardian can be contacted and has the opportunity to intervene.**
- 2. Designate specific individuals to be promptly contacted, including:**
  - a. The school counselor, psychologist, nurse, and/or principal**
  - b. The student's parent/guardian**
  - c. As necessary, local police or counseling agencies**
- 3. Set forth one or more plans by which the school can transfer responsibility for the student's welfare to the parent/guardian and/or the appropriate support agent or agency.**
- 4. Provide for the timely follow-up by designated school staff regarding the parent/ guardian and student's contact with an appropriate support agent or agency.**

*From California School Boards Association: West Sacramento, CA*

**Used by permission.**

**Copyright 2004.**



## Attachment 2

### **School District Crisis Response:**

#### ***District Office Responsibilities***

- \_\_\_ Develop and implement District Crisis Intervention policies and procedures.
- \_\_\_ Establish a District Crisis Support Team.
- \_\_\_ Train administrators and District Crisis Support Team in policies and procedures
- \_\_\_ Facilitate establishment of crisis teams at all district school and non-school sites.
- \_\_\_ Involve the District Crisis Support Team in all district-wide crisis planning, exercises and prevention program evaluation.
- \_\_\_ Coordinate district-wide crisis suicide prevention gatekeeper skill training
- \_\_\_ Establish and maintain district-wide suicide intervention and postvention protocols and procedures.
- \_\_\_ Establish and maintain a call list of District Crisis Support Team members.
- \_\_\_ Establish and maintain close relationships with community mental health agencies and secure their involvement in planning, training, and mutual aid...
- \_\_\_ Request mutual aid assistance when necessary from neighboring school districts.
- \_\_\_ Establish and maintain list of names and phone numbers for community support systems needed to provide additional services when required.
- \_\_\_ Collect and maintain current psycho-educational materials and other supplies that may be needed to support crisis intervention activities.
- \_\_\_ Respond to requests for assistance 24/7 from local schools and support by mobilizing district resources and providing on-site support to School/Site Crisis Team Leader and District Crisis Guidance Manager
- \_\_\_ Support mutual aid system for neighboring school crisis intervention teams.

Adapted with permission from *School Mental Health Crisis Intervention Teams* – Los Angeles County Office of Education.

### **Attachment 3**

#### **DISTRICT CRISIS GUIDANCE MANAGER RESPONSIBILITIES**

The District Crisis Guidance Manager functions as a general consultant to all schools and departments in the district for issues pertaining to suicide, crisis, death, and grief.

***District Crisis Guidance Manager is contacted:***

1. For support and resources whenever an issue of suicide, crisis, death, or grief impacts student or staff members.
2. In consultation with School Crisis Team, to determine if the help of the District Crisis Support Team is needed at the site.
3. To further decide if any other support staff or outside resources need to be called to offer assistance.
4. To mobilize outside resources if appropriate.

***District Crisis Guidance Manager Responsibilities:***

1. Make sure school district employees know the identity and function of the District Crisis Guidance Manager and how to access that person.
2. Introduce self and role to parents of students via PTA newsletters and meetings.
3. Contact and enlist the cooperation of community resources.
4. Be available for on-going review and debriefing post-crisis.
5. Maintain a file of letters, memos, and other exchanges developed to communicate the crisis information.
6. Prepare and provide resource materials on grief, loss, traumatic stress reactions and make them available for loan to individuals and sites needing them.
5. Develop, coordinate and lead prevention program plan evaluations.

Adapted with permission from: School Crisis Response Teams: Lessening the Aftermath Training Manual. 1993 by Mary Schoenfeldt. Third Edition.

## Attachment 4

### **DISTRICT CRISIS SUPPORT TEAM: MEMBERSHIP & ROLES**

The District Crisis Support Team functions as a back up support system to all schools and departments in the district for issues pertaining to suicide, crisis, death, and grief. The District Crisis Support Team responds when asked by a school site administrator or the crisis guidance manager. They are not immediately deployed into service for every crisis.

#### ***District Crisis Support Team Membership:***

1. District Crisis Guidance Manager should serve on teams at all levels.
2. Team should also include building principals, psychologists, nurse, counselors, social workers, and other support staff.
3. Volunteers commit to a minimum of two years and perform this duty in addition to regularly assigned roles.
4. All members should have crisis intervention experience and be provided training.
5. Team members share information with co-workers as appropriate.
6. Team members make arrangements with supervisors so they can respond if needed to other sites in the school district.
7. There should be 4-5 members of a team for each level (elementary, junior high/middle school, senior high).
8. Ad-hoc members may be added if necessary to respond to a crisis, (to include any staff member deemed helpful to a given situation by virtue of background, skill, or relationship to the people in crisis).
9. Volunteers from departments of Student Services, Special Education, E.S.L./Bilingual and other departments as appropriate.

#### **District Crisis Support Team Role:**

1. As a **BACK-UP** when a School Crisis Team is unable to handle a situation according to their site plan in situations such as:
  - a. Number of students traumatized so large it is not possible to support them without extra help.
  - b. School staff members so shocked or saddened by event they want assistance to provide crisis intervention.
  - c. The School Crisis Team is unsure of appropriate way to handle the particular situation and asks for help.
2. Help facilitate establishment of crisis teams at school and non-school district sites if needed.
3. Help with district-wide training, exercises, and prevention program evaluation.

Adapted with permission from: *School Crisis Response Teams: Lessening the Aftermath Training Manual*. 1993 by Mary Schoenfeldt. Third Edition.

## Attachment 5

### DISTRICT MEDIA GUIDELINES

- 1. General Guidelines for School Administrators /Designated Media Liaison.**
  - a. Designated media liaison is individual who interacts with reporters, nobody else.
  - b. Assist news professionals in reporting responsibly and accurately.
  - c. Avoid "no comment", instead using media request as an opportunity to influence content of story, always being sure to give information about community crisis resources and other services available.
  - d. Develop a district policy that gives media professionals seeking to report on sensitive stories related to school some clear and fair guidelines in advance of a tragedy.
  - e. Crisis Manager annually prepares media kits for local media professionals annually, sharing the research on media and suicide contagion and name designated media liaison and designee.
  - f. Strive to have a positive relationship with media professionals as it benefits both parties--this creates more opportunities for positive messages and helpful information to be shared.
- 2. Specific Guidelines for School Administrators /Designated Media Liaison in event of suicide.**
  - a. Explain to media professionals the potential for suicide contagion linked with certain types of reporting and ways to minimize risk:
    - (1) Presenting simplistic explanations for suicide is a problem.
    - (2) Suicide usually a result of complex interaction of factors.
    - (3) Most people have a history of problems, often unknown in aftermath of suicide.
    - (4) Detailed descriptions of suicide not needed, but acknowledgement of problems is recommended.
    - (5) Engaging in repetitive, prominent, and excessive reporting of suicide prolongs preoccupation with suicide among those at risk and is associated with suicide contagion.
  - b. Reinforce with media professionals that sensationalized news coverage of a suicide also raises people's preoccupation with suicide and has been linked to suicide contagion.
    - (1) Limit morbid details to tone the sensationalism down.
    - (2) Decrease prominence of suicide story.
    - (3) Avoid using dramatic images related the suicide.
  - c. Reporting technical details of the means that a suicide was completed is also related to suicide contagion and is known to encourage imitation.
  - d. Stories that paint suicide as an acceptable coping mechanism for stress or problems may encourage that as a solution for at-risk people. Instead:
    - (1) Use the suicide story as opportunity to present alternative, more positive ways to cope, which has been related to prevention of further suicide attempts.
    - (2) Encourage reporters to report about times that other people who, when challenged, chose a more pro-life option.
  - e. Community expressions of grief over a suicide should not be focused on in a large way by the media.

- (1) Such actions exacerbate suicide contagion by the suggestion the suicidal behavior is being honored, rather than the dead person mourned.
- (2) Limit reporting of large memorial gestures, (public eulogies, half-mast flags, permanent memorials).
- f. Only focusing on the positive characteristics of a suicide completer in the news and in public forums may elicit contagion if not accompanied by acknowledgement of the deceased person's problems. A person vulnerable to suicide contagion may interpret such press as positive reinforcement for imitating the suicidal act.
- g. News stories dealing with suicide are given an important opportunity to feature information about seeking help and illuminating that suicide can be prevented.
- h. Be proactive with media professionals around difficult issues. Contact editor before reporters come to site and re-iterate media guidelines and threat of suicide contagion.

**Attachment 6**

**ESTABLISHING SCHOOL CRISIS TEAM:**

*SCHOOL SITE RESPONSIBILITIES*

The School Crisis Team is a group of employees from the school site who are knowledgeable enough about their school community, its students and staff, to make necessary decisions when a crisis occurs in the district for issues pertaining to suicide, crisis, death, and grief. The school crisis team leader (usually the principal) should convene the school crisis team in the event of a completed suicide or a serious suicide attempt by a student or staff member.

\_\_\_\_\_ **Establish School Crisis Team Leader/Designee** (Typically the Principal):

\_\_\_\_\_ **Team Leader: Assigned:** \_\_\_\_\_  
Call team together to plan response; summarize crisis situation; facilitate development of School Crisis Response Plan; respond to media personally or through District Office; act as District Office liaison; conduct debriefing/evaluation of response plan.

**School Crisis Team Membership:**

Team leader should select the team from willing staff volunteers who agree to a one-year term or longer. Size of team varies with school size. Try to include volunteers from these ranks:

- |                                  |                                |
|----------------------------------|--------------------------------|
| _____ Asst. Principal            | _____ School Psychologist      |
| _____ Campus Security Supervisor | _____ School Secretary         |
| _____ School Counselor           | _____ School Teacher           |
| _____ School Nurse               | _____ Ad-hoc, as needed: _____ |

The Team Leader should request the following information be assembled and distributed to Crisis Team:

- \_\_\_\_\_ Names and phone numbers of members of the District Crisis Support and School Crisis Teams (home and work).
- \_\_\_\_\_ Name and phone number of Team Leader and at least two back-up leaders.
- \_\_\_\_\_ Responsibilities of each member depending on the crisis.
- \_\_\_\_\_ Determine person (and back-up) responsible to call team together following Roles of School Crisis Team

**Suggested Team Member Roles:**

- \_\_\_\_\_ **Community Liaison:** Assigned: \_\_\_\_\_  
Network with community agencies involved in plan; coordinate community involvement with school psychologist; keep records of all community agency involvement at site; update other team members as needed.
- \_\_\_\_\_ **Parent Liaison:** Assigned: \_\_\_\_\_  
Contac family of deceased; contact parents as needed; coordinate any communication to be sent home; conduct parent meeting if appropriate.

\_\_\_\_\_ **Medical Liaison:** Assigned: \_\_\_\_\_  
Typically school nurse; plan to ensure student health following the crisis and act as a liaison with medical personnel.

\_\_\_\_\_ **Crisis Intervention Coordinator:** Assigned: \_\_\_\_\_  
Typically school counselor/psychologist; acts as referral resource for students and staff; serve as in-take counselor for highly traumatized students; consult with staff; help screen students at risk; develop support systems as needed; conduct group meetings with parents/staff as needed.

\_\_\_\_\_ **Security Liaison:** Assigned: \_\_\_\_\_  
Typically school administrator; plans to ensure student safety after crisis and acts as liaison with law enforcement; responsible for crowd control and monitoring common areas to find students who might need help or are out of class.

\_\_\_\_\_ **Staff Liaison:**  
Assigned: \_\_\_\_\_  
Keep staff informed of action plan; coordinate classroom activities dealing with trauma; coordinate referral system for highly traumatized students; conduct staff debriefing.

**Suggested Training for School Crisis Team:**

- |  |                                   |
|--|-----------------------------------|
| _____ Critical incident stress debriefing                | _____ Violence/bully prevention   |
| _____ Team building                                      | _____ Drug abuse prevention       |
| _____ Active Listening                                   | _____ Effective media relations   |
| _____ Small group techniques for children and adults     | _____ Grief and loss intervention |
| _____ First aid and how to call for medical assistance   | _____ Suicide intervention skills |
| _____ Crowd control and evacuation of students and staff | _____ Anger management            |

Adapted with permission from: School Crisis Response Teams: Lessening the Aftermath Training Manual. 1993 by Mary Schoenfeldt. Third Edition.

## Attachment 7

### SCHOOL-BASED SUICIDE PREVENTION GATEKEEPER TRAINING CHECKLIST

Gatekeepers in the school setting are adult employees who have contact with students and colleagues as part of their usual routine. Gatekeeper training provides these gatekeepers information that will help them identify signs that may indicate suicidality or other mental health issues. It and instructs them on who they should inform and how to make a referral when they have concerns. It gives research-based information about risk and protective factors. District Crisis Guidance Manager will organize this training and manage data about training attendance and frequency.

#### School-Based Suicide Prevention Gatekeeper Training Checklist:

- \_\_\_\_\_ Recruit school or community mental health professional(s) knowledgeable in suicide prevention to conduct training.
- \_\_\_\_\_ Plan for trainer(s) to receive continuing education in suicide prevention methodology and research to remain current.
- \_\_\_\_\_ Institute mandatory school gatekeeper training and repeat the training every two years for all school and non-school site personnel, certified and classified.
- \_\_\_\_\_ Training to develop and increase knowledge, attitude, and skills in:
  - \_\_\_\_\_ The nature of suicidal behavior
  - \_\_\_\_\_ De-stigmatizing mental illness
  - \_\_\_\_\_ Depression and biological basis of some mental illness.
  - \_\_\_\_\_ Identification of students/staff who may be at risk for suicide or mental illness, warning signs, and contagion.
  - \_\_\_\_\_ Knowledge of school's policy and procedures when a student or staff member threatens suicide and/or exhibits suicidal behavior either verbally, in writing, or by action.
  - \_\_\_\_\_ Steps used in determining levels of risk and who does that.
  - \_\_\_\_\_ Referral procedures and school and community referral resources.
  - \_\_\_\_\_ Schools role in suicide prevention.
  - \_\_\_\_\_ Means restriction.
  - \_\_\_\_\_ Understanding content of suicide prevention programs that will be taught to students.
  - \_\_\_\_\_ Understanding of roles and responsibilities of District Crisis Guidance Manager, District Crisis Support Team, and School Crisis Team, and identification of the members of each.
  - \_\_\_\_\_ Knowledge about how to interact with and assist family and friends in the aftermath of a suicidal event.



## Attachment 8

### **SCHOOL-BASED SUICIDE PREVENTION PROGRAM RECOMMENDATIONS: CURRICULUM-BASED AND SCREENING PROGRAMS**

There is tension in the literature about the usefulness and efficacy of curriculum-based suicide prevention/awareness programs. The National Strategy of Suicide Prevention (DHHS, 2000) put out a call for evidence-based suicide prevention programs and early mental health screenings, assessments, and referral services. With a sudden increase in the number of suicide prevention programs being developed, the Suicide Prevention Resources Center (SPRC), together with the American Foundation of Suicide Prevention (AFSP), began to develop a registry of effective programs, the National Registry of Effective Programs, (NREP). These effective programs had to meet certain standards in order to be found "evidence-based". These reviewed programs were classified as "effective", "promising", or "unrated" based on reviews using ten scoring criteria that were heavily weighted toward integrity and utility (SPRC, 2005). The NREP registry has been discontinued since January 2005 and its duties are being subsumed by the newly formed National Registry of Evidence-Based Programs and Practices (newly labeled the NREPP), which is under the Substance and Mental Health Services Administration (SAMSHA). The NREPP is a new rendition of the former NREP. All programs that had qualified as evidence-based suicide prevention programs under the NREP will have to undergo new review to be eligible for the new NREPP effective, evidence-based program registry.

#### **Summary of the reviewed programs found to be either “effective” or “promising”:**

It is highly likely new programs will be continue to be reviewed and added as effective or promising. The list below identifies the programs currently identified as either effective or promising, giving a school district a good place to begin as it seeks to employ research-based suicide prevention methods in its pursuit to help students and staff prevent suicide. Schools have been asked to participate in suicide prevention and screening efforts by the National Strategy for Suicide Prevention and the President’s New Freedom Commission on Mental Health: Achieving the Promise: Transforming Mental Health Care in America.

#### **A. C-Care/CAST (Thompson, Eggert, Randell, & Pike, 2001):**

This program is a school-based intervention for students at risk for suicide. It combines one-to-one (C-Care for Counselors Care) with a series of small group sessions (CAST for Coping and Support Training).

1. SPRC Classification: Effective.
2. Target Age: 14-18.
3. Gender: Male & Female.
4. Ethnicity/Race: Multiple.
5. Level of Intervention: Selective, Indicated.
6. Increased Protective Factors: Skills in problem solving, conflict resolution, and nonviolent handling of disputes.

7. Decreased Risk Factors: Mental disorders, particularly mood disorders, schizophrenia, anxiety disorders and certain personality disorders, hopelessness.
8. Cost: Price not specified, but indicated cost involved.
9. Contact info:  
Reconnecting Youth Prevention Research Program  
Phone: 206-543-8555;  
Email: [elainet@u.washington.edu](mailto:elainet@u.washington.edu);  
Elaine Thompson, PhD, RN  
Psychosocial & Community  
Health -Campus Box 357263  
UW School of Nursing - Seattle, WA 98195-7263

#### **B. SOS: Signs of Suicide (Aseltine & DeMartino, 2004)**

This program uses a two-prong approach by combining a suicide awareness curriculum and a brief depression screening. It clearly teaches suicide as directly related to mental illness, and not a reaction to stressful circumstances. The Signs of Suicide (SOS) program has been designated as a “promising program” by the federal Substance Abuse and Mental Health Services Administration (SAMHSA), becoming the only suicide prevention program selected for its National Registry of Effective Prevention Programs (the old NREPP) (Screening for Mental Health, Inc., 2002) and the first to empirically demonstrate significant reductions in self-reported suicide attempts.

1. SPRC Classification: Promising
2. Target Age: 14-18.
3. Gender: Male & Female.
4. Ethnicity/Race: Multiple.
5. Level of Intervention: Universal.
6. Increased Protective Factors: Easy access to variety of clinical interventions & support for help seeking, cultural and religious beliefs that discourage suicide and support self-preservation, strong family and community support.
7. Decreased Risk Factors: Stigma associated with help-seeking behaviors.
8. Language Availability: Spanish language versions available.
9. Cost: Approximately \$200
10. Contact Info: **Screening for Mental Health, Inc.**  
SOS Suicide Prevention Program for Secondary Schools  
One Washington Street, Suite 304  
Wellesley Hills, MA 02481  
Web site: [www.mentalhealthscreening.org/sos\\_highschool](http://www.mentalhealthscreening.org/sos_highschool)  
Email: [highschool@mentalhealthscreening.org](mailto:highschool@mentalhealthscreening.org)  
Phone: 781-239-0071; Fax: 781-431-7447

#### **C. Columbia TeenScreen (Shaffer, et al., 2004):**

This program seeks to identify youth at risk for suicide and are suffering from mental illnesses yet to be diagnosed using screening instruments.

1. SPRC Classification: Promising.
2. Target Age: 11-18.
3. Gender: Male & Female.
4. Ethnicity/Race: Varied.
5. Level of Intervention: Universal.
6. Increased Protective Factors: Easy access to clinical interventions and support for help-seeking.
7. Decreased Risk Factors: Alcohol and substance abuse disorders; Mental disorders, particularly mood disorders, schizophrenia, anxiety disorders and certain personality disorders, impulsive and/or aggressive tendencies; Relational or social loss.
8. Decreased Risk Factors: Barriers to accessing health care, particularly mental health and substance abuse treatment.
9. Cost: ColumbiaTeenScreen© is currently offering materials and consultation for free.
10. Contact Info: Columbia TeenScreen© Program  
1775 Broadway, Suite 715  
NY, NY 10019  
Phone: 1-800-TEENSCREEN (833-6727)  
Email: [teenscreen@childpsych.columbia.edu](mailto:teenscreen@childpsych.columbia.edu)  
Website: [www.teenscreen.org](http://www.teenscreen.org)

**D. Lifelines** (Kalafat & Elias, 1994)

Lifelines is a school-based program made up of four 45-lessons of suicide prevention awareness material. It also includes model policies and procedures for schools to adopt, as well as gatekeeper presentations for educators and parents/guardians.

1. SPRC Classification: Promising.
2. Target Age: 12-17.
3. Gender: Male & Female.
4. Ethnicity/Race: Multiple.
5. Level of Intervention: Universal
6. Increased Protective Factors: Easy access to variety of clinical interventions & support for help-seeking; Strong family and community support connections.
7. Decreased Risk Factors: Stigma associated with help-seeking behavior.
8. Cost: Varies depending on number staff/students. Manual \$40.
9. Contact Info: John Kalafat, PhD  
Rutgers University  
Graduate School of Applied and Professional Psychology  
152 Frelinghuysen Road  
Piscataway, NJ 08854  
Phone: 732-445-2000 x121; FAX: 732-445-4888  
Email: [kalafat@rci.rutgers.edu](mailto:kalafat@rci.rutgers.edu)

**E. Reconnecting Youth Class (RY)** (Eggert & Nicholas, 2004)

This program targets students in grades 9-12 who are achieving poorly, potential for drop out, and other at-risk type behaviors. It endeavors to build resiliency skills against risk factors, give social support and life skills training, and reduce substance abuse and depression/aggression issues.

1. SPRC Classification: Promising.
2. Target Age: 14-18.
3. Gender: Male & Female.
4. Ethnicity/Race: Multiple.
5. Level of Intervention: Selective, Indicated.
6. Increased Protective Factors: Strong family and community support connections; Skills in problem solving, conflict resolution, and nonviolent handling of disputes.
7. Decreased Risk Factors: Alcohol & substance abuse disorders; Mental disorders, particularly mood disorders, schizophrenia, anxiety disorders, and certain personality disorders, impulsive and/or aggressive tendencies; Relational or social loss.
8. Costs: Curriculum Guide - \$300; 10 Student Workbooks - \$212; RY Training (8 participants/1 trainer) \$3000; Other varies.
9. Contact Info: RY Program Information & Scheduling  
Ms. Beth McNamara  
Phone: 425-861-1177  
FAX: 206-726-6049  
Email: [beth.mcnamara@comcast.net](mailto:beth.mcnamara@comcast.net)

#### **F. Zuni Life Skills Development (LaFromboise, 1995)**

This program is based on social cognition theory, is culturally tailored to American Indian youth, and seeks to develop competency in a range of life skills.

1. SPRC Classification: Promising
2. Target Age: 14-18
3. Gender: Male & Female
4. Ethnicity/Race: American Indian
5. Level of Intervention: Selective
6. Increased Protective Factors: Easy access to a variety of clinical interventions & support for help-seeking; Skills in problem-solving, conflict resolution, and nonviolent handling of disputes.
7. Decreased Risk Factors: Hopelessness.
8. Cost: Life Skills Curriculum Text - \$30.00; Other materials vary.  
Contact info:  
Teresa D. LaFromboise, PhD, Associate Professor of Education  
Stanford University - Cubberly 216, 3096  
Stanford, CA 94305-3096  
Phone: 650-723-1202; FAX: 650-725-7412  
Email: [lafrom@stanford.edu](mailto:lafrom@stanford.edu)

- G. The programs listed below were noted in the SPRC Registry as not yet proven, but SPRC gave information about them and indicated they will continue to see how the evidence for them does or does not continue to build.

1. **ASIST**

This program is a 2 day workshop designed to impart gatekeeper skills & knowledge.

- a. SPRC Classification: Unrated at this time
- b. Target Age: 18+
- c. Gender: Male & Female
- d. Ethnicity/Race: Multiple
- e. Level of Intervention: Selective, Universal
- f. Increased Protective Factors: Easy access to a variety of clinical interventions & support for help-seeking; Strong family and community support connections.
- g. Decreased Risk Factors: Lack of social support & sense of isolation; Barriers to accessing health care, particularly mental health & substance abuse treatment.
- h. Cost: Approximately \$100.00.
- i. Contact Info: LivingWorks Education, Inc.  
Website: [www.livingworks.net](http://www.livingworks.net)  
Email: [info@livingworks.net](mailto:info@livingworks.net)  
Phone: 403-3009-0242

2. **Yellow Ribbon**

This program is based on the model of promoting help-seeking behavior, gatekeeper training, and public awareness of suicide prevention.

- a. SPRC Classification: Unrated at this time
- b. Target Age: All Ages.
- c. Gender: Male & Female.
- d. Ethnicity/Race: Multiple.
- e. Level of Intervention: Selective, Universal.
- f. Increased Protective Factors: Easy access to a variety of clinical interventions & support for help-seeking; Strong family and community support connections.
- g. Decreased Risk Factors: Stigma associated with help-seeking behavior.
- h. Costs: Variable; offered on a sliding scale.
- i. Contact info: Yellow Ribbon Program  
Website: [www.yellowribbon.org](http://www.yellowribbon.org)  
Email: [Ask4help@yellowribbon.org](mailto:Ask4help@yellowribbon.org)  
Phone: 303-429-3530

**Attachment 9**

**SCHOOL CRISIS IMMEDIATE RESPONSE: BRIEF CHECKLIST**

**I. IMMEDIATE SCHOOL SITE RESPONSE: Team Leader/ Designee.**

- \_\_\_\_ A. Identify problem/event and determine degree of impact on school.
- \_\_\_\_ B. Be certain appropriate “alarms” have been sounded.
- \_\_\_\_ C. Notify District Superintendent or Designee.

- \_\_\_\_\_D. Assemble Site Crisis Team, contact Superintendent (or designee), and District Crisis Guidance Manager.
- \_\_\_\_\_E. Designate individual coordinating responsibilities & locations.
- \_\_\_\_\_F. Clarify additional resources to be called in:
  1. American Red Cross.
  2. Children's Welfare - CPS.
  3. Fire Department.
  4. Police.
  5. Hospitals.
  6. Poison Control Center: 800-
  7. Suicide Prevention & Crisis Services.
  8. County Mental Health.
  9. Other: \_\_\_\_\_.
- \_\_\_\_\_G. Alter daily/weekly schedule as needed.

**II. COMMUNICATION: Team Leader/ Designee**

- \_\_\_\_\_A. Designate Crisis Coordinating Center
- \_\_\_\_\_B. Establish visitor sign-in Procedures (at all campus entry sites as possible/feasible)
- \_\_\_\_\_C. Principal or designee:
  1. Review facts & determine what information should be shared.
  2. Consider police investigation parameters.
  3. Notify family with sensitivity and dispatch. (Identify person to contact family).
- \_\_\_\_\_D. Develop and disseminate bilingual FACT SHEET (written bulletin) for
  1. Staff.
  2. Students.
  3. Parents/Community.
- \_\_\_\_\_E. Identify media liaison.
  1. Principal and media spokesperson confer on information to be shared.
  2. Designate a location for media representatives.
- \_\_\_\_\_F. Contact other district schools, (priority to schools of affected students' siblings).
- \_\_\_\_\_G. Principal and Crisis Counseling Coordinator coordinate counseling efforts:
  1. Classroom presentations/discussions.
  2. Parent/community meetings.
  3. School staff meeting.
- \_\_\_\_\_H. Provide for RUMOR CONTROL.
  1. Keep a TV set or radio tuned to a news station.
  2. Verify ALL facts heard.
  3. Update Fact Sheet as needed.
  4. Utilize student leaders, as appropriate: (as sources knowledgeable of student rumors; as peer leaders to convey factual info; as runners (written bulletins should be sealed when necessary)).

**III. FIRST AID AND EMERGENCY RELEASE PLAN: Nurse**

- \_\_\_\_\_A. Initiate First Aid Team procedures

- \_\_\_\_\_ B. Designate Emergency Health Office Location: \_\_\_\_\_.
- \_\_\_\_\_ C. Initiate Emergency Release Plan procedures if necessary
- \_\_\_\_\_ D. Designate student checkout location \_\_\_\_\_

**IV. PSYCHOLOGICAL FIRST AID: Crisis Intervention Coordinator**

- \_\_\_\_\_ A. Logistics: Designate rooms/locations/areas and assign responsibilities.
  - 1. Individual counseling-Location: \_\_\_\_\_ Whom: \_\_\_\_\_
  - 2. Parents- Location: \_\_\_\_\_ Whom: \_\_\_\_\_
  - 3. Staff: \_\_\_\_\_ Whom: \_\_\_\_\_
  - 4. Sign-in for Support Services: \_\_\_\_\_ Whom: \_\_\_\_\_
- \_\_\_\_\_ B. Initiate referral process, including procedures for self-referral.
  - 1. Identify location of crisis team members.
  - 2. Provide bilingual services as needed.
  - 3. Distribute appropriate forms to staff for student counseling referrals.
  - 4. Distribute summary of students who were referred to all staff daily.
- \_\_\_\_\_ C. Disseminate appropriate supportive information to staff and parents.
- \_\_\_\_\_ D. Identify and coordinate contact of high-risk students.
- \_\_\_\_\_ E. Identify and coordinate contact other highly affected students & staff.
- \_\_\_\_\_ F. Plan & initiate appropriate interventions (individual counseling, group counseling, parent/community meetings, meetings with ENTIRE staff, classroom activities/presentations, referrals to community agencies)
- \_\_\_\_\_ G. Schedule support and time out breaks for crisis workers.


**V. DAILY DE-BRIEFING OF SCHOOL CRISIS-**

**Team Leader, Team, District Crisis Guidance Manager:**

- \_\_\_\_\_ A. Crisis Intervention Activities.
  - 1. Review actions of each day.
  - 2. Identify weaknesses & strengths of crisis responses.
  - 3. Review status of referred students.
  - 4. Prioritize needs/personnel needed next day & plan follow-up activities.
  - 5. Allow time for emotional debriefing of each crisis team member.

**Attachment 10**

**SCHOOL CRISIS TEAM QUICK CHECKLIST FOLLOWING A SUICIDE**

Check if  done.	Tasks to be Done	Person Responsible for Task (Identify before a crisis)
	Verify Facts	
	Contact School Superintendent & District Crisis Guidance Manager	
	Contact Staff (phone tree)	
	Convene School Crisis Team	
	Identify Family Contact Person	
	Arrange for Substitute Teachers	
	Write Announcement to Students	
	Morning Staff Meeting	
	Set up Safe Rooms (places for very upset individuals to de-brief privately)	
	Distribute Suggestions for Classroom Discussion	
	Notify Students	
	Provide List of Readings & Materials to Teachers	
	Write & Send Letter to Parents/Legal Guardians	
	After School Staff Meeting	
	Parent, Family, Community Meeting	
	Carefully decide on a school memorial/remembrance not glorifying the act of suicide	
	Post Intervention Debriefing	
	Follow up with Students	

Adapted with permission from: Canter, A. S., & Carroll, S. A. (Eds.). (1999). Crisis Prevention & Response: A Collection of NASP Resources. Bethesda, MD: National Association of School Psychologists



**Attachment 11**

**SUICIDE/CRISIS PREVENTION AND RESPONSE:  
COMMUNITY COORDINATION:  
Community Mutual Aid Checklist**

*Every school district should have a formalized, coordinated agreement and understanding about mutual aid in the event of a suicide that requires help from one or more of the following type of community providers and neighbors. This Checklist should be continually updated by the District Crisis Guidance Manager, (annually at the minimum).*

<b>AGENCY/NAME</b>	<b>PHONE NUMBER</b>	<b>CONTACT PERSON</b>
American Red Cross		
City Fire Department		
Clergy/Religious Leaders		
Community/County Mental Health		
Coroner's Office		
County Juvenile Probation		
County Office of Education		
County Office of Emergency Services		
County Sheriff		
District Attorney		
Hospice		
Local Bus		
Medical Hospital/Clinic		
Neighboring School District(s)		
Newspaper		
Police Department		
Psychiatric Hospital Facilities:		
Radio Stations		
Sexual Assault & Domestic Violence Center		
Suicide Prevention Agency		
Survivors' Group		
TV Stations		

## Attachment 12

### **SAMPLE LETTERS AND FORMS FOR PARENT/GUARDIAN AND STAFF**

**In a crisis situation it is important that the voice of the individual site be heard in correspondences, but sometimes in times of crisis it helps to have a prompt. There are several sample letters/announcements below intended to help site administrators and Crisis Teams get started as they compose their own thoughts and words in the following situations. They serve as a rough template and it is expected that they will be tailored to your individual site and its situation and on your school site's own letterhead. (Adapted with permission from: School Crisis Response Teams: Lessening the Aftermath Training Manual. 1993 by Mary Schoenfeldt. Third Edition).**

#### **1. Parent/Guardian Letters**

- a. Sample parent/guardian letter inviting to Gatekeeper Training
- b. Sample parent/guardian letter in death by accident
- c. Sample parent/guardian letter in death by suicide.
- d. Sample parent/guardian letter in death by murder
- e. Sample parent/guardian letter after a natural disaster
- f. Sample parent/guardian informational evening guidelines

#### **2. Staff Letters/Announcements/Documentation/ Forms**

- a. Sample staff letter advising of meeting to introduce Crisis and Suicide Prevention Handbook.
- b. Sample staff letter seeking School Site Crisis Team volunteers.
- c. Sample staff application for School Site Crisis Team volunteers.
- d. Sample staff letter seeking District Crisis Support Team volunteers.
- e. Sample application for District Crisis Support Team volunteers
- f. Sample staff letter advising of schedule of mandatory gatekeeper meetings.
- g. Sample staff script to inform students of death of another student or employee.
- h. Sample staff announcement of death of a staff member or his/her family.
- i. Sample form for documentation of suicidal behavior or threat.
- j. Sample Guidelines after Assessing Suicidal Risk - Checklist
- k. Sample documentation & referral form for parent/guardians.

## 1. Parent/Guardian Letters

### a. Invitation to parents/guardians-Gatekeeper Training

Dear \_\_\_\_\_:

Our school district has made a commitment to school-based suicide prevention education for our students, which includes gatekeeper training for staff and interested parents/guardians. Gatekeepers are people who see the children routinely and may notice when they are having difficulty emotionally. Gatekeeper training serves to teach gatekeepers to identify signs that may signal emotional difficulty and equip the gatekeepers with information about referral and getting the student the appropriate help. Parent/Guardian Gatekeeper Training will occur:

Date:

Time:

Place:

Child care will be provided by school employees who have volunteered to help make this opportunity available to more people. This presentation will be jointly facilitated by \_\_\_\_\_, one of our school psychologists; \_\_\_\_\_, our District Crisis Guidance Manager; and \_\_\_\_\_, of our county Suicide Prevention Agency. We anticipate this information will help us all be more aware of individuals who may be considering suicide, who to contact if we are concerned, and what resources are available in the schools and the community. Thank you for considering this important opportunity.

Sincerely,

**(Principal name inserted)**

**b. Sample parent/guardian letters in death by accident of student or staff member**

Dear Parents/Guardians:

It is with great sadness that I share the following news with you. (i.e. This morning one of our **(insert grade number)** grade students, **(insert name)**, hit by a car outside of **his/her** home in \_\_\_\_\_. According to **his/her** family, **he/she** ran out into the street and was seriously injured. **He/she** died at the hospital. We are profoundly saddened by **his/her** death).

We have shared this information and had discussions with all of our students at **(insert school's name)** School so they know what has happened. Counselors, teachers, and other support personnel have been on hand to assist and will continue to be available to students, staff, and parents as long as needed. Please call the school **(insert phone number)** if you want assistance.

As parents and guardians, we urge you to talk to your children about this sad event, also. The death of a (student/teacher) may affect a child in a variety of ways, depending on the age of the student, how well the child knew Sally, and the child's prior experience with death and grief. Typical reactions children have to a death include:

- Appear to be unaffected
- Think about it privately
- Fright
- Sadness & withdrawal
- Agitation & anger
- Ask many questions
- Attempts to be extra good

May I suggest that you listen to your children intently? If your child seems to need to talk, answer their questions simply, honestly, and possibly over and over again. (optional) A Parent Informational Night is planned for:

Date:

Time:

Place:

At that time we can talk further about helping children through grief.

Our thoughts are with the family and friends of **(insert child's name)**.

Sincerely,

**(Principal name inserted)**

**c. Sample guardian/parent letter in death by suicide-*ONLY share this information if the coroner has confirmed the death was a suicide and if the site administrator has so advised.***

Dear Parents and Guardians:

This letter is sadly to inform you a student, **(insert student's name)**, has died. The coroner has reported the cause of death to be a completed suicide

We have shared this information and had discussions with all of our students at **(insert school's name)** School so they know what has happened. Counselors, teachers, and other support personnel have been on hand to assist and will continue to be available to students, staff, and parents as long as needed. Please call the school **(insert phone number)** if you want assistance.

We have had counselors and extra staff on campus today to help meet the emotional needs of students and staff who have been upset by this tragedy. We will continue to have extra mental health assistance for students and staff as long as it appears necessary. I urge you to discuss this event with your child and explain that suicide is not a positive response to life's challenges. Please explain that experts tell us most people who attempt or complete suicide have many emotional problems for which there is help available. Emphasis the importance of people seeking help when needed. This death of a **(student/teacher)** may affect children in a variety of ways, depending on the ages of the students, how well they knew Sally, and the children's prior experiences with grief and death, (particularly a suicide). Typical reactions children have:

- Appear to be unaffected
- Think about it privately
- Fright
- Sadness & withdrawal
- Agitation & anger
- Ask many questions
- Attempts to be extra good
- Guilt
- Wanting to be with peers

This is an important time to listen to your children intently. If your children seem to need to talk, answer their questions simply, honestly, and possibly over and over again.

I have enclosed a flyer of community mental health resources that are available should this event trigger any undue or exaggerated reactions to your child or someone else you know. Please advise us if you have any serious concerns about the effect this event is having on your child or any of your child's friends. This is a time we must be eyes and ears for each other as we seek to nurture and protect the children in our midst and teach them how to seek help when it is needed.

Please join us in mourning the **(insert name) that** has died, but not glamorizing, and thereby positively reinforcing, the method by which **he/she** died. Suicide contagion is always an issue we must be mindful of in matters such as these. A positive way to honor this child is to use this tragic suicide as a teachable moment. We must reinforce positive options our children can employ when life is difficult and how to access those options.

*(Optional)* A Parent Informational Night is planned for:

Date:

Time:

Place:

At that time we can talk further about helping children through the grief and unsettled feelings they might be having in the wake of this tragedy.

Our thoughts are with the family and friends of **(insert name of deceased)**.

Sincerely,

**(Principal name inserted)**

**d. Sample parent letter in death by murder**

Dear Parents/Guardians:

A very sad thing happened today that I need to share with you. A student, **(insert child's name)**, an **(insert grade number)** grade student at **(insert name of school)** was murdered **(insert when & where if known)**. We are profoundly saddened by **his/her death.**

We have had counselors and extra staff on campus today to help meet the emotional needs of students and staff who have been upset by this tragedy. We will continue to have extra mental health assistance for students and staff as long as it appears necessary. This death may affect children in a variety of ways, depending on the ages of the students, how well they knew **(insert child's name)**, and the children's prior experiences with grief and death, (particularly a violent one such as this). Typical reactions children have:

- Appear to be unaffected
- Think about it privately
- Fright
- Sadness & withdrawal
- Agitation & anger
- Ask many questions
- Attempts to be extra good
- Wanting to be with peers, especially older students

This is an important time to listen to your children intently. If your children seem to need to talk, answer their questions simply, honestly, and possibly over and over again. I have enclosed a flyer of community mental health resources that are available should this event trigger any undue or exaggerated reactions to your child or someone else you know. Please advise us if you have any serious concerns about the effect this event is having on your child or any of your child's friends. This is a time we must be eyes and ears for each other as we seek to nurture and protect the children in our midst and teach them how to seek help when it is needed.

*(Optional)* A Parent Informational Night is planned for:

Date:

Time:

Place:

At that time we can talk further about helping children through grief.

Our thoughts are with the family and friends of **(insert child's name)**.

Sincerely,

**(Principal name inserted)**

**e. Sample parent/guardian letter about coping with child's reaction to disaster natural or other)**

Dear Parents/Guardians:

The **(insert name of disaster)** we have experienced is profoundly troubling to everyone. Below are some coping strategies we offer that we hope will help you respond positively to your child.

**FEAR AND ANXIETY:**

1. Normal reaction to any danger that threatens one's life or well-being
2. What triggers the fear after a disaster?
  - ✓ Fear of reoccurrence
  - ✓ Fear of separation from family
  - ✓ Fear of being left alone
3. First priority as parents is to understand the kinds of fears and anxieties children experience.

**ADVICE TO PARENTS:**

1. Important for the family to be together
2. Children need reassurance by parents' words and actions
3. LISTEN to
  - ✓ What your children tell you about their fears
  - ✓ How they feel and what they think about what has happened—validate them.
4. Explain the disaster and known facts to children & listen carefully to what they say and ask—encourage them to talk.
5. Children's fear does not need to completely disrupt their own and the family's activities.
6. Communicate and work cooperatively with the school crisis team at your children's school.

**SETTLING DOWN:**

1. Parents need to communicate they are maintaining control; they need to be understanding, but firm, supportive, and make decisions for children.
2. Bedtime problems
  - ✓ Children may refuse to go to bed alone
  - ✓ They may have difficulty sleeping when they do go to bed
  - ✓ They may wake up repeatedly in the night and have difficulty falling back to sleep.
3. It is natural for children to want to be close to their parents and for parents to want their children near them.
4. Parents should be aware of their own fears, their own uncertainty, and of the effect these have upon their children.
5. Children may demonstrate regressive behavior such as:
  - ✓ Bedwetting
  - ✓ Clinging to parents
  - ✓ Thumb sucking
6. Children respond to praise; parents should make a deliberate effort not to focus on the child's regressed or immature behavior.
7. Specific fears



- ✓ Refusal to go to school
- ✓ Fear of the dark
- ✓ Fear of going to bed
- ✓ Fear of “monsters”

**WHEN IT IS TIME FOR PARENTS TO SEEK PROFESSIONAL HELP:**

1. If sleeping problem continues for more than a few weeks, if the clinging behavior does not diminish, or if the fears become worse, it is time to ask for professional advice.
2. Mental health professionals are specially trained to help people in distress. They can help parents cope with and understand the unusual reactions of the child. By talking to the parents and child, either individually or in a group, a therapist can help a child overcome fears more easily.

By working with the school crisis team at their child’s school, parents can gain access to resources and obtain recommendations.

We hope these suggestions are helpful to you. Please contact me or our school support staff (school counselor, nurse, or psychologist) if we can we of further help.

Sincerely,

**(Principal name inserted)**

**f. Sample parent/guardian invitation to informational evening guidelines**

**PARENT INFORMATIONAL EVENING GUIDELINES**

- INVITE:** Send a note home to inform parents of the Parent Informational Night. Send an e-mail if your school has a list of parents, also.
- FACILITATE:** Have enough support person for every 15 parents estimated to attend.
- INTRODUCE:** Introduce the evening as a time for parents to express their concerns and ask questions of the school and school crisis team in regards to the (death or disaster) that has impacted their children and families.
- SHARE:** Divide into circles of no more than 15 participants with a facilitator for each group.
- ✓ Go around circle, one at a time, starting with facilitator
  - ✓ Ask that each person share his/her concerns, feelings, questions and experiences in regards to the (death or disaster)
  - ✓ Do not encourage discussion at this time. Facilitator takes notes to be addressed at end of sharing circle.
- BREAK:** Take a break after sharing circle.
- REJOIN:** Rejoin as one group for feedback session.
- FEEDBACK:** A designated facilitator addresses the questions and concerns raised by parents. This may include material about:
- ✓ Family dynamics during grieving process
  - ✓ Adult grief
  - ✓ Neighborhood and school safety
  - ✓ Suicide and violence prevention
  - ✓ Information on how school is responding
  - ✓ Other topics
- ENCOURAGE:** The facilitator closes the evening with a statement of encouragement and faith in the family's ability to respond to the individual needs of the family members and with appreciation for the great difficulties facing the families.

**Staff Letters/Announcements/Documentation/ Forms**

**a. Sample staff letter advising of meeting to introduce Crisis and Suicide Prevention Handbook and the District Crisis Guidance Manager.**

Dear Staff Member:

Members of our school district have worked very hard on the creation of a **(insert school district name)** Crisis and Suicide Prevention Handbook. It is important that each staff member know the contents of this handbook and where it is located in the building. The **(insert name of Crisis Guidance Manager)**, will come to our school to present and explain the handbook and answer questions you might have. Please attend this important meeting.

Date:

Time:

Place:

As our school forms its own school site crisis team, this handbook will serve as an important tool. I look forward to seeing you there.

Sincerely,

**(Principal name inserted)**

**b. Sample staff letter seeking School Site Crisis Team volunteers.**

Dear Staff:

With the introduction of the DJUSD Crisis and Suicide Prevention Handbook to help facilitate a district-wide crisis plan this year, we are going to formalize a school site crisis team. I have attached the form, *Establishing School Crisis Team: School Site Responsibilities*, (from the Handbook, Section 6, Attachment 6). Please read it over and consider if you would be interested in applying.

We are looking for school crisis team members: comfortable dealing with crisis situations; knowledgeable about community and district-wide resources; able to work well with others on a team; and would be willing to spend the time to meet regularly for training and coordination. If you are interested, please submit the attached application to me before **(insert date)**.

See me if you have any questions or would like more information.

Sincerely,

**(Principal name inserted)**

**b. Sample staff application for School Site Crisis Team volunteers. Application for School Site Crisis Team**

**Name:** \_\_\_\_\_

**Assignment:** \_\_\_\_\_

**I am interested in being assigned to the School Site Crisis Team. I have experience in:**

- |   |  |
|---|--|
| <input type="checkbox"/> Critical incident stress debriefing                | <input type="checkbox"/> Violence/bully prevention   |
| <input type="checkbox"/> Team building                                      | <input type="checkbox"/> Drug abuse prevention       |
| <input type="checkbox"/> Active Listening                                   | <input type="checkbox"/> Effective media relations   |
| <input type="checkbox"/> Small group techniques for children and adults     | <input type="checkbox"/> Grief and loss intervention |
| <input type="checkbox"/> First aid and how to call for medical assistance   | <input type="checkbox"/> Suicide intervention skills |
| <input type="checkbox"/> Crowd control and evacuation of students and staff | <input type="checkbox"/> Anger management            |

Please comment below on items checked. If you need more space use the back or attach another paper):

Thanks for applying. Please turn into **(insert principal's name)** by **(insert date)**.

**d. Sample staff letter seeking District Crisis Support Team volunteers.**

Dear Staff:

With the introduction of the DJUSD Crisis and Suicide Prevention Handbook to help facilitate a district-wide crisis plan this year, we are going to formalize a school site crisis team. I have attached the form, *District Crisis Support Team: Membership & Roles*, (from the Handbook Section 6, Attachment 8). Please read it over and consider if you would be interested in applying.

We are looking for District Crisis Support Team members: comfortable dealing with crisis situations; knowledgeable about community and district-wide resources; able to work well with others on a team; and would be willing to spend the time to meet regularly for training and coordination. If you are interested, please submit the attached application to me before **(insert date)**. You must have your supervisor's approval to apply as there is a possibility your presence on the team could take you from your building periodically.

See me if you have any questions or would like more information.

Sincerely,

**(Crisis Guidance Manager name inserted)**

**e. Sample staff application for District Crisis Support Team volunteers Application for District Crisis Support Team**

Name: \_\_\_\_\_ School Site: \_\_\_\_\_

Assignment \_\_\_\_\_

Supervisor Signature approving application: \_\_\_\_\_

**I am interested in applying to become a member of the District Crisis Support Team. I have experience in:**

- |   |  |
|---|--|
| <input type="checkbox"/> Critical incident stress debriefing                | <input type="checkbox"/> Violence/bully prevention   |
| <input type="checkbox"/> Team building                                      | <input type="checkbox"/> Drug abuse prevention       |
| <input type="checkbox"/> Active Listening                                   | <input type="checkbox"/> Effective media relations   |
| <input type="checkbox"/> Small group techniques for children and adults     | <input type="checkbox"/> Grief and loss intervention |
| <input type="checkbox"/> First aid and how to call for medical assistance   | <input type="checkbox"/> Suicide intervention skills |
| <input type="checkbox"/> Crowd control and evacuation of students and staff | <input type="checkbox"/> Anger management            |

Please comment below on items checked. If you need more space use the back or attach another paper):

Thanks for applying. Please turn into (**insert name of Crisis Guidance Manager**) at (**insert location**) by (**insert date**).

**f. Sample staff letter advising of schedule of mandatory gatekeeper meetings.**

Dear Staff,

There have been many requests for staff in-service on suicide prevention and a delineation of the roles and responsibilities we have to students and colleagues when we work for the school district. In a sense we are all the eyes and ears of the school community. Adults who see members of the school community on a routine basis are “gatekeepers”, people who help others be safe. Beginning this year we will be providing gatekeeper training so our employees know what to do when they suspect someone in the school community is contemplating suicide or is in some other kind of profound danger. We all need to know the protocol of who to inform, what resources are available, and how to best help. These are all issues of compassion as well as liability. The gatekeeper training for our school will be:

Date:

Time:

Place:

I look forward to joining each of you at this important meeting.

Sincerely,

**(Principal’s name inserted)**



**g. Sample staff script to inform students of death of another student or employee, (after they have been told personally by Crisis Team Leader/Designee).**

Dear Staff;

It is a hard task to tell your students about the death of (insert name of deceased).

**FIRST AND FOREMOST:** If you do not feel you want to be the one to tell your students, then do not! The school crisis team will provide someone who can lead the discussion for you, or take over your class while you seek the support you need. Please take advantage of this resource. We care about you and your needs in this sad time and want you to feel our support.

If you would like to lead the discussion yourself, here are the facts and some suggestions for procedures; please make appropriate developmental adjustments according to the age of students in class:

“I have something very sad I want to share with you. **Here you will read the factual information (agreed upon by the school crisis team), e.g.:**

*This morning a student who attends our school in (insert grade number), (insert name), was hit by a car outside of his/her home in \_\_\_\_\_. According to his/her family, he/she ran out into the street and was seriously injured. He/she died at the hospital.*

**(Then in your own words continue on with the following or something similar):**

*I am feeling very sad about what has happened. I would like to spend some time together now to share with each other. Maybe we could help each in expressing how we feel about (insert name). Maybe we could help each other in expressing how we feel about (insert name) and how he/she died.”*

- ✓ Take some time for discussion
- ✓ See the two attached handouts: “Guidelines for Teachers on How to Tell Students about a Death” and “Guidelines for Teachers on How to Lead a Discussion with Grieving Students”
- ✓ After discussion, tell students there are counselors in the building if they need to talk further and arrange a procedure for them going to see the counselor.
- ✓ After your discussion you may want to: take a break or recess; playground play; do some drawing, art project or other projects-subject matter student choice; journal writing-any thoughts about whatever is on the child’s mind.
- ✓ These activities may be useful to continue to do at intervals during the day and to intersperse throughout curriculum in coming days.

**If you need support, please call the office. Do not hesitate to ask.**

**THANK YOU FOR BEING THE WONDERFUL STAFF THAT YOU ARE.....**

**h. Sample staff announcement of death in a staff member's family.**

Dear Colleagues:

Many of you are dealing with a number of very difficult emotions around the death of **(insert name of staff member's family member)**.

**(Give accurate information.) e.g.: (Insert staff member's name)** was informed that **(his/her) (insert family member)** died of **(insert condition)** while **(insert location)**.

This loss has left **(insert staff member's name)** and **(his/her)** family with a void that can only be understood by those of you who have experienced such a loss. Your love and support for **(insert staff member's name)** will be an important part in **(his/her)** ability to cope with this loss and at the same time continue on with **(his/her)** personal and professional life.

I have talked with **(staff member)** and shared the fact that so many of you are concerned and that you wished to do something for **(her/him)** to show your feelings and send comfort. **(He/She)** was touched by this and appreciated each of you and your caring ways.

In discussing ways people could demonstrate their caring, **(staff member)** and **(his/her)** family has decided that **(he/she)** would like people to make contributions to **(insert name of charity or cause)** in **(insert deceased's name)**. **(He/She)** had a special connection to that work.

Funeral services for **(deceased)** will be held at **(time)** on **(day of the week)** at **(location)**. Everyone is invited to **(place)** following the service. The address is **(insert address)**.

Your caring ways and support has been, and will continue to be, important to **(staff member)**. **(School name)** is fortunate to be made up of so many wonderful human beings who take such good care of one another.

Sincerely,

**(Principal's name inserted)**

## I. Sample form for documentation of suicidal behavior or threat.

### SUICIDAL ASSESSMENT CHECKLIST\*\*

Name of person at risk: \_\_\_\_\_ Date: \_\_\_\_\_

Interviewer: \_\_\_\_\_

(Suggested points to cover with person at risk & later with family member of at-risk person)

#### A. PAST ATTEMPTS, CURRENT PLANS, & VIEW OF DEATH

- |  |   |   |
|--|---|---|
| 1. Does the individual have frequent suicidal thoughts?  | Y | N |
| 2. Have the individual or a significant other in his or her life ever attempted suicide?                                   | Y | N |
| 3. Does the individual have a detailed, feasible suicide plan?   | Y | N |
| 4. Has the individual made special arrangements such as giving away prized possessions?                                    | Y | N |
| 5. Does the individual fantasize about suicide as a way to make others feel guilty or as a way to get a happier afterlife? | Y | N |

#### B. REACTIONS TO PRECIPITATING EVENTS

- |  |   |   |
|--|---|---|
| 1. Is the individual experiencing severe psychological distress?                               | Y | N |
| 2. Have there been major changes in recent behavior along with negative feelings and thoughts? | Y | N |

*(Such changes are often related to recent loss or threat of loss of significant others or of positive status and opportunity. They may also stem from sexual, physical, or substance abuse. Negative feelings & thoughts are often expressions of a sense of extreme loss, abandonment, failure, sadness, hopelessness, guilt, and sometimes anger directed inward).*

#### C. PSYCHOSOCIAL SUPPORT

- |   |   |   |
|---|---|---|
| 1. Is there a lack of significant other to help individual survive? | Y | N |
| 2. Does the individual feel alienated?                              | Y | N |

#### D. HISTORY OF RISK-TAKING BEHAVIOR

- |   |   |   |
|---|---|---|
| 1. Does the individual take life-threatening risks or display poor impulse control? | Y | N |
|---|---|---|

*\*\*Use this checklist as an exploratory guide with individuals you are concerned about. Each yes raises the level of risk, but there is no single score indicating high risk. A history of suicide attempts is a sufficient reason for action. High risk is also associated with very detailed plans (when, where, how) that specify a lethal and readily available method, a specific time, and location where it is unlikely the act would be disrupted. Further high risk indicators include individual making final arrangements and information about a recent, critical loss. Because of informal and confidential nature of this type of assessment, it should NOT be filed as part of a student's regular school records. (Adapted and used with permission of UCLA: Center for Mental Health in Schools, May 2004)*

## **j. Sample Guidelines after Assessing Suicidal Risk - Checklist**

### **FOLLOW-THROUGH STEPS AFTER ASSESSING SUICIDAL RISK CHECKLIST**

- \_\_\_ (1) In assessment process, efforts will have been made to discuss problem openly and non-judgmentally with individual. Remember how seriously devalued individual feels. Avoid saying anything demeaning; convey warmth, empathy, and respect. If individual resists talking about the matter, give another effort because the more the individual shares, better change of engaging individual in problem-solving.
- \_\_\_ (2) Explain to individual importance of your responsibility to break confidentiality in matters of suicidal risk. Ask if individual would prefer to take the lead or at least be present during process of informing parents and/or other concerned parties.
- \_\_\_ (3) Try to contact parents by phone to:
  - a. Inform about concern
  - b. Gather more info to assess risk
  - c. Provide info about problem and available resources
  - d. Offer help in connecting with appropriate resources++
- ++*Note: In case of a minor, if family is uncooperative, it may be necessary to report child endangerment after taking the following steps*
- \_\_\_ (5) If an individual is considered to be in danger, only release him/her to parent (in case of child) or adult family member (in case of adult) who is equipped to provide help. In high risk cases, if parent or family are unavailable or uncooperative, and no one else is available to help, you must contact the A local public agency, (children's services, local law enforcement, services for emergency hospitalization). These agencies will need and want the following info:
  - a. Individual's name, address, birth date, Social Security number
  - b. Data about individual's suicide risk (see Suicide Risk Checklist)
  - c. Stage of parent (child) or family (adult) notification
  - d. Language spoken by individual and family
  - e. Health coverage plan if there is one
  - f. Location of at-risk individual
- \_\_\_ (6) Follow-up with individual and adult family member to determine what steps have been taken to minimize risk.
- \_\_\_ (7) Document all steps taken and outcomes. Plan for aftermath intervention & support.
- \_\_\_ (8) Report child endangerment if necessary.

**k. Sample documentation & referral form for parent/guardians.**

**Advise of At-Risk Individual and Referral Form**

I, **(insert name of parent/guardian for child or adult family member for adult)**, am **(state relationship to at-risk person)** of **(insert name of at-risk individual)**.

I have been advised by **(insert name of school employee)** of the **(insert name of school district)** that **(name of at-risk person)** is considered to be at-risk for suicide based on the following: **(School mental health person should briefly write in space below what was said, done, or written to cause this concern)**.

I acknowledge that I have been advised to seek mental health care for my child via my family doctor, the local mental health agency, or a private mental health therapist. **(In case of child)** I have been advised and understand I have the right to ask for a special education assessment if I deem that necessary. I have been given the name and phone number of **(insert name(s) of District Crisis Guidance Manager, school psychologist, nurse, or counselor)** should I have further questions of concerns.

Signed: \_\_\_\_\_

Relationship to At-Risk Individual: \_\_\_\_\_

Date: \_\_\_\_\_

## Section VII. Bibliography

### References

- Adams, D. M., Overholser, J. C., & Lehnert, K. L. (1994). Perceived family functioning and adolescent suicidal behavior. *Journal of the American Academy of Child and Adolescent Psychiatry, 33*(4), 498-507.
- Adelman, H. S., & Taylor, L. (1991). Early school adjustment problems: Some perspectives and a project report. *American Journal of Orthopsychiatry, 61*, 468-474.
- Adelman, H. S., & Taylor, L. (2000). Moving prevention from the fringes into the fabric of school improvement. *Journal of Educational and Psychological Consultation, 11*, 7-36.
- Agerbo, F., Nordentoft, M., Mortensen, P. B. (2002). Familial, psychiatric, and socioeconomic risk factors for suicide in young people: Nested case-control study. *British Medical Journal, 325*, 74.
- Allberg, W. & Chu, L. (1990). Understanding adolescent suicide: Correlates in a developmental perspective. *The School Counselor, 37*, 343-349.
- Allen, B.P. (1987). Youth suicide. *Adolescence, 22*(86), 271-290.
- Allen, J. P., Aber, J. L., & Leadbeater, B. J. (1990). Adolescent problem behaviors: The influence of attachment and autonomy. *Psychiatric Clinics of North America, 13*, 455-467.
- American Association of Suicidology, Prevention Division (1999). *Guidelines for school-based suicide prevention programs*. Washington, DC: Author.

- American Psychiatric Association (APA). (1980). *Diagnostic and statistical manual of mental disorders* (3rd ed.). Washington, DC: Author.
- American Psychiatric Association (APA). (1987). *Diagnostic and statistical manual of mental disorders* (3rd ed.-rev). Washington, DC: Author.
- Anderson, B. S. (1996). *The counselor and the law* (4th ed.). Alexandria, VA: American Counseling Association.
- Anderson, R. N. (2002). Deaths: Leading causes for 2000. *National Vital Statistics Reports*, 50(16). Hyattsville, MD: National Center for Health Statistics.
- Annie E. Casey Foundation. (2003) *KIDS COUNT Data Book: 2003*, Annie E. Casey
- Arias, E. & Smith, B. (2003). *Deaths: Preliminary data for 2001*, 51(5).  
National Center for Health Statistics. Hyattsville, MD: Table 6. URL available:  
[http://www.cdc.gov/nchs/data/nvsr/nvst51/nvsr51\\_05.pdf](http://www.cdc.gov/nchs/data/nvsr/nvst51/nvsr51_05.pdf).
- Arango, V., Underwood, M. D., Boldrini, M., et al. (2001). Serotonin 1A receptors, serotonin transporter binding and serotonin transporter mRNA expression in the brainstem of depressed suicide victims. *Neuropsychopharmacology*, 25, 892-903.
- Arias, E., Anderson, R. N., Kung, H. S., & Murphy, S. L. (2003). *Deaths: Final data for 2001*. National Vital Statistics Report. Hyattsville, MD: U.S. Department of Health and Human Services.
- Arias, E., & Smith, B. L. (2003). *Deaths: Preliminary data for 2001*. National Vital Statistics Report 51(5). Hyattsville, MD: National Center for Health Statistics.
- Armacost, R. L. (1990). High school student stress and the role of the counselors. *School Counselor*, 38, 197-203.

- Asarnow, J., Carlson, G., & Guthrie, D. (1987). Coping strategies, self-perceptions, hopelessness and perceived family environments in depressed and suicidal children. *Journal of Consulting and Clinical Psychology, 55*, 361-366.
- Aseltine, R. H. (2002). *An evaluation of the SOS suicide prevention program: Final report of findings from the 2001-2002 school year*. Farmington, CT: University of Connecticut, Health Center.
- Aseltine, R. H. (2003). An evaluation of a school-based suicide prevention program. *Adolescent Family Health, 2*, 81-88.
- Aseltine, R. H., & DeMartino, R. (2004). An outcome evaluation of the SOS suicide prevention. *American Journal of Public Health, 94*(3), 446-452).
- Aseltine, R. H., Gore, S., & Colten, M. (1994). Depression and the social developmental context of adolescence. *Journal of Personality & Social Psychology, 67*, 252-263.
- Aseltine, R. H., Gore, S., & Colten, M. E. (1998). The co-occurrence of depression and substance abuse in late adolescence. *Development and Psychopathology, 10*, 549-570.
- Ashworth, S., Spirito, A., Colella, A., & Benedict-Drew, C. (1986). A pilot suicidal awareness, identification, and prevention program. *Rhode Island Medical Journal, 69*, 457-461.
- Ayyash-Abdo, H. (2001). Individualism and collectivism: The case of Lebanon. *Social Behavior and Personality, 29*, 503-518.
- Ayyash-Abdo, H. (2002). Adolescent suicide: An ecological approach. *Psychology in the Schools, 39*(4), 459-470.



- Bailey, J.E., Kellerman, A. L., Somes, G. W., Banton, J. G., Rivara, F. P. & Rushforth, N. P. (1997). Risk factors for violent death of women in the home. *Archives of Internal Medicine*, 157(7), 777-782.
- Ballard v. Polly, 387 F. Supp. 895 (1975).
- Beautrais, A. L. (2001). Child and young adolescent suicide in New Zealand. *Australian and New Zealand Journal of Psychiatry*, 125, 355-373.
- Beautrais, A. L., Coggan, C. A., Fergusson, D. M., & Rivers, L. (1997). *The prevention, recognition and management of young people at risk of suicide: Development of guidelines for schools*. Wellington, New Zealand: New Zealand Ministry of Education.
- Beautrais, A. L., Joyce, P. R., & Mulder, R. T. (1996). Risk factors for serious suicide attempts among youths aged 13 through 24 years. *Journal of the American Academy of Child and Adolescent Psychiatry*, 35, 1174-1182.
- Beautrais, A. L., Joyce, P. R. & Mulder, R. T. (1998). Psychiatric contacts among youths aged 13 through 24 years who have made serious suicide attempts. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(4), 504-511.
- Beautrais, A. L., Joyce, P. R., & Mulder, R. T. (1999). Personality traits and cognitive styles as risk factors for serious suicide attempts among young people. *Suicide and Life-Threatening Behavior*, 29, 37-47.
- Beck, A. T., Steer, R., Kovacs, M. & Garrison, B. (1985). Hopelessness and eventual suicide: A 10-year prospective study of patients hospitalized with suicidal ideation. *American Journal of Psychiatry*, 142, 559-563.

- Bell, C. C., & Clark, D. C. (1998). Adolescent suicide. *Paediatric Clinics of North America*, 45, 365-380.
- Belsky, J. (1980). Child maltreatment: An ecological integration. *American Psychologist*, 35, 320-335.
- Berliner, D.C., & Biddle, B. J. (1995). *The manufactured crisis: Myths, fraud, and attack on America's public school*. Reading, MA: Addison-Wesley Publishing Company.
- Berman, A. L. (1996). Dyadic death: A typology. *Suicide and Life-Threatening Behavior*, 26, 342-350.
- Berman, A. L., & Jobes, D. A. (1991). *Adolescent suicide: Assessment and intervention*. Washington, DC: American Psychological Association.
- Berman, A. L. & Jobes, D. A. (1995). Suicide prevention in adolescents (age 12-18). *Suicide and Life-Threatening Behavior*, 23, 143-154.
- Berman, A. L., & Schwartz, R. H. (1990). Suicide attempts among adolescent drug users. *American Journal of Diseases of Children*, 144, 310-314.
- Berman, P., & McLaughlin, M. W. (1978). *Federal programs supporting educational change*. Santa Monica, CA: Rand Corporation.
- Birckmayer, J., & Hemenway, D. (2001). Suicide and firearm prevalence: Are youth disproportionately affected? *Suicide and Life-Threatening Behavior*, 31, 303-310.

- Blake, S. M., Ledsky, R., Lehman, T., Goodenow, C., Sawyer, R. & Hack, T. (2001). Preventing sexual risk behaviors among gay, lesbian, and bisexual adolescents: The benefits of gay-sensitive HIV instruction in schools. *American Journal of Public Health, 91*, 940-946.
- Bogust v. Iverson, 102 N.W.2d 228 (1960).
- Boivin, M., Poulin, F. & Vitaro, F. (1994). Depressed mood and peer rejection in childhood. *Development and Psychopathology, 6*, 483-498.
- Bongar, B., Maris, R. W., Berman, A. L., & Litman, R. E. (1992). Outpatient standards of care and the suicidal patient. *Suicide and Life-Threatening Behavior, 22*, 453-478.
- Bongar, B., Maris, R. W., Berman, A. L. & Litman, R. E. (1992). Outpatient standards of care and the suicidal patient. *Suicide & Life-Threatening Behavior, 22*, 453-477.
- Bonnie, R. J., Fulco, C. E., & Liverman, C. T. (Eds.). (1999). *Reducing the burden of injury: Advancing prevention and treatment*. Washington, DC: Institute of Medicine, National Academy Press.
- Borowsky, I. W., Ireland, M., & Resnick, M. D. (2001). Adolescent suicide attempts: Risk and protectors, *Pediatrics, 107*(3), 485-493.
- Borowsky, I. W., Resnick, M.D., Ireland, M., & Blum, R.W. (1999). Suicide attempts among American Indian and Alaska Native youth: Risk and protective factors. *Archives of Pediatrics and Adolescent Medicine, 153*(6), 573-580.

- Botvin, G. J., Baker, E., Dusenbury, L., Tortu, S., & Botvin, E. M. (1990). Preventing adolescent drug abuse through a multimodal cognitive-behavioral approach: Results of a 3-year study. *Journal of Consulting and Clinical Psychology, 58*, 437-446.
- Brand, E. F., King, C. A., Olson, E., Ghazuddin, N., & Naylor, M. (1996). Depressed adolescents with a history of sexual abuse: Diagnostic comorbidity and suicidality. *Journal of the American Academy of Child and Adolescent Psychiatry, 35*(1), 34-41.
- Brener, N. D., Krug, E. G., & Simon, T. R. (2000). Trends in suicidal ideation and behavior among high school students in the United States, 1991-1997. *Suicide and Life-Threatening Behavior, 30*, 304-312.
- Brent, D. A. (1995). Risk factors for adolescent suicide and suicidal behavior: Mental and substance abuse disorders, family environmental factors and life stress. *Suicide and Life-Threatening Behavior, 25*(Suppl.), 52-63.
- Brent, D. A., Baugher, M., Bridge, J., Chen, T., & Chiappetta, L. (1999). Age- and sex-related risk factors for adolescent suicide. *Journal of the American Academy of Child and Adolescent Psychiatry, 38*, 1497-1505.
- Brent, D. A., Johnson, B., Perper, J., Connolly, J., Bridge, J., Bartle, S., et al. (1994). Personality disorder, personality traits, impulsive violence, and completed suicide in adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry, 33*, 1080-1086.
- Brent, D. A., Kalas, R., Edelbrock, C., Costello, A. J., Dulcan, M. K., & Conover, N. (1986). Psychopathology and its relationship to suicidal ideation in childhood and

- adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry*, 25, 666-673.
- Brent, D. A., Kolko, D. J., Birhamer, B., et al. (1998). Predictors of treatment efficacy in a clinical trial of three psychosocial treatments for adolescent depression. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37, 906-914.
- Brent, D. A., Perper, J. A., Goldstein, C. E., Kolko, D. J., Allan, M. J., Allman, C. J., et al. (1998). Risk factors for youth suicide: A comparison of youth suicide victims with suicidal inpatients. *Archives of General Psychiatry*, 45, 581-588.
- Brent, D. A., Perper, J. A., Moritz, G., Allman, C., Friend, A., Schweers, J., et al. (1992). Psychiatric effects of exposure to suicide among the friends and acquaintances of adolescent suicide victims. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 629-639.
- Brent, D. A., Perper, J. A., Moritz, G., Allman, C., Friend, A., Roth, C., et al. (1993). Psychiatric risk factors for adolescent suicide: A case control study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 32, 521-529.
- Brent, D. A., Perper, J. A., Moritz, G., Baugher, M., & Allman, C. (1993). Suicide in adolescents with no apparent psychopathology. *Journal of the American Academy of Child and Adolescent Psychiatry*, 32, 494-500.
- Brent, D. A., Perper, J. A., Moritz, G. M., Baugher, M., Schweers, J., & Ross, C. (1993). Firearms and adolescent suicide: A community case control study. *American Journal of Diseases of Children*, 147, 1066-1071.
- Brent, D. A., Perper, J. A., Moritz, G. M., Liotus, L., Schweers, J. & Canobbio, R. (1994). Major depression or uncomplicated bereavement? A follow-up study of

- youth exposed to suicide. *Journal of the Academy of Child and Adolescent Psychiatry*, 33, 231-239.
- Breton, J.-J., Boyer, R., Bilodeau, H., Raymond, S., et al. (2002). Is evaluative research on youth suicide programs theory-driven? The Canadian experience. *Suicide and Life-Threatening Behavior*, 32(2), 176-190.
- Brickmayer, J., & Hemenway, D. (1999). Minimum-age drinking laws and youth suicide, 1970-1990. *American Journal of Public Health*, 89, 1365-1368.
- Bridge, J. A., Brent, D. A., Johnson, B. A., & Connolly, J. (1997). Familial aggregation of psychiatric disorders in a community sample of adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36(5), 628-637.
- Brock, S. (2002a). Crisis theory: A foundation for the comprehensive crisis prevention and intervention team. In S. Brock, P. Lazarus, & S. Jimerson (Eds.), *Best practices in school prevention and intervention* (pp.5-18). Bethesda, MD: National Association of School Psychologists.
- Brock, S. E. (2002b). School suicide postvention. In S. E. Brock, P. J. Lazarus, & S. R. Jimerson (Eds.), *Best practices in school crisis prevention and intervention* (pp. 211-224). Bethesda, MD: National Association of School Psychologists.
- Brock, S., Lazarus, P., & Jimerson, S. (1994). *Health, mental health and safety guidelines for schools: References best practices in school crisis prevention and intervention*. New York: Cambridge University Press.
- Brock, S. E., & Sandoval, J. (1997). Suicidal ideation and behaviors. In G. C. Bear, K. M. Minke, & A. Thomas (Eds.), *Children's needs II: Development, problems, and*

- alternatives* (pp. 361-374). Bethesda, MD: National Association of School Psychologists.
- Bronfenbrenner, U. (1979). *The ecology of human development*. Cambridge, MA: Harvard University Press.
- Brooks v. Logan, 903 p2d. 73 (1995).
- Brown, D. R. & Blanton, C. J. (2002). Physical activity, sports participation, and suicidal behavior among college students. *American College of Sports Medicine, 34*, 1087-1096.
- Burge, V., Felts, M., Chenier, T. & Parillo, A. V. (1995). Drug use, sexual activity, and suicidal behavior in high school students. *Journal of School Health, 65*, 222-227.
- Burns, B. J., Costello, E. J., Angold, A., Tweed, D., Stangl, D. Farmer, E. M., & Errantly, A. (1995). Children's mental health service use across sectors. *Health Affairs, 14*, 147-159.
- Burns, J. M., & Patton, George C. (2000). Preventive interventions for youth suicide: a risk factor-based approach. *Australian and New Zealand Journal of Psychiatry, 34*, 388-407.
- California Department of Education (CDE). (1987). *Suicide prevention program for California public schools*. ISBN # 0-8011-0682-6. Sacramento, CA: Author.
- Callahan, J. (1996). Negative effects of a school suicide postvention program: A case example. *Crisis, 17*, 108-115.
- Callahan, J. (2000). Predictors and correlates of bereavement in suicide support group participants. *Suicide and Life-Threatening Behavior, 30*(2), 104-124.

- Canter, A. S., & Carroll, S. A. (Eds.). (1999). *Crisis prevention and response: A collection of NASP resources*. Bethesda, MD: National Association of School Psychologists.
- Cantor, C.H. & Slater, P. J. (1995). The impact of firearm control legislation on suicide in Queensland: Preliminary findings. *Medical Journal of Australia*, 162,583-585.
- Capuzzi, D. (1994). *Suicide prevention in the schools: Guidelines for middle and high school settings*. Alexandria, VA: American Counseling Association.
- Capuzzi, D. (1998). Addressing the needs of at-risk youth: Early prevention and systemic intervention. In C. C. Lee & G. R. Walz (Eds.), *Social action: A mandate for counselors*. Alexandria, VA: American Counseling Association.
- Capuzzi, D. (2004). From Seneca to suicidology: A history of suicide. In D. Capuzzi (Ed.), *Suicide across the life span: Implications for counseling*. Alexandria, VA: American Counseling Association.
- Capuzzi, D., & Golden, L. (1988). *Preventing adolescent suicide*. Muncie, IN: Accelerated Development, Inc.
- Capuzzi, D., & Gross, D. (2000). *Youth at risk: A prevention resource for counselors, teachers, and parents*. (3<sup>rd</sup> ed.). Alexandria, VA: American Counseling Association.
- Capuzzi, D., & Gross, D. (2004). *Youth at risk: A prevention resource for counselors, teachers, and parents*. Alexandria, VA: American Counseling Association.
- Carnegie Corporation. (1989). Turning points: Preparing American youth for the 21st century. *The report of the task force on education of young adolescents*. New York: Author.



- Carrington, P. J., & Moyer, S. (1994). Gun control and suicide in Ontario. *Journal of Psychiatry, 151*, 606-608.
- Caruano, R. (2004). *Columbia University TeenScreen program*. Paper presented at the Spring Meeting of the California Suicide Prevention Action Network meeting, Sacramento, CA, May 2005.
- Celotta, B. (1995). The aftermath of suicide: Postvention in a school setting. *Journal of Mental Health Counseling, 17*(4), 397-412.
- Cellota, B., Jacobs, G., Keys, S. G., & Cannon, G. (1989). A model prevention program. In D. Capuzzi & L. Golden (Eds.), *Preventing adolescent suicide*. Muncie, IN: Accelerated Development.
- Center for Mental Health in Schools at UCLA. (2003). A technical assistance sampler on school interventions to prevent youth suicide. Los Angeles, CA: Author.
- Center for Mental Health in Schools at UCLA. (2004). Integrating agendas for mental health in schools into the recommendations of the President's New Freedom Commission on Mental Health. *Addressing Barriers to Learning, 9*(1), 1-12.
- Center for Disease Control and Prevention (1998). Youth risk **behavior** surveillance- United States, 1997. *Morbidity and Mortality Weekly Report, 47*(SS-3), 1-89.
- Centers for Disease Control & Prevention (CDC). (1992). *Youth suicide prevention programs: A resource guide*. Atlanta, GA: Author.
- Centers for Disease Control & Prevention (CDC). (1994). Programs for the prevention of suicide among adolescents and young adults: Suicide contagion and the reporting

of suicide: Recommendations from a national workshop. *Morbidity and Mortality Weekly Report*, 43(RR-6), 1-18.

Centers for Disease Control & Prevention. (1998). *Youth risk behavior surveillance-United States, 1997*. *Morbidity and Mortality Weekly Report*, 43(SS-3), 1-89.

Centers for Disease Control & Prevention (CDC). (1999a). Wonder, Mortality (compressed data set). National Center for Health Statistics. [Online database]. Retrieved October 10, 2003 from: <http://wonder.cdc.gov>

Centers for Disease Control and Prevention (CDC). (1999b). Framework for program evaluation in public health. *MMWR*, 48 (No. RR-11): [1-40].

Centers for Disease Control & Prevention (CDC). (2001). *Healthy people 2000 final review*. National Center for Health Statistics. Hyattsville, MD: Public Health Service.

Centers for Disease Control & Prevention (CDC). (2002). CDC surveillance summaries: Youth risk behavior surveillance-United States, 2001. *MMWR* 51(SS-4), 1-64. Atlanta, GA: U.S. Department of Health and Human Services.

Centers for Disease Control & Prevention (CDC). (2003). *Web-based injury statistics query and reporting system (WISQARS)*. [Online]. National Center for Injury Prevention & Control. Retrieved July 10, 2004 from:  
<http://www.cdc.gov/ncipc/wisqars/>  
<http://www.cdc.gov/nchs/datawh/statab/unpubd/mortabs/gmwk292a.htm>

Child Trends. (2002). *Child Trends databank indicator: Suicidal teens. Tables 1 and 22, Homicide, suicide, and firearm death among youth ages 15-19*. Original data from the National Vital Statistics System administered by the National Center for

- Health Statistics, Centers for Disease Control & Prevention. Retrieved February 14, 2003, from [http://www.childtrendsdatbank.org/tables/70\\_Table\\_1.htm](http://www.childtrendsdatbank.org/tables/70_Table_1.htm)
- Children's Safety Network. (2000). *Fact sheets: Youth suicide prevention plans*. Newton, MA: Education Development Center.
- Ciffone, F. (1993). Suicide prevention: A classroom presentation to adolescents. *Social Work, 98*, 196-203.
- Collaborative Management Services, Inc. (2004). Benefits of collaborative group-decision technologies. Retrieved August 28, 2004, from <http://cmsi-web/cmsi/Benefits.htm>.
- Cole, D. A. (1989). Psychopathology of adolescent suicide: Hopelessness, coping beliefs, and depression. *Journal of Abnormal Psychology, 98*, 248-255.
- Collins, J., Small, M. L., Kann, L., Pateman, B. C., Gold, R. S. & Kolbe, L. J. (1995). School health education. *Journal of School Health 65*,(8): 302-311.
- Committee on Prevention of Mental Disorders. (1994). *Reducing the risks for mental disorders: Frontiers for preventive intervention research* (1st ed.) Washington, DC: National Academy Press.
- Cosand, B. J., Bourque, L. B., & Kraus, J. F. (1982). Suicide among adolescents in Sacramento County, California 1950-1979. *Adolescence, 17*(68), 917-930.
- Crumley, F. E. (1990). Substance abuse and adolescent suicidal behavior. *Journal of the American Medical Association, 280*, 471-472.
- Cummings, R., Koepsell, T. D., Grossman, D. C., Savarino, J. & Thompson, R. S. (1997). The association between the purchase of a handgun and homicide or suicide. *American Journal of Public Health, 87*, 974-978.

- Curran, D. F. (1987). *Adolescent suicidal behavior*. Washington, DC: Hemisphere Publishing.
- Davidson, L. E., Rosenberg, M. L., Mercy, J. A., et al. (1989). An epidemiologic study of risk factors in two teenage suicide clusters. *Journal of the American Medical Association, 262*, 2687-2692.
- Davidson, M., & Range, L. M. (1997). Practice teachers' responses to a suicidal student. *Journal of Social Psychology, 137*, 530-532.
- Davidson, M. W., & Range, L. M. (1999). Are teachers of children and young adolescents responsive to suicide prevention training manuals? Yes. *Death Studies, 23*(1), 61-71.
- Davis, J. M., & Brock, S. E. (2002). Suicide. In J. Sandoval (Ed.), *Handbook of crisis counseling, intervention, and prevention in the schools* (2nd ed.; pp 273-299). Hillsdale, NJ: Earlbaum.
- Davis, J. M., & Sandoval, J. (1991). *Suicidal youth: School-based intervention and prevention*. San Francisco: Jossey-Bass.
- Davis, J. M., Sandoval, J. & Wilson, M. P. (1988). Strategies for the primary prevention of adolescent suicide. *School Psychology Review, 17*, 559-569.
- DeMan, A. F., & Leduc, C. P. (1993). Correlates of suicidal ideation in French-Canadian adolescents: Personal variables, stress, and social support. *Adolescence, 28*, 112-123.
- DeMan, A. F., & Leduc, C. P. (1995). Suicidal ideation in high school students: Depression and other correlates. *Journal of Clinical Psychology, 51*, 173-181.
- Demetriades, D., Murray, J., Myles, D., Chand, L., Sathyaragiswaran, L., Noguchi, T.,

- Bongard, F. S., Vryer, G. H., & Gaspard, D. J. (1998). Epidemiology of major trauma and trauma deaths in Los Angeles County. *Journal of the American College of Surgery*, 187, 373-383.
- Dishion, T. J., Capaldi, D., Spracklen, K. M., & Li, F. (1995). Peer ecology of male adolescent drug use. *Development and Psychopathology*, 7, 803-824.
- Dryfoos, J. (1994). Full-service schools: A revolution in health and social services for children, youth, and families. San Francisco: Jossey-Bass.
- Dubow, E. F., Kausch, D. F., Blum, M. C., Reed, J., & Bush, E. (1989). Correlates of suicidal ideation and attempts in a community sample of junior high and high school students. *Journal of Clinical Child Psychology*, 18, 158-166.
- Durkheim, E. (1951). Suicide: A study in sociology. In J. A. Spalding (Trans.) & K.E. Early (Eds.), *Suicide, a study in sociology* (pp. 305-321). New York: Free Press. (Original work published in 1897).
- Durlak, J. A., & Wells, A. M. (1997). Primary prevention mental health programs for children and adolescents: a meta-analytic review. *American Journal of Community Psychology*, 25, 115-152.
- Early, K. E., & Akers, R. L. (1993). "It's a White thing": An exploration about beliefs about suicide in the African-American community. *Deviant Behavior*, 14, 227-296.
- Eckersley, R. (1993). Failing a generation: The impact of culture on the health and well being of youth. *Journal of Pediatrics and Child Health*, 29, 516-519.

- Eckert, T. L. , Miller, D. N., DuPaul, G. J., & Riley-Tillman, T. C. (2003). Adolescent suicide Prevention: School psychologists' acceptability of school-based programs. *School Psychology Review, 32*(1), 57-76.
- Eggert, L. L., Thompson, E. A., Herting, J. R. (1994). A Measure of Adolescent Potential for Suicide (MAPS): Development and preliminary findings. *Suicide and Life-Threatening Behavior, 24*(4), 359-381.
- Eggert, L. L., Thompson, E. A., Herting, J. R., & Nicholas, L. J. (1994). A prevention research program: Reconnecting at-risk youth. *Issues in Mental Health Nursing, 15*(2), 107-135.
- Eggert, L. L., Thompson, E. A., Herting, J. R., & Nicholas, L. J. (1995). Reducing suicide potential among high-risk youth: Tests of a school-based prevention program. *Suicide and Life-Threatening Behavior, 25*(2), 276-296.
- Eggert, L. L., Thompson, E. A., Herting, J. R., & Randell, B., & Mazza, J. (1998). Identifying and assessing youth at risk for suicidal behaviors. Paper presented at workshop on Suicidality in Youth: Development of the Knowledge Base for Youth at Risk, National Institute of Mental Health, Bethesda, MD, November 1998.
- Eggert, L. L., Thompson, E. A., Randell, B. P., & McCauley, E. (1995). Youth Suicide Prevention Plan for Washington State. Olympia, WA: Washington Department of Health .
- Eisel v. Board of Education of Montgomery County, 597 A.2d 447 (Md. Ct. App. 1991).
- Elias, M. J. (1997). Reinterpreting dissemination of prevention programs as widespread Implementation with effectiveness and fidelity. In R.P. Weissberg, T. P. Gullotta,

- R. L. Hampton, B. A. Ryan, & G. R. Adams (Eds.), *Enhancing children's wellness* (pp.253-289), Thousand Oaks, CA: Sage.
- Elias, M. J., Gara, M. A., Schuyler, T. F., Branden-Muller, L. R., & Sayette, M.A. (1991). The promotion of social competence: Longitudinal study of a preventive school-based program. *American Journal of Orthopsychiatry*, *61*, 409-417.
- Emery, P. E. (1983). Adolescent depression and suicide. *Adolescence*, *18*(70), 245-258.
- Emslie, G. J., & Mayes, T. L. (2001). Mood disorders in children and adolescents: Psychopharmacological treatment. *Biological Psychiatry*, *49*, 1082-1090.
- Eshun, S. (2000). Role of gender and rumination in suicide ideation. *Cross-Cultural Research*, *34*, 250-264.
- Evans, W., Smith, M., Hill, G., Albers, E., & Neufeld, J. (1996). Rural adolescent views of risk and protective factors associated with suicide. *Crisis Intervention* *3*, 1-12.
- Farberow, M. L. (1991). Preparatory and prior suicidal behavior factors. In L. Davidson & M. Linnoila (Eds.), *Risk factors for youth suicide*. New York: Hemisphere.
- Farberow, N., Atman, S., & Thorne, A. (1985). *Report of the National Conference on Youth Suicide: June 19-20, 1985*. Washington, DC: Administration for Children, Youth and Families.
- Felner, R. D., & Felner, T. W. (1989). Primary prevention programs in the educational context: A transactional-ecological framework and analysis. In L. A. Bond & B. E. Compas (Eds.), *Primary prevention and promotion in the schools* (pp. 13-49). Newbury Park, CA: Sage.
- Felner, R. D., Felner, T. Y., & Silverman, M. M. (2000). Prevention in mental health and social intervention: Conceptual and methodological issues in the evolution of the

- science and practice of prevention. In J. Rapaport & E. Seidman (Eds.), *Handbook of community psychology* (pp. 9-42). New York: Plenum Press.
- Fergusson, D. M., Woodward, L. J., & Horwood, L. J. (2000). Risk factors and life processes associated with the onset of suicidal behavior during adolescence and early adulthood. *Psychological Medicine, 30*, 23-39.
- Fernquist, R. M. (2000). Problem drinking in the family and youth suicide. *Adolescence, 35*(139), 551-558.
- Fetro, J. V. (1998). Implementing coordinated school health programs in local schools. In E. Marx, S. F. Wooley, & D. Northrop (Eds.), *Health is academic: A guide to coordinated school health programs*. New York, NY: Teachers College Press.
- Flynn, L., Craft, L., McGuire, L. C., & Crandall, D. T. (2002). The Columbia TeenScreen Program: Providing mental health checkups to youth. *Emotional and Behavioral Disorders in Youth, 2*(4), 83-86.
- Forman, S. & Kalafat, J. (1998). Substance abuse and suicide: Promoting resilience against self-destructive behavior in youth. *The School Psychology Review, 27*(3), 398-406.
- Foster, T. (2001). Dying for a drink: Global suicide prevention should focus more on alcohol use disorders. *British Medical Journal, 323*(7317), 817-818.
- Frances, A., & Clarkin, J. F. (1985). Considering family versus other therapies after a teenager's suicide attempt. *Hospital and Community Psychiatry, 36*(10), 1041-1046.
- Frederick, C. J. & Resnick, A. L. P. (1971). How suicidal behaviors are learned. *American Journal of Psychotherapy, 25*, 37-55.



- Freeman, D. G. (1998). Determinants of youth suicide: The Easterlin-Holinger cohort Hypothesis re-examined. *American Journal of Economics and Sociology*, 57, 183-98.
- Gallup, G. (1991). *The Gallup Survey on Teenage Suicide*. Princeton, NJ: George H. Gallup International Institute.
- Garfinkel, B. D., Crosby, E., Herbert, M. R., Matus, A. L., Pfeifer, J. K., & Sheras, P. L. (1988). *Responding to adolescent suicide*. Bloomington, IN: Phi Delta Task Force on Adolescent Suicide, Phi Delta Kappa Educational Foundation. (ERIC Document Reproduction Service No. ED 301 813).
- Garfinkel, B. D., & Golombek, H. (1983). Suicidal behavior in adolescence. In H. Golombek & B. D. Garfinkel (Eds.), *The adolescent and mood disturbance*. New York: International University Press.
- Garland, A. F., Shaffer, D. & Whittle, B. (1989). A national survey of school-based, adolescent suicide prevention programs. *Journal of the American Academy of Child & Adolescent Psychiatry*, 28(6), 931-934.
- Garland, A. F. & Zigler, E. (1993). Adolescent suicide prevention: Current research and social policy implications. *American Psychologist*, 48(2), 169-182.
- Garlow, S. J., Purselle, D., & Heninger, M. (2005). Ethnic differences in patterns of suicide across the life cycle. *American Journal of Psychiatry*, 162:319-323.
- Garret Lee Smith Memorial Act, 42 U.S.C. § 201 (2004).
- Gibbs, J. T. (1997). African-American suicide: A cultural paradox. *Suicide and Life-Threatening Behavior*, 27(1), 68-79.

- Glowinski, A. L., Bucholz, K. K., Nelson, E. C., Quiang Fu, P., Madden, A. F., Reich, W., et al. (2001). Suicide attempts in an adolescent female twin sample. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40, 1300-1308.
- Gould, M. S. (2001). Suicide and the media. *Annals of the New York Academy of Sciences*, 932, 200-224.
- Gould, M. S., Fisher, P., Parides, M., Flory, & Shaffer, D. (1996). Psychosocial risk factors of child and adolescent completed suicide. *Archives of General Psychiatry*, 53(12), 1155-1162.
- Gould, M. S., Greenberg, T., Velting, D. M., & Shaffer, D. (2003). Youth suicide risk and factors: A review of the past 10 years. *Journal of the Academy of Child and Adolescent Psychiatry*, 42(4), 386-405.
- Gould, M. S., & Kramer, R. A., (2001). Youth suicide prevention. *Suicide & Life-Threatening Behavior*, 31, 6-31.
- Gould, M. S., Petrie, K., Kleinman, M., & Wallenstein, S. (1994). Clustering of attempted suicide: New Zealand national data. *International Journal of Epidemiology*, 23, 1185-1189.
- Gould, M. S., & Shaffer, D. (1991). Epidemiology and child psychiatry. In R. Michels, A.M. Cooper, S. B. Guze, L. L. Kudd, G. L. Klerman, A. J. Solnit, & A. J. Stunkard (Eds.), *Psychiatry*. New York: Lippincott.
- Gould, M. S., Shaffer, D., Fisher, P., & Garfinkel, R. (1998). Separation/divorce and child and adolescent completed suicide. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(2), 155-163.

- Gould, M. S., Wallenstein, S., Kleinman, M. H., O'Carroll, P. W., & Mercy, J. A. (1990). Suicide clusters: An examination of age-specific effects. *American Journal of Public Health, 80*, 211-212.
- Greenberg, M. T., Domitrovich, C., & Bumbarger, B. (2001). The prevention of mental disorders in school-aged children: Current state of the field. *Prevention & Treatment, 4*(1).
- Grunbaum, J., Kann, L., Kinchen, S., Williams, B., Ross, J., Lowry, R, et al. (2002). Youth risk behavioral surveillance. *MMWR: Morbidity and Mortality Weekly Report, 51*(SS04), 1-62.
- Guo, G., & Harstall, C. (2002). *Efficacy of suicide prevention programs for children and youth*. HTA 26: Services A Health Technology Assessment (available from Director, Health Technology). Alberta, Canada: Alberta Heritage Foundation for Medical Research.
- Gutkin, T. (Ed.). (2002). Evidence-based interventions in school psychology: The state of the art and future directions [Special issue]. *School Psychology Quarterly, 17*(4), 339-341.
- Hammond, W. R. (2001). Suicide prevention: Broadening the field toward a public health approach. *Suicide and Life-Threatening Behavior, 32*, 1-2. Retrieved April 19, 2003 from <http://proquest.umi.com/pqdlink?Ver-1&Exp=02-15-2008&FMT-FT&DID-0000001144>
- Harry, J. (1991). Sexual identity issues. In L. Davidson & M. Linnoila (Eds.), *Risk factors for youth suicide*. New York: Hemisphere.

- Hawkes, M., & Romiszowski, A. (2001). Examining the reflective outcomes of asynchronous computer-mediated communication on in-service teacher development. *Journal of Technology and Teacher Education*, 9(2), 285-308.
- Hawkins, J. D. & Catalano, R. F. (1992). *Communities that care: Action for drug abuse Prevention*. San Francisco: Jossey-Bass.
- Hawton, K. (1986). *Suicide and attempted suicide among children and adolescents*. Beverly Hills, CA: Sage.
- Hawton, K., Fagg, J., & Simkin, S. (1996). Deliberate self-poisoning and self-injury in children and adolescents under 16 years of age in Oxford, 1976-1993. *British Journal of Psychiatry*, 69(2), 202-208.
- Hawton, K., Simkin, S., Deeks, J. J., O'Connor, S., Keen, A., Altman, D. G., et al. (1999). Effects of a drug overdose in a television drama on presentations to hospital for self-poisoning: Times series and questionnaire study. *British Medical Journal*, 318, 972-977.
- Hawton, K., Townsend, E., Deeks, J., Appleby, D., Bennewith, O., & Cooper, J. (2001). Effects of legislation restricting pack sized of paracetamol and salicylate on self-poisoning in the United Kingdom: before and after study. *British Medical Journal*, 322, 1203-1207.
- Hayden, D. C. & Lauer, P. (2000). Prevalence of suicide programs in schools and roadblocks to implementation. *Suicide & Life-Threatening Behavior*, 30(3), 239-251.
- Hazell, P. (1993). Adolescent-suicide clusters: Evidence, mechanisms, and prevention. *Australian and New Zealand Journal of Psychiatry*, 30, 633-642.

- Hazell, P., & King, R. (1996). Arguments for and against teaching suicide prevention in schools. *Australian and New Zealand Journal of Psychiatry*, 30, 633-642.
- Helsel, D. C. (2001). Does your school track the suicidal student? *The Clearing House*, 75(2), 92-95.
- Henning, C. W., Crabtree, C. R., & Baum, T. (1998). Peer contracting as a response to potential suicide in adolescents. *Archive of suicide research*, 4, 169-187.
- Henry, C. (1996). Adolescent suicide and families: An ecological approach. *Adolescence*, 28, 110, 291-308.
- Henry, C. S., Stephenson, A. L., Hanson, F., & Hargett, W. (1993). Adolescent suicide and families: An ecological approach. *Adolescence*, 28, 291-308.
- H. R. No. 212, 105<sup>th</sup> Cong., 143 Cong. Rec. 57 (1997) (enacted).
- H R. Rep. No. 99-4650 (1986).
- Hoberman, H. M., & Garfinkel, B.D. (1988). Completed suicide in children and adolescents. *Journal of the Academy of Child and Adolescent Psychiatry*, 27, 689-695.
- Hoff, L. (1991). Crisis intervention in schools. In A. A. Leenaar & Wenckstern (Eds.). *Suicide prevention in the schools* (pp.123-14). New York: Hemisphere Publishing Corp.
- Holinger, P. C., & Offer, D. (1981). Perspectives on suicide in adolescence. In R. G. Simmons (Ed.), *Research in community and mental health* (Vol. 2; pp. 139-157). Greenwich, CT: JAI Press.
- Huberman, A. M., & Miles, M. G. (1984). *Innovation up close: How school improvement works*. New York: Plenum Press.

- Indian Health Service (IHS). (1997). *Trends in Indian health 1997*. Retrieved September 9, 2003 at <http://www.ihs.gov/PublicInfo/Publications/trends97/trends97.asp>
- Indian Health Service (IHS). (1999). *Indian health focus: Injuries, 1989-99*. Washington, DC: U.S. Department of Health and Human Services, Indian Health Service.
- Institute of Medicine. (1994). *Reducing risks for mental disorders: Frontiers for Preventive intervention research*. Washington, DC: National Academy Press.
- Ishii, K. (1991). Measuring mutual causation: Effect of suicide news on suicide in Japan. *Social Science Research, 20*, 188-195.
- Jackson, H. P., Hess, P. M., & van Dalen, A. (1995). Preadolescent suicide: How to ask and how to respond. *Families in Society: The Journal of Contemporary Human Services, (May)*, 267-79.
- Jacobs, D. G., Brewer, M., & Klein-Benham, M. (1999). Suicide assessment: An overview and recommended protocol. In D. G. Jacobs (Ed.), *The Harvard Medical School guide to assessment and intervention* (pp. 3-39). San Francisco: Jossey-Bass.
- James, S. H., & DeVaney, S. B. (1995). Preparing to testify: The school counselor as court witness. *School Counselor, 42*, 261-267.
- Jamison, K. R., & Baldessarini, R. J. (Eds.). (1999). Effects of medical interventions on suicidal behavior. *Journal of Clinical Psychiatry, 60* (Suppl. 2), 1-122.
- Jennings, J., Pearson, G., & Harris, M. (2000). Implementing and maintaining school based mental health services in a large, urban school district. *Journal of School Health, 70*(5), 201-205.

- Jessor, R., Van Den Bos, J., Vanderryn, J., Costa, F., & Turbin, M. S. (1995). Protective factors in adolescent problem behavior: Moderator effects and developmental change. *Developmental Psychology, 31*(6), 923-933.
- Jobes, D. A., Berman, A. L., O'Carroll, P. W., Eastgard, S., & Knickmeyer, S. (1996). The Kurt Cobain suicide crisis: Research, public health, and news media perspectives. *Suicide and Life-Threatening Behavior, 26*, 260-271.
- Johnson, J. G., Cohen, P., Gould, M. S., Kasen, S., Brown, J., & Brook, J. S. (2002). Childhood adversities, interpersonal difficulties, and risk for suicide attempts during late adolescence and early adulthood. *Archives of General Psychiatry, 59*, 741-749.
- Joiner, T.E., Pfaff, J.J., & Acres, J. G. (2002). A brief screening tool for suicidal symptoms in adolescents and young adults in general health settings: Reliability and validity data from the Australian National General Practice Youth Suicide Prevention Project. *Behavior Research and Therapy, 40*, 471-481.
- Jones, G. D. (1997). The role of drugs and alcohol in urban minority adolescent suicide attempts. *Death Studies, 21*, 189-203.
- Kalafat, J. (1990). Adolescent suicide and the implications for school counselors. *The School Counselor, 37*, 359-69.
- Kalafat, J. (1997) The prevention of youth suicide. In R. P. Weissberg, T. P. Gullatta, B. A. Ryan, & G. R. Adams (Eds.), *Healthy children 2010: Enhancing children's wellness* (pp.175-213) Thousand Oaks, CA: Sage.
- Kalafat, J. & Elias, M. (1992). Adolescents' experience with and response to suicidal peers. *Suicide and Life-Threatening Behavior, 22*, 315-321.

- Kalafat, J. & Elias, M. (1994). An evaluation of a school-based suicide awareness intervention. *Suicide and Life-Threatening Behavior*, 22, 315-321.
- Kalafat, J. & Elias, M. (1995). Suicide prevention in the education context. *Suicide and Life-Threatening Behavior*, 25, 125-133.
- Kalafat, J. & Gagliano, C. (1996). The use of simulations to assess the impact of an adolescent suicide response curriculum. *Suicide and Life-Threatening Behavior*, 26(4), 359-364.
- Kalafat, J., & Lazarus, P. J. (2002). Suicide Prevention in Schools. In S. E. Brock, P. J. Lazarus, S. R. Jimerson (Eds.), *Best practices in school crisis prevention and intervention* (pp. 211-224). Bethesda, MD: National Association of School Psychologists.
- Kalafat, J., & Ryerson, D. M. (1999). The implementation and institutionalization of a school-based youth suicide prevention program. *The Journal of Primary Prevention*, 19(3), 157-175.
- Kalafat, J., & Underwood, M. (1989). *Lifelines: A school-based adolescent suicide response program*. Dubuque, IA: Kenal/Hunt.
- Kaslow, N. J., McClure, E., & Connell, A. (2002). Treatment for depression in children and adolescents (pp. 441-464). In I. H. Gotlib & C. L. Hammen (Eds.), *Handbook of depression*. New York: Guilford Press.
- Kazdin, A. E. (1991). Prevention of conduct disorder. *The prevention of mental disorders: Progress, problems, and prospects*. Washington, DC: National Institute of Mental Health.



- Kellermann, A. L., Rivara, F. P., Somes, G., Reay, D. T., Francisco, J., Gillentine Banton, J., et al. (1992). Suicide in the home in relation to gun ownership. *New England Journal of Medicine*, 327, 467-472.
- Kelson v. City of Springfield, 767 F.2d 651 (1985).
- Kessler, R. C., & Walters, E. (1998). Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Comorbidity Survey. *Depression and Anxiety*, 7, 3-14.
- King, K. A. (2001). Developing a comprehensive school suicide prevention program. *The Journal of School Health*, 71(4), 132-37.
- King, K. A., Price, J. H., Telljohann, S. K., & Wahl, J. (1999). High school health teachers' knowledge of adolescent suicide. *American Journal of Health Studies*, 15(3), 156-163.
- King, K. A., & Smith, J. (2000). Project SOAR: A training ground to increase school counselors' knowledge and confidence regarding suicide prevention and intervention. *Journal of School Health*, 70, 402-407.
- King, K. A., Vidourek, R. A., & Davis, B. (2002). Increasing self-esteem and school connectedness through a multidimensional mentoring program. *The Journal of School Health*, 72(7), 294-299.
- Kirby, L. M., & Fraser, M. (1997). Risk and resilience in childhood. In M. Fraser (Ed.). *Risk and resilience in childhood: An ecological perspective*. Washington, DC: NASW Press.
- Kleespies, P. M. (Ed.). (1998). *Emergencies in mental health practice: Evaluation and management*. New York: Guilford Press.

- Knickmeyer, S. (1996). Commentary-the media perspective. *Suicide and Life-Threatening Behavior, 26*, 269-271.
- Kosky, R., Silburn, S., & Zubrick, S. R. (1990). Are children and adolescents who have suicidal thoughts different from those who attempt suicide? *Journal of Nervous Mental Disorders, 178*, 38-43.
- Krug, E. G., Dahlberg, L. L., Mercy, J. A., Zwi, A., Lozano, R. (Eds.). (2002). *World report on violence and health*. Geneva: World Health Organization.
- Kush, F. A. (1991). A descriptive study of school-based adolescent suicide prevention/intervention programs: Program components and the role of the school counselor. *Dissertation Abstracts International, 52*, 1692A.
- Lacourse, E., Claes, M., & Villeneuve, M. (2001). Heavy metal music and adolescent suicidal risk. *Journal of Youth and Adolescents, 30*(3), 321-332.
- LaFramboise, T., & Howard Pitney, B. (1995). The Zuni life skills development and curriculum: Description and evaluation of a suicide prevention program. *Journal of Counseling Psychology, 42*(4), 479-486.
- Landau-Stanton, J., & Stanton, M. D. (1985). Treating suicidal adolescents and their families. In M. P. Mirkin & S. L. Koman (Eds.), *Handbook of adolescent and family therapy* (pp.309-328). New York: Gardner Press.
- Leane, W., & Shute, R. (1998). Youth suicide: The knowledge and attitudes of Australian teachers and clergy. *Suicide and Life-Threatening Behavior, 28*, 165-73.
- Leenaars, A. A., & Wenckstern, S. (1990). Suicide prevention in schools: An introduction. *Death Studies, 14*, 297-302.

- Leenaars, A. A., & Wenckstern, S. (1998). Principles of postvention: Applications to suicide and trauma in schools. *Death Studies, 22*, 357-391.
- Lester, D. (1997). The role of shame in suicide. *Suicide and Life-Threatening Behavior, 27*, 352-361.
- Lewis, M. W., & Lewis, A. C. (1996). Peer helping programs: Helper role, supervisor training, and suicidal behavior. *Journal of Counseling Development, 74*, 307-313.
- Lewis, M. W., & Lewis, A. C. (1998). Peer helping programs: Helper role, supervisor training suicidal behavior. *Peer-Facilitator Quarterly, 15*, 64-74.
- Lewinsohn, P. M., Rohde, P., & Seeley, J. R. (1993). Psychosocial characteristics of adolescents with a history of suicide attempt. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*, 60-68.
- Lewinsohn, P. M., Rohde, P., & Seeley, J. R. (1994). Psychosocial risk factors for future adolescent suicide attempts. *Journal of Consulting and Clinical Psychology, 62*, 297-305.
- Lieberman, R., & Davis, J. (2002). Suicide intervention. In S. E. Brock, P. J. Lazarus, & S. R. Jimerson (Eds.), *Best practices in school crisis prevention and intervention*. Bethesda, MD: National Association of School Psychologists.
- Loeber, R., Green, S. M., & Lahey, B.B. (1990). Mental health professionals' perception of the utility of children, mothers, and teachers as informants on childhood psychopathology. *Journal of Clinical Child Psychology, 19*, 136-143.
- Ludwig, J. & Cook, P. J. (2000). Homicide and suicide rates associated with implementation of the Brady Handgun Violence Prevention Act. *Journal of the American Medical Association, 284*, 585-591.

- Lyon, M. E., Benoit, M., O'Donnell, R. M., Getson, P.R., Silber, T., & Walsh, T. (2000). Assessing African American adolescents' risk for suicide attempts: Attachment theory. *Adolescence, 35*(137), 121-134.
- MacKay, A. P., Fingerhut, L. A., & Duran, C. R. (2000). Adolescent Health Chartbook: Health, United States, 2000. Hyattsville, MD: National Center for Health Statistics.
- Mackesy-Amiti, M.E., Fendrich, M., Libby, S., Goldenberg, D., & Grossman, J. (1996). Assessment of knowledge gains in pro-active training for postvention. *Suicide and Life-Threatening Behavior, 26*, 161-174.
- Malley, P.B., Kush, F., & Bogo, R.J. (1994). School-based adolescent suicide and prevention programs: A survey. *School Counselor, 42*, 130-136.
- Malone, K.M., Oquendo, M. A., Haas, G. L., Ellis, S. P., Li, S., & Mann, J. J. (2000). Protective factors against suicidal acts in major depression: Reasons for living. *American Journal of Psychiatry, 157*, 1084-1089.
- Maris, R. W. (2002). Suicide. *The Lancet, 360*, 319-326.
- Markus, H. R., Kitayama, S., & Heiman, R. J. (1996). Culture and "basic" psychological principles. In E. T. Higgins & A. W. Kruglanski (Eds.), *Social psychology: Handbooks of basic principles* (pp. 857-913). New York: Guilford Press.
- Marttunen, M. J., Aro, H. M., & Lonnqvist, J. K. (1993). Precipitant stressors in adolescent suicide. *Journal of the American Academy of Child and Adolescent Psychiatry, 32*(6), 1178-1183.

- Marttunen, M. J., Hillevi, M. A., Hendriksson, M. M., & Lönnqvist, J. K. (1991). Mental disorders in adolescent suicide: DSM-III-R Axes I and II diagnoses in suicides among 13-19- year olds in Finland. *Archives of General Psychiatry*, 48, 834-839.
- Marzuk, P. M., Leon, A. C., Tardiff, K., & Morgan, E. B. (1992). The effects of access to lethal methods of injury on suicide rates. *Comprehensive textbook of suicidology*. New York: Guilford Press.
- Mazza, J. J. (1997). School-based suicide prevention programs: Are they effective? *School Psychology Review*, 26, 382-396.
- McDaniel, J. S., Purcell, D. W., & D'Augelli, A. R. (2001). The relationship between sexual orientation and risk for suicide: Research finding and future directions for research and prevention. *Suicide and Life-Threatening Behavior*, 31(1) (Suppl.), 84-105.
- McEvoy, M. L., & McEvoy, A. W. (1994). *Preventing youth suicide. A handbook for educators and human service professionals*. Holmes Beach, FL: Learning Publications, Inc.
- McGuire, L C., & Flynn, L. (2003). The Columbia TeenScreen Program: Screening youth for mental illness and suicide. *Trends in Evidenced-Based Neuropsychiatry*, 5(2), 56-62.
- McKee, P.W., Jones, R. W., & Barbe, R. H. (1993). *Suicide and the school: A practical guide to suicide prevention*. Horsham, PA: LRP Publications.
- McKeown, R. E., Garrison, C. Z., Cuffe, S. P., Waller, J. L., Jackson, K. L., & Addy, C. L. (1998). Incidence and predictors of suicidal behaviors in a longitudinal sample

- of young adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37, 612-619.
- McLaughlin, J., Miller, R., & Warwick, H. (1996). Deliberate self-harm in adolescents: Hopelessness, depression, and problem solving. *Journal of Adolescence*, 19, 523-532.
- Metha, A., Weber, B. & Webb, L. D. (1998). Youth suicide prevention: A survey and analysis of policies and efforts in the 50 states. *Suicide and Life-Threatening Behavior*, 28(2), 150-164.
- Miller, D. N., Eckert, T. L., DuPaul, G. J., & White, G. P. (1999). Adolescent suicide prevention: Acceptability of school-based programs among secondary school principals. *Suicide and Life Threatening Behavior*, 29, 72-85
- Milsom, A. (2002). Suicide prevention in schools: Court cases and implications for principals. *National Association for Secondary School Principals (NASSP) Bulletin*, 86, 24-33.
- Minkler, M. & Roe, K. M. (1993). *Grandmothers as caregivers: Raising children of the crack cocaine epidemic*. Newbury Park, CA: Sage Publications.
- Mishara, B. L., & Daigle, M. S. (2001). Help lines and crisis intervention services: Challenge for the future. In D. Lester (Ed.), *Suicide Prevention: Resources for the Millennium* (pp.153-171). Philadelphia: Brunnel/Mazel.
- Molin, R. S. (1986). Covert suicide and families of adolescents. *Adolescence*, 21(90), 469-475.

- Morgan, G. (1993). Long term risks after attempted suicide: Identifying risk factors should help to reduce subsequent suicide. *British Medical Journal*, 306, 1626-1628.
- Moscicki, E. K. (1997). Identification of suicide risk factors using epidemiological studies. *Psychiatric Clinics of North America*, 20, 499-517.
- Motes, P. S., Melton, G., Simmons, W. E. W. , & Pumariega, A. (1999). Ecologically oriented school-based mental health services: Implications for service system delivery. *Psychology in the Schools*, 36(5), 391-401.
- National Adolescent Health Information Center (NAHIC). (2000). Fact sheet on mortality: Adolescents and young adults. San Francisco: University of California, San Francisco, NAHIC.
- National Institutes of Mental Health (NIMH). (2004). Pragmatic considerations of culture in preventing suicide. – *Proceedings of NIH Office of Rare Diseases, National Institute of Mental Health, Indian Health Service, Annenberg Foundation Trust at Sunnylands, and Emerging Scholars Interdisciplinary Network*, 1-6.
- URL: <http://www.nimh.nih.gov/scientificmeetings/suicideprevention2004.pdf>.
- Retrieved from internet November 28, 2005.
- National Institutes of Mental Health (NIMH). (2000). Child and adolescent violence research at NIMH. Washington, DC: Author.
- New Freedom Commission on Mental Health (NFCMH). (2003). *Achieving the promise: Transforming mental health care in America*. Final Report. DHHS Pub. No. SMA-03-3832. Rockville, MD: Author.

- O'Carroll, P.W., Berman, A. L., Maris, R., Moscicki, E., Tanney, B., & Silverman, M. (1998). Beyond the Tower of Babel: A nomenclature of suicidology. In R. J. Kosky, et al. (Eds.). *Suicide prevention: The global context* (pp. 23-40). New York: Plenum Press.
- O'Carroll, P. W., & Silverman, M. M. (1994). Community suicide prevention: The effectiveness of bridge barriers. *Suicide and Life-Threatening Behavior*, 24, 89-99.
- Offer, D., Howard, K. I., Schonert, K. A., & Ostrov, E. (1991). To whom do adolescents turn for help? Differences between disturbed and non-disturbed adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 30(4), 623-630.
- Olfson, M., Marcus, S. C., Weissman, M. M., & Jensen, P. S. (2002). National trends in the use of outpatient psychotropic medications by children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 514-521.
- Ollendick, T. H., Greene, R. W., Werst, M. D., & Oswald, D. P. (1990). The predictive validity of teacher nominations: A five-year follow-up of at-risk youth. *Journal of Abnormal Child Psychology*, 18, 699-713.
- Orbach, I., & Bar-Joseph, H. (1993). The impact of a suicide prevention program for adolescents with suicidal tendencies, hopelessness, ego identity, hopelessness, ego identity, and coping. *Suicide and Life-Threatening Behavior*, 23, 120-129.
- Overholser, J. C., Hemstreet, A. H., Spirito, A., & Vyse, S. (1989). Suicide awareness programs in the schools: Effects of gender and personal experience. *Journal of the American Academy of Child and Adolescent Psychiatry*, 28, 925-930.



- Panak, W. F., & Garber, J. (1992). Role of aggression, rejection, and attributions in the prediction of depression in children. *Development and Psychopathology, 4*, 145-165.
- Patton, M. Q. (1990). *Qualitative Evaluation and Research Methods*. Newbury Park, CA: Sage Publications.
- Paykel, E. S. (1991). Stress and life events. In L. Davidson & M. Linnoila (Eds.), *Risk factors for youth suicide*. New York: Hemisphere.
- Perr, I. N. (1985). Right to refuse treatment and delayed medical care for psychiatric patients. *American Journal of Psychiatry, 142*, 1127.
- Petersen, G. J., Pietrzak, D., & Speaker, K. M. (1998). The enemy within: A national study on school violence and prevention. *Urban Education, 33* (3), 331-359.
- Petersen, G. J., & Speaker, K. M. (1999). An early intervention model for violence and suicide reduction. *Contemporary Education, 71*(1), 14-18.
- Piacentini, J., Rotheram-Borus, M. J., Gillis, J. R., Graae, F., Trautman, P., Cantwell, C., et al. (1995). Predictors of treatment and attendance among adolescent suicide attempters. *Journal of Consulting and Clinical Psychology, 63*, 469-473.
- Pinto, A., & Whisman, M. A. (1996). Negative affect and cognitive biases in suicidal and nonsuicidal hospitalized adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry, 35*, 158-165.
- Poland, S. (1989). *Suicide intervention in the schools*. New York: The Guilford Press.
- Poland, S., & McCormick, J. S. (1999). *Coping with crisis: Lessons learned*. Longmont, CO: Sopris West.

- Popenhagen, M. P., & Qualley, R. M. (1998). Adolescent suicide: Detection, intervention, and prevention. *Professional School Counseling, 1*, 30-35.
- Portner, J. (2000). Suicide: Many Schools Fall Short on Prevention. *Education Week, 19*(32), 1, 20-22.
- Prothrow-Stith, D. (1994, April). Building violence prevention into curriculum: A physician-administrator applies a public health model to schools. *The School Administrator, 4*(51), 8-12.
- Queralt, M. (1993). Psychosocial risk factors associated with suicide in a small community sample of Latino adolescent attempters. *Social Work in Education, 15*(2), 91-103..
- Randell, B. P., Eggert, L. L., & Pike, K. C. (2001). Immediate post-intervention effects of two brief youth suicide prevention interventions. *Suicide and Life-Threatening Behavior, 31*, 41-61.
- Range, M. (1993) Suicide prevention: Guidelines for schools. *Educational Psychology Review, 5*, 135-154.
- Reid, W. H. (1998). Promises, promises: Don't rely on patients' no-suicide/no-violence "contracts." *Journal of Practicing Psychiatry and Behavioral Health, 4*, 316-318.
- Remley, T. P., Jr., & Herlihy, B. (2001). *Ethical, legal, and professional issues in counseling*. Upper Saddle River, NJ: Prentice Hall.
- Resnick, M. D., Berman, P. S., & Blum, R. W., Bauman, K. E., Harris, K. M., Jones, J. et al. (1997). Protecting adolescents from harm: Findings from the National Longitudinal Study on Adolescent Health. *Journal of the American Medical Association, 278*(10), 823-832.

- Resnick, M. D., Harris, L. J., & Blum, R. W. (1993). The impact of caring and connectedness on adolescent health and well-being. *Journal of Paediatrics and Child Health*, 29(Suppl 1), S3-9.
- Rew, L., Thomas, N., Horner, S. D., et al. (2001). Correlates of recent suicide attempts in a triethnic group of adolescents. *Journal of Nursing Scholars*, 33, 361-367.
- Reynolds, W. M. (1988). *Suicidal ideation questionnaire: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Reynolds, W. M. (1991). A school-based procedure for the identification of adolescents at-risk for suicidal behaviors. *Family and Community Health*, 14, 64-75.
- Reynolds, W. M., & Mazza, J. (1994). Suicide and suicidal behavior in children and adolescents. In W.M. Reynolds & H.F. Johnston (Eds.), *Handbook of depression in children and adolescents* (pp.525-580). New York: Plenum Press.
- Rich, C. L., Young, J. G., Fowler, R. C. Wagner, J. & Black, N.A. (1990). Guns and suicide: Possible effects of some specific legislation. *American Journal of Psychiatry*, 147, 342-346.
- Roberts, R. E., Roberts, C. R. & Chen, R. Y. (1998). Suicidal thinking among adolescents with a history of attempted suicide. *Journal of the American Academy of Child and Adolescent Psychiatry*, 37(12) 1294-1300.
- Roberts, R. E., Roberts, C. R. & Chen, R. Y. (1997). Ethnocultural differences in prevalence of adolescent depression, *American Journal of Community Psychology*, 25: 1,95-105

- Roberts, R. L., Lepkowski, W. J., Davidson, K. (1998). After a student suicide, the team approach. *The Education Digest*, 64(2), 50-55. Retrieved April 19, 2003 from <http://webspirs3.silverplatter.com/cgi-bin/waldo.cgi>
- Roeser, R. W., & Midgley, C. (1997). Teacher's view of issues involving students' mental health. *The Elementary School Journal*, 98, 115-33.
- Rose, G. (1992). *The strategy of preventive medicine*. Oxford: Oxford University Press.
- Ross, C. P. (1980). Mobilizing schools for suicide prevention. *Suicide and Life-Threatening Behavior*, 10, 239-244.
- Rotheram-Borus, M. J., Trautman, P. D., Dopkins, S. C., & Shrout, P.E. (1990). Cognitive style and pleasant activities among female adolescent suicide attempters. *Journal of Consulting and Clinical Psychology*, 58, 554-561.
- Rowling, L., & Holland, J. (2000). Grief and school communities: The impact of social context, a comparison between Australia and England. *Death Studies*, 24, 34-50.
- Rubenstein, J. L., Halton, A., Kasten, L., Rubin, C., & Stechler, G. (1998). Suicidal behavior in adolescents: Stress and protections in different family contexts. *American Journal of Orthopsychiatry*, 68, 274-284.
- Rubenstein, J. L., Heeren, T., Housman, D., Rubin, C., & Stechler, G. (1989). Suicidal behavior in "normal" adolescents: Risk and protective factors. *American Journal of Orthopsychiatry*, 59, 59-71.
- Rudd, M. D. (2000). Integrating science into the practice of clinical suicidology: A review of the psychotherapy literature and a research agenda for the future. In R. W. Martis, S. S. Canetto, J. L. McIntosh, & M. M. Silverman (Eds.), *Review of Suicidology* (pp. 47-83). New York: Guilford Press.

- Rudd, M. D., & Joiner, T. (1998). The assessment, management, and treatment of suicidality: Toward clinically informed and balanced standards of care. *Clinical Psychology: Science and Practice*, 5, 135-150.
- Ruof, S. R., & Harris, J. M. (1988). How to get suicide prevention training and a crisis team for your schools. *Communique*, 16(10), 28-29.
- Russell, S. T., & Joyner, K. (2001). Adolescent sexual orientation and suicide risk: Evidence from a national study. *American Journal of Public Health*, 91 1276-1281.
- S. 2175, 108th Cong., (2004).
- S. Res. 84, 105<sup>th</sup> Cong., 143 Cong. Rec. 57 (1997) (enacted).
- Sadowski, C., & Kelley, M. L. (1993). Social problem-solving in suicidal adolescents. *Journal of Consulting and Clinical Psychology*, 61, 121-127.
- Salk, L., Lipsitt, L. P., Sturner, W. Q., Reilly, B. M., & Levat, R. H. (1985). Relationship of maternal and prenatal conditions to eventual adolescent suicide. *The Lancet*, 16(1), 624-627.
- Sanford, M. N., Offord, D. R., Boyle, M. H., Peace, A., & Racine, Y. A. (1992). Ontario child health study: Social and school impairments in children aged 6 to 16 years. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31, 60-67.
- Sarason, S. B. (1982). *The culture of the school and the problem of change*. Boston: Allyn and Bacon.
- Satcher, D. (1998). Bringing the public health approach to the problem of suicide. *Suicide and Life-Threatening Behavior*, 28, 325-327.

- Scherff, A. R., Eckert, T. L., Miller, D. N. (2005). Youth suicide prevention: A survey of public school superintendents' acceptability of school-based programs. *Suicide and Life-Threatening Behavior*, 2, 154-169.
- Schmidtke, A., & Schaller, S. (2000). The role of mass media in suicide prevention. In K. Hawton & K. van Heeringen (Eds.), *The international handbook of suicide and attempted suicide* (pp.121-137). New York: John Wiley & Sons.
- Schoenfeldt, M. (1993). *School crisis response teams: Lessening the aftermath* (3rd ed.). Marysville, WA: Schoenfeldt & Associates.
- School Mental Health Project/Center for Mental Health in Schools. (2004). Integrating agendas for mental health in schools in to the recommendations of the president's New Freedom Commission on Mental Health. *Addressing Barriers to Learning*, 9(1).
- Scouller, K. M., & Smith, D. I. (2002). Prevention of youth suicide: How well informed are the potential gatekeepers of adolescents in distress? *Suicide & Life-Threatening Behavior*, 32(1), 67-79.
- Screening for Mental Health, Inc. (2002). *High school suicide prevention program reduces suicide attempts*. Wellesley Hills, MA: Author.
- Shaffer, D. (1974). Suicide in childhood and early adolescence. *Journal of Child Psychology and Psychiatry*, 15, 275-291.
- Shaffer, D., & Craft, L. (1999). Methods of adolescent suicide prevention. *Journal of Clinical Psychiatry*, 60, 70-74.