

PRIMATES IN PERIL

The world's 25 most endangered primates
2018-2020

Edited by

Christoph Schwitzer, Russell A. Mittermeier, Anthony B. Rylands, Federica Chiozza,
Elizabeth A. Williamson, Dirck Byler, Serge Wich, Tatyana Humle, Caspian Johnson,
Holly Mynott and Gráinne McCabe



Bristol Zoological
Society
Saving Wildlife Together



GLOBAL
WILDLIFE
CONSERVATION



PRIMATES IN PERIL

The world's 25 most endangered primates
2018-2020

Edited by

Christoph Schwitzer, Russell A. Mittermeier, Anthony B. Rylands,
Federica Chiozza, Elizabeth A. Williamson, Dirck Byler, Serge Wich,
Tatyana Humle, Caspian Johnson, Holly Mynott and Gráinne McCabe

Illustrations by

Stephen D. Nash

IUCN SSC Primate Specialist Group (PSG)
International Primatological Society (IPS)
Global Wildlife Conservation (GWC)
Bristol Zoological Society (BZS)

CONTENTS

Published by: IUCN SSC Primate Specialist Group (PSG), International Primatological Society (IPS), Global Wildlife Conservation (GWC), Bristol Zoological Society (BZS)

Copyright: ©2019 Global Wildlife Conservation
All rights reserved. No part of this report may be reproduced in any form or by any means without permission in writing from the publisher.
Inquiries to the publisher should be directed to the following address:
Russell A. Mittermeier, Chair, IUCN SSC Primate Specialist Group, Global Wildlife Conservation, c/o WeWork, 718 7th Street, N.W., Washington, DC 20001, USA.

Citation (report): Schwitzer, C., Mittermeier, R.A., Rylands, A.B., Chiozza, F., Williamson, E.A., Byler, D., Wich, S., Humle, T., Johnson, C., Mynott, H., and McCabe, G. (eds.). 2019. *Primates in Peril: The World's 25 Most Endangered Primates 2018–2020*. IUCN SSC Primate Specialist Group, International Primatological Society, Global Wildlife Conservation, and Bristol Zoological Society, Washington, DC. 130pp

Citation (species): Perkin, A. 2019. Rondo Dwarf Galago *Paragalago rondoensis* (Honest in Kingdon, 1997). In: C. Schwitzer, R.A. Mittermeier, A.B. Rylands, F. Chiozza, E.A. Williamson, D. Byler, S. Wich, T. Humle, C. Johnson, H. Mynott and G. McCabe (eds.), *Primates in Peril: The World's 25 Most Endangered Primates 2018–2020*, pp. 24–27. IUCN SSC Primate Specialist Group, International Primatological Society, Global Wildlife Conservation, and Bristol Zoological Society, Washington, DC.

Layout: Sarah Clark, Bristol Zoological Society, Clifton, Bristol, UK, BS8 3HA

Illustrations: © Stephen D. Nash, Global Wildlife Conservation, Washington DC, and Department of Anatomical Sciences, Health Sciences Center, State University of New York at Stony Brook, NY, USA

Available from: Jill Lucena, Global Wildlife Conservation, c/o WeWork, 718 7th Street, N.W., Washington, DC 20001, USA
E-mail: jlucena@globalwildlife.org
Website: www.primates-sg.org

Printed by: Gemini West, Bristol, UK

ISBN: 978-1-5272-4806-9

Cover photo: Tapanuli orangutan *Pongo tapanuliensis* © Maxime Aliaga

Acknowledgements 2
The World's 25 Most Endangered Primates: 2018–2020 3

Madagascar	4	Neotropics	76
Bemansy mouse lemur (<i>Microcebus manitatra</i>)	6	Buffy-tufted-ear marmoset (<i>Callithrix aurita</i>)	78
Lake Alaotra gentle lemur (<i>Hapalemur alaotrensis</i>)	9	Pied tamarin (<i>Saguinus bicolor</i>)	82
James' sportive lemur (<i>Lepilemur jamesorum</i>)	12	Ecuadorian white-fronted capuchin (<i>Cebus aequatorialis</i>)	86
Indri (<i>Indri indri</i>)	15	Olalla Brothers' titi monkey (<i>Plecturocebus olallae</i>)	90
Aye-aye (<i>Daubentonia madagascariensis</i>)	19	Brown howler monkey (<i>Alouatta guariba</i>)	94
		Central American spider monkey (<i>Ateles geoffroyi</i>)	98
Africa	22		
Rondo dwarf galago (<i>Paragalago rondoensis</i>)	24	Other Species Considered	102
Roloway monkey (<i>Cercopithecus roloway</i>)	28	Grauer's gorilla (<i>Gorilla beringei graueri</i>)	102
Kipunji (<i>Rungwecebus kipunji</i>)	32	Crested macaque (<i>Macaca nigra</i>)	102
White-thighed colobus (<i>Colobus vellerosus</i>)	36	Bornean banded langur (<i>Presbytis chrysomelas</i>)	102
Niger Delta red colobus (<i>Ptilocolobus epieni</i>)	40	Tonkin snub-nosed monkey (<i>Rhinopithecus avunculus</i>)	103
Tana River red colobus (<i>Ptilocolobus rufomitratu</i>)	42	Siau Island tarsier (<i>Tarsius tumpara</i>)	103
Western chimpanzee (<i>Pan troglodytes verus</i>)	46		
Asia	50		
Javan slow loris (<i>Nycticebus javanicus</i>)	52		
Pig-tailed snub-nose langur (<i>Simias concolor</i>)	56		
Golden-headed langur or Cat Ba langur (<i>Trachypithecus poliocephalus</i>)	59		
Golden langur (<i>Trachypithecus geei</i>)	62		
Purple-faced langur (<i>Semnopithecus vetulus</i>)	66		
Gaoligong hoolock gibbon (<i>Hoolock tianxing</i>)	70		
Tapanuli orangutan (<i>Pongo tapanuliensis</i>)	73		

Appendix	104
References	106
Editors' addresses	120
Contributors' addresses	121

ACKNOWLEDGEMENTS

Here we present the 2018–2020 iteration of the World’s 25 Most Endangered Primates list, drawn up during an open meeting held during the XXVII Congress of the International Primatological Society (IPS), Nairobi, 22 August 2018.

We have updated the species profiles from the 2016–2018 edition (Schwitzer *et al.* 2017) for those species remaining on the list and added additional profiles for newly listed species.

This publication is a joint effort of the IUCN SSC Primate Specialist Group, the International Primatological Society, Global Wildlife Conservation, and the Bristol Zoological Society.

We are most grateful to the Margot Marsh Biodiversity Foundation for providing significant support for research and conservation efforts on these endangered primates through the direct provision of grants and through the Primate Action Fund, administered by Ms. Ella Outlaw, of the Chief Conservation Officer’s Office at Global Wildlife Conservation. Over the years, the foundation has provided support for the training workshops held before the biennial congresses of the International Primatological Society and helped primatologists to attend the meetings to discuss the composition of the list of the world’s 25 most endangered primates.

We would like to thank all of the authors who contributed to the final 2018–2020 version:

Laura V. Allen, Yves Rostant Andriamalala, Tsarafilamatra R. Andriamisedra, Herizo T. Andrianandrasana, Andie Ang, P. C. Bhattacharjee, Júlio César Bicca-Marques, Christophe Boesch, Giovanna Bonadonna, Andrew E. Bowkett, Gerson Buss, Thomas M. Butynski, Rodrigo S. Carvalho, Laura Cervera, Óscar M. Chaves, Dilip Chetry, Rekha Chetry, Susan M. Cheyne, Fanny Cornejo, Tim R. B. Davenport, Renata B. de Azevedo, Chiara De Gregorio, Yvonne de Jong, Stella de la Torre, Daniela W. De Luca, Andrea Dempsey, Giuseppe Donati, William Dreyer, Nicholas J. Dunn, Pengfei Fan, Izeni P. Farias, Hanlan Fei, Sally J. Fransen, Gabriella Fredriksson, Marco Gamba, Cristina Giacoma, Sery Gonedelé Bi, Marcelo Gordo, Harry Hilser, Zelinda B. Hirano, Tomas Hrbek, Iwan Hunowu, Cláudia A. Igayara-Souza, Rachel Ashgbofe Ikemeh, Leandro Jerusalinsky, Caspian L. Johnson, Daphne Kerhoas, Stanislaus M. Kivai, Juliet King, Inza Koné, Hjalmar Kühn, Wagner R. Lacerda, Diogo Lagroteria, Neahga Leonard, Richard Lewis, Edward E. Louis Jr., L. C. Loyola, Sophy J. Machaga, Jesus Martinez, Reiko Matsuda Goodwin, W. Scott McGraw, Erik Meijaard, Fabiano R. Melo, Sérgio L. Mendes, Pedro G. Méndez-Carvajal, Longondraza Miaretsoa, Gilberto Pozo Montuy, Noah E. Mpunga, Tilo Nadler, Andrianjaka Rijaniaina Jean Nary, K. Anna I. Nekarís, Leonardo G. Neves, Vincent Nijman, Stuart Nixon, Matthew G. Nowak, John F. Oates, Luciana I. Oklander, Lisa M. Paciulli, Anna P. Pannick, Richard J. Passaro, Alfons Patandung, Andrew Perkin, Colin Peterson, Gabriela Ponce, Márcio Port-Carvalho, Héctor Portillo-Reyes, Phan Duy Thuc, Wulan Pusparini, Martina Raffel, Brigitte M. Raharivololona, Rivo Rahasia, Jean-Baptiste Ramanamanjato, Doménico R. Randimbiharinarina, Rose Marie Randrianarison, Fidy Ralainasolo, Jonah Ratsimbazafy, Lucile Mialisoa Raveloarimalala, Lena M. Reibelt, Guopeng Ren, Melissa E. Rodríguez, Bonarge A. Rodríguez-Beitia, Fábio Röhe, Noel Rowe, Rasanayagam Rudran, Anthony B. Rylands, Agus Salim, Daniela Schrudde, Timothy M. Sefczek, Myron Shekelle, Alessandro Silva, María Fernanda Solórzano, Tenekwetché Sop, Roswitha Stenke, Jatna Supriatna, Nelson Ting, Valeria Torti, Mônica M. Valença-Montenegro, Daria Valente, Marcelo D. Vidal, Patrick O. Waeber, Robert Wallace, Edward D. Wiafe, Julie Wieczkowski, Serge A. Wich, Dominic Wormell, Lu Zhang.

References

Schwitzer, C., Mittermeier, R.A., Rylands, A.B., Chiozza, F., Williamson, E.A., Macfie, E.J., Wallis, J. and Cotton, A. (eds.). 2017. *Primates in Peril: The World’s 25 Most Endangered Primates 2016–2018*. IUCN SSC Primate Specialist Group (PSG), International Primatological Society (IPS), Conservation International (CI), and Bristol Zoological Society, Arlington, VA. 99 pp.

THE WORLD’S 25 MOST ENDANGERED PRIMATES: 2018–2020

Here we report on the tenth iteration of the biennial listing of a consensus of the 25 primate species considered to be among the most endangered worldwide and the most in need of conservation measures.

The 2018–2020 list of the world’s 25 most endangered primates has seven species from Africa, five from Madagascar, seven from Asia, and six from the Neotropics (Appendix: Table 1). Indonesia, Brazil, Ghana and Cote d’Ivoire each have three, Nigeria and Tanzania two, and China, Myanmar, India, Bhutan, Sri Lanka, Vietnam, Argentina, Ecuador, Peru, Mexico, Guatemala, Nicaragua, Honduras, El Salvador, Costa Rica, Panama, Bolivia, Guinea-Bissau, Liberia, Republic of Guinea, Senegal, Sierra Leone, Togo, Benin and Kenya each have one.


Twelve of the primates were not on the previous (2016–2018) list (Appendix: Table 3). Eight of them are listed as among the world’s most endangered primates for the first time. The Rondo dwarf galago, kipunji, Tana River red colobus and indri had already been on previous iterations, but were subsequently removed in favour of other highly threatened species. The 2018–2020 list contains two members each of the genera *Ptilocolobus* and *Trachypithecus*, thus particularly highlighting the severe threats that large primates are facing in all of the world’s primate habitat regions.

The changes made in this list compared to the previous iteration (2016–2018) were not because the situation of the twelve species that were dropped (Appendix: Table 2) has improved. In some cases, the situation has in fact worsened. By making these changes we intend rather to highlight other, closely related species enduring equally bleak prospects for their survival.






During the discussion of the 2018–2020 list at the XXVII Congress of IPS in Nairobi in 2018, a number of other highly threatened primate species were considered for inclusion (page 102). For all of these, the situation in the wild is as precarious as it is for those that finally made it on the list, thus they have been included as ‘*Other Species Considered*’, a new category in the Top 25 Most Endangered Primates’ series.




MADAGASCAR



Madagascar Primates

- | | |
|---|---|
|  <i>Daubentonia madagascariensis</i> |  <i>Indri indri</i> |
|  <i>Hapalemur alaotrensis</i> |  <i>Microcebus manitatra</i> |
|  <i>Lepilemur jamesorum</i> | |



BEMANASY MOUSE LEMUR

Microcebus manitatra Hotaling et al., 2016

Madagascar
(2018)

Giuseppe Donati, Andrianjaka Rijaniaina Jean Nary & Jean-Baptiste Ramanamanjato

The Bemanasy mouse lemur, *Microcebus manitatra*, was recently described from specimens collected in April 2007 in Bemanasy (also called Petite Lavasoa), one of the forest fragments in the Nouvelle Aire Protégée of Ambatotsirongorongo in the south-eastern corner of Madagascar (Hotaling et al. 2016). It was distinguished as a separate species to *M. murinus* using its nuclear and mtDNA, differences in which were diagnosed using species discovery delimitation criteria of mtDNA gene tree monophyly, distinct nuclear population structuring, and validation using coalescent-based Bayesian species delimitation tests (Hotaling et al. 2016). This recent description confirmed the findings of previous comprehensive surveys of mtDNA of *M. murinus* from across the south-eastern region, which revealed divergent geographic clades in close geographical proximity with no evidence of haplotype sharing (Weisrock et al. 2010; Hapke et al. 2012). This species takes its name from the Malagasy 'manitatra,' meaning 'range expansion,' to reflect the divergent distribution of this mouse lemur from the primarily western distribution of *M. murinus*.

Microcebus manitatra is a relatively large mouse lemur characterized by its size (total length of 270–276 mm and body mass of 58 g), long tail, dense, short fur (150 mm), relatively short hind feet (33 mm) and long ears (25–26 mm). The dorsal pelage is uniformly greyish brown on the back and tail and the underside is a greyish beige with dark grey underfur. This contrasts with *M. murinus* which has a variable greyish-brown to brownish-grey back and tail with a dull reddish-brown or cinnamon mid-dorsal stripe, as well as a mixed beige and grey underside (Hotaling et al. 2016).

The greatest threats to the Bemanasy mouse lemur are habitat loss and degradation driven by wood extraction, slash-and-burn cultivation, and fires.

The type locality, a forest called Bemanasy, is a 31.1 ha forest patch averaging 617 m a.s.l. located around 25 km south-west of the town of Tolagnaro (Fort Dauphin). It is one of a complex of three fragments (the other two are named Grand Lavasoa and Ambatotsirongorongo) today formally protected as Special Reserve of Ambatotsirongorongo (since 2015 according to the decree 2015-792). These fragments are remnants of one large forest block in the southern part of the Lavasoa Mountains that was positioned in a steep ecological gradient between spiny bush habitat to the west, the humid lowland forest to the north, and the littoral humid forest to the east (Donque 1972). This is reflected by its mixed floristic composition, which contains both humid forest plants and representatives of drier formations (Ramanamanjato et al. 2002; Andrianarimisa et al. 2009). Historical land cover maps indicate that these forests have not been directly connected to other forests for more than 40 years (Foiben-Taosarintanin'i Madagasikara 1979). The ecological complexity of the Lavasoa–Ambatotsirongorongo

mountains and the close geographical proximity of several mouse lemur species in the area (*M. griseorufus* in the west, *M. tanosi* in the north, and *M. ganzhorni* in the east) is remarkable. According to Hapke et al. (2012), this can be regarded as a micromodel of the retreat and expansion of various forest types and their resident mouse lemur species.

Neither population estimates or eco-ethological data are available for *M. manitatra*. Currently its distribution has only been confirmed in the 31.1 ha Bemanasy fragment, thus its population size is assumed to be very small. However, it is likely that

LAKE ALAOTRA GENTLE LEMUR

Haplemur alaotrensis Rumpler, 1975

Madagascar
(2000, 2014, 2016, 2018)

Lena M. Reibelt, Herizo T. Andrianandrasana, Fidy Ralainasolo, Lucile Mialisoa
Raveloarimalala, Richard Lewis, Jonah Ratsimbazafy & Patrick O. Waeber



the *M. cf. murinus* present in the other two fragments in the Special Reserve of Ambatotsirongorongo is also *manitatra*, which would result in a total area of occupancy of around 136 ha. MtDNA analyses indicate that the population of *M. cf. murinus* in Petriky, a 300-ha littoral forest 5 km east of Ambatotsirongorongo, clusters in the same clade of *M. manitatra* (Hapke *et al.* 2012). If confirmed by future analyses, this would increase the population size of this highly threatened mouse lemur species by several times.

The greatest threats to *M. manitatra* are habitat loss and degradation driven by wood extraction, slash-and-burn cultivation, and fires (Hapke *et al.* 2012). Although a recent report indicates that most fields near forest edges are abandoned, the fragments are surrounded by fields and cattle pastures, so slash-and-burn cultivation remains a threat (Ramanamanjato *et al.* 2002; Andrianjaka and Hapke 2015). The largest pressure currently seems to be wood extraction to meet the demand of construction timber (Andrianjaka and Hapke 2015). The forests in the Special Reserve of Ambatotsirongorongo also play a key role as sources of irrigation and drinking water (Andrianjaka and Hapke 2015).

Since 2017, the forest has been co-managed by local communities through an association of neighboring villages, called FIMPIA (Forest Police Association for Ambatotsirongorongo), and by the DREF (Direction Régionale de l'Environnement et des Forêt Anosy). Several aspects of forest management such as land use, sanctions for violations of these rules, and fees for visitors and researchers have now been implemented in the DINA (the community set regulations) as a result (Andrianjaka and Hapke 2015). The conservation value of this area has long been recognized and several conservation projects involving habitat restoration and local livelihood improvement have been conducted over recent years. However, the extremely small size of the remaining forest coupled with the increasing anthropogenic pressures from the town of Fort Dauphin call for further and immediate conservation actions. Because of its location, the Special Reserve of Ambatotsirongorongo contains an unusual and highly diverse mixture of lemurs, including one species of dwarf lemur previously listed (2014-16) in the World's 25 most endangered primates publication, *Cheirogaleus lavasoensis* (see Schwitzer *et al.* 2015). This makes these tiny forest fragments in the south-eastern corner of Madagascar a very high conservation priority.



The Critically Endangered Lake Alaotra gentle lemur, also known as the Lac Alaotra bamboo lemur (*Haplemur alaotrensis*), is the only primate taxon living constantly in a wetland. While all other bamboo lemurs are forest dwellers occupying a variety of forest habitats across Madagascar, *H. alaotrensis* is confined to the marshlands surrounding Lake Alaotra. It is this limited geographical range and the ever-increasing pressures on its shrinking habitat that have brought this lemur close to extinction.

This lemur is a cathemeral and a highly specialized herbivorous feeder. It targets 11 plant species and 16 distinct plant parts such as shoots, stem, pith, leaves, seeds, buds, and flowers (Mutschler 1999). The species is characterized by an average adult body length of about 30 cm (12 in), with a 30–40 cm (12–16 in) tail and a weight between 1 and 1.4 kg (2–3 lb) (Mutschler 1999). Its home range varies from 2 to 5 ha depending on group size (Nievergelt *et al.* 1998). Reibelt *et al.* (2017a) provide a more detailed account of the lemur's biology.

Its habitat, the marsh of Lake Alaotra, is located in the Alaotra-Mangoro region, where it covers about 20,000 ha and acts as a buffer zone between the lake (~20,000 ha of open water) and the agricultural zone (~120,000 ha of ricefields). Due to the high importance of the wetland for biodiversity and agro-economy, it was designated as a Ramsar site in 2003 (722,500 ha covering the whole watershed). Throughout the 2000s, marsh management was delegated from the state to environmental associations based in the villages adjoining the marshes, and in 2015 this mosaic formed the creation of the Alaotra Protected Area (46,432 ha encompassing the lake and the marshes) under the Regional Forestry Department.

Despite legal protection efforts, the lemur population has been declining continuously since the first published census in the 1990s. From the first estimate (7,500–11,000 individuals; Mutschler and Feistner 1995) it has shrunk to an estimated 2,500 individuals (Ralainasolo *et al.* 2006). Even the high priority conservation zone and tourist focal area, Park Bandro, recorded losses in recent years. While 170 individuals were estimated there in 2013 (Ratsimbazafy *et al.* 2013), only half of the original 85 ha park remained three years later, resulting in a reduced carrying capacity of 40–80 individuals (Raveloarimalala and Reibelt 2016). However, no recent sightings have been confirmed for this northern subpopulation. Subpopulations also exist in the southwestern marsh of Lake Alaotra, which is more protected due to its inaccessibility.

The principal threats to *H. alaotrensis* are habitat loss, habitat degradation, and hunting. While lemurs were regularly hunted for food in the 1990s, this is no longer the case due to long-term conservation public awareness work. Marsh burning to establish irrigated rice fields and to access fishing ponds is the main driver of habitat conversion (Copsey *et al.* 2009a, 2009b; Guillera-Arroita *et al.* 2010; Ralainasolo *et al.* 2006), which caused a nearly 30% loss of total marsh area between 2000 and 2016 (Andrianandrasana 2017). Furthermore, recent declines in rice harvests and fish catches have resulted in counter-season rice (*vary jeby*) production; grown in shallow lake water, or by converting the marshes at the lake-edge. More recently, the establishment of counter-season rice fields has been the main cause of suitable habitat loss in Park Bandro. Fires are also problematic for the lemurs, as marsh vegetation needs three years to regenerate to provide suitable habitat for the lemur again (Andrianandrasana 2002, 2009).

The conservation and management of the Alaotra Protected Area is challenging. Four permanent technical agents from the Ministry of Environment, Ecology and Forests are responsible for more than 50,000 ha of marshes. In close partnership with the authorities, Durrell Wildlife Conservation Trust (Durrell) and Madagascar Wildlife Conservation (MWC) have been collaborating for more than ten years on community-based efforts around the

lake, focusing on supporting the local resource-management associations, participatory ecological monitoring, environmental education, and eco-tourism (Andrianandrasana *et al.* 2005; Rendigs *et al.* 2015; Waeber *et al.* 2017a).

Durrell Wildlife Conservation Trust has recently installed more than 200 posts around core biodiversity zones to properly delineate the Protected Area, and local authorities approved a law prohibiting the utilization of the marshes inside this area. Durrell further supports the Regional Ministry of Fisheries in implementing a fishing ban, and works with the Regional Ministries of the Environment, Ecology and Forests, and the National Gendarmerie to ensure law enforcement, i.e. detect and follow-up on illegal activities in the Protected Area. In 26 villages around the lake, 96 CFL (marsh patrollers) monitor the Protected Area and report to Durrell for follow-up actions. Fires are monitored by sight and using the system 'Global Fire' to ensure timely detection and control of fires.

The total population of the Lake Alaotra gentle lemur has shrunk to an estimated 2,500 individuals.

MWC engages in research on human dimensions of lemur conservation for a better understanding of local livelihood strategies as well as values and perceptions to inform conservation actions (Reibelt *et al.* 2017b, 2017c; Stoudmann *et al.* 2017; Waeber *et al.*

2017b; Reibelt and Waeber 2018). MWC also develops alternative income opportunities such as compost from invasive species and ecotourism through Camp & Park Bandro. Environmental education activities include a comic book and poster (Maminirina *et al.* 2006; Richter *et al.* 2015), local restoration events in Park Bandro, and a new educational table-top game that explores ecosystem links and facilitates social learning (cf. Reibelt *et al.* 2018a, 2018b).

The annual Bandro festival (World Lemur Festival) is celebrated to value the lake and its biodiversity. It involves local authorities, MWC, Durrell and GERP (Groupe d'étude et de recherche sur les primates), and is accompanied by restoration events with the community. The festival is held in late October, when the marshes are dry, accessible and easy to burn, hence when protection activities are most critical. The Bandro Festival thus reminds everyone in the Alaotra region of its unique biodiversity and the need to protect it.

JAMES' SPORTIVE LEMUR

Lepilemur jamesorum (Louis et al., 2006)

Madagascar
(2016, 2018)

Tsarafilamatra R. Andriamisedra, Jonah Ratsimbazafy, William Dreyer,
Colin Peterson & Edward E. Louis, Jr.

James' sportive lemur (*Lepilemur jamesorum*) is a medium-sized nocturnal sportive lemur of Madagascar. It is found only in Manombo Special Reserve and Vevembe Classified Forest in the southeastern coastal region of the island, and is currently listed as Critically Endangered on the IUCN Red List (Andriaholinirina et al. 2014).

James' sportive lemur is similar to all other sportive lemurs in being folivorous (Ganzhorn 1993). However, it supplements its diet by feeding on gums. The species weighs approximately 780 g, with a total length of 56 cm, a head-body length of 26 cm, and a tail length of 30 cm. The pelage is short and smooth, primarily brown on the body and lightly grayish-brown on the belly and ventral portion of the extremities. The face is distinguished by the whitish-gray marking along the jaw and throat from the chin to the ears, forming a mask. The upper part of the head is brown with a black midline stripe that is continuous for almost the entire length of the body. The ears are large and cup-shaped, gray dorsally with black borders and a small cream-colored patch on the region beneath (Louis et al. 2006; Mittermeier et al. 2010). In general, the tail is uniformly brown, but several individuals have been noted to have a whitish tip.

Lepilemur jamesorum is primarily known from the Manombo Special Reserve, although it has been documented, through molecular genetic data, to exist in the Vevembe Classified Forest inland near Vondrozo. Although James' sportive lemur was first described in 2006 (Louis et al., 2006), there has been limited field research before or

since. Nocturnal surveys were carried out for the species in 2017 and 2018 by the authors. Low population densities were found, with only two individuals occupying an 800 ha survey plot in the special reserve. There were also few signs of their presence, such as tree holes and feeding traces. Sustained detection of the animal through censuses is planned to continue before research starts on behaviour and parasitism.

The primary threats to James' sportive lemur are habitat loss and hunting for bushmeat. Anthropogenic pressure is a significant problem in the Manombo area due to high levels of poverty,

limited job opportunities and inflation, which increase dependence of local communities on resources inside protected areas to survive.

James' sportive lemurs are hunted with traditional traps or simply by cutting into trees and taking them directly from their tree holes. Deforestation in the

Manombo Special Reserve is a significant problem not only for the sportive lemurs but also for the local communities which are gradually losing their timber, firewood and charcoal. This utilization of the forest also reduces natural food availability for James' sportive lemur and fragments their habitat. Fires started by humans destroy habitat and food resources. The distribution of James' sportive lemur in coastal littoral forests in southeastern Madagascar makes it especially vulnerable to stochastic events such as cyclones, e.g., cyclone Gretelle in 1997. This powerful cyclone in eastern Madagascar created difficulties for local people, and many animals died including lemurs. A census

James' sportive lemur is hunted with traditional traps or simply by cutting into trees and taking them from tree holes.



is required to quantify the impact on *L. jamesorum* as the population appeared to have declined.

The exact distribution and population numbers for James' sportive lemur are not known. Field surveys and research are required to ascertain population size and to establish baseline population parameters for the species.

Reforestation is important, not only to increase the habitat available and reconnect forest fragments, but also to provide resources for the local communities that they can harvest outside of the two protected areas. Complementary to this will be engaging local communities in alternative livelihoods to encourage sustainable practices and conservation in the Manombo region. In addition, reinforcing environmental education in the school system and local surrounding communities is needed to safeguard the future of James' sportive lemur.

INDRI

Indri indri Gmelin, 1788

Madagascar
(2012, 2018)

Valeria Torti, Longondraza Miaretsoa, Daria Valente, Chiara De Gregorio,
Giovanna Bonadonna, Rose Marie Randrianarison, Jonah Ratsimbazafy,
Marco Gamba & Cristina Giacoma



Variation in pelage colouration has resulted in two indri subspecies being identified (Groves 2001), but results from recent research (Brenneman *et al.* 2016) provide no support for this hypothesis. Molecular analysis of the mitochondrial sequences underline no significant genetic differentiation of the indri populations that would allow a two-subspecies classification system. The two major color forms are merely part of a clinal variation and not indicative of distinct taxa.

This highly distinctive lemur is endemic to the island of Madagascar where it inhabits the eastern rainforests from Anjanaharibe-Sud and Antohaka Lava (15 km SE of Andapa) in the north, south to Anosibe Anala Classified Forest. It has not been found on the Masoala Peninsula or in Marojejy (Mittermeier *et al.* 2008). It usually occurs at low elevations, but ranges up to 1,800 m (Goodman and Ganzhorn 2004).

Few studies on indri population densities are available. Betampona (2228 ha) has an estimated population of 77–147 indris (Glessner and Britt 2005). In Torotorofotsy (9900 ha) and Analamazaotra Special Reserve (700 ha), 21–32 indris have been reported (Junge *et al.* 2011). Population densities are presumably low, typically ranging from 5.2–22.9 per km² or 6.9–13.2 per km² (C. Golden pers. comm.). A recent survey in the Maromizaha New Protected Area (GERP 2017) suggested an indri population density of 82.32 individuals per km². A reasonable total population estimate, based on few studies (Glessner and Britt 2005; Junge *et al.* 2011; Brenneman *et al.* 2016; Nunziata *et al.* 2016; Bonadonna *et al.* 2017), would be 1,000–10,000 individuals. Population figures are in decline due to habitat destruction and hunting.

The indri is a mainly folivorous lemur (Powzyk and Mowry 2003), inhabiting tropical moist lowland



and montane forests. Their diet consists primarily of immature leaves supplemented by flowers, fruit, seeds and bark, which vary in proportion according to season (Randrianarison *et al.* 2018). They occasionally descend to the ground to eat earth, perhaps to detoxify seeds that have also been eaten (Powzyk 1997; Britt *et al.* 2002; Powzyk and Thalmann 2003; Randrianarison *et al.* 2016).

The indri lives in socially monogamous family groups, composed of one adult pair and their offspring (Pollock 1979). Group size is reported to vary between two to six individuals (Torti *et al.* 2013). The limited number of adult individuals in a group suggests that intrasexual dominance is age-related (Pollock 1977). Females give birth every two years. Reproduction is highly seasonal, with the birth of a single offspring in May or June. Reproductive maturity is reached between seven and nine years of age (Pollock 1977), but individuals may disperse earlier (V. Torti pers. obs). Both males and females disperse (C. Giacoma pers. obs; Torti *et al.* 2018) and the sex ratio at birth is approximately 1:1 (Kappeler 1997). Groups in fragmented landscapes tend to be larger than those in more extensive, undisturbed areas (Pollock 1979; Powzyk 1997; Bonadonna *et al.* 2017). Indri groups occupy relatively small territories, the sizes of which vary according to the study and the site: 17.6 to 25.9 ha in Analamazaotra Reserve (Bonadonna *et al.* 2017), 34 to 40 ha in Mantadia (Powzyk 1997), 27 ha in Betampona (Glessner and Britt 2005) and 5 to 17.47 ha in Maromizaha New Protected Area (Bonadonna *et al.* 2017). Indri movements in the territory are evenly distributed. Mean daily path lengths are 243 m in Maromizaha NPA (Torti, unpublished data) and 350 m per day in Andasibe-Mantadia NP (J. A. Powzyk unpubl. data).

Group encounters are rare; Bonadonna and colleagues (2017) observed only 30 encounters of 16 different indri groups in 36 months. However, when two groups meet, they defend their territory ownership by using long vocal interactions or territorial song (Torti *et al.* 2013), and eventually starting physical fights (Pollock 1979; Powzyk and Mowry 2006). After a dispute resolution, the adult

males reunite with their females and resume their normal ranging and feeding activities.

The indri has a rich vocal repertoire (Sorrentino *et al.* 2013) and is the only lemur that communicates through songs. Their songs are long sequences of vocal units that are organized in phrases (Thalmann *et al.* 1993; Gamba *et al.* 2011). They have the form of a chorus in which all the adults and subadults of a group utter their contribution in a precise and coordinated manner (Gamba *et al.* 2016a). Songs have various functions depending on the context in which they are emitted and they are used for both inter- and intra-group communication (Torti *et al.* 2013, 2018). Furthermore, songs are likely to provide information about the group composition and mediate the formation of new groups (Giacoma *et al.* 2010; Gamba *et al.* 2016b; Torti *et al.* 2017).

The principal threat to the indri is habitat destruction for slash-and-burn agriculture, logging and fuelwood gathering, all of which also take place in protected areas.

The principal threat to the indri is habitat destruction for slash-and-burn agriculture, logging and fuelwood gathering.

Illegal hunting is a major problem for the indri in certain areas (Jenkins *et al.* 2011). Although long thought to be protected by local *fadys* (traditional taboos), these do not appear to be universal and

the animals are now hunted even in places where such tribal taboos exist, with observed increases in hunting since the political crisis. In many areas, these taboos are breaking down with cultural erosion and immigration, and local people often find ways to circumvent taboos even if they are still in place. For example, a person for whom eating the indri is forbidden may still hunt the animals to sell to others, while those who may be forbidden to kill indris can purchase them for food. In 2018, in the Commune of Lakato (Alaotra Mangoro Region), nine indris were killed by poachers in the Antavolobe forest (J. Ratsimbazafy pers. obs.). Recent studies of villages in the Makira Forest indicate that indris have been hunted for their skins (which are worn as clothing), and show that indri meat is prized, fetching a premium price, and that current levels of indri hunting are unsustainable (Golden 2009; Jenkins *et al.* 2011; R. Dolch pers. comm.). The indri is listed on Appendix I of CITES.

It occurs in three national parks (Mananara-Nord, Andasibe-Mantadia and Zahamena), two nature reserves (Betampona and Zahamena) and five special reserves (Analamazaotra, Mangerivola, Ambatovaky, Anjanaharibe-Sud, and Marotandrano) (Mittermeier *et al.* 2008). It is also present in several of the 27 new protected areas declared in 2015 (e.g., Anosibe An'Ala, Anjozorobe-Angavo, Makira, Maromizaha). The corridors between Mantadia and Zahamena are an important Conservation Site, where widespread conservation education and capacity building should be implemented to eliminate hunting, with the indri as the flagship species. This species has never been successfully kept in captivity and thus the success of a captive breeding program is difficult to predict. In the next years it will be critical to support local forest management by improving the existing community-based approach (Randrianarison *et al.* 2015). Actions should include expansion of protected habitats to increase population connectivity (e.g., the Ankeniheny-Zahamena corridor) and to decrease lemur disturbance by rural communities.

AYE-AYE

Daubentonia madagascariensis Gmelin, 1788

Madagascar
(2016, 2018)

Doménico R. Randimbiharirina, Timothy M. Sefczek, Brigitte M. Raharivololona, Yves Rostant Andriamalala, Jonah Ratsimbazafy & Edward E. Louis, Jr.



The aye-aye, *Daubentonia madagascariensis*, is the only surviving representative of the Daubentoniidae, the most anciently diverged family of primates known to date (Martin 1990; Simons 1995; Catlett *et al.* 2010; Perry *et al.* 2012). Aye-ayes have the widest distribution of any extant lemur, ranging from Montagne d'Ambre in the north of Madagascar, to Parc National d'Andohahela in the south, and Parc National Tsingy de Bemaraha in the west (Ganzhorn and Rabesoa 1986; Simons 1993; Schmid and Smolker 1998; Rahajanirina and Dollar 2004). Aye-ayes were also introduced to Nosy Mangabe, an island located off the Masoala Peninsula in northeast Madagascar (Petter 1977; Sterling 1993).

The aye-aye is the largest nocturnal primate, with a body mass of 2.5–2.6 kg and a total length of 74–90 cm, of which 30–37 cm is head-body length and 44–53 cm is tail length (Oxnard 1981; Glander 1994; Feistner and Sterling 1995). Aye-ayes have several unusual, derived traits including an elongated, thin, highly-flexible middle finger with a metacarpophalangeal ball and socket joint, continuously-growing fused canines and incisors, the greatest encephalization quotient of any strepsirrhine or nocturnal primate, a relatively long gestation period for lemurs, and a slow life history, including late weaning and a protracted learning period (Owen 1863; Jouffroy 1975; Martin 1990; Simons 1995; Barrickman and Lin 2010; Catlett *et al.* 2010).

Aye-ayes live in several forest types, from primary rainforest to dry undergrowth forest, and occupy habitats of varying qualities, including disturbed forests and heavily degraded forests near plantations (Pollock *et al.* 1985; Ganzhorn and Rabesoa 1986; Ancrenaz *et al.* 1994; Andriamasimanana 1994). Their distribution across various habitats is partially due to their




dietary adaptability. Though aye-ayes primarily eat insect larvae and *Canarium* seeds, they may expand their diet to include nectar from *Ravenala madagascariensis*, coconut, banana, mango, litchi, breadfruit, sugar cane and cankers (Petter and Petter 1967; Petter 1977; Iwano and Iwakawa 1988; Sterling 1993; Andriamasimanana 1994; Simons and Meyers 2001; Randimbiharirina *et al.* 2018).

Despite the aye-aye's distribution and dietary flexibility, their huge individual home ranges and long interbirth intervals may translate to low population densities (Perry *et al.* 2012). As aye-ayes are solitary and can have large home ranges upwards of 973.12 ha, reliable population estimates remain elusive (Randimbiharirina *et al.* 2018). Most locality records are based on feeding traces. However, at any given site, traces may be made by one aye-aye or multiple individuals (Aylward *et al.* 2018).

The greatest threats to aye-ayes are habitat destruction (forest degradation, fragmentation and slash and burn agriculture) and persecution by some local populations who believe aye-ayes to be an evil omen (Petter and Peyrieras 1970; Simons and Meyers 2001). Although there are multiple scientific and popular articles focusing on the aye-aye, Randimbiharirina and colleagues' (2018) research from the Kianjavato Classified Forest currently represents the only long-term field study conducted on naturally-occurring individuals. Therefore, further long-term studies are needed to understand the aye-aye's ecology across its distribution before accurate population estimates and conservation initiatives can be achieved. Furthermore, novel studies utilizing molecular genomic platforms, such as in Aylward *et al.* (2018), should be further explored as a means to provide accurate population estimates. The reclusive nature of the aye-aye provides a cautionary facet of this species' status, since its extirpation from forest habitat will only be documented long after it has been unknowingly eliminated.



AFRICA



African Primates

- | | | |
|--|--|---|
| ■ <i>Cercopithecus roloway</i> | ■ <i>Pan troglodytes verus</i> | ■ <i>Rungwecebus kipunji</i> |
| ■ <i>Colobus vellerosus</i> | ■ <i>Piliocolobus epieni</i> | ■ <i>Piliocolobus rufomitratu</i> |
| ■ <i>Paragalago rondoensis</i> | | |



RONDO DWARF GALAGO

Paragalago rondoensis Honess in Kingdon, 1997

Tanzania
(2006, 2008, 2010, 2012, 2014, 2018)

Andrew Perkin

Weighing approximately 60 g, the Rondo dwarf galago, *Paragalago rondoensis*, is the smallest of all galago species (Perkin and Honess 2013). It is distinct from other dwarf galagos in its diminutive size, a bottle-brush-shaped tail, its reproductive anatomy, and its distinctive "double unit rolling call" (Perkin and Honess 2013). Current knowledge indicates that *P. rondoensis* is endemic to the coastal forests of Tanzania. There are three spatially distinct sub-populations. One is in southeast Tanzania near the coastal towns of Lindi and Mtwara. The second is approximately 400 km north, in pockets of forest around the capital city of Dar es Salaam. The third sub-population is in Sadaani National Park, approximately 100 km north of Dar es Salaam. However, there is emerging data (vocal and penile morphological) that suggests the northern and southern populations may be phylogenetically distinct.

Rondo dwarf galagos have a mixed diet of insects and fruit. They often feed close to the ground, and move by vertical clinging and leaping in the shrubby understory.

They build daytime sleeping nests, usually in the canopy (Bearder *et al.* 2003). As with many small primates, *P. rondoensis* is probably subject to predation from owls and other nocturnal predators, such as genets, palm civets and snakes. The presence of these predators invokes intense episodes of alarm calling (Perkin and Honess 2013).

Across its known range, the Rondo galago can be found sympatrically with a number of other galagos, including two much larger species in the genus *Otolemur*: Garnett's galago *O. garnettii* (Least Concern, Butynski *et al.* 2008a), the thick-tailed galago, *O. crassicaudatus* (Least Concern, Bearder 2008). In the northern parts of its range (for example, in Zaraninge forest, Pugu/Kazimzumbwi

Forest Reserve (FR) and Pande Game Reserve (GR), the Rondo galago is sympatric with the Zanzibar galago, *P. zanzibaricus* (Least Concern, Butynski *et al.* 2008b) and in the southern parts of its range (for example, in Rondo, Litipo and Noto), it is sympatric with Grant's galago, *P. granti* (Least Concern, Honess *et al.* 2008).

P. rondoensis, classed as Critically Endangered, (Perkin *et al.* 2008), has an extremely limited and fragmented range in a number of remnant patches of Eastern African Coastal Dry Forest (Burgess and Clarke 2000) in Tanzania. These are at Zaraninge forest (06°08'S, 38°38'E) in Sadaani National Park (Perkin 2000), Pande GR (06°42'S, 39°05'E), Pugu/Kazimzumbwi (06°54'S, 39°05'E) (Perkin 2003, 2004), Rondo Nature Reserve (NR) (10°08'S, 39°12'E), Litipo (10°02'S, 39°29'E) and Ziwani (10°20'S, 40°18'E) FRs (Honess 1996b; Honess and Bearder 1996). New sub-populations were identified in 2007 near Lindi town in Chitoo FR (09°57'S, 39°27'E) and Ruawa FR (09°44'S, 39°33'E), and in 2011 in

Noto Village FR (09°53'S, 39°25'E) (Perkin *et al.* 2011, 2013) and in the northern population at Ruvu South FR (06°58'S, 38°52'E). Specimens of *P. rondoensis*, originally described as *Galagoides demidoffi phasma*, were collected by Ionides from the Rondo Plateau, SE Tanzania in 1955, and by Lumsden from Nambunga, near Kitangari, (approximately 10°40'S, 39°25'E) on the Makonde Plateau, Newala District in 1953. There are doubts as to the persistence of the species on the Makonde Plateau, which has been extensively cleared for agriculture. Surveys there in 1992 failed to detect any extant populations (Honess 1996). Distribution surveys have been conducted in the southern (Honess 1996; Perkin *et al.* in prep.) and northern coastal forests of Tanzania (29 surveyed)

The major threat facing the Rondo dwarf galago is loss of habitat.



and Kenya (seven surveyed) (Perkin 2000, 2003, 2004; Perkin *et al.* 2013). Absolute population sizes remain undetermined, but recent surveys have provided density estimates of: 3–6/ha at Pande GR (Perkin 2003) and 8/ha at Pugu FR (Perkin 2004). Relative abundance has also been estimated from encounter rates: 3–10/hr at Pande GR and Pugu/Kazimzumbwi FR (Perkin 2003, 2004), and 3.94/hr at Rondo FR (Honest 1996b). There is a clear and urgent need for further surveys to determine population sizes in these dwindling forest patches.

The major threat facing the Rondo dwarf galago is loss of habitat. All sites are subject to some level of agricultural encroachment, charcoal manufacture and/or logging. In 2008, the known area of *P. rondoensis* occurrence was 101.6 km^2, but new data on forest area change indicates this figure has fallen to 87.4 km^2 . In Pande GR (2.4 km^2), Chitoo FR (5 km^2) and Rondo FR (25 km^2), forest cover remained the same between 2008 and 2014. However, forest cover between 2008 and 2014 fell in Zaraninge forest from 20 km^2 to 15 km^2 , in Pugu/Kazimzumbwi FR from 33.5 km^2 to 8 km^2 , in Ruawa FR and Litipo FR from 4 km^2 to 3 km^2 , and in Ziwani FR from 7.7 km^2 to 1 km^2 . Two newly discovered areas of occupancy are Ruvu S, in which forest cover fell from 20 km^2 to 5 km^2 , and Noto, in which forest cover fell from 21 to 20 km^2 , in the same time period (Burgess and Clarke 2000; Doggart 2003; Perkin *et al.* in prep).

As habitat availability decreases, the population trend must also be assumed to be declining, the rate varying according to the level of protection of each forest fragment. All sites, except Pande GR, Zaraninge (in Saadani National Park) and Rondo NR forest, are national or local authority forest reserves and as such nominally, but in practice minimally, protected. Since 2008, protection of two forests has increased: the Noto plateau forest, formerly open village land, is part of a newly created village forest reserve, and the Rondo Forest Reserve has now been declared a new nature reserve. Both are important for Rondo dwarf galago conservation given their relatively large size. Given current trends in charcoal production for nearby Dar es Salaam, the forest reserves of Pugu and Kazimzumbwi were predicted to disappear over the next 10–15 years (Ahrends 2005). Recorded forest loss in Pugu/Kazimzumbwi and Ruvu South has been attributed to the rampant charcoal trade. Pande, as a Game Reserve, is relatively secure, and Zaraninge forest,

being in a national park, is the most protected part of the range of *G. rondoensis*. In the south, the Noto, Chitoo and Rondo populations are the most secure, as they are buffered by tracts of woodland. The type population at Rondo is buffered by woodland and *Pinus* plantations managed by the Rondo Forestry Project, and is now a nature reserve. Litipo and Ruawa FRs are under threat from bordering village lands. Ziwani is now mostly degraded scrub forest, thicket and grassland.

The following conservation actions are recommended to safeguard the future of this species: 1) continued monitoring of habitat loss rates, 2) surveying new areas for remnant populations, 3) implementation of community-based conservation and awareness, 4) assessment of population status and phylogenetic relationships between the sub-populations and confirmation of suspected phylogenetic distinctions. Until such time that the latter has been carried out, each subpopulation must be considered to be of high conservation value.

ROLOWAY MONKEY

Cercopithecus roloway Schreber, 1774

Ghana, Côte d'Ivoire
(2002, 2006, 2008, 2010, 2012, 2014, 2016, 2018)

Sery Gonedelé Bi, Andrea Dempsey, Inza Koné, W. Scott McGraw & John F. Oates

Cercopithecus roloway and its close relative *Cercopithecus diana* are highly attractive, arboreal monkeys that inhabit the Upper Guinean forests of West Africa. The roloway monkey, which once occurred in many of the southern forests of Ghana and central and eastern Côte d'Ivoire, is distinguished from the Diana monkey by its broad white brow line, long white beard and yellow thighs. Because individuals with intermediate coat patterns are known from near the Sassandra River in Côte d'Ivoire, some scientists treat the roloway and Diana as subspecies of one species, *C. diana* (for example, Oates 2011). Of the two forms, the roloway is the more seriously threatened, and it is now rated as Critically Endangered on the IUCN Red List (Koné et al. 2019).

Roloway monkeys are upper-canopy specialists that prefer undisturbed forest. Destruction and degradation of their habitat and relentless hunting for the bushmeat trade have reduced their population to small, isolated pockets. Miss Waldron's red colobus (*Piliocolobus waldroni*) once inhabited many of the same forests as the roloway, but is now almost certainly extinct (Oates 2011). Unless much more effective conservation action is taken very quickly, there is a strong possibility that the roloway monkey will also disappear in the near future.


Over the last 50 years, roloway monkeys have been steadily extirpated in Ghana. In southwestern Ghana, once a stronghold of *C. roloway*, an ornithological study showed a 600% increase in both legal and illegal logging between 1995 and 2008 (Arcilla et al. 2015). Illegal logging, which makes up 80% of timber harvested in Ghana,

is particularly devastating; because it is wholly unregulated or monitored, there are no limits on number, size or species of trees taken. One third of illegal logging is by companies that take more than their quota, expand into protected areas, and/or continue to log after their permit has expired. The remaining two thirds are rogue illegal chainsaw operators (Arcilla et al. 2015). Additional factors causing the roloway monkey's decline are clearing for agriculture, charcoal production and bushmeat hunting. Hunting has very likely been the major cause in the recent crash in roloway populations; bushmeat is a major food source for Ghanaians, with an estimated 80% of the rural population dependent on it as their main source of protein (Dempsey 2014; Trench 2000).

The captive population of the roloway monkey is now so small that extinction in captivity is also a strong possibility.

Several recent surveys have failed to confirm the presence of roloway monkeys in any reserves in western Ghana, including the Ankasa Conservation Area, Bia National Park, Krokosua Hills Forest Reserve, Subri River Forest Reserve and Dadieso Forest Reserve

(Oates 2006; Gatti 2010; Buzzard and Parker 2012; Wiafe 2013). Community-owned forests along the Tano River (referred to as the "Kwabre Community Rain Forest") in the far southwestern corner of the country are the only localities in Ghana where any roloways have been recorded by scientists or conservationists in the last decade. Kwabre consists of patches of swamp forest along the lower Tano River, adjacent to the Tanoé forest in Côte d'Ivoire. Surveys of these forests have been conducted under the auspices of the West African Primate Conservation Action organization since 2011, and several sightings of roloway groups have been made, along with mona monkeys,



spot-nosed monkeys, white-naped mangabeys and olive colobus (WAPCA 2014; Dempsey 2014; Osei *et al.* 2015). WAPCA has supported a community-based conservation project with villages around these forests, establishing a Kwabre Community Resource Management Area, which works to protect the forest through the sustainable management of natural resources. Meanwhile, further efforts should be made to ascertain whether any roloway monkeys still survive in Ankasa, because this site has significant conservation potential and roloways have been reported there in the relatively recent past, as well as in the Amazuri Wetlands Area.

In neighbouring Côte d'Ivoire, the roloway monkey's status is dire as well. Less than twelve years ago, roloways were known or strongly suspected to exist in three forests: the Yaya Forest Reserve, the Tanoé forest adjacent to the Ehy Lagoon, and Parc National des Îles Ehotilé (McGraw 1998, 2005; Koné and Akpatou 2005; Gonedelé Bi *et al.* 2013). Surveys of eighteen areas between 2004 and 2008 (Gonedelé Bi *et al.* 2008, 2012) confirmed the presence of roloways only in the Tanoé forest, suggesting that the roloway monkey may have been eliminated from at least two forest areas (Parc National des Îles Ehotilé, Yaya Forest Reserve) in the last dozen years. Subsequent surveys carried out in southern Côte d'Ivoire suggest a handful of roloways may still survive in two forest reserves along the country's coast. In June 2011, Gonedelé Bi observed one roloway individual in the Dassioko Sud Forest Reserve (Gonedelé Bi *et al.* 2014, in review; Bitty *et al.* 2013). In 2012, Gonedelé Bi and A. E. Bitty observed roloways in Port Gauthier Forest Reserve, and in October 2013, Gonedelé Bi obtained photographs of monkeys poached inside this reserve, including an image purported to be a roloway. The beard on this individual appears short for a roloway, raising the possibility that surviving individuals in this portion of the interfluvial region may in fact be hybrids. In any case, no sightings of roloways have been made in the Dassioko Sud or Port Gauthier Forest Reserves since 2012, including during the most recent patrols (February 2017). These reserves are described as coastal evergreen forests, and both are heavily degraded due to a large influx of farmers and hunters from the northern portion of the country (Bitty *et al.* 2013). Gonedelé Bi and colleagues, in cooperation with SODEFOR (Société de Développement des

Forêts) and local communities, have organized regular surveys aimed at removing illegal farmers and hunters from both reserves. However, surveys made in August 2015 revealed that a logging company (SIDB) had begun clearing a portion of the Port Gauthier reserve. Efforts are underway to work with SODEFOR to stop logging and other illegal activities in these reserves (Gonedelé Bi 2015).

Thus, the only forest in Côte d'Ivoire where roloways are confirmed to exist is the Tanoé forest adjacent to the Ehy Lagoon, and immediately across the Tano River from the Kwabre forest in Ghana. This wet forest also harbours one of the few remaining populations of white-naped mangabeys in Côte d'Ivoire. Efforts led by I. Koné and involving several organizations (WAPCA, ACB-CI, Mulhouse Zoo) helped stop a large palm oil company from causing further habitat degradation, and a community-based conservation effort has helped slow poaching in this forest (Koné 2015). Unfortunately, hunting still occurs in Tanoé, and the primate populations there are undoubtedly decreasing (Gonedelé Bi *et al.* 2013).

As the potential last refuge for roloways and one of the last refuges for white-naped mangabeys, the protection of the Tanoé Forest in Côte d'Ivoire and the adjacent Kwabre Forest in Ghana should be the highest conservation priority. By any measure, the roloway monkey must be considered one of the most Critically Endangered monkeys in Africa and is evidently on the verge of extinction (Oates 2011). In addition, the captive population is now also so small that extinction in captivity is a strong possibility (Lefaux and Montjardet 2016).

KIPUNJI

Rungwecebus kipunji Ehardt, Butynski, Jones and Davenport in Jones *et al.* 2005

Tanzania
(2006, 2008, 2018)

Tim R. B. Davenport, Noah E. Mpunga, Sophy J. Machaga & Daniela W. De Luca

The discovery of the kipunji (*Rungwecebus kipunji*), a monkey endemic to southern Tanzania, on the forested flanks of Mt Rungwe in 2003, (Davenport 2005), demonstrated how much there was still to learn about Africa's primate fauna. Originally thought to be a mangabey (Jones *et al.* 2005), it was later placed in a new monotypic genus *Rungwecebus* by Davenport *et al.* (2006) on the basis of molecular and morphological data, making it the first new genus of African monkey to be described in 83 years. Subsequent research supported this phylogenetic position and taxonomic status (Olson *et al.* 2008; Roberts *et al.* 2009).

The kipunji is one of the world's most threatened primates. This was first demonstrated by a census that provided the initial systematically-derived data on the animal's abundance and distribution (Davenport *et al.* 2008). Whilst there is growing evidence that a decade of conservation has ameliorated the situation (Davenport and Marques 2018), the kipunji remains Critically Endangered, with the species and genus facing a high risk of extinction in the wild (Davenport & Jones 2008).

The kipunji is known from only two populations, separated by approximately 350 km of non-forested, agricultural land. One population is at 1,750–2,450 m in 12.4 km² of Rungwe-Kitulo Forest (Davenport *et al.* 2008) in Tanzania's Southern Highlands (09.12°–09.18°S, 33.67°–33.92°E). The Rungwe-Kitulo Forest includes the Mt Rungwe Nature Reserve (150 km²) and the Livingstone Forest (191 km², found in the 412 km² Kitulo National Park) (Davenport 2002; Davenport and Bytebier 2004; Davenport *et al.* 2005, 2006, 2008). Mt Rungwe and Livingstone Forest (in Kitulo)

are connected by the Bujingijila Corridor, a 2-km-wide degraded forest connection (Davenport 2005, 2006; Mwakilema and Davenport 2005) that is currently being re-wilded. The kipunji inhabits the wetter forest of southern Mt Rungwe, and isolated groups are scattered in the north and south of Livingstone Forest (Davenport *et al.* 2006, 2008; Bracebridge *et al.* 2011), with five groups in Madehani forest a few kilometres further south.

The other population of *R. kipunji* is located at 1,300–1,750 m in the Vikongwa Valley, Ndundulu Forest (07.67°–07.85°S, 35.17°–36.83°E; ca. 180 km² of closed forest), in the Kilombero Nature Reserve of the Udzungwa Mountains (Jones *et al.* 2005). The area occupied by the species in Ndundulu is estimated to be approximately 25 km² (Davenport pers. obs. in Marshall *et al.* 2015) although its density is low. The kipunji has not been recorded from the contiguous Udzungwa Mountains National Park (Davenport *et al.* 2008; De Luca *et al.* 2010; Marshall *et al.* 2015). The species' extent of occurrence (EOO) is estimated to be 42.4 km² in Rungwe-Kitulo (Bracebridge *et al.* 2011) and ~25 km² in Ndundulu (Davenport pers. obs. in Marshall *et al.* 2015), with the combined total EOO (species range) at just 67.5 km².

The kipunji is sparsely distributed (Davenport *et al.* 2006, 2008; Bracebridge *et al.* 2001, 2012). The small extent of occurrence in Rungwe-Kitulo (42.4 km²) and even smaller estimate for Ndundulu (~25 km²) give grounds for conservation concern. The total extent of occurrence is much less than the 100 km² required to meet the threshold for listing as Critically Endangered under criterion B of the IUCN Red List.

Previous surveys estimated the total population of kipunji to be 1,117 animals.



Surveys totalling 2,864 hours and covering 3,456 km of transect were undertaken by Davenport *et al.* (2008) to determine distribution and group numbers. In addition, 772 hours of simultaneous multi-group observational follows in Rungwe-Kitulo and Ndundulu forests enabled 209 total counts to be carried out. Davenport *et al.* (2008) estimated some 1,042 individuals in Rungwe-Kitulo, ranging from 25 to 39 individuals per group ($\mu = 30.65$; SE = 0.62; n = 34), and 75 individuals in Ndundulu, ranging from 15 to 25 individuals per group ($\mu = 18.75$; SE = 2.39; n = 4). The total kipunji population was thus estimated to be 1,117 animals in 38 groups ($\mu = 29.39$; SE = 0.85; n = 38). The Ndundulu population is restricted and the Rungwe-Kitulo population is fragmented with isolated subpopulations remaining in degraded habitat (Davenport *et al.* 2008, 2010). Kipunji densities appeared to be three times greater in the secondary forests of Rungwe-Kitulo than in the primary forests of the Udzungwas (Davenport *et al.* 2008), possibly because competition with other diurnal primates is not present in Rungwe-Livingstone (Bracebridge *et al.* 2011). There was also a statistically significant difference in mean group size between the Rungwe-Kitulo and the Ndundulu populations (Davenport *et al.* 2008), possibly related to the small total population in Ndundulu. A new census by the Wildlife Conservation Society (WCS) is now underway (Davenport and Markes 2018).

Rungwecebus kipunji occurs entirely within protected areas (PAs); Kitulo National Park, Mt Rungwe Nature Reserve, Kilombero Nature Reserve (Davenport *et al.* 2008) as well as Nkuka and Madehani forests managed under lease by WCS. Of the state-run PAs, however, only Kitulo National Park has ongoing management activities and these remain limited in the forest. Neither Mt Rungwe nor Kilombero Nature Reserves have financial resources, and the management of Mt Rungwe has been dependent on funds from WCS since 2008. Forest loss and degradation has been much reduced over the last decade, especially compared with the situation in 2003 when unmanaged resource extraction was commonplace (Machaga *et al.* 2004; Davenport 2005, 2006). However, challenges remain in some of the more remote areas.

The Bujingijila corridor linking Mt Rungwe Nature Reserve to Livingstone Forest in Kitulo National

Park and the areas joining the northern and southern sections of Livingstone are encroached (Davenport 2005; Mwakilema and Davenport 2005). However, a decade of protected area and community conservation work by WCS has meant that solid recovery is underway (Davenport and Markes 2018). Despite this, some subpopulations remain fragmented even within the same protected area. Kipunji are killed by log traps, mainly as retribution for the raiding of maize, beans and potatoes, and mostly from January to April. In Ndundulu Forest the species is present in low numbers (Davenport *et al.* 2008) but the reason for this remains unclear (Marshall *et al.* 2015).

The focus of current kipunji conservation work is the protection and restoration of the montane habitats of Mt Rungwe, especially the forest connections such as the Bujingijila corridor (Davenport 2006; Bracebridge *et al.* 2011). Protecting connections is a high priority for the conservation of this genus/species and restoring Bujingijila could provide habitat for an additional 88 kipunji (8% population increase), using density estimates from the 2008 census (Bracebridge *et al.* 2013). This would also reconnect the Mt Rungwe and Kitulo subpopulations.

The kipunji is being used as a 'flagship species' by WCS's long-term Southern Highlands Conservation Program in and around Rungwe-Kitulo, especially in education and awareness raising activities, and as part of a long-term monitoring program. A section of forest contiguous with Mt Rungwe that contains groups of kipunji is now being leased to and managed by WCS. A habituated group is monitored daily for research, especially on aspects of the kipunji's social and reproductive behaviour, feeding ecology, home range dynamics, predation and demography. A new group is now being habituated for tourism. The focus of applied kipunji conservation work is the protection and restoration of its montane forest habitats, widespread environmental education, and support to both management authorities and local communities across its range.



WHITE-THIGHED COLOBUS

Colobus vellerosus I. Geoffroy Saint-Hilaire, 1830

Côte d'Ivoire, Ghana, Togo, Benin, possibly Nigeria
(2016, 2018)

Reiko Matsuda Goodwin, Sery Gonedelé Bi, Edward D. Wiawe & John F. Oates

The white-thighed colobus (*Colobus vellerosus*) is one of five *Colobus* species that inhabit sub-Saharan Africa. As the name suggests, it has white-silvery thighs and a white ruff around the face. It is a middle-to-upper canopy species, but it sometimes goes down to the forest floor to cross tree gaps, play, and drink water. This colobine feeds mostly on the leaves of trees and lianas (Wong *et al.* 2006). It lives in lowland rainforests, swamp forests, seasonally inundated semi-deciduous forests, savannah-woodlands and gallery forests from the coastal areas to 10° 6' N (Oates 2011). The fact that it occurs in wet, dry, and moderately disturbed habitats and that groups occupy small home ranges (Wong and Sicotte 2007) attests to its ecological flexibility.

This colobus monkey has an extremely fragmented distribution from the area between the Sassandra-Bandama rivers in Côte d'Ivoire to Benin, traversing Ghana and Togo. It is not clear if it still occurs in western Nigeria. The species' broad range may give a false impression that it is not as threatened as some primates with more restricted ranges. In each range country, however, *C. vellerosus* occurs in only a limited number of protected areas and community forests.

Habitat destruction and degradation have significantly reduced *C. vellerosus* populations, but the primary cause of their decline in all range countries is uncontrolled hunting (Oates 2011). Large primates, like the other six colobine species included in this volume, have been particularly decimated by illegal hunting in Africa and Asia.

Our current estimate of the population size of *C. vellerosus* at sites from which there are some data is only about 1,200 (Matsuda Goodwin *et al.* unpublished). Although this number excludes several groups in some forest areas where surveys have not yet been conducted or have been incompletely performed, we estimate that the total population is now less than 1,500. Apart from the Boabeng-Fiema Monkey Sanctuary (BFMS) in Ghana, which has a population of about 370 white-thighed colobus (Kankam *et al.* 2010), the average number of individuals known or inferred to be present at all other sites is 23, and these sites are typically separated from each other by about 60 km. This means that there are considerable impediments to gene flow between the populations and a high risk of inbreeding depression.

The encounter rate in seven protected areas has declined by 87.2% in the last four decades.

In Côte d'Ivoire, the white-thighed colobus used to be commonly observed in many protected areas, but in the Bandama-Sassandra interfluvial zone, Grébouo 1 sacred forest is the only place where this species still occurs (Gonedelé Bi *et al.* 2014). East of this zone, this species is known to occur at only three sites: Tanoé-Ehy swamp forest, Comoé National Park, and Dinaoudi sacred forest (McGraw *et al.* 1998; Gonedelé Bi *et al.* 2014). The successive periods of civil unrest in Côte d'Ivoire between 1999 and 2012 exacerbated the threats to *C. vellerosus*. Many protected areas suffered from weakened protection and heavy hunting, some were converted into plantations, and the settlement of people inside protected areas was widespread (Campbell *et al.* 2008; Bitty *et al.* 2015).



Dinaoudi and Grébouo 1 sacred forests have only two groups of *C. vellerosus* each (S. Gonedélé Bi pers. obs.). A survey conducted in the Tanoé-Ehy swamp forest in 2007 had an encounter rate of 0.05 groups/km and the observed group size was extremely small (mean = 2) (Gonedélé Bi *et al.* 2010). In the Comoé National Park, camera images and *ad libitum* sightings by J. Lapuente (pers. comm.) in the last four years indicate that several groups still occur there.

Three Critically Endangered taxa are found in the Tanoé-Ehy swamp forest—the rolaway monkey and the white-naped mangabey, besides the white-thighed colobus—and the Comoé Monkey Project is now taking an initiative to survey and conserve these species there, alongside local non-governmental organizations (NGOs) from both Côte d'Ivoire and Ghana (RASAP-CI, CEPA, WAPCA). While logging (to create an oil palm plantation) and hunting continue, these NGOs have been engaging with local communities to promote conservation activities (Gonedélé Bi *et al.* 2013; McGraw *et al.* 2017).

The white-thighed colobus used to be widely distributed in Ghana (Grubb 1998). However, it has been extirpated from at least 14 of 26 sites where surveys were conducted between 1976 and 2018. It is still present in two national parks (Kakum, Mole), several forest reserves (Ayum, Bia Tributaries North, Bonsam Bepo, Bonkoni, Cape Three Points, Subin, Mpameso, Krokosua Hills, Atewa Range), Bia Resource Reserve (RR), and at BFMS (Oates 2006; Burton 2010; Gatti 2010; Wiafe 2013, 2016, 2018; Osei *et al.* 2015; Akom 2015). The species also occurs in the Kwabre swamp forest, where a transboundary project linked to the Tanoé-Ehy project in Côte d'Ivoire that focusses on community-based conservation for the rolaway monkey has been in place since 2014 (Osei *et al.* 2015; McGraw *et al.* 2017; WAPCA 2018).

Only one to a few groups remain at each of these sites, however, and hunting and logging continue in most of them. The species may still occur in the Digya and Kyabobo national parks, the Yoyo River Forest Reserve, the Kalakpa Resource Reserve, and the Amanzule swamp forest, but it is very unlikely that large populations survive at any of these sites given ongoing high levels of hunting and forest destruction (Owusu-Ansah 2010). Atewa Range Forest Reserve, which is also home to the white-

naped mangabey, still harbours the white-thighed colobus, but hunting, snaring, mining, logging and farming threaten their sustained existence there (Kusimi 2015; Wiafe 2018). A campaign against a joint China-Ghana bauxite mining at Atewa Range here is ongoing (Environmental Justice Atlas 2018). The only site in Ghana where there is a stable *C. vellerosus* population is the BFMS, but deforestation and habitat degradation have been increasing there (E. Wiafe pers. obs.). At BFMS, group size varies from 9 to 38 with a mean of 13 in forest fragments and 15 in the larger forest (Wong and Sicotte 2007). Past survey data from seven protected areas indicate that the encounter rate of this species has declined on average by 87.2% in the last four decades, suggesting an equivalent population decline (Matsuda Goodwin *et al.* unpublished).

The white-thighed colobus has probably been extirpated from Burkina Faso (Ginn and Nekaris 2014). In Togo, recent surveys found a few groups of this species at each of Togodo Faunal Reserve, Fazao-Malfakassa National Park and Yikpa Community Forest (Segniagbeto *et al.* 2017, 2018). *Colobus vellerosus* is on the verge of extinction in Benin; only 1–2 groups live in the Lama Forest Reserve and Kikélé sacred forest, and the species appears to have been extirpated from the forest reserves of Pénésoulou, Mt. Kouffé, Wari-Marou, Ouémé Supérieur, and the Lokoli swamp forest where it used to occur (Matsuda Goodwin *et al.* 2016). The chance of this species still being present at Bonou swamp forest and Ouémé Boukou Forest Reserve in Benin is slim. In Nigeria, there is little credible information that this species still occurs in the Old Oyo and Kianji Lake national parks, where it was said to occur more than 30 years ago (Happold 1987).

Urgent measures are required to stop further local extirpations of the white-thighed colobus. We urge range countries to enact tough laws against the hunting of threatened species, similar to the Wildlife Conservation and Management Law (2013) in Kenya, and to strictly enforce the laws against hunting. Moreover, systematic surveys should be conducted in Mole and Digya national parks, and the Atewa Forest Reserve in Ghana, and the Old Oyo and Kainji Lake national parks in Nigeria. We also encourage the European Association of Zoos and Aquaria to consider establishing a captive-breeding programme for

this species at Zoo Duisburg (Germany) where such a programme for the closely-related king colobus (*Colobus polykomos*) already exists, or at Zoo Barcelona (Spain) where the white-naped mangabey is housed.

NIGER DELTA RED COLOBUS

Piliocolobus epieni Grubb and Powell, 1999

Nigeria
(2008, 2010, 2016, 2018)

Rachel Ashegbofe Ikemeh & John F. Oates



The Niger Delta red colobus (*Piliocolobus epieni*) is endemic to the marsh forests in the central part of the Niger Delta of Nigeria (Oates 2011). Its species name is derived from its name in the Ijaw language of the people who inhabit the limited area of about 1,500 km² in Bayelsa State where it occurs. *Piliocolobus epieni* only became known to science in 1993 in the course of a biodiversity survey coordinated by C. Bruce Powell (Powell 1994). Studies of vocalizations and mitochondrial DNA suggest that this population is not closely related to its closest geographic relatives, the Bioko red colobus (*Piliocolobus pennantii*) or Preuss's red colobus (*Piliocolobus preussi*) of eastern Nigeria and western Cameroon, leading Ting (2008) to treat this monkey not as a subspecies of *P. pennantii* (see Groves 2001; Grubb *et al.* 2003) but as a distinct species, *Procolobus epieni*. Groves (2007) regarded almost all the different red colobus monkeys, including *epieni*, *pennantii* and *preussi*, as separate species in the genus *Piliocolobus* – a taxonomy that we follow here. Since 2008, the IUCN Red List of Threatened Species has regarded *P. epieni* as Critically Endangered (Oates and Struhsaker 2016).

The marsh forests where the Niger Delta red colobus is found have a high water table all year round, but do not suffer deep flooding or tidal effects. The most intensive ecological study of this monkey, by Lodjewijk Werre (1994–1996), suggested that the clumped distribution of food species in the marsh forest is a key factor restricting *P. epieni* to its limited range, which is demarcated by the Forcados River and Bomadi Creek in the northwest, the Sagbama, Osiana and Apoi creeks in the east, and the mangrove belt to the south (Werre 2009). At the time of its discovery in the mid 1990s, this red colobus was locally common, especially in forests

near the town of Gbanraun, but it was beginning to come under intense pressure from degradation of its habitat and commercial hunting. Important colobus food trees, especially *Fleroya ledermannii*, were being felled at a high rate by artisanal loggers, and the logs were floated out of the Delta on rafts to processing centers in Lagos and elsewhere. In addition, large canals dug as part of oil extraction activities, as well as smaller canals dug by loggers into the interior swamps, were changing local hydrology (Werre and Powell 1997; Grubb and Powell 1999). The Ijaw people are traditionally fishermen, but outside influences introduced by the oil industry have encouraged commercial bushmeat hunting and logging throughout the Niger Delta. The most recent range-wide assessment of *P. epieni* conducted in 2013 suggests that, as a result of habitat destruction and hunting, the population has declined significantly since the 1990s, and that it may now be around 90% lower than the previous estimate of <10,000 (Ikemeh 2015).

In the 2013 survey, the presence of *P. epieni* was confirmed only in four forest areas, and it was considered extirpated from 11 other forests where it had been reported in the 1990s by Werre (2009). Cumulative survey data indicate that the current number of individuals surviving in the wild may be only a few hundred (Ikemeh 2015). The two most important remaining areas for *P. epieni* conservation are thought to be the Apoi Creek Forests, flanked by the communities of Gbaraun, Apoi and Kokologbene, and forests near Kolotoro. Insecurity in the region and the consequences of corrupt governance, amongst other factors, have exacerbated the major threats of habitat degradation and commercial hunting. Because red colobus monkeys are known to be sensitive to habitat disturbance and hunting in other parts of Africa (Struhsaker 2005), it is feared that the Niger Delta red colobus, with its restricted range, is at risk of extinction.

The red colobus monkeys are probably more threatened than any other taxonomic group of primates in Africa (Oates 1996; Struhsaker 2005). *Piliocolobus badius temminckii* (Senegal to Guinea or Sierra Leone) (Galat-Luong *et al.* 2016), *Piliocolobus badius badius* (Sierra Leone to western

Côte d'Ivoire) (Oates *et al.* 2016a), *Piliocolobus preussi* (western Cameroon and eastern Nigeria) (Oates *et al.* 2016b), and *Piliocolobus pennantii* (Bioko Island, Equatorial Guinea) (Oates and Struhsaker 2008) are all now Critically Endangered from different combinations of habitat loss and hunting, while *Piliocolobus waldroni* (eastern Côte d'Ivoire and western Ghana) may already be extinct (Oates *et al.* 2019).

At present, no areas in the Niger Delta are formally protected for wildlife, even though the region has great ecological significance and supports many rare, unique and/or threatened taxa (Ikemeh 2015). The Niger Delta red colobus shares its marsh forest habitat with two other threatened primates; the Nigerian white-throated guenon (*Cercopithecus erythrogaster pococki*) and the red-capped mangabey (*Cercocebus torquatus*) (Ikemeh 2015). Also found in these forests are the putty-nosed monkey (*Cercopithecus nictitans*), the mona monkey (*Cercopithecus mona*) and the olive colobus (*Procolobus verus*) (Efenakpo *et al.* 2018). However, political instability in the Delta, related in the most part to disputes over the allocation of oil revenues, has prevented any progress in biodiversity conservation during the last decade (Ikemeh 2015).

It is feared that the Niger Delta red colobus, with its restricted range, is at risk of extinction.

Because security in the Niger Delta continues to be challenging, undertaking effective conservation actions remains difficult. Despite these challenges, and with the urgent need to save this species from extinction, a locally driven research, awareness-raising and conservation initiative is currently underway in Bayelsa State, coordinated by the SW/ Niger Delta Forest project (directed by R. Ikemeh). Plans exist to intensify this effort, with concerted focus on ensuring that *P. epieni* is protected by law in Bayelsa State and that one or more protected areas are established for the fragile remaining populations. Two areas in particular have already been suggested for special protected status to conserve red colobus monkeys: the Otolo-Kolotoro-Ongoloba area and the Apoi Creek area (Ikemeh 2015).

Note: To date, photographs of this species have been difficult to obtain. Thus, it is represented here only by the illustration.



TANA RIVER RED COLOBUS

*Piliocolobus rufomitratu*s Peters, 1879

Kenya
(2002, 2004, 2006, 2008, 2012, 2014, 2018)

Stanislaus Kivai, Laura Loyola, Julie Wieczkowski, Juliet King, Yvonne de Jong,
Laura Allen, Nelson Ting & Tom Butynski

The taxonomy of the Tana River red colobus *Piliocolobus rufomitratu*s has been disputed (Grubb *et al.* 2013; Oates and Ting 2015); it has been considered a species (Groves 2007; Zinner *et al.* 2013; Groves and Ting 2013, Butynski and Hamerlynck 2015) and a subspecies *P.r.rufomitratu*s (Napier 1985; Grubb *et al.* 2003; Ting 2008; Struhsaker and Grubb 2013). The classification used here follows the taxonomy adopted at the IUCN SSC Primate Specialist Group African Primate Red List Assessment Workshop in Rome in April 2016. The species is listed as Endangered on the IUCN Red List of Threatened Species based on its extremely limited and fragmented geographic range, continued habitat fragmentation, and the unsustainable exploitation of forest products (Butynski *et al.* in press). The Tana River red colobus is listed as Class B in the African Convention on the Conservation of Nature and Natural Resources.

The Tana River red colobus, as well as the Endangered Tana River mangabey *Cercocebus galeritu*s, are endemic to the riparian forests of the lower Tana River and Tana Delta of the north coast of Kenya. They are broadly sympatric with the Vulnerable *Cercopithecus mitis albotorquatu*s, as well as with *Paragalago cocos*, and *Otolemur garnettii lasiotis*, and narrowly sympatric on the forest edge with *Papio cynocephalus ibleanus*, *Chlorocebus pygerythrus hilgerti*, and *Galago senegalensis braccatus* (De Jong and Butynski 2009, 2012; Butynski and De Jong 2019). The gallery forests are part of the East African Coastal Forests Biodiversity Hotspot. The evergreen riparian forest patches are narrow, often restricted to less than 500 m from

the river or water channels. They provide habitat in this, otherwise, expansive arid and semi-arid landscape (Baker *et al.* 2015).

The Tana River red colobus occupies about 34 riverine and flood-plain forests that range in size (1-500 ha) along a 60-km stretch of the lower Tana River and in the upper Tana Delta (Butynski and Mwangi 1995; Mbora and Meikle 2004; Hamerlynck *et al.* 2012). The area of occupancy is extremely small (<13 km²; Butynski and Mwangi 1994; Hamerlynck *et al.* 2012; Butynski and Hamerlynck 2015). Fewer than 1000 individuals remain (Mbora and Butynski 2009). Primate surveys in 2000 found two groups in Onkolde forest (Tana Delta) but a survey in December 2017 recorded only one male and one female (Nature Kenya, unpubl. data).

Population densities of the diurnal, arboreal, and folivorous Tana River red colobus range from 33 to 253 individuals/km², varying according to habitat quality (Mbora and Meikle 2004;

Loyola 2015). As all remaining forests inhabited by the Tana River red colobus are small, fragmented, degraded, and, therefore, seriously threatened, the long-term survival of this species seems bleak.

With the Tana River red colobus population in a continual state of decline (Marsh 1978, 1986; Butynski and Mwangi 1995), mean group sizes have fallen by about 50% since the 1970s (Marsh 1978; Karere *et al.* 2004). Many factors account for these declines, most (if not all) of which are associated with the rapidly increasing human population, including poverty, insecurity, poor

Continuing deforestation, forest fragmentation and invasive plants challenge the survival of the Tana River red colobus.



conservation leadership among local communities and weak governance by government authorities. Seventy-five percent of the population lives below the poverty line (Baker *et al.* 2015). This requires rigorous community awareness and outreach on the importance of human population control strategies, initiation of sustainable livelihood alternatives, and adoption of agricultural technologies that improve food production (Mбора and Allen 2011).

The construction of hydroelectric power dams along the upper Tana River, and several big irrigation projects, have altered the water table, river flow volume, and flood cycle, leading to drastic vegetation changes in the lower catchment (Butynski 1995; Maingi and Marsh 2002). The situation is likely to worsen with the upcoming High Grand Falls Dam, which is expected to be completed in 2031 (Hamerlynck *et al.* 2012; Mwangasha 2018). This will be the second largest dam in Africa. It will be accompanied by large-scale irrigation schemes, and water transfer to the Lamu Port and to the Southern Sudan-Ethiopia Transport Corridor (LAPSSET). Additionally, the High Grand Falls Dam will have negative impacts on the floods and groundwater recharge required for the maintenance of the lower Tana River and Tana Delta forests. It is expected to have a negative impact on flood-dependent livelihoods (crop farmers, pastoralists, and fishermen) and the biodiversity of this biologically-sensitive region (Njiru 2011; Duvail *et al.* 2012; Hamerlynck *et al.* 2012).

Continuing deforestation, habitat degradation, fragmentation, and spread of invasive plant species, further threaten the Tana River red colobus, as do agricultural encroachment and unsustainable forest exploitation (e.g., building materials, palm wine, medicinal plants, wood for canoe-making, firewood collection) (Butynski and Mwangi 1995; Mбора and Meikle 2004; Moinde-Fockler *et al.* 2007; De Jong & Butynski 2009; Duvail *et al.* 2012; Hamerlynck *et al.* 2012; Butynski & De Jong 2019). The invasive mesquite *Propopis juliflora*, facilitated by forest clearing by people and elephants *Loxodonta africana*, is steadily spreading into indigenous forests and inhibiting regeneration of native tree species. Stringent habitat protection, restoration of abandoned farmlands and degraded forests, and research on the control of the invasive species are required.

Proposed and failed irrigation schemes continue to threaten Tana River red colobus habitat. The Tana Delta Irrigation Project (TDIP), a failed rice growing development [financed by the Japan International Cooperation Agency (JICA) and managed by the Tana and Athi Rivers Development Authority (TARDA)] led to the loss of some of the most important forests for the Tana River red colobus and Tana River mangabey (Butynski and Mwangi 1994; Moinde-Fockler *et al.* 2007). Similarly, the Bura Maize and Cotton Irrigation Scheme cleared 350 km² and diverted river water through furrows, but failed (Horta 1994; Christensen *et al.* 2012). A proposed 10,000 km² irrigation scheme in the lower Tana River has also been initiated by the Kenyan government, which will lead to further habitat loss.

The entire Tana River red colobus population lies in a politically insecure area and much unplanned, unregulated, and unsustainable exploitation of natural resources. Conservation efforts, as well as monitoring, are impeded by insecurity and regional conflicts (Duvail *et al.* 2012). Resettlement of pastoralist communities from the Tana Delta and Tana River Primate National Reserve (TRPNR; Baker *et al.* 2015), by Kenya Wildlife Service (KWS) led to unrest in the region. A court case to degazette the only protected area in the geographic range of the Tana River red colobus (and Tana River mangabey), the TRPNR (171 km²) was won by the local community. In January 2007, the High Court of Kenya ordered the annulment of the Reserve, citing lack of proper involvement of the local people during its gazette. In 2012, the Tana Delta became a Ramsar Site, otherwise, the Tana Delta has no protected area or protection status. To secure the long-term survival of the Tana River red colobus, conservation initiatives that aim to enable local people to conserve biodiversity, improve livelihoods, rehabilitate degraded areas, and establish sustainable income generating projects, including ecotourism, are required (Butynski & De Jong 2019).

Despite the many challenges of the lower Tana River and Tana Delta, there is some good news. Various unsustainable commercial developments have aborted their plans, and conservation initiatives have started in the region, including the Ndera Community Conservancy (NCC) (Northern Rangeland Trust; Mбора and Allen 2011; Butynski and De Jong 2019). In addition,

plans to form the Ngwano Community Conservancy are on-going and the Tana River County Government is supporting various community conservation efforts.

WESTERN CHIMPANZEE

Pan troglodytes verus Schwarz, 1934

Côte d'Ivoire, Ghana, Guinea-Bissau, Liberia, Mali, Republic of Guinea,
Senegal, Sierra Leone
(2018)

Tenekwetché Sop, Christophe Boesch & Hjalmar Kühl

The western chimpanzee (*Pan troglodytes verus*) is one of the four currently recognised subspecies of chimpanzees. Since 2016, it has been classified as Critically Endangered on the IUCN Red List of Threatened Species (Humble *et al.* 2016).

Historically, the western chimpanzee ranged widely from eastern Senegal to Benin, across 11 countries. Currently, it only occurs in eight countries. It is believed to have disappeared from Benin, Togo, and Burkina Faso (Ginn *et al.* 2013; Campbell and Hounbedji 2015), and is close to extinction in Ghana. It is possible that remnants of its historic geographic range extend into western Nigeria, but genetic evidence is still needed as confirmation (Humble *et al.* 2016).

Twenty years ago, knowledge about the population status of the western chimpanzee was limited (Kormos and Boesch 2003), but extensive large-scale surveys since have increased knowledge (e.g., Brncic *et al.* 2010; WCF 2012; Tweh *et al.* 2014). The most comprehensive estimation suggests a wild population of 35,000–55,000 chimpanzees (Kühl *et al.* 2017; Heinicke *et al.* 2019). Guinea alone has been estimated to host more than 20,000 western chimpanzees, with the largest remaining population – more than 17,000 individuals – found in the Fouta Djallon region (WCF 2012). The other strongholds for this taxon are Liberia with 7,000 individuals (Tweh *et al.* 2014), and Sierra Leone with 5,000 individuals (Brncic *et al.* 2010). Five to six hundred are estimated for Senegal (Wessling and Pruetz, pers. comm.) while in Guinea-Bissau a population of 1,000–1,500 individuals was recently reported in the Boe area (A. Goedmakers pers. comm. 2017). Côte d'Ivoire was once a stronghold

for this subspecies, with an estimated population of 8,000–12,000 individuals in the 1990s (Marchesi *et al.* 1995). The number of chimpanzees in Côte d'Ivoire has declined by 80% since 1990 (Kühl *et al.* 2017).

Between 1990 and 2014, the western chimpanzee's geographic range declined by 20% from 657,600 to 524,100 km² (Kühl *et al.* 2017). At the same time, populations declined annually by 6.5%, with a total population reduction of 80%. This period corresponds to a single chimpanzee generation (23 years) (Langergraber *et al.* 2012). If this decline continues, 99% of the remaining population will be lost by 2060. This drastic population decline and

range reduction provided the quantitative basis for uplisting the western chimpanzee to Critically Endangered (Humble *et al.* 2016).

The western chimpanzee is thought to have been lost from Benin, Togo and Burkina Faso, and is close to extinction in Ghana.

The major causes of population decline are illegal hunting, habitat loss (Kormos *et al.* 2003; Carvalho *et al.* 2012; Tweh *et al.* 2014; Brncic *et al.* 2015; Humble *et al.* 2016) and infectious disease (Köndgen *et al.* 2008). Throughout its range,

the western chimpanzee is losing its natural rainforest and savanna woodland habitat. Even though hunting chimpanzees is illegal, it is widespread, mainly for food, but also in retaliation for crop raiding, and occasionally, infants are captured and sold to wildlife traffickers (e.g., Pruetz and Kante 2010). Large spatial variation, however, exists in the importance and magnitude of threats and their underlying drivers across the subspecies' range.

Several range countries of the western chimpanzee (Liberia, Sierra Leone, and Côte d'Ivoire) have suffered in past decades from civil unrest and acute political



crises that have undermined their capacity to protect their biological resources. This has had direct and indirect impacts on forest resources and wildlife protection, including the occupation of protected areas. In Marahoué National Park in Côte d'Ivoire, for example, the chimpanzee population disappeared within a few years only because of human immigration, and the Mount Peko and Mount Sangbe National Park were occupied by migrants in the 2000s. The influx of humans resulted in unsustainable exploitation of natural resources and the illegal circulation of weapons, exacerbated by the absence of law enforcement in protected areas. Indeed, the dramatic decline of western chimpanzees is a consequence of the combined effect of increasing levels of threats and lack of political, financial and conservation commitments (Kühl *et al.* 2017). In Côte d'Ivoire, where up to 12,000 chimpanzees were estimated to occur in the 1990s, a 90% decline was recorded from 1990 - 2007. This was mostly driven by increasing human population density and the associated pressures of poaching, massive deforestation in classified forests, immigration from the Sahelian belt, and political instability in the country (Campbell *et al.* 2008; Kühl *et al.* 2017). As a consequence, chimpanzees became locally extinct in several protected areas including Marahoué National Park and several classified forests, including Nizoro, Dassikro, Haute Dodo, Niouniourou, Okromodou, Port Gauthier, Monogaga, Mount Kope, Tamin and likely Mount Peko National Park (based on data in the IUCN SSC A.P.E.S. database).

Nowadays, the majority of chimpanzees that remain in Cote d'Ivoire are found in Tai National Park (Tiedoué *et al.* 2018), likely because of the regular anti-poaching patrols, long-term research activities, eco-tourism and awareness-raising activities conducted there by OIPR (Office Ivoirien des Parcs et Reserves), as well as by national and international NGOs over many decades.

Liberia holds the largest remaining rainforest population of western chimpanzees (Tweh *et al.* 2014), but this is threatened by bushmeat hunting that also occurs in protected areas (Greengrass 2016) and rapidly developing mining, forestry and agro-industrial sectors (Junker *et al.* 2015). The 5,000 chimpanzees of Sierra Leone (Brncic *et al.* 2010) are mostly threatened by mining expansion, industrial agriculture and hunting for bushmeat and in retaliation to crop foraging.

In Mali, Senegal and Guinea-Bissau, at the northern limit of the western chimpanzee range, they are

particularly susceptible to habitat loss due to mining concessions and unregulated artisanal mining. In Senegal, for example, nearly the entirety of the chimpanzee range has been licensed for gold mining exploration and exploitation. This chimpanzee population is also facing immediate threats of habitat loss as it is vulnerable to aridity and environmental change (Kühl *et al.* 2017). The chimpanzees of Guinea-Bissau occur predominantly in the Boé area which has important Bauxite reserves, and future mining projects will considerably impact this fragile population.

Heinicke *et al.* (2019) report that chimpanzees persist in higher densities under three social-ecological conditions: rainforest habitats with low human impact, rough terrain that is difficult to access, and regions where cultural taboos such as hunting prohibitions prevail and there is a low human footprint. In Guinea, where the largest remaining and most stable population is found, western chimpanzees live predominantly in the unprotected area of the Fouta Djallon, which meets the third category (Boesch *et al.* 2017). The majority of local people belong to the Fulani ethnic group, who neither hunt nor eat chimpanzees for cultural reasons and are mostly pastoralists or practitioners of small-scale traditional agriculture (Ham 1998). Although currently hosting a stable chimpanzee population, Fouta Djallon is rich in bauxite. Anticipated large-scale mining in the future is expected to cause further population decline and threaten the main stronghold of *Pan t. verus*. Furthermore, several large infrastructure projects planned by the Guinean authorities across the country (e.g., dams, roads, and railway construction) are threats to chimpanzees in Guinea at large.

Chimpanzees are protected by national and international laws throughout their range, but enforcement of laws is almost nonexistent. Seventy percent of the current population occurs outside protected areas (IUCN SSC A.P.E.S. database 2016). With exponential human population growth in range countries, global demand for natural and mineral resources, and poor law enforcement in and outside protected areas in most range countries, it is very likely that the western chimpanzee will continue to decline in the future unless conservation efforts are made to effectively protect the remaining populations.

Following the uplisting of *P. t. verus* to Critically Endangered on the IUCN Red List of Threatened Species, a multi-stakeholder meeting was held in Monrovia (December 2017) to discuss necessary


conservation measures. A new Regional Action Plan for *P. t. verus*, currently in preparation, has identified 9 conservation strategies. These include improvement of regional law enforcement coordination, improved legal frameworks and landuse planning, capacity for protected area management, disease surveillance and conservation finance mechanisms (IUCN, unpublished).

A crucial first step would be to include conservation priorities into development and land use planning (LUP) across sectors. This requires increased commitment by the conservation community and relevant political bodies (ministries for resources and environmental affairs), to conduct cross-disciplinary planning of infrastructure, development, and resource extraction projects that affect chimpanzee habitat. This will also require increased funding of conservation initiatives on the ground, and additional engagement from political authorities to improve the management of ape habitat, including existing protected areas. Protected areas are important in conserving biodiversity (Lovejoy 2006), hence it is also essential that range countries "expand [their] protected area system to at least 17% of land surface (...) by 2020", as agreed upon in the Aichi Biodiversity Targets (CBD 2010), so that a greater number of chimpanzees are protected. There are some promising initiatives in Liberia, with the creation of the new Grebo-Krahn National Park in the chimpanzee range, and in Guinea, where the Moyen Bafing National Park is in the process of being created. If finalized, this new national park would protect ca. 5,000 chimpanzees of the Fouta Djallon. In June 2017, the Government of Guinea Bissau officially created the Boé and Dulombi national parks in the south-east of the country, which if properly governed, should protect several chimpanzees in the area.

Apart from immediate actions, long-term, sustainable protection of the region's biodiversity will likely depend on interventions which address critical problems specific to range countries of the region, such as poverty, regional immigration, poor governance, unsustainable agricultural practices and lack of land-use planning. The Fouta Djallon landscape, the main stronghold of the western chimpanzee, deserves special attention. It is particularly important to understand how chimpanzees and humans co-exist in agricultural landscapes, how factors contributing to this coexistence can be maintained over time, and how they can eventually be transferred to other regions inhabited by great apes.



ASIA



Asian Primates

- | | |
|---|---|
|  <i>Pongo tapanuliensis</i> |  <i>Trachypithecus poliocephalus</i> |
|  <i>Trachypithecus geei</i> |  <i>Semnopithecus vetulus</i> |
|  <i>Nycticebus javanicus</i> |  <i>Simias concolor</i> |
|  <i>Hoolock tianxing</i> | |



JAVAN SLOW LORIS

Nycticebus javanicus É. Geoffroy Saint-Hilaire, 1812

Indonesia
(2008, 2010, 2012, 2014, 2016, 2018)

K. Anna I. Nekaris & Vincent Nijman

Devastating habitat loss throughout southeast Asia threatens all nine species of slow loris with extinction, i.e. greater *Nycticebus coucang*, pygmy *N. pygmaeus*, Bengal *N. bengalensis*, Philippine *N. menagensis*, Bornean *N. borneanus*, Kayan *N. kayan*, Sody's *N. bancanus*, Sumatran *N. hilleri*, and Javan *N. javanicus* (Munds *et al.* 2013; Pozzi *et al.* 2014; Rowe and Myers 2016). Slow lorises exhibit numerous unique traits including slow life history, locomotion and digestion, the ability to enter torpor and hibernate, and being the only venomous primates (Nekaris 2014). Still, wild slow lorises have seldom been studied for more than a year (Malaysia *N. coucang*, Wiens *et al.* 2006; Cambodia *N. pygmaeus*, Starr *et al.* 2011; India, *N. bengalensis*, Das *et al.* 2014), with only *N. javanicus* being the focus of a long-term study (Rode-Margono *et al.* 2014). Many researchers and conservationists have only ever seen a slow loris in the illegal wildlife trade, either dried on bamboo sticks in preparation for traditional medicine, paraded as a photo prop on a tourist beach, or sold as a pet (Schulze and Groves 2004; Das *et al.* 2009; Nijman *et al.* 2015; Osterberg and Nekaris 2015). The extreme popularity of viral slow loris internet videos is a double-edged sword, to some extent making the public aware of their decline, but also causing the public to perceive that they are not threatened (Nekaris *et al.* 2013a). The extent of trade raised international concern, resulting in the transfer of the genus *Nycticebus* to CITES Appendix I in 2007 (Nekaris and Nijman 2007).

Javan slow lorises are now listed by IUCN as Critically Endangered, thus here we use the Critically Endangered Javan slow loris as the flagship for slow loris conservation (Nekaris *et al.*

2013b). Since being re-recognised as a species by the IUCN in 2006, work on the Javan slow loris has increased and provides a sound example of understanding and mitigating the threats to a highly threatened species. Both morphologically and genetically distinct, the Javan slow loris weighs about 1 kg, and exhibits a facial mask comprised of bold fork marks leading from the eyes and ears to the crown of the head, revealing a white diamond pattern on the forehead (Nekaris and Jaffe 2007).

Capturing individuals to meet the demand for pets is the most severe threat to the survival of Javan slow lorises. Despite being legally protected in Indonesia since 1973, with their striking coloration and presence on Java, Indonesia's commercial centre, it is no wonder that Indonesian pet traders in the 1990s targeted Javan slow lorises above other endemic slow loris species. Since 2012, the number of Javan slow lorises openly traded in markets has decreased, with a stark rise in numbers of greater slow lorises from Sumatra, a species with a

threat status that must also be carefully monitored. Indeed, over three years of market surveys on Java between 2012 and 2015, four times more greater slow lorises than Javan slow lorises were counted, with traders claiming that Javan slow lorises are increasingly difficult to obtain (Nijman *et al.* 2015). In November 2013, nearly 300 greater slow lorises were confiscated in two raids. Following the smaller of these raids, 31 out of the 76 slow lorises confiscated died in the next few weeks. Death rates of the larger raid are unknown. Successful prosecution of lawbreakers buying or selling slow lorises in Indonesia is a very rare occurrence, so much so that we are not aware of a single slow loris trader having been sentenced in the last

To avoid being bitten by venomous slow lorises, traders habitually cut or pull out an animal's lower front teeth.



decade, despite hundreds of slow lorises having been confiscated from traders. Wildlife traders in Indonesia have increasingly turned to social media to advertise their illegal stock, including Javan slow lorises. The huge rise in Facebook and WhatsApp means that many are sold via social media without ever being seen in a wildlife market. An ongoing online monitoring programme by the Little Fireface Project suggests that in 2018, an average of 43 Javan slow lorises are offered per month in online forums, an increase of 13 individuals per month since 2017.

To avoid being bitten by venomous slow lorises, traders habitually cut or pull out an animal's lower front teeth prior to selling them (Nekaris *et al.* 2013c). Traders may also cut teeth prior to packing slow lorises tightly into crates, as they often damage each other with their venomous bites during transport. Indeed, Fuller *et al.* (2017) showed that in a single confiscation of 77 slow lorises by Cikananga Wildlife Rescue Centre, nearly 30% died in the first 6 months, with morbidity from wounds, mainly bites, being the main cause of death. Other causes of death due to dental removal include dental abscess or pneumonia (Nekaris and Starr 2015). Those that do survive are no longer able to eat their preferred food (gum) (Das *et al.* 2014), or engage in the important behavior of social grooming with the toothcomb, meaning that any confiscated animals are unlikely to survive if released to the wild.

Reintroduction itself is a threat to the Javan slow loris. In the major markets in Java, at least four of the other six Indonesian species are traded along with the Javan slow loris, and in the markets in Sumatra at least three species are regularly traded, including ones that do not occur naturally on the island. The similar appearance of slow lorises, to the untrained eye, results in release of slow loris species from Sumatra and Borneo into Java and vice versa, causing potential for hybridization or even displacement of native species by introduced ones (Nekaris and Starr 2015). The ability of slow lorises to persist in human habitat if left undisturbed means that well-meaning people may translocate animals to habitat that is unknown to the animals, exacerbating these problems (Kumar *et al.* 2014).

Moore *et al.* (2014) assessed the success of Javan slow loris reintroductions, finding a death rate of up to 90%. Illness, hypothermia and exhaustion were all implicated in the death of slow lorises.

Reintroductions were started before the basics were known about the Javan slow loris' behaviour, ecology or distribution. No habitat suitability assessment could be made, since details were lacking on the type of habitat the species preferred and what it avoided. It has recently been reported by rescue centers that the success rates of Javan slow loris reintroductions are improving, but unfortunately no published data are available to verify these claims. Newspaper reports show that up to 30 slow lorises are released in one site at one time, but the highly territorial and venomous nature of slow lorises means that such releases are destined to have a high failure rate. A related study of pygmy slow lorises in Vietnam found that the season of release and age should be considered to increase the likelihood of survival (Kenyon *et al.* 2014).

To obtain vital information on the Javan slow lorises, in 2011 the Little Fireface Project instigated a study of the species' behavioural ecology in Garut District of West Java, Indonesia (Rode-Margono *et al.* 2014). This multi-disciplinary project has obtained data on home range size, social organisation, infant dispersal and feeding ecology. It was found that both sexes disperse from their natal range at about 20 months old, dispersal distances are 1–3 km from the natal range, home range sizes are large relative to the size of the animal (5–10 ha), the species goes into torpor, and the diet comprises mainly gum, supplemented with nectar and insects (Cabana *et al.* 2017). Several initiatives have been put into place to conserve slow lorises in the area and in other parts of Java. National workshops have been held for law enforcement officers and rescue center employees to provide essential data for a national slow loris action plan. At the local level, slow lorises are often totally dependent on local people for their protection, feeding on human planted tree species and residing in human farmlands. Thus, a major conservation program combining empowerment activities, conservation education and village events has been launched, and it is hoped that it can be used as a model for other key slow loris sites in Indonesia (Nekaris and Starr 2015).

For a long time, slow lorises were thought to be common throughout Indonesia, and the presence of animals in trade was believed to be an indicator of their abundance. We are only beginning to unravel the complexity of their taxonomy and distribution, leading to a bleak overall picture. While Java has an impressive and comprehensive

protected area network, encompassing over 120 terrestrial conservation areas and covering 5,000 km², enforcement of environmental laws and active protection in most of these parks is lacking. Besides curbing the illegal trade, it is paramount that these conservation areas, and indeed all other remaining forest areas on the island, be effectively protected.

PIG-TAILED SNUB-NOSE LANGUR

Simias concolor G.S. Miller, 1903

Indonesia

(2002, 2004, 2006, 2008, 2010, 2012, 2014, 2016, 2018)

Lisa M. Paciulli & Anna P. Pannick

The pig-tailed snub-nose langur (*Simias concolor*) is serving as the representative species for the six threatened and endemic Mentawai Islands primates. The other primates inhabiting this 7,000 km², ~70 island archipelago west of Sumatra are Kloss's gibbon (*Hylobates klossii*), the Pagai surili (*Presbytis potenziani*), the Siberut surili (*Presbytis siberu*), the Pagai macaque (*Macaca pagensis*), and the Siberut macaque (*Macaca siberu*). *Simias* is a monotypic genus with two subspecies: *S. c. concolor* / masepsep (Miller 1903) that inhabits Sipora, North Pagai Island, and South Pagai Island; and *S. c. siberu* / simakobu (Chasen and Kloss 1927), which is restricted to Siberut Island (Zinner *et al.* 2013). Threatened mainly by commercial logging, human encroachment, and hunting (Whittaker 2006), *Simias concolor* is classified as Critically Endangered on the IUCN Red List (Whittaker and Mittermeier 2008).

The populations of *S. concolor* on Pagai Island are threatened by forest conversion to oil palm plantations, forest clearings, product extractions by local people (Whittaker 2006), and opportunistic hunting (Paciulli 2004). On Siberut Island, a 1,000 km² oil palm plantation development was planned in 2014, and a 200-km² timber plantation for biomass energy production was planned in 2016 (Gaworecki 2016). Although both plans were cancelled as a result of local opposition, protests, and environmental assessments, attempts to exploit the Mentawai Islands' natural resources are likely to continue since the national government has designated

these biodiverse tropical islands as production forests (Gaworecki 2016).


Timber removal on a large scale is a concern as *S. concolor* has significantly lower densities in forests logged ~20 years previously – 2.54 individuals/km², compared to 5.17 individuals/km² in unlogged forests (Pagai Islands) (Paciulli 2004). It is estimated that on the Pagai Islands, there are approximately 3,347 pig-tailed snub-nosed langurs, 1,049 Kloss's gibbons, 1,545 Pagai surilis, and 7,984 Pagai macaques (Paciulli and Viola 2009). All of the primate species seem to reach

their highest known densities in the Peleonan Forest, an underused research base (Whitten 2009) and site of the Siberut Conservation Project in northern Siberut (Waltert *et al.* 2008). In Peleonan peat-swamp forests, *S. c. siberu* has densities as high as 65.5 individuals/km² (Quinten *et al.* 2010). However, in other Siberut peat swamp forests located in mangroves, the roots of many mangrove trees

are cut for the construction of ephemeral kitchen shelters. This human-inflicted soil and tree damage has long-term impacts on the regeneration of plants and the morphology of the area (Guillaud and Burgos 2018), likely making some of the mangroves unsuitable as primate habitat.

Logging facilitates hunting by providing easier access to forested areas, leaving primates more exposed and vulnerable (Febrianti 2015). Where hunting occurs on the Mentawai Islands, it has devastating effects on *S. concolor*, as it is the

Locals consider pig-tailed snub-nose langur meat to be a delicacy, and entire groups can be eliminated in a single hunting event.



preferred game species (Mitchell and Tilson 1986; Fuentes 2002; Paciulli and Sabbi 2017). The locals consider pig-tailed snub-nose langur meat to be a delicacy (Febrianti 2015), and entire groups can be eliminated in a single hunting event (Hadi *et al.* 2009). On the Pagais, few men report actively hunting (Paciulli 2004), but on Siberut, 24% of the men still hunt, with 77% targeting pig-tailed snub-nose langurs (Quinten *et al.* 2014). On Siberut, hunting reduces pig-tailed snub-nose langur group size, significantly impacts adult sex ratios, and affects the number of immature individuals in groups (Erb *et al.* 2012).



The uncertainty of Indonesian government land use means that land function and protection levels on the Mentawai Islands can change at any time with little notice, putting the species further at risk. There is only one large protected area for *S. concolor*: the 190,500 ha Siberut National Park, a UNESCO Biosphere Reserve that covers 47% of Siberut. Although the park serves as the main reserve for ~51,000 primates (Quinten *et al.* 2015), hunting is much more prevalent there than elsewhere, with ~4,800 primates being removed each year (min. 6.4 % of the population) (Quinten *et al.* 2014). Drastic measures need to be taken to ensure that the Peleonan Forest on Siberut is truly protected. The same goes for forests on the southern islands, where the Pagai macaque and Pagai surili are not represented in any protected areas (Supriatna *et al.* 2017).



Whittaker (2006) suggested protecting areas in the Pagai Islands by cooperating with a logging corporation that has practiced sustainable logging there since 1971, as well as increased protection for Siberut National Park, which currently lacks enforcement. The Peleonan Forest in North Siberut, which is home to unusually high primate populations and easily accessible, also needs to be safe-guarded. In addition, the Mentawai people could benefit from conservation education, especially regarding hunting, and the development of alternative economic models to reduce the likelihood that land will be sold to logging companies (Whittaker 2006). Although the World Wildlife Fund, Asian Development Bank (1992-2000 loan project), and Phase I of the World Bank-implemented Critical Ecosystem Partnership Fund have poured over \$1 million into Siberut, Whitten (2009) noticed that little has changed on the remote islands in thirty years.

GOLDEN-HEADED LANGUR


or Cat Ba Langur

Trachypithecus poliocephalus Trouessart, 1911

Vietnam

(2000, 2002, 2004, 2006, 2008, 2010, 2012, 2014, 2016, 2018)

Neahga Leonard, Richard J. Passaro, Daniela Schrudde, Roswitha Stenke,
Phan Duy Thuc & Martina Raffel



The Cat Ba langur (also known as the golden-headed langur), *Trachypithecus poliocephalus*, is probably the most endangered of the Asian colobines, and is assessed as Critically Endangered (Bleisch *et al.* 2008). This species occurs only on Cat Ba Island in the Gulf of Tonkin off the north-eastern Vietnamese shore (Stenke and Chu 2004). The Cat Ba Archipelago is adjacent to the world-famous Ha Long Bay, a spectacular karst formation that was invaded by the sea following the last major glaciation. The favoured habitat of the Cat Ba langur is tropical moist forest on limestone karst hills, a habitat preference it shares with the other six to seven taxa of the *T. francoisi* group.

While there are no systematic and reliable data available on the historic density of the langur population on Cat Ba Island, reports by indigenous people suggest the entire island of Cat Ba (140 km²) and some smaller offshore islands were previously densely populated by langurs. Hunting has been identified as the sole cause for the dramatic and rapid population decline from an estimated 2,400–2,700 in the 1960s to approximately 50 individuals by 2000 (Nadler and Ha 2000). The langurs were poached mainly for trade in traditional medicines and for sport. Since the implementation of strict protection measures in 2000, the langur population on Cat Ba Island has stabilized (Nadler *et al.* 2003) and since 2003 has been on the increase (Leonard 2016). In the latter half of 2015 numbers fell from the mid-high 60s to the low 50s and have since been slowly recovering. This has raised concerns that as langur numbers recover, interest in poaching by people from adjacent regions may also revive (Leonard 2016).

Although the growth of the population is encouraging, the overall status of the species



remains critical and the total population is worryingly small. Habitat fragmentation and hunting has divided the remaining population into several isolated sub-populations, some of which are non-reproducing social units. A surplus of young males is a cause for concern as take-over attempts can lead to infanticide and inadvertent infant deaths, both of which were recorded in early 2018 (N. Leonard pers. comm.).

The total reproductive output of *Trachypithecus poliocephalus* has been low due to the small population and the long inter-birth cycle, but records indicate that the birth rate is increasing, with 49% of the total births recorded between 2000 and 2018 having taken place from 2014–2018. Births occur throughout the year, with a peak in January–April, just prior to the rainy season (Leonard et al. 2016, N. Leonard pers. comm.).

In 2012, after many years of planning and preparation, one group of two females was successfully translocated from a small off-shore islet where they had become stranded to the relative safety of the strictly protected core zone of Cat Ba National Park. Here they quickly assimilated into existing groups containing males, thus allowing them the opportunity to reproduce for the first time ever. It is hoped that continued protection efforts and additional population management interventions such as these will enhance the rebound of this species.

The Cat Ba Archipelago and adjacent Ha Long Bay are nationally and internationally recognized for their importance to biodiversity conservation. Cat Ba National Park was established in 1986. It presently covers more than half of the main island. Ha Long Bay was established as a World Heritage site in 1994, and the combined archipelago includes ~1,500–2,000 large and small islands, cliffs and rocks. In 2004, the Cat Ba Archipelago was designated a UNESCO Man and the Biosphere Reserve. Despite the conservation designations and laws to protect the region, nature and wildlife protection on Cat Ba Island is deficient. Environmental awareness and commitment

among the local communities is slowly increasing, and hunting/trapping of all animals is illegal on Cat Ba Island. Unfortunately, efforts to effectively conserve the langurs and their habitat continue to face major obstacles from increasing tourism development, increasing human population and severe deficiencies in law enforcement (Stenke 2005; Leonard 2018). As is common elsewhere in the region, poaching by local people is driven by livelihood issues, brought about by low incomes and lack of employment opportunities. Immense local and regional demand for wildlife and animal parts for food and dubious traditional medicines provide a market for poached animals and plants. Although langur hunting ostensibly stopped years ago, the 2015 decline in numbers raises doubts as to the permanence of the hunting cessation. Regardless, hunters continue to poach other animals and plants in langur areas, thus jeopardising langur habitat. Strict enforcement of

the established protections is therefore necessary for the survival of all species on Cat Ba Island that are targeted by the illegal Asian wildlife trade.

A conservation program for the Cat Ba langur is supported by Zoo Leipzig, Zoological Society for the Conservation of Species and Populations (ZGAP), and the Allwetterzoo Münster in Germany. The

project was initiated on Cat Ba Island in November 2000 by Allwetterzoo Münster and ZGAP. The aim of the Cat Ba Langur Conservation Program is to provide protection for the langurs and their habitat, to conduct research that will help inform future population management decisions, and to help contribute to the conservation of the overall biodiversity of the Cat Ba Archipelago, all in collaboration with Vietnamese authorities.

A surplus of young male Cat Ba langurs is a cause for concern as group takeover attempts can lead to infanticide.

GOLDEN LANGUR

Trachypithecus geei Khajuria, 1956

India, Bhutan
(2018)

Rekha Chetry, Dilip Chetry & P. C. Bhattacharjee

The Golden langur (*Trachypithecus geei*) is an attractive, arboreal and diurnal primate endemic to India and Bhutan in south-east Asia. It was first discovered by E. P. Gee in 1953. As the name suggests, its coat color is golden orange, but only during the breeding season – in the rest of the year it becomes creamy or dirty white. Ventral coat color is comparatively lighter, and the golden orange color is brighter in females than males. The species is predominantly arboreal, spending 99% of its active time in trees, foraging primarily in the top and middle strata of forest canopies (Biswas 2004). However, in degraded habitats, they descend to the ground (Chetry and Chetry 2009).

Golden langurs mature sexually after 5–7 years for males and 4 years for females. The breeding season is between June and January and the birth season is January to June. The gestation period is 168–180 days with one infant being born at a time, and the inter-birth interval is two years (Chetry and Chetry 2009). Sexual harassment is an integral part of the species' reproductive behaviour. Infants and, to some extent, females, have been identified as harassers.

The golden langur is highly social and maintains diverse forms of societies: (1) uni-male, multi-female troops/harem troops (3–9 members with 1:2.68 male–female ratio), (2) bi-male, multi-female troops (8–15 members with 1:1.94 male–female ratio), (3) multi-male, multi-female troops, (4) all male bands (2–5 individuals) and (5) lone males (Chetry and Chetry 2009). However, uni-male, multi-female troops or societies are the most stable and common form of social structure, followed by bi-male, multi-female societies (Biswas

2004). Troops are cohesive and both intra- and inter-troop interactions are mostly peaceful. The annual home range is between 10 and 58 ha for diverse social troops in different habitat conditions (Chetry and Chetry 2009) and day path lengths vary from 200 to 700 m.

The golden langur is diurnal. They spend, on an annual basis, 12.8–33% of time feeding, 40–63.1% resting, 6.3–19% in locomotion, 5–11.5% monitoring, 2–3.7% playing and 0.3–6% grooming (Mukherjee 1996; Chetry 2002; Medhi and Chetry 2003; Biswas 2004; Medhi 2004). At night, golden langurs select tall trees of few selected species to sleep in (Biswas 2004).

Golden langur deaths are being attributed to electrocution on power lines, road accidents, and attacks by domestic dogs.

Green leaves (both young and mature) form the major constituent of the folivorous golden langur's diet. Other dominant food items include fruits, seeds, flowers, stem cortex and twigs from >200 plant and tree species. Gum, soil, algae, snails and alcoholic effluence are also important supplements in the diet of golden langurs (Medhi 2004;

Biswas 2004). Their primary predators are leopard, wild dog and python (Chetry *et al.* 2005; Chetry *et al.* 2018). Their anti-predator response varies according to the predator (Chetry *et al.* 2007).

In India, its distribution extends over an area greater than 2,500 km², bordered by the rivers Manas in the east, Sankosh in the west and the Brahmaputra in the south (Srivastava 1999). Remarkably, Ram *et al.* (2016) found that the Aie and Champabati rivers are also acting as natural barriers to migration between golden langur populations in Assam. Its distribution in Bhutan is limited between the Sankosh river and Chamkhar-Mangde-Manas river complex and



covers a range of 4,782 km² (Wangchuk 2005, Lhendup *et al.* 2018).

In India, the estimated available habitat for the golden langur is 1,255 km². While it primarily inhabits wet evergreen and tropical semi-evergreen forests, it also thrives in sal (*Shorea robusta*) dominated forests and secondary forests. In Bhutan, available habitat is 3,475 km², out of an estimated 4,782 km² potential habitat (Wangchuk 2005). The preferred habitat here is warm broad-leaved forests between 1,000 m and 2,600 m asl, and subtropical forests between 200 m and 1,000 m asl. The golden langur shares its habitat with three other primate species: Assamese macaque (*Macaca assamensis*), rhesus macaque (*Macaca mulatta*) and slow loris (*Nycticebus bengalensis*).

The estimated population size of the golden langur in Bhutan is 6,637 (Wangchuk 2005). In India, Ghosh (2009) and Biswas *et al.* (2010) observed 5,141 individuals in 566 troops. Thus, the global population of the species is estimated to be >12,000 (Chetry and Chetry 2009; Horwich *et al.* 2013).

The conservation status of the golden langur, according to the IUCN Red List, is Endangered (Das *et al.* 2008). The Wildlife (Protection) Act of India (1972) and the Forest and Nature Conservation Act of Bhutan (1995) have classified the animal as a Schedule-I species, and it is an Appendix-I species in CITES. In India, protected habitat is limited to Manas National Park and the Chakrashila Wildlife Sanctuary. The combination of habitat loss and fragmentation have already taken their toll on golden langur populations in India where it is also threatened by encroachment, illegal tree felling, fuel wood collection and cattle grazing (Chetry *et al.* 2018). Srivastava *et al.* (2001) reported a 50% loss of original habitat of the species in India.

Due to severe shrinkage and fragmentation, eight isolated populations were wiped out from eight forest patches between 1970 and 1990, all in Assam, India (Choudhury 2002). Moreover, at the human-wildlife interface, golden langur deaths are being attributed to electrocution on power lines and road accidents (Chetry and Chetry 2009) and they are increasingly attacked by dogs.

In Bhutan, the species is better protected, with 50% of its habitat situated within the protected area network, including Royal Manas National

Park, Black Mountain National Park and Phipsoo Wildlife Sanctuary.

Nevertheless, the species remains under pressure from a myriad of anthropogenic threats including fluctuating land tenure systems, infrastructure development, shifting cultivation and commercial logging (Wangchuk 2005). However, the most severe threat has arisen from the hybridization of the Golden langur with capped langurs as a result of the recently built suspension bridges over the Chamkhar river (Wangchuk *et al.* 2005; Choudhury 2008; Ram *et al.* 2016). Alarming, 15% of the golden langur population is now hybridised (Wangchuk 2005).

Conservation challenges are likely to increase despite current conservation initiatives. There is no doubt that the local administration and NGOs are working at different levels to address the threats faced by the golden langur. Yet the current situation calls for more effective and continuous conservation action in order to safeguard the golden langur and prevent it from becoming Critically Endangered.

To secure the future of the species, several actions are recommended. Firstly, we advocate for a reform in the protected area network. Specifically, we propose that Bhumeswar Proposed Reserve Forest (PRF), Bamungaon PRF, Nadangiri Reserve Forest (RF), Kakoijana RF and the Sankarhola area under the Bhairabchura PRF should be declared as Community Reserves and directly involve the local community in the conservation of the golden langur. In Assam, the proposed Ripu Chirang Wildlife Sanctuary should be instated as early as possible. Finally, Chakrashila Wildlife Sanctuary along with its adjacent reserve forests such as Sreegram, Katrigacha, Buxamara and Nadangiri Hill should be upgraded to a National Park at the earliest opportunity to ensure the preservation of the golden langur's habitats in India.

Secondly, we propose forest corridors to connect these forest fragments. Specifically, forested corridors should be created between (1) Chirang RF to Bengtol RF to Manas RF in the northern range of its population, (2) Chakrashila to Abhaya rubber garden to Naddengri RF, (3) Bamingaon to Khoragaon PRF, and (4) Nakkati to Kakoijana RF in the southern range, to provide larger areas for the long-term survival of these populations.

These efforts would be complemented by habitat restoration to reverse habitat loss. In instances where it is not possible to connect fragments by forested corridors, a translocation management plan should be considered, to reinforce remnant populations.

Thirdly, the re-assessment of the current status of the species across its entire distribution is urgently needed. Finally, a recognised and state-sponsored species action plan is needed, which includes the recommendations included here.

PURPLE-FACED LANGUR

Semnopithecus vetulus Erxleben, 1777

Sri Lanka
(2004, 2006, 2008, 2010, 2012, 2014, 2016, 2018)

Rasanayagam Rudran

After Sri Lanka's twenty-six year civil war ended in 2009, extensive deforestation occurred, which escalated conflicts between humans and monkeys. This conflict undermined the long-term survival of all three Sri Lankan primate species (*Semnopithecus vetulus*, *S. priamthersites* and *Macaca sinica*), which are not only endemic, but also threatened with extinction. As public outcry and political pressure mounted to resolve these conflicts, several government institutions and non-governmental organizations, led by SPEARS Foundation, helped the country's Department of Wildlife Conservation (DWC) to develop an action plan for people to conserve and coexist with all species of monkeys. The plan was submitted to the country's government in March 2016 for cabinet approval.


While awaiting approval, SPEARS Foundation used funds from foreign donors to implement some key elements of the plan. One was to develop Community Conservation Areas (CCAs), which, when established, would be administered and managed sustainably by local communities under DWC supervision. To find suitable sites for CCAs, the SPEARS team analysed complaints of human-monkey conflicts received by the DWC between 2007 and 2015. The analysis indicated that conflicts occurred throughout the country, but their frequency varied between localities (Cabral *et al.* 2018). Therefore, thirteen field surveys were conducted from 2016 to 2018 to locate sites best suited for the establishment of CCAs. Information from these surveys and other relevant data on all four purple-faced langur subspecies are presented below.

Urbanisation poses a serious threat to the long-term survival of the Critically Endangered and endemic subspecies of the western purple-faced langur.

The range of the western purple-faced langur (*Semnopithecus vetulus nestor*) includes the most densely populated region around Colombo, the country's capital. Therefore, urbanisation poses a serious threat to the long-term survival of this Critically Endangered and endemic subspecies (Dittus *et al.* 2008; Rudran *et al.* 2009). A survey conducted in 2007 (Rudran 2007) indicated that 81% of *S. v. nestor*'s historical range (Hill 1934; Phillips 1935) had been deforested and converted to human altered landscapes. Due to this habitat reduction, much of *S. v. nestor*'s current population subsists mainly on fruit from home gardens (Dela 2007; Rudran 2007). Nutritional consequences of feeding on a low diversity diet of cultivated fruits are unclear but considered detrimental to the folivorous *S. v. nestor* (Rudran 2015).

Besides depleting natural food sources, deforestation causes habitat fragmentation, forcing animals to travel on the ground and along power lines to move between fragments. These movements increase mortality by exposing them to attacks by dogs, speeding vehicles, and electrocution (Parker *et al.* 2008). In some parts of its range, *S. v. nestor* is occasionally shot and killed as a pest while feeding in home gardens (Dela 2004). Such human-induced fatalities reduce group sizes and appear to lead to local extinctions in *S. v. nestor*'s range (Rudran 2007).

To promote environmental awareness among school children, classroom lectures were presented to 1,360 students, and about 400 of them participated in nature walks. To enhance economic stability and reduce unsustainable dependence on natural resources, 90 adults



were trained to cultivate pepper, a crop with considerable demand in world markets. They also received pepper plants to grow in their home gardens to generate income. Another initiative trained young adults as nature guides to earn income by accompanying visitors to the species-rich lowland rainforest around Thummodara. Some stakeholders were also given loans to produce food, drinks and local handicrafts to sell to visitors. A brochure was prepared to advertise the ecotourism program to hotel visitors, and a website is under construction to promote it via the internet.

The highland purple-faced langur (*S. v. monticola*), also known as the bear monkey, was investigated for two years at Horton Plains by Rudran (1973a, 1973b) nearly fifty years ago. When the area was surveyed again in 2016, Rudran noted appreciable changes to the vegetation. Many species previously recorded as important food plants of the bear monkey were dead or dying. This appeared to be primarily due to debarking of the adult trees and feeding of saplings by the sambar (*Cervus unicolor*) population, which had increased in population because of the invasive soft grass introduced to Horton Plains with the fertilizer used by a now defunct potato farm (Adikaram *et al.* 1999). The death and lack of regeneration of food plants appear to have undermined bear monkey survival. A census was not conducted in 2016, but early morning loud calls of harem males were considerably less frequent than before, indicating a population decline. The area was surveyed again in 2017 to collect data on crop damage and human attitudes towards monkeys.

Three surveys were conducted in the range of the southern purple-faced langur (*S. v. vetulus*) in 2017. Data from these surveys are still being analysed but a long-term study of *S. v. vetulus* (Roscoe *et al.* 2013) reported several threats to the future survival of this subspecies. These threats were the same as those experienced by *S. v. nestor*. Additionally, a major highway constructed through *S. v. vetulus*'s range is expected to create a permanent barrier to gene flow between the populations found along the coast and the interior of the country.

The northern purple-faced langur (*S. v. philbricki*) was investigated for two years in the late 1960s (Rudran 1973a, 1973b) when conflicts with humans were not a serious issue. In the late

1970s, however, the impact of the Accelerated Mahaweli Development Program (AMDP) on wildlife in *S. v. philbricki*'s range became a serious concern. To mitigate this concern, an Environmental Impact Assessment (EIA) of AMDP recommended the establishment of four new National Parks around the development area (Tippetts-Abbott-McCarthy-Stratton 1980). While these areas provided protection to *S. v. philbricki*, serious threats such as habitat fragmentation and hunting for food, medicinal purposes and rituals still remained in other areas (Wickremasinghe *et al.* 2016). Similar findings have also been reported by other investigators (Nahallage and Huffman 2013). Two surveys conducted in 2018 by SPEARS Foundation staff found that populations of the highly arboreal *S. v. philbricki* were fewer than that of the other two subspecies in the area due to habitat fragmentation.

In conclusion, although Sri Lanka's monkeys face a perilous future (Rudran 2013), there is hope that they can be conserved. One reason for hope is that most Sri Lankans follow the Buddhist doctrine of compassion towards all living things. Therefore, promoting this doctrine and Buddha's own reverence of the forest present opportunities to deter deforestation in a country steeped in cultural traditions but ignorant of the detrimental effects of habitat destruction. Another reason for optimism stems from a decision by successive governments to increase Sri Lanka's forest cover from 27% to 36% using native plants, to achieve the country's economic development goals (Yatawara 2011). The political will to increase forest cover augurs well for the future protection of wildlife. It is important that the Sri Lankan government approves the 2016 action plan in order to ensure a steady flow of financial support to conserve Sri Lanka's monkeys.

GAOLIGONG HOOLOCK GIBBON

or Skywalker hoolock gibbon

Hoolock tianxing Fan et al., 2017

China, Myanmar
(2018)

Pengfei Fan, Hanlan Fei, Lu Zhang, Guopeng Ren & Susan M. Cheyne

Hoolock gibbons were first described scientifically by Harlan (1834) under the name *Simia hoolock*. They were subsequently transferred to the genus *Hylobates*, and then assigned to their own distinct subgenus (later elevated to genus), first *Bunopithecus* (later restricted to an extinct Quaternary gibbon from China) (Prouty et al. 1983; Groves 2001) and then *Hoolock* Mootnick and Groves 2005. Taxonomic variation between different hoolock populations was first recognized by Groves (1967), who identified a major east-west morphological division and described *Hylobates hoolock leuconedys* to distinguish eastern hoolock populations from those in the west, geographically isolated by the Chindwin River. Both subspecies were latterly elevated to full species: the western (*Hoolock hoolock*) and eastern hoolock (*H. leuconedys*) gibbons. Fan et al. (2017) assessed the morphological and genetic characteristics of wild animals and museum specimens to evaluate the taxonomic status of the hoolock population in China. The results suggested that hoolocks distributed to the east of the Irrawaddy and Nmai Hka rivers, which were previously assigned to *H. leuconedys*, are morphologically and genetically distinct from those to the west of the rivers, resulting in them now being recognized as a new species: the Gaoligong hoolock gibbon or skywalker hoolock gibbon, *Hoolock tianxing* Fan et al., 2017.

In 2009, the Skywalker hoolock gibbon population was estimated to be less than 200 individuals; now, it is less than 150.

Hoolock tianxing was once widely distributed around the west bank of the Salween River, west of Yunnan, China, but >90% of its habitat was lost by 1994 (Fan et al. 2017). In 2009, the population was estimated to be <200 individuals (Fan et al. 2011). Now, this figure is less than 150 individuals, made up of 34 family groups and 10 solitary individuals across 17 subpopulations. The largest subpopulation has seven groups. Five subpopulations have only one group. The population has remained relatively stable from 2009 to 2017, but is isolated from other populations by distance, villages and roads, and has a low birth rate. For example, between 2008 and 2018 the reproductive rates of three mature females were tracked. One of these females produced two offspring in this time (November 2008 and December 2012), whilst the others produced just one (2008 and 2012). *Hoolock tianxing* is listed as Critically Endangered under criterion A4a,c,d.

Agricultural encroachment, commercial logging, habitat fragmentation and isolation, and hunting (for bushmeat and pet trade) are major threats to *H. tianxing*. Additionally, the population is threatened by stochastic loss, in which subpopulations are reduced to 1-2 groups with no opportunity for dispersal or gene-flow. Population linking, protection and habitat restoration are urgently needed, and the translocation of non-viable subpopulations may also be required. There is also a hoolock population in Myanmar. While unstudied, it

TAPANULI ORANGUTAN

Pongo tapanuliensis Nurcahyo, Meijaard, Nowak, Fredriksson
& Groves in Nater et al., 2017

Indonesia (Sumatra)
(2018)

Gabriella Fredriksson, Matthew G. Nowak, Jatna Supriatna & Serge Wich

is likely that the population faces similar difficulties to those in China, (i.e. habitat loss and poaching) but there is comparatively less conservation action and law enforcement in Myanmar. As the population in China decreases, the importance of the Myanmar population increases. Therefore, although demand for conservation intervention in Myanmar is high, a careful approach is advised to safely navigate the recent political unrest.

The following actions are needed: (1) Raise awareness of this species, especially in China through targeted campaigns, (2) Determine population status in Myanmar through population surveys, (3) Address threats at a local scale through an ethnographic approach, (4) Investigate possibilities for connecting populated forest fragments and/or translocation of isolated groups/individuals.



The Tapanuli orangutan, *Pongo tapanuliensis*, was only formally described in 2017, when it was shown that an isolated orangutan population in the Batang Toru region, which used to be considered the southernmost range of extant Sumatran orangutans (*Pongo abelii*), south of Lake Toba, is distinct from other Sumatran and Bornean populations (Nater et al. 2017). Through a comparison of cranio-mandibular and dental characters from an orangutan killed during human-orangutan conflict to a comparative sample of adult male orangutans of similar developmental stage, Nater et al. (2017) found consistent differences between the Batang Toru individual and other extant Ponginae. Similarly, comparisons of adult male long calls from two Tapanuli males with those of a large sample of Bornean and Sumatran males also revealed a unique mix of long call characteristics. Model-based approaches based on the analyses of 37 orangutan genomes supported the morphological results, revealing that the deepest split in the evolutionary history of extant orangutans occurred ~3.38 mya between the Batang Toru population and those to the north of Lake Toba. In comparison, the Bornean orangutan and Sumatran orangutan separated much later at about 674 ka. The analyses show that there was some gene flow between the Sumatran and Tapanuli orangutan species until 10–20 ka. Combined, these analyses support a new classification of orangutans into three extant species.

Due to high levels of habitat conversion and fragmentation, along with illegal hunting and poaching, the Tapanuli orangutan is estimated to have experienced a significant population reduction in recent decades (Nowak et al. 2017). With a population estimate of 767 (95% confidence intervals 231-1,597) individuals (Wich et al. 2019), the Tapanuli orangutan is the least numerous of all great ape species. Its distribution is separated by around 100 km from the closest population of the



© Maxime Aliaga

Sumatran orangutan to the north. A combination of small population size and geographic isolation is of particularly high conservation concern, as it may lead to inbreeding depression (Hedrick and Kalinowski 2000) and threaten population persistence (Allendorf *et al.* 2013). Nater *et al.* (2017) recorded extensive runs of homozygosity in the genomes of two Tapanuli orangutan individuals, pointing at the occurrence of recent inbreeding.

The only known population of Tapanuli orangutans occurs in the uplands of the Batang Toru Ecosystem, an area of roughly 1,500 km² consisting of three forest blocks, of which 1,023 km² is suitable orangutan habitat (Wich *et al.* 2016, 2019). Most of this is upland forest (>500 m asl, up to 1800 m asl), covering the upper watersheds of eight river systems and providing fresh water for over 100,000 people across Tapanuli. Forest loss data indicate that orangutan habitat below 500 m asl was reduced by 60% between 1985 and 2007 for both the Tapanuli and the Sumatran orangutan (Wich *et al.* 2008, 2011). It is thought that more Tapanuli orangutan habitat will be lost as significant areas

of forest in its range remain under considerable threat (Wich *et al.* 2016, 2019; Sloan *et al.* 2018) from habitat conversion for small-scale agriculture, mining exploration and exploitation, a large-scale hydroelectric scheme, geothermal development and agricultural plantations. Only about 10% of its geographic range is in an area recognized by the 'World Database of Protected Areas'. Another 76% is in Hutan Lindung (Protection Forest), and 14% does not have any 'forest status' in the spatial plans. The area without any 'forest status' consists of rugged primary forest with the highest densities of Tapanuli orangutans in the Batang Toru Ecosystem (SOCP unpublished data). The protected areas are not immune from the above threats (Wich *et al.* 2008, 2011, 2016) and orangutans in these areas are also hunted (Wich *et al.* 2012). Due to their slow life history, with a generation time of at least 25 years, orangutans on Sumatra are unable to sustain substantial and continual loss of individuals (Wich *et al.* 2004, 2009; Marshall *et al.* 2009).

The Tapanuli orangutan was more widespread until relatively recently, with sightings further south in

the lowland peat swamp forests in the Lumut area (Wich *et al.* 2003) and several nests encountered during a rapid survey in 2010 (G. Fredriksson pers. obs.). However, the forests in the Lumut area have been almost completely converted to oil-palm plantations in recent years. Observations were also made of a male orangutan in the Adiankoting subdistrict in North Tapanuli, north of the Batang Toru West forest block, during a human conflict situation where the orangutan was shot at with an air rifle when it was found foraging on durian fruits (G. Fredriksson pers. obs.). The persistence of viable subpopulations in these areas is currently not known.

Tapanuli orangutans have been observed feeding on a number of tree species that have not previously been recorded as orangutan food species. These unique species include *Gymnostoma sumatranum* from the Casuarinaceae family, and *Dacrydium imbricatus*, *Dacrydium beccarii*, *Dacrydium comosum*, and *Podocarpus neriifolius* from the Podocarpaceae family. At SOCP's long-term monitoring station in the Batang Toru Ecosystem, 21.9% of all feeding observations recorded between 2011 and 2015 were represented by five conifer species (Araucariaceae and Podocarpaceae) and one non-conifer evergreen species (Casuarinaceae). Seeds of *Agathis borneensis* from the Araucariaceae family have been considered a 'fallback' fruit, frequently consumed when few other fruits are available (Nater *et al.* 2017; SOCP unpublished data). Thus, a significant proportion of the dietary profile of Tapanuli orangutans is markedly different from that of previously studied orangutan populations.

Due to the extremely rugged terrain, external threats have been primarily limited to illegal clearing of protected forests, hunting and killing during crop conflict, and trade in young orangutans (Wich *et al.* 2012, 2016). Encroachment and hunting have increased in recent decades, due to an influx of migrants from Nias Island, west of Sumatra, who settle on protected forest land on Batang Toru's forest edge where no land claims exist at present (Wich *et al.* 2012). In addition, despite land


status changes from Production Forest to that of Protection Forest in 2014 (Ministry of Forestry of the Republic of Indonesia 2014), one company still maintains a controversial 300 km² logging permit located in primary forest in the current range of the Tapanuli orangutan. In the southwest corner of the Batang Toru Ecosystem, a large gold and silver mine has converted key lowland habitat of the Tapanuli orangutan and retains controversial mining permits overlapping parts of the remaining Tapanuli orangutan range. Land speculation related to the company's exploration is further threatening the primary forest. More recently, the development of a hydroelectric project has started in the area of the highest orangutan density, which could impact roughly 100 km² of Tapanuli orangutan habitat, or nearly 10% of the entire species' population (Sloan *et al.* 2018). This controversial hydroelectric scheme, located at a hotspot of seismic activity on the Great Sumatran Faultline,

also jeopardizes the chances of maintaining and restoring habitat corridors between the western and eastern Tapanuli orangutan ranges and a strict nature reserve with a small population of Tapanuli orangutans (Wich *et al.* 2019). If the connectivity between these populations is not restored, and the last core high-density habitat of the Tapanuli orangutan is bisected by infrastructure


development related to the hydro dam (roads, tunnel, high electricity power lines), the long-term survival of the Tapanuli orangutan will be severely threatened (Wich *et al.* 2019).

In order to safeguard the future of the most endangered great ape species in the world, all possible efforts must now be made to prevent any further degradation of Tapanuli orangutan habitat, and to reconnect its three habitat fragments to restore genetic exchange. As it currently stands, two of the three habitat fragments do not contain viable populations, leaving only one viable and highly threatened population as the future of the species. Lastly, field management activities need to be established to prevent further hunting and encroachment, with clear and enforced boundary demarcation, and active human-orangutan conflict mitigation efforts put in place.

With a population estimate of fewer than 800 individuals, the Tapanuli orangutan is the least numerous of all great ape species.



NEOTROPICS



Neotropical Primates

- | | |
|--|--|
|  <i>Alouatta guariba</i> |  <i>Saguinus bicolor</i> |
|  <i>Cebus aequatorialis</i> |  <i>Callithrix aurita</i> |
|  <i>Ateles geoffroyi</i> |  <i>Plecturocebus olallae</i> |



BUFFY-TUFTED-EAR MARMOSET

Callithrix aurita É. Geoffroy Saint-Hilaire, 1812

Brazil
(2018)

Rodrigo S. Carvalho, Sally J. Fransen, Mônica M. Valença-Montenegro, Nicholas J. Dunn, Cláudia A. Igayara-Souza, Márcio Port-Carvalho, Dominic Wormell, Fabiano R. Melo, Alessandro Silva, Wagner R. Lacerda & Leandro Jerusalinsky

There are six species in the genus *Callithrix*, all endemic to Brazil, but only two, *Callithrix aurita* (VU) and *Callithrix flaviceps* (EN), are listed as Threatened on the IUCN Red List (IUCN 2019). *Callithrix aurita* is under consideration for uplisting from Vulnerable to Endangered in its most recent reassessment (IUCN SSC Primate Specialist Group, in prep.). *Callithrix aurita* is under extreme and increasing threat from habitat loss, habitat fragmentation and competition and hybridization with invasive *Callithrix* species, and the recent outbreak of yellow fever in its range adds significantly to these threats.

Remarkable for its face, which resembles a "little skull", the buffy-tufted-ear marmoset is a small (~420g) primate from the montane region of south-eastern Brazil. *Callithrix aurita*'s habitat is in the mountain chains of the Atlantic Forest of the south-eastern states of Brazil (São Paulo, Rio de Janeiro and Minas Gerais) (Rylands *et al.* 1993). Their presence there has been related to the lower temperatures found in the highlands, with most populations encountered at altitudes between 600 and 1300 m (Norris *et al.* 2011).

Of all Brazilian biomes, the Atlantic Forest is the most heavily populated and a substantial part of it is now an archipelago of small islands of vegetation embedded in a matrix of degraded areas, pasture, agriculture, forestry and urban areas (Joly *et al.* 2014).

Another serious issue is the presence of

marmosets introduced from distant and different Brazilian ecosystems into *C. aurita*'s range. Ecological research has shown that *C. aurita* faces significant competition from invasive marmosets, *C. penicillata*, *C. jacchus* and hybrids between them (Pereira 2006, 2010; Rylands *et al.* 2008; Melo and Rylands 2008; Pereira *et al.* 2008, 2014; Port-Carvalho and Kierulff 2009; Nogueira *et al.* 2011; Bechara 2012; Carvalho *et al.* 2013; Carvalho 2015; Melo *et al.* 2015; Nunes 2015; Gonçalves 2016; Silva *et al.* 2018), and genetic research also demonstrates hybridization between *C. aurita* and

their introduced congeners (Pereira 2010; Nogueira *et al.* 2011; Carvalho *et al.* 2013; Carvalho 2015). Groups of invasive and hybrid marmosets are replacing native *C. aurita* populations (Pereira 2006, 2010; Bechara 2012; Oliveira 2012; Carvalho *et al.* 2013; Carvalho 2015; Melo *et al.* 2015; Nunes 2015), and hybridization is undoubtedly affecting

the genetic integrity of small, pure *C. aurita* populations through genetic introgression.

The speed and potential consequences of the invasive process that *C. aurita* faces is well exemplified by the situation in the Serra dos Órgãos National Park in Rio de Janeiro State, where five years of observations of two pure *C. aurita* groups in the park found no contact between these native groups and any invasive marmosets. However, field observations in 2015 demonstrated, for the first time, the arrival of invasive marmosets competing

Groups of invasive and hybrid marmosets are replacing native buffy-tufted-ear marmoset populations.



with the native study groups. The replacement of one of the pure *C. aurita* groups by a mixed group of the invasive and native species was observed in the same year. A recent genetic study has also evidenced ongoing hybridization on the edge of the Serra dos Órgãos National Park (Carvalho 2015). This area is emblematic of the invasive processes that are happening in other parts of *C. aurita*'s native range (C. Knogge, W. Lacerda, R. Carvalho, L. Oliveira, D. Pereira, J. Malukiewicz, pers. obs.).

From 2016 to 2017, the yellow fever virus emerged in eastern Brazil, predominantly in the states of Minas Gerais, Espírito Santo, São Paulo and Rio de Janeiro. This was the largest outbreak of yellow fever observed in recent history. The re-emergence of the virus had great impact on the country's non-human primate (NHP) populations, affecting more than 7,000 NHPs, with 1,412 confirmed epizootics, and 777 confirmed human cases with 261 deaths (MinSaude 2017). *Callithrix aurita*'s range contained the highest number of confirmed cases during the 2016-2017 outbreak. In *Callithrix* alone, 257 yellow fever cases have been confirmed so far (2511 reports with final analysis and outcomes still in process), demonstrating the susceptibility of the genus to the disease. Numbers are likely to be far higher than reported as, because of their small size, marmoset cadavers are very difficult to find in the dense forests.

The species' situation has concerned primatologists since 1971 (Coimbra-Filho 1971), but it was only in 2014 that deeper questions were raised and the severity of the threats *C. aurita* faces were widely embraced. These concerns resulted in action being taken to establish an international conservation program to save the species (the Mountain Marmosets Conservation Program - MMCP). *The National Action Plan for the Atlantic Forest Primates and the Maned Sloth* (ICMbio 2018) was launched in 2018, covering *C. aurita* and 12 other threatened primates.

One of the first actions of the collaborative conservation initiative (MMCP) has focussed on increasing surveys to identify regions with pure *C. aurita* populations, hybrid groups and invasive *Callithrix* spp. (Lacerda *et al.* 2015; Melo *et al.* 2015; Nunes 2015; Carvalho *et al.* 2018). The NGOs PREA and Muriqui Institute of Biodiversity (Minas Gerais), the Environmental Secretariat of

São Paulo and the research team and managers of the Serra dos Órgãos National Park began surveys in October 2015. The surveys over the species range show alarming results. In approximately 100 different locations, 85 groups of *C. aurita* and 47 groups of invasive species or hybrids were found (Carvalho and Lacerda, unpubl. data).

Due to the alarming results from the field, the MMCP recognized the urgency of organizing and reinvigorating the captive population of *C. aurita*. In 2017, the MMCP executed a masterplan for movements and pairings for *C. aurita* in captivity. Using SPARKS and PMX software it showed that, with adequate management, the *ex situ* population is expected to reach 60 individuals in 3 years, kept in at least 5 institutions, with the formation and maintenance of at least 12 reproductive pairs. To maintain a genetically healthy population in captivity, the target number for the population was identified to be 350-400 individuals. The program therefore requires more institutions to join, with the appropriate technical capacity and long-term commitment to hold the species.

Recently, an important political initiative was approved by the office of the Environmental Secretary of São Paulo. This was the legislative proposal to establish reproductive restrictions for non-native *Callithrix* legally held in captivity in the state, in order to reduce surplus legally-captive marmosets and reduce undue releases of invasive *Callithrix* into the wild.

Considering that more than 83% of the Atlantic forest comprises fragments smaller than 50 ha (Ribeiro *et al.* 2009), it is suggested that most remaining *C. aurita* sub-populations number less than 1,000 individuals (Rylands 2008), and populations are both fragmented and isolated (Norris *et al.* 2011; Carvalho *et al.* 2018). The minimum viable population size to ensure the long-term survival of a species is thought to be 3,000-5,000 individuals (Traill *et al.* 2007). Based on density estimates and review of the literature, no sub-population of *C. aurita* can be considered viable in the long term (Norris *et al.* 2011).

In this context, the conservation of *C. aurita* will depend on several actions. Firstly, surveys are required to fully comprehend how the species is being affected by invasive congeneric marmosets and yellow fever. Secondly, effective techniques

and protocols must be developed to control the invasive *Callithrix* populations. Thirdly, monitoring and management is required within a meta-population framework, with individual movements facilitated by human interventions. Continued legal protection of the species and its habitats must be guaranteed, and finally, the involvement of more institutions and social awareness towards *C. aurita* conservation must be promoted.

In addition to the scientific and environmental communities, the general public will also be indispensable in *C. aurita* conservation. For example, we have received important records of the occurrence of *C. aurita* and invasive marmosets via social media, cell phone applications and birdwatchers. Such information highlights the notion that efficient communication with the general public and the involvement of citizen-science strategies is an important conservation tool. Accordingly, one future priority of the *C. aurita* Conservation Plan will be to amplify the outreach efforts to educate and communicate with the public, and to build upon current achievements, motivating local communities to commit to the conservation of native species.

The first steps of raising awareness for *C. aurita*'s plight and development of a conservation plan have been taken, but future conservation efforts will require partnerships with other researchers and institutions in order to synergise efforts to face the various challenges of saving *C. aurita* as an evolutionarily and ecologically unique Neotropical primate.

PIED TAMARIN

Saguinus bicolor Spix, 1823

Brazil
(2018)

Marcelo Gordo, Diogo Lagroteria, Fábio Röhe, Leandro Jerusalinsky, Renata B. de Azevedo, Marcelo D. Vidal, Tomas Hrbek, Izeni P. Farias & Anthony B. Rylands

The distinctive pied tamarin, *Saguinus bicolor*, a member of the family Callitrichidae, has a black, hairless face, ears and crown, contrasting with white fur on the back of the head, the mantle, the chest and arms. The back, flanks, abdomen, and legs have pale (sometimes quite dark) greyish brown fur, and the tail and outer surface of the thighs are variably a more reddish brown or have reddish tinges (Hershkovitz 1977; Egler 1986; Gordo *et al.* 2008, 2017). It weighs between 450 and 550 g, occasionally reaching 600 g. The length of the head and body is 28–32 cm, and the long, thin tail is 38–42 cm (Gordo 2008; Gordo *et al.* 2017). With its slim body and claw-like nails, the pied tamarin can move with great agility through thick vegetation, and even climb broad vertical trunks. The sexes are the same in their appearance.

Like other tamarins, *S. bicolor* gives birth to twins once a year, rarely twice (Gordo 2012). Gestation ranges from 180 to 219 days (Hershkovitz 1977; Egler 1992; Baker *et al.* 2009). All group members carry the infants for the few weeks after birth until they become independent. They reach sexual maturity at about two years of age (Gordo 2008). Group sizes range from 2 to 13 (Gordo 2012). *Saguinus bicolor* is diurnal, and groups are extremely territorial. It is found in dense primary forest, secondary forest and *campinarana* (forest on white sand). Its diet includes ripe fruits, small animal prey – invertebrates and small vertebrates – eggs, tree exudates (gums) and nectar (Egler 1986, 1991, 1992; Gordo 2008; Gordo *et al.* 2017).

An endemic primate of the Brazilian Amazon, its geographic distribution is approximately 7,500

km², with a large portion of this taken up by the city of Manaus, capital of the state of Amazonas, and its metropolitan region. The adjacent regions suffer strong anthropogenic effects because of urban expansion, roads, colonisation, agriculture and cattle ranching (Emmons 1990; Röhe 2006; Gordo *et al.* 2013; Coelho *et al.* 2017, 2018). Population densities of the pied tamarin are low throughout their range. Gordo (2012; Gordo *et al.* 2017) has recorded densities of 1 group/km² in extensive tracts of mature forest, and slightly higher numbers, about 2 groups/km², in isolated forest patches in and near the city of Manaus. These fragmented areas, despite having higher densities, support small populations that are not viable in the medium to long term (Gordo 2012; Campos *et al.* 2017). In urban areas, pied tamarins are run over, electrocuted when using power lines, attacked by cats and dogs, captured as pets and generally mistreated (Gordo 2012; Gordo *et al.* 2013, 2017).

Pied tamarins are run over, electrocuted when using power lines, attacked by cats and dogs, captured as pets and generally mistreated.

Away from urban areas, the pied tamarin is threatened by deforestation, habitat degradation and fragmentation, and displacement by encroaching red-handed (or Midas) tamarins, *Saguinus midas*, a closely related species which otherwise occurs over a large part of the Guiana Shield, in Brazil, French Guiana, Suriname and Guyana (Hershkovitz 1977; Röhe 2006; Gordo *et al.* 2017). Interspecific interactions and competition between *S. bicolor* and *S. midas* were reported by Ayres *et al.* (1980, 1982) and Egler (1983) and studied later in detail by Röhe (2006). The documented, apparent displacement of the pied tamarin by red-handed tamarins contributed



to its classification as Critically Endangered (CR) (Vidal *et al.* 2018) on the Brazilian List of Threatened Wildlife (Brazil, ICMBio MMA 2018), and, while still pending review and confirmation by the Species Survival Commission (SSC), it has also been ranked as CR by the IUCN SSC Primate Specialist Group.

The rapid demographic expansion of *S. midas* at the expense of *S. bicolor* is very worrying. Genetic studies, however, have demonstrated that in addition to genetic bottlenecks observed in urban fragments, *S. bicolor* has been experiencing population declines for some 10,000 years (Farias *et al.* 2015). Hybrids, which have a predominantly *S. midas* phenotype, have been recorded in the contact zone of the two species (Jeferson Oliveira pers. obs. 2014; Diogo Lagroteria pers. obs. 2016). The southernmost part of the range of *S. bicolor*, which in principle could serve as a natural refuge for the species, encompasses precisely the urban area of Manaus and its zone of expansion.

Another threat is disease. Little is known about the effects of parasites and pathogens in free-living or even captive animals, something that has been of major concern for *in situ* and *ex situ* conservation measures (Baker *et al.* 2009; Brazil, ICMBio, MMA 2011; Jerusalinsky *et al.* 2017; Lagroteria *et al.* 2017a). This concern has stimulated upcoming research (Maia da Silva *et al.* 2008; Solorio *et al.* 2015). Current projects are investigating the relation of this primate to the Zika, Dengue and Chikungunha viruses, filariasis, and digestive tract parasites.

A key initiative for the conservation of the species was the creation of the action plan *Plano de Ação Nacional para a Conservação do Sauim-de-Coleira (National Action Plan for the Conservation of the Pied Tamarin)* in 2011 (executive summary) and 2017 (the final document), coordinated by the National Center for Research and Conservation of Brazilian Primates (ICMBio) (Brazil, ICMBio, MMA 2011; Jerusalinsky *et al.* 2017). An environmental education program linked to the action plan was implemented with great success, disseminating an understanding of the threatened status of the species, promoting the adoption of the species as the symbol of Manaus, and also cultivating new allies for conservation and research (Lagroteria *et al.* 2017b). The *ex situ* management program is strongly supported by European zoos and a few North American zoos, and participation from

Brazilian institutions is growing (Baker *et al.* 2009; Lagroteria *et al.* 2017b). The financial support provided by the European and US zoos for *in situ* and *ex situ* conservation projects has been, and remains, extremely important for pied tamarin conservation efforts.

Considering that there are currently just two protected areas for the species – one municipal and one state, both located in urban areas, and each smaller than 50 ha – the creation of a reserve that could support a viable population is essential. A concrete output of the action plan was two formalized proposals for protected areas: one an environmental protection area (APA) of the government of the state of Amazonas, the other a federal biological reserve of about 16,000 ha in the east of the pied tamarin's range. The proposal for the APA was accepted, and the APA Sauim-de-Manaus, 1,050 ha, was established in June 2018.

Recent research and conservation measures have also taken a positive turn with the participation of the local population (Santos *et al.* 2017a, 2017b), the scientific community, and diverse institutions including the Federal Public Ministry (Brazil, ICMBio, MMA 2011; Gordo 2012; Barr 2016; Campos *et al.* 2017; Coelho *et al.* 2017, 2018; Jerusalinsky *et al.* 2017) resulting in the reforestation of degraded areas and the creation of ecological corridors vital for the maintenance and connectivity of viable populations in the more urbanized areas.

ECUADORIAN WHITE-FRONTED CAPUCHIN

Cebus aequatorialis J.A. Allen, 1914

Ecuador, Peru
(2018)

Stella de la Torre, Fanny Cornejo, Laura Cervera & María Fernanda Solórzano

The Ecuadorian white-fronted capuchin *Cebus aequatorialis* is a Critically Endangered (Tirira 2011; Cornejo and de la Torre 2015) primate found in western Ecuador and northwestern Peru from 0 to 2000 m asl (Cornejo and de la Torre 2015; Rylands *et al.* 2013; Tirira 2017). At the north of its range, most records now are south of the Guayllabamba River, and its southernmost distribution appears to be the Cerros de Amotape National Park in Peru (Cornejo and de la Torre 2015; Cervera *et al.* 2018; Tirira *et al.* 2018). The species is included in CITES Appendix 2 and it is illegal to hunt or trade *C. aequatorialis* in both Ecuador (Tirira 2011) and Peru (SERFOR 2015).

Cebus aequatorialis is a medium-sized monkey (body length: 35-51 cm, tail: 39.5-50 cm) with males slightly larger and heavier than females (Rylands *et al.* 2013). There is considerable individual variation in fur color and length. The upperparts, from the nape to the back, are usually pale cinnamon rufous, darker along the midline of the back. Front and sides of the head are pale, yellowish white, with a narrow black transverse line on the forehead forming the cap, from which a narrow median black line descends to the nose. The outsides of the limbs are similar in color to the body. Hands and feet are more brownish than the arms and legs. The ventral surface is paler than the flanks. The chest is lighter than the belly. The dorsal surface of the tail is dull wood-brown and is darker than the body. The undersurface of the tail is paler (Rylands *et al.* 2013).

Distribution of the white-fronted capuchin has been reduced to less than 1% of its original range in the last few decades.

The species inhabits tropical and subtropical forests of the Chocó and Tumbes eco-regions (Albuja 2002; Albuja and Arcos 2007). It is diurnal and arboreal, using all forest strata including the ground of primary and secondary forests and orchards (Campos and Jack 2013; Tirira 2017). It feeds mainly on mature fruits, complementing its diet with animal prey (insects, eggs and small vertebrates). In Ecuador, about 30 different plant species are known to be part of its diet (Albuja *et al.* 2018). Groups vary in size from 5 to 20 individuals. The adult sex ratio appears to be 1:1 or mildly skewed towards females 0.8:1 (Albuja 1992; Jack and Campos 2012; Rylands *et al.* 2013). Group home range size appears to be large, at about 500 ha (Jack and Campos 2012; Rylands *et al.* 2013), but more studies are needed to confirm this estimation.

The reproductive biology of *C. aequatorialis* is unknown. In other species of the genus, sexual maturity in females occurs when they are 4-7 years old and one year later in males. However, both sexes only reach adult body size at about 15 years old (Rylands *et al.* 2013). Considering that successful reproduction usually occurs only when animals have attained adult body size, the generation time is estimated to be about 15-16 years (Tirira *et al.* 2018).

The main threats to *C. aequatorialis* are forest loss and fragmentation, which have been particularly severe in western Ecuador, reducing its habitat. About 70% of the original forest cover in this



region has been converted to other uses, mainly agriculture and ranching (Cervera *et al.* 2018; Ecuador 2012; Sierra 2013; Gonzalez-Jaramillo 2016). It is estimated that the species distribution has been reduced to less than 1% of its original range in the last few decades (Albuja and Arcos 2007), and different modelling methods have indicated that only 5,000 km² (Campos and Jack 2013) or 8,600 km² (Albuja *et al.* 2018) of suitable habitat remains. However, the presence of *C. aequatorialis* in most of this area has yet to be confirmed.

Cebus aequatorialis is considered a pest in plantations of corn (mainly), bananas, plantain and cacao, and hence is persecuted and hunted. In some areas of mangrove, local people see it as a competitor in crab hunting and persecute it. Captive animals have been observed in villages in western Ecuador and in the Huaquillas market on the Ecuador-Perú boundary (Tirira *et al.* 2018).

In Ecuador, *C. aequatorialis* has mainly been reported to occur in public and private protected areas (see below), which are the only sites with sufficient forest to support the species. In disturbed areas, i.e., most localities in Ecuador, the species is elusive, tending to flee upon sighting. In a census of four species of western Ecuadorian primates carried out from October 2016 – March 2017, in 83 localities of 13 provinces, only 13 out of 260 records (5%) were of *C. aequatorialis* (Cervera *et al.* 2018). Surveys from previous years evidenced a relatively wide variability in local abundance (Tirira *et al.* 2018). In central western Ecuador, Jack and Campos (2012) estimated densities of 2–22 ind/km² (mean 2.4 ind/km²). In central and southwestern Ecuador, Albuja and Arcos (2007) estimated densities of 3.5 and 3.9 ind/km², while in southwestern Ecuador, De la Torre *et al.* (in prep.) estimated a density of 0.1 ind/km².

Some information about the species demography and distribution in the west of Ecuador has been provided (Albuja and Arcos 2007; Jack and Campos 2012; Campos and Jack 2013), and was updated by the most recent census of western Ecuadorian primates (Cervera *et al.* 2018). Other, smaller, short-term studies have provided information about local abundance and conservation threats (Cervera *et al.* 2015; Moscoso-Silva 2013; Solórzano 2014), but the species remains poorly known in most of its potential area of distribution.

In Peru, studies on *C. aequatorialis* are scarce. It is known to occur only in government protected areas that provide a certain degree of protection. However, there is very little information on its status in these protected areas and limited capacity to monitor them. In these areas, Hurtado *et al.* (2016) reported a group size of 3–12 individuals and an encounter rate of 0.3 ind/km (based on 7 sightings during 112 km transects). Previously, group sizes of 3–5 individuals were reported in 1980 (Saavedra and Velarde 1980) and 1994 (Encarnación and Cook 1998). Improving forest connectivity along Ecuador and Peru's border is imperative to maintain the species in both countries (Hurtado *et al.* 2016).

Given the degree of fragmentation across the species' range, targeted efforts are required to better understand its relationship with humans and how the disturbed landscape affects its demography, ecology and behaviour. The Peruvian population in the protected areas can be used as a baseline for comparisons. This information is imperative for conservation planning by decision makers, so that actions such as habitat corridor creation, conservation education, and resolution of human-wildlife conflicts can be undertaken. Additionally, successful conservation efforts must include collaboration between Ecuadorian and Peruvian authorities, stakeholders, and scientists.

The species has been reported to occur in various public and private protected areas. In Ecuador (Tirira *et al.* 2018) these are: Chocó Andino de Pichincha, Parque Nacional Machalilla, Reserva Ecológica Los Ilinizas, Reserva Ecológica Mache-Chindul, Reserva Ecológica Manglares Churute, Refugio de Vida Silvestre Manglares Estuario Río Muisne, Refugio de Vida Silvestre Marino y Costera Pácoche, Área importante para las Aves Tito Santos, Bosque Protector Puyango, Bosque Protector Bellavista, Bosque Protector Buenaventura, Bosque Protector Cambugán, Bosque Protector Cerro de Hayas, Bosque Protector Cerro Blanco, Bosque Protector Jama-Coaque, Bosque Protector Jauneche, Bosque Protector La Hesperia, Bosque Protector La Otonga, Bosque Protector Lalo Loor, Bosque Protector Maquipucuna, Bosque Protector Mashpi, Bosque Protector Mindo-Nambillo, Bosque Protector Río Guajalito. In Peru, it is reported from Cerros de Amotape National Park and Tumbes National Reserve. Finally, it is also reported from Bosques de Paz, a bi-national reserve of Ecuador and Peru.

OLALLA BROTHERS' TITI MONKEY

Plecturocebus olallae Lönnberg, 1939

Bolivia
(2018)

Jesus Martinez & Robert Wallace

The Olalla brothers' titi monkey was described in 1939 (Lönnberg 1939) and is one of two primates endemic to Bolivia. The original description was made from just one specimen, but it has since been validated as a species by a series of taxonomic assessments (*Callicebus olallae*; Hershkovitz 1990; Kobayashi 1995; van Roosmalen *et al.* 2002). The last review of callicebine taxonomy updated the species name to *Plecturocebus olallae*, maintaining it as a member of the *P. donacophilus* species group (Byrne *et al.* 2016). Its extremely restricted range (267 km²) in a naturally fragmented forest threatened by deforestation, together with a falling population size (<2,000 individuals remain) make *P. olallae* a Critically Endangered species (Wallace *et al.* 2013; López-Strauss and Wallace 2015; Martinez and Wallace 2016).

No information on wild *P. olallae* populations was available for more than 60 years after its description. It was first observed in 2002 around the Río Yacuma (Felton *et al.* 2006). Subsequent distribution and demography studies showed that the population is small and occurs only in the forests around the upper part of the Río Yacuma in the western part of the Beni Department, confirming its endemism to Bolivia (Martinez and Wallace 2007, 2013; Wallace *et al.* 2013). This area corresponds to the Moxos savannah ecosystem where the landscape, sculpted by flooding regimes, consists of a grassland matrix in which gallery forest and naturally fragmented forest patches are immersed (Martinez and Wallace 2007, 2010).


No information on wild Olalla Brothers' titi monkey populations was available for more than 60 years after its description.

Plecturocebus olallae is monogamous, living in family groups of up to five individuals (Martinez and Wallace 2007, 2010). Group home range size is around 7 ha and they cover a daily distance of approximately 500 m (Martinez 2014; Martinez and Wallace 2016). Their diet is mainly frugivorous (48.5%), with leaves also being important (38.9%), while flowers, insects, and other foods are consumed in much smaller proportions (Martinez and Wallace 2016). Gestation lasts for around 4 months, but offspring are not produced every year (Martinez and Wallace 2010). The monkeys reach sexual maturity in two years, when they look

for opportunities to leave their natal groups (Martinez and Wallace 2010; Bicca-Marques and Heymann 2013).

Habitat loss is the main threat to *P. olallae*, especially considering the fragmented forest coverage that characterizes the region. Habitat loss is linked to cattle ranching, the main economic activity in the

region, in which grasslands are burned annually to promote their regeneration as pasture. Unfortunately, this technique often results in uncontrolled fires that can affect forest patches inhabited by titi monkeys. The smoke from nearby fires can result in loss of territory (Martinez and Wallace 2011), showing that alternative methods for grassland management are urgently required. The increasingly fragmented forest promotes unusual ground travel and risky displacements for titi monkeys (Martinez and Wallace 2007, 2010, 2011).



Another important risk for *P. olallae* is the ongoing improvement of the Northern Corridor, a major road that passes around 10 km from the area it inhabits (Martinez and Wallace 2007, 2010). The dirt road is currently being upgraded to an asphalt highway, which could result in more numerous and sizeable human settlements as well as an increase in the intensity of cattle ranching and agricultural activities, both of which would cause forest loss (Martinez and Wallace 2007, 2010; Wallace *et al.* 2013; Porter *et al.* 2013). Although most of the land adjacent to the road belongs to private properties, which reduces the chances of the establishment of new human settlements, encroachment is still a risk due to proximity.

Several measures have been taken for the conservation of *P. olallae*. Initially, outreach activities included talks with local authorities of the Reyes and Santa Rosa municipalities. These authorities were already interested in consolidating tourism, the second largest economic activity in the region. Information on the presence of *P. olallae* and other wildlife was very important in promoting the creation of two municipal protected areas where titi monkeys are a conservation priority (Martinez and Wallace 2010; Wallace 2013). To involve local people in *P. olallae* conservation, posters were designed and distributed to schools and public offices. From 2011 to 2012, an intensive outreach project took place, supported by the municipal authorities, consisting of talks to students in the main towns and several communities, as well as contests and fairs oriented to raise awareness and support the conservation of *P. olallae* as a unique symbol of the local natural patrimony (Martinez *et al.* 2015). These efforts were very successful because they shared information about *P. olallae* and promoted interest in biodiversity conservation.

More recently, efforts have focused on creating management plans for the Santa Rosa and Reyes municipal protected areas, Pampas del Yacuma and Los Santos Reyes, which represent the majority of the distributional range for *P. olallae*. In 2017, a management plan was developed for Pampas del Yacuma (Santa Rosa), in which *P. olallae* conservation is linked to protection of forest and other wildlife in the region (GAMSR 2016). A similar plan is being created for Los Santos Reyes (Reyes), which covers a large portion of the species' range. Currently, conservation action focuses on

reinforcing the management of the municipal protected areas and the development of conservation plans that include monitoring and protection actions (MMyA 2009, 2014). Working with the protected areas of Reyes and Santa Rosa represents the best way to work with cattle ranchers towards implementing improved grassland and cattle management to reduce negative effects on *P. olallae* and other wildlife. A second priority is to work with the National Road Authority to implement mitigation measures along the Northern Corridor improvement project, involving both municipalities and local stakeholders. The complete establishment of the municipal protected areas will also enhance protection of *P. olallae* and the areas' wildlife and landscapes, which will also help establish ecotourism as a more sustainable option for local development.

Research is still required to fill the biological and ecological knowledge gaps concerning *P. olallae*. This knowledge will aid in the development of an appropriate monitoring program for *P. olallae* populations to determine how the distinct pressures derived from human activities affect these monkeys and to develop better protection measures.

BROWN HOWLER MONKEY

Alouatta guariba Humboldt, 1812

Brazil, Argentina
(2012, 2014, 2016, 2018)

Gerson Buss, Luciana I. Oklander, Júlio César Bicca-Marques, Zelinda B. Hirano, Óscar M. Chaves, Sérgio L. Mendes, Leonardo G. Neves, Fabiano R. Melo, Anthony B. Rylands & Leandro Jerusalinsky

Alouatta guariba is endemic to the Atlantic Forest in eastern Brazil and northeastern Argentina. In the south, its range is limited by the Camaquã river basin in the state of Rio Grande do Sul (Printes *et al.* 2001) and to the north the limit is Boa Nova, southern Bahia, south of the Rio de Contas (Neves *et al.* in prep.), although it occurred north as far as the right (south) bank of the Rio Paraguaçu in the past (Gregorin 2006). The western boundary is marked by the limits of the Atlantic Forest. In Argentina, the species occurs in the province of Misiones (Agostini *et al.* 2014). Although with some uncertainty, two subspecies are recognized: the southern brown howler, *A. guariba clamitans*, and the northern brown howler, *Alouatta g. guariba*, north of the rios Jequitinhonha or Doce (Rylands *et al.* 2000; Glander 2013).

The brown howler is a folivore-frugivore, including more or less fruit in its diet according to seasonal availability (Neville *et al.* 1988; Chaves and Bicca-Marques 2013). As such, brown howlers are important seed dispersers for numerous plant species (Chaves *et al.* 2018). Home range size varies between study sites but averages 13 ha (Fortes *et al.* 2015). Ranges of 15 groups studied varied from 1.8 to 33 ha (Miranda and Passos 2011). Day range varies from 50 m to 1,677 m (Fortes *et al.* 2015). Groups average 4 to 6 individuals, but can be as large as 13 (Jardim 2005; Miranda and Passos 2005; Ingberman *et al.* 2009). Unimale-unifemale and multimale-multifemale groups have been reported (Glander 2013). The size of an adult male is 50–60 cm (head-body) and 52–67 cm (tail), while an adult female is 44–54

cm (head-body) and 48–57 cm (tail). Adult males weigh 5.3–7.2 kg and adult females weigh 4.1–5.0 kg (Glander 2013). Longevity is estimated at 15–20 years (Strier 2004). Females have single offspring, with an interbirth interval of 9–22 months (Strier *et al.* 2001).

As for all the Atlantic Forest primates, the brown howler has suffered extensive habitat loss since European arrival in South America more than five centuries ago. During the colonization process, the forest cover was broadly devastated due to exploitation of natural resources (e.g., Brazil wood, gold), extensive agriculture (e.g., sugar cane, coffee) and cattle ranching. This region today concentrates around 70% of the Brazilian population (~150 million people) and the principal capital cities, with corresponding industrial activity and urbanization (Scarano and Ceotto 2015). In Brazil, the Atlantic Forest has been reduced to 11.7% of its original coverage (Ribeiro *et al.* 2009). The remaining

forest is immensely fragmented into hundreds of thousands of patches, of which the great majority are 50 ha or less (Ribeiro *et al.* 2009), hence unsuitable to support viable populations in the long term. Being one of the largest primates in the Atlantic Forest, the species has been extensively hunted, and also suffers to some extent from the pet trade.

Disease epidemics are an additional and very serious threat. Howlers are highly susceptible to yellow fever, and two recent outbreaks (2008/2009, 2016/2018), have severely affected

Howlers are highly susceptible to yellow fever, and two recent outbreaks have severely affected their numbers.



their numbers throughout the Atlantic Forest (Holzmann *et al.* 2010; Almeida *et al.* 2012; Bicca-Marques *et al.* 2017). Due to misinformation and the dissemination of the fear that humans could be infected directly through contact or proximity with monkeys, howlers were persecuted, with many injured and killed during the outbreaks (Bicca-Marques *et al.* 2017). In the next few decades, pathogen exposure could act synergistically with other threats such as habitat loss, putting populations at high extinction risk.

Southern brown howlers (*Alouatta guariba clamitans*) occur in lowland forests along Brazil's coast, as well as in higher elevation submontane and montane forests and seasonal semi-deciduous forests inland (Bicca-Marques *et al.* 2018). In southern Brazil and northeast Argentina, they also occupy a transition of mixed Upper Paraná Atlantic Forest and Araucaria Moist Forest (Miranda and Passos 2005; Agostini *et al.* 2014). Aguiar *et al.* (2007) recorded the species in periodically flooded and semi-deciduous forests in the Paraná river floodplains.

The primary threats are widespread forest loss and fragmentation throughout the subspecies' range due to logging, agriculture and cattle-ranching (Bicca-Marques *et al.* 2018). Attacks by domestic dogs, traffic accidents, and electrocution are serious threats to howlers living close to urban areas (Printes *et al.* 2010; Chaves *et al.* in prep.). The design and implementation of conservation strategies for the southern brown howlers in urban and suburban regions are crucial for the long-term survival of these animals (Jerusalinsky *et al.* 2010).

Although some local population census data are available for Brazil, the total remaining population is unknown, but certainly declining. In Argentina, the situation is even worse; only a few populations persist with no more than 20–50 adult individuals (Agostini *et al.* 2014).

The southern brown howler is listed as Vulnerable on the Brazilian list of threatened fauna (Brazil, MMA 2014) and the IUCN Red List (Buss *et al.* in prep), but may be a candidate for the Endangered category after the 2016–2018 yellow fever outbreak (Bicca-Marques *et al.* 2017). It is considered Critically Endangered in Argentina (Agostini *et al.* 2012).

The northern brown howler monkey (*Alouatta guariba guariba*) inhabits lowland, submontane and montane Brazilian Atlantic forest. It has a considerably more restricted range than *A. g. clamitans* and is classified as Critically Endangered both in the Brazilian list of threatened fauna (Brazil MMA 2014) and the IUCN Red List (Mendes *et al.* 2008). It has been listed as one of the world's 25 most endangered primates since 2012 (Neves *et al.* 2017). Adding the locations in the lower reaches of the Jequitinhonha basin reported by Rylands *et al.* (1988) and the small populations of *A. g. guariba* discovered in the last few years, the total population is unlikely to sum more than 250 mature individuals, and no subpopulation is believed to have more than 50 mature individuals (Neves *et al.* 2018). Overall, the main threats to the wild populations of this subspecies are habitat fragmentation, hunting, and the very small sizes of the scattered populations (Neves *et al.* 2017).

There are protected areas in the northern brown howler's range in the state of Bahia and northeastern Minas Gerais, all created since 1980. Nevertheless, the only strictly protected area where the species has been confirmed is the Mata Escura Biological Reserve (51,046 ha), just north of the middle Rio Jequitinhonha (Melo 2005).

The two subspecies of *Alouatta guariba* are included in the *Brazilian National Action Plan for Conservation of the Atlantic Rainforest Primates and Maned Sloth* (Brazil MMA 2018), and will be part of the Primate Conservation Action Plan of Argentina which will be produced in 2019. These plans provide measures to identify important areas for conservation in order to (a) restore, maintain and increase habitat and its connectivity, (b) mitigate the impact of roads and power lines, and (c) assess and mitigate the impact of epizootics on the species.

CENTRAL AMERICAN SPIDER MONKEY

Ateles geoffroyi Kuhl, 1820

Mexico, Guatemala, Nicaragua, Honduras, El Salvador, Costa Rica, Panama (2016, 2018)

Pedro G. Méndez-Carvajal, Melissa E. Rodríguez, Gilberto Pozo Montuy, Óscar M. Chaves, Gabriela Ponce, Bonarge A. Rodríguez-Beitia & Héctor Portillo-Reyes

Central American spider monkeys *Ateles geoffroyi* are distributed in Mexico, Guatemala, Nicaragua, Honduras, El Salvador, Costa Rica and Panama (Rylands *et al.* 2006). Because of habitat loss and fragmentation, severe hunting pressure, and the pet trade, they are considered Critically Endangered (Smith 2005; Cuarón *et al.* 2008). *Ateles geoffroyi* was subdivided into nine subspecies by Kellogg and Goldman (1944). Since then three have been synonymized (*A. g. pan*, *A. g. panamensis*, and *A. g. yucatanensis*) (Rylands *et al.* 2006), and Groves (2005) considered *A. g. azuerensis* to be a synonym of *A. g. ornatus*.

Recent taxonomic studies using mitochondrial DNA have validated other subspecies which are mentioned here, but more information on identification and sample locations is needed to corroborate these conclusions (Morales-Jiménez *et al.* 2015; Ruiz-García *et al.* 2016).

The genus *Ateles* has long been considered the most threatened in the Neotropics (Mittermeier *et al.* 1989). *A. geoffroyi* has a long gestation period (226-232 days) compared to other Atelidae, such as *Alouatta*, *Brachyteles* and *Lagothrix* (Campbell 2000). They also spend a larger proportion of their time foraging compared to other Central American primate species (Chapman *et al.* 1989), with a major dietary requirement of 69-91% fruit (Campbell 2000). With their highly frugivorous diet, spider monkeys need large expanses of

forest and are less able to adapt to fragmentation than *Alouatta* (Méndez-Carvajal 2013). In addition to its ecological requirements, it is one of the main species hunted in indigenous regions (Smith 2005). This species has a large range compared to other non-human primates in the Mesoamerican region, but it is threatened by high rates of deforestation. Narco-effect rates (deforestation related to illegal drug trade) are 20-60% per year, sometimes also affecting natural parks and reserves (McGrath 2014).

Ateles geoffroyi azuerensis (CR) was initially described as *Ateles azuerensis* Bole, 1937, and was studied for the first time in La Vaca, Coto Region, in the Chiriqui Province (Carpenter 1935). The actual distribution and total population have been assessed by the Fundación Pro-Conservación de los Primates Panameños (FCPP), a Panamanian NGO that has been monitoring this primate since 2001. *Ateles g. azuerensis* has been extirpated in Chiriqui

Province, west and north Veraguas and Herrera Province. It is present now only in southwestern Veraguas and Los Santos Province, on the Azuero Peninsula, in the southern areas near the Cerro Hoya National Park, and in the fragmented landscape between Punta Duarte, La Barra, Guanico, Quema, La Tronosa Forest Reserve, La Miel and Pedasi. Only 10 subgroups and five complete groups have been detected, with a mean of 3.8 individuals/subgroup, SE ± 0.6 (range 2-7) and a mean of 12.5 individuals/

Narco-effect rates (deforestation related to illegal drug trade) are 20-60% per year in parts of the spider monkey range.



group, SE ± 3.7 (range 10–22), with densities of 1.4 individuals/km² (for fragmented habitats), and an approximate total population of <150 individuals (Méndez-Carvajal and Ruiz-Bernard 2009; Méndez-Carvajal 2013). Conservation measures led by FCPP involve community volunteers from Azuero, environmental education and the distribution of an educational *Azuero's Primate Guide*, as well as monitoring biodiversity and surveying the Azuero Peninsula (Méndez-Carvajal *et al.* 2013).

Ateles geoffroyi frontatus (EN) ranges from northern Nicaragua to the northwestern parts of Costa Rica, including the basins of the ríos Principolca, Tuma and Uluce, and is also found in Metagalpa and the Nicaraguan highlands (Allen 1914; Rylands *et al.* 2006; Cuarón *et al.* 2008).

Ateles geoffroyi geoffroyi (CR) inhabits San Juan del Norte, Martina Bay and southeastern Nicaragua, and the population probably extends into northern Costa Rica (Rylands *et al.* 2006).

The subspecies *Ateles geoffroyi grisescens* (DD) was reported by Kellogg and Goldman (1944) from the valley of the Río Tuira, Serranía del Sapo, Pirre, Tucuti in Darien Province, Panama (Elliot 1913; Gray 1865; Sclater 1875). It also occurs in Baudó, northwestern Colombia (Rylands *et al.* 2006). Recent studies have reported that *A. g. grisescens* is no longer in its original area (Tuira River), nor in Chucanti or the Maje Mountain Chain (Méndez-Carvajal 2012). However, the presence/absence of this primate from Panama is still being studied (Méndez-Carvajal *et al.* 2016). A documentary related to the expedition to find *A. g. grisescens* has been filmed for Barbara Réthoré and Julien Chapuis from Conserv-action and NatExplorers, in support of FCPP projects and the re-discovery of this subspecies.

The natural range of *A. g. ornatus* (CR) is in Costa Rica and Panama. In Costa Rica it is known to be in the Osa Peninsula, Carara Biological Reserve, Corcovado National Park (Matamoros and Seal 2001), and Cerro Chirripo, Cantón de Pérez Zeledón, at 1700 m asl, with a density of 0.012 individuals/km² (Rodríguez-Beitia pers. obs.). In Panama, it is present on the northern side of the Caribbean coast, in the lowlands and highlands of Bocas del Toro, the northern coast of Veraguas Province, Coclé (rare in Coclé

and Donoso; Méndez-Carvajal, pers. obs.), Portobelo National Park, and San Blas mountain chain (Méndez-Carvajal *et al.* 2016). An isolated population was introduced onto Barro Colorado Island (Campbell 2000). In Panama, FCPP started a long-term monitoring project in 2010 in the San Blas mountain chain to understand the actual distribution and population densities for this subspecies (Méndez-Carvajal 2014).

Ateles geoffroyi vellerosus (CR) is present in Mexico, Belize, Guatemala, Honduras and El Salvador (Cuarón *et al.* 2008). The population density of *A. g. vellerosus* is between 2.9 individuals/km² – 9.3 individuals/km² at Montes Azules Biosphere Reserve in Marqués Comillas ejido, Chiapas, Mexico (Chaves *et al.* 2011). It also occurs in northern Veracruz, Oaxaca, Tamaulipas, Chiapas, Tabasco, Campeche, Quintana Roo and some other regions on the Yucatan Peninsula (Chaves *et al.* 2011). It occurs in densities of 2–12 individuals/km² (Pozo-Montuy *et al.* 2015). In Guatemala, it is reported at Petén, Alta Verapaz, Baja Verapaz, Izabal, Sololá, Huehuetenango and Quiché (Ponce-Santizo *et al.* 2009). It is reported in El Salvador at Chaguantique Natural Protected Area (NPA), El Tercio, El Nacascolo, Normandía NPA (Usulután Department), and a group was recently rediscovered in Olomega lagoon (San Miguel and La Unión Departments) (Pineda-Peraza *et al.* 2017). Some historical records were made in Montecristo, Cerro el Mono y Conchagua (Rodríguez-Menjivar 2007). *A. g. vellerosus* is threatened by forest fires, the pet trade, habitat fragmentation due to farming activities such as palm oil plantations, and road construction (McGrath 2014). Some conservation activities to protect this taxon include environmental education, and setting up canopy bridges to facilitate canopy connection and reduce the number of animals killed on the roads. These activities have been implemented by the Mexican Primates Regional Monitoring System led by the project Conservación de la Biodiversidad del Usumacinta A.C. since 2013 (Pozo-Montuy *et al.* 2015). The Maya Biosphere Reserve (MBR) in the north of Guatemala, with 2.2 million ha, constitutes the largest and most important habitat for the subspecies (68.6% is its original forests). Conservation actions are maintained by several organizations with the aim to preserve this important forest block in Guatemala (Ponce-Santizo *et al.* 2009).

OTHER SPECIES CONSIDERED



AFRICA GRAUER'S GORILLA

Gorilla beringei graueri Matschie, 1914
(2010, 2012, 2014, 2016)

Stuart Nixon

The Critically Endangered Grauer's gorilla is endemic to the eastern Democratic Republic of Congo (DRC) and distributed discontinuously throughout the forests east of the Lualaba River to the western Albertine Rift escarpment. Long-term insecurity since 1996 has had a devastating effect on their populations. Surveys completed in 2015 identified a catastrophic decline of 77% in just a single generation since the 1990s (Plumptre *et al.* 2016a, 2016b) primarily due to illegal hunting. Today, an estimated 3,800 Grauer's gorillas remain across their 19,700 km² range. However, bushmeat hunting remains the single largest threat to Grauer's gorilla, followed by habitat loss, disease such as ebola, and the unmitigated effects of global climate change. Targeted conservation action in priority sites will be vital to slow the further demise of this subspecies.



ASIA CRESTED MACAQUE

Macaca nigra Desmarest, 1822
(2016)

Caspian L. Johnson, Rivo Rahasia, Wulan Pusparini, Iwan Hunowu, Alfons Patandung, Andrew E. Bowkett, Harry Hilser & Daphne Kerhoas

Macaca nigra is the only Critically Endangered species of Sulawesi's seven endemic macaques (Fooden 1969). A forest-dwelling macaque, *M. nigra* is endemic to the northern peninsula of Sulawesi from the tip to the Onggak-Dumoga River, where it meets the boundary with *M. nigrescens* (Johnson *et al.* 2019). Dominant threats to the species include hunting and habitat loss, caused by agricultural expansion, human-induced fires (for cattle grazing) and illegal logging. Between 2001 and 2017, 9% of tree cover was lost in the species' range (Hansen *et al.* 2013). Continued habitat loss outside protected areas is therefore expected to have severe implications for the species if left unchecked. Confounding this is the fragmentation of remaining habitat. *Macaca nigra* is being hunted at a potentially unsustainable rate.



ASIA BORNEAN BANDED LANGUR

Presbytis chrysomelas Müller, 1838
(2018)

Andie Ang, Erik Meijaard, Vincent Nijman & Noel Rowe

The Bornean banded langur (*Presbytis chrysomelas*) is a Critically Endangered primate endemic to the northwestern corner of Borneo (Nijman *et al.* 2008). Two subspecies are recognised: *P. c. chrysomelas* and *P. c. cruciger*. In the past it could be found in protected areas such as Tanjung Datu National Park, Similajau National Park, Maludam National Park, Niah National Park, Lanjak-Entimau Wildlife Sanctuary and Samunsam Wildlife Sanctuary, which are all in Sarawak and Danau Sentarum Wildlife Sanctuary in West Kalimantan (Groves *et al.* 2013; Phillipps and Phillipps 2018). However, the most recent records of their occurrence in several of these parks date back more than a decade. While Nijman *et al.* (2008) conservatively estimated that 200-500 individuals remained, more recent research suggested this might have been an underestimate. However, habitat is declining, partially because of cash crop plantations and forest fires.



ASIA TONKIN SNUB-NOSED MONKEY

Rhinopithecus avunculus Dollman, 1912
(2000, 2002, 2004, 2006, 2008, 2010, 2012, 2014, 2016)

Tilo Nadler

The Tonkin snub-nosed monkey, *Rhinopithecus avunculus*, is a Critically Endangered species endemic to Vietnam, confined to a few areas of the far northwest (Nadler and Brockman 2014). Its distribution has been drastically reduced in recent decades due to massive deforestation and intensive hunting. As a result, the population has become severely fragmented (Nadler *et al.* 2003; Nadler and Brockman 2014). The species was thought to be extinct until its rediscovery near the town of Na Hang, Tuyen Quang Province in 1989. Conservation activities there were unsuccessful and it is likely now extirpated. A population of 20 to 40 individuals was estimated for Cham Chu Nature Reserve, Tuyen Quang Province, but subsequent surveys provided no sightings. In 2001, a population was discovered in Khau Ca, close to Du Gia Nature Reserve, Ha Giang Province. A census in 2015 confirmed 125-130 individuals. Subsequently the area was declared as Tonkin snub-nosed monkey Species/Habitat Conservation Area. It is the only population which is not immediately threatened. In 2007, a population of about 20 Tonkin snub-nosed monkeys was discovered in Tung Vai, Ha Giang Province, close to the border with China. This population is threatened through hunting and habitat loss (Le and Covert 2010). The total population of the Tonkin snub-nosed monkey is currently believed to be fewer than 250 individuals (Xuan *et al.* 2008).



ASIA SIAU ISLAND TARSIER

Tarsius tumpara Shekelle *et al.*, 2008
(2006, 2008, 2010)

Myron Shekelle & Agus Salim

The Siau Island tarsier (*Tarsius tumpara*) is Critically Endangered and first attracted the notice of the scientific community when it was added to the list of the 25 Most Endangered Primates in 2006, on which it remained until 2012 (Shekelle and Salim 2011). The main threats to *T. tumpara* are its very restricted range, limited remaining habitat in its range, and hunting. The species is known only from Siau Island, Sulawesi, Indonesia (116 km²), on which Mt. Karengetang, a highly active volcano, comprises 55% of the land. Satellite imagery and field surveys indicate that no primary forest remains on the island and that secondary forest might make up as little as 17%. The island lacks protected areas, other than some green areas set aside as water catchments. Most alarmingly, tarsiers are regularly eaten as a snack food, with as many as 10 individuals consumed at one sitting (Shekelle and Salim 2009). It is possible that *T. tumpara* exists on other islands nearby to Siau, which would bring its total extent of occurrence to 125 km² (Shekelle and Salim 2009). *Badan Pusat Statistik Kabupaten Kepulauan Sangihe* (2018) reports the total population for the islands of Siau, Tagulandang, and Biaro in 2017 as 65,976. We expect about 66% (or 43,500) to be from Siau.

APPENDIX

Table 1. The World's 25 Most Endangered Primates: 2018–2020.

MADAGASCAR		
<i>Microcebus manitatra</i>	Bemanasy mouse lemur	Madagascar
<i>Hapalemur alaotrensis</i>	Lake Alaotra gentle lemur	Madagascar
<i>Lepilemur jamesorum</i>	James' sportive lemur	Madagascar
<i>Indri indri</i>	Indri	Madagascar
<i>Daubentonia madagascariensis</i>	Aye-aye	Madagascar
AFRICA		
<i>Paragalago rondoensis</i>	Rondo dwarf galago	Tanzania
<i>Cercopithecus roloway</i>	Roloway monkey	Côte d'Ivoire, Ghana
<i>Rungwecebus kipunji</i>	Kipunji	Tanzania
<i>Colobus vellerosus</i>	White-thighed colobus	Côte d'Ivoire, Ghana, Togo, Benin, possibly Nigeria
<i>Piliocolobus epieni</i>	Niger Delta red colobus	Nigeria
<i>Piliocolobus rufomitratu</i>	Tana River red colobus	Kenya
<i>Pan troglodytes verus</i>	Western chimpanzee	Côte d'Ivoire, Ghana, Guinea-Bissau, Liberia, Mali, Republic of Guinea, Senegal, Sierra Leone
ASIA		
<i>Nycticebus javanicus</i>	Javan slow loris	Indonesia
<i>Simias concolor</i>	Pig-tailed snub-nose langur	Indonesia
<i>Trachypithecus poliocephalus</i>	Golden-headed langur or Cat Ba langur	Vietnam
<i>Trachypithecus geei</i>	Golden langur	India, Bhutan
<i>Semnopithecus vetulus</i>	Purple-faced langur	Sri Lanka
<i>Hoolock tianxing</i>	Skywalker hoolock gibbon	China, Myanmar
<i>Pongo tapanuliensis</i>	Tapanuli orangutan	Indonesia
NEOTROPICS		
<i>Callithrix aurita</i>	Buffy-tufted-ear marmoset	Brazil
<i>Saguinus bicolor</i>	Pied tamarin	Brazil
<i>Cebus aequatorialis</i>	Ecuadorian white-fronted capuchin	Ecuador, Peru
<i>Plecturocebus olallae</i>	Olalla Brothers' titi monkey	Bolivia
<i>Alouatta guariba</i>	Brown howler monkey	Brazil, Argentina
<i>Ateles geoffroyi</i>	Central American spider monkey	Mexico, Guatemala, Nicaragua, Honduras, El Salvador, Costa Rica, Panama

Table 2. Primate species included on the 2016–2018 list that were removed from the 2018–2020 list.

MADAGASCAR		
<i>Lemur catta</i>	Ring-tailed lemur	Madagascar
<i>Microcebus gerpi</i>	Gerp's mouse lemur	Madagascar
<i>Propithecus perrieri</i>	Perrier's sifaka	Madagascar
AFRICA		
<i>Paragalago orinus</i>	Mountain galago	Tanzania
<i>Gorilla beringei graueri</i>	Grauer's gorilla	Democratic Republic of Congo
ASIA		
<i>Macaca nigra</i>	Crested macaque	Indonesia
<i>Nomascus hainanus</i>	Hainan gibbon	China
<i>Pongo pygmaeus</i>	Bornean orangutan	Indonesia
<i>Rhinopithecus avunculus</i>	Tonkin snub-nosed monkey	Vietnam
NEOTROPICS		
<i>Ateles fusciceps</i>	Brown-headed spider monkey	Ecuador, Colombia, Panama
<i>Cebus kaapori</i>	Ka'apor capuchin	Brazil
<i>Plecturocebus caquetensis</i>	Caquetá titi monkey	Colombia

Table 3. Primate species that were added to the 2018–2020 list. The Rondo dwarf galago, kipunji, Tana River red colobus and indri were added to the list after previously being removed. The other eight species are new to the list.

MADAGASCAR		
<i>Microcebus manitatra</i>	Bemanasy mouse lemur	Madagascar
<i>Indri indri</i>	Indri	Madagascar
AFRICA		
<i>Paragalago rondoensis</i>	Rondo dwarf galago	Tanzania
<i>Rungwecebus kipunji</i>	Kipunji	Tanzania
<i>Piliocolobus rufomitratu</i>	Tana River red colobus	Kenya
<i>Pan troglodytes verus</i>	Western chimpanzee	Côte d'Ivoire, Ghana, Guinea-Bissau, Liberia, Mali, Republic of Guinea, Senegal, Sierra Leone
ASIA		
<i>Hoolock tianxing</i>	Gaoligong hoolock gibbon	China, Myanmar
<i>Pongo tapanuliensis</i>	Tapanuli orangutan	Indonesia
NEOTROPICS		
<i>Callithrix aurita</i>	Buffy-tufted-ear marmoset	Brazil
<i>Saguinus bicolor</i>	Pied tamarin	Brazil
<i>Cebus aequatorialis</i>	Ecuadorian white-fronted capuchin	Ecuador, Peru
<i>Plecturocebus olallae</i>	Olalla Brothers' titi monkey	Bolivia

EDITORS' ADDRESSES

Schwitzer, Christoph

Bristol Zoological Society, c/o Bristol Zoo Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: cschwitzer@bristolzoo.org.uk

Mittermeier, Russell A.

Global Wildlife Conservation, c/o WeWork, 718 7th Street, N.W., Washington, DC 20001, USA
E-mail: rmittermeier@globalwildlife.org

Rylands, Anthony B.

Global Wildlife Conservation, c/o WeWork, 718 7th Street, N.W., Washington, DC 20001, USA
E-mail: arylands@conservation.org

Chiozza, Federica

Global Mammal Assessment Program, Department of Biology and Biotechnology, Sapienza Università di Roma, Viale dell'Università 32, 00185 Roma, Italy
E-mail: federica.chiozza@uniroma1.it

Williamson, Elizabeth A.

School of Natural Sciences, University of Stirling, Stirling FK9 4LA, UK
E-mail: e.a.williamson@stir.ac.uk

Byler, Dirck

Global Wildlife Conservation, c/o WeWork, 718 7th Street, N.W., Washington, DC 20001, US
E-mail: dbyler@globalwildlife.org

Wich, Serge

Natural Sciences and Psychology, Faculty of Science, Liverpool John Moores University, Tom Reilly Building, Byrom Street, Liverpool L3 3AF, UK
E-mail: sergewich@gmail.org

Humle, Tatyana

Senior Lecturer, University of Kent, Canterbury, Kent, CT2 7NR, UK
E-mail: t.humle@kent.ac.uk

Johnson, Caspian

Bristol Zoological Society, c/o Bristol Zoo Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: cjohnson@bristolzoo.org.uk

Mynott, Holly

Bristol Zoological Society, c/o Bristol Zoo Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: top25primates@gmail.com

McCabe, Gráinne

Bristol Zoological Society, c/o Bristol Zoo Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: gmccabe@bristolzoo.org.uk

CONTRIBUTORS' ADDRESSES

Allen, Lara V.

Centre for Global Equality
Cambridge, CB2 1RR, UK
Email: lara.allen@centreforglobalequality.org

Andriamisedra, Tsarafilamatra Rotsi

GERP, Logement 34, Cité des Professeurs, Fort Duchesnes, Antananarivo 101, Madagascar
E-mail: andriamisedrar@gmail.com

Andriamalala, Yves Rostant

Anthropobiologie et Développement Durable, Faculté des Sciences, Université d'Antananarivo, PO Box 906, Antananarivo 101, Madagascar
E-mail: rossrajh@gmail.com

Andrianandrasana, Herizo T.

Department of Zoology, University of Oxford, UK
E-mail: herizo.andrianandrasana@zoo.ox.ac.uk

Ang, Andie

Department of Anthropology, University of Colorado, 1350 Pleasant St, 233 UCB, Boulder, CO 80309-0233
E-mail: andie.ang@colorado.edu

Bhattacharjee, P.C.

Department of Zoology, Gauhati University, Guwahati, 781014, Assam, India
E-mail: bhattapc@gmail.com

Bicca-Marques, Júlio César

Laboratório de Primatologia, Escola de Ciências, PUCRS, Av. Ipiranga, 6681 Prédio 12A, Porto Alegre, RS 90619-900, Brazil
E-mail: jcbicca@pucrs.br

Boesch, Christophe

Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6 04103 Leipzig, Germany
E-mail: boesch@eva.mpg.de

Bonadonna, Giovanna

Department of Biology, Monmouth University, United States
E-mail: gbonadon@monmouth.edu

Bowkett, Andrew E.

Field Conservation & Research Programmes Manager, Whitley Wildlife Conservation Trust, Totnes Road, Paignton, Devon, UK
E-mail: Andrew.bowkett@paigntonzoo.org.uk

Buss, Gerson

Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio), Centro Nacional de Pesquisa e Conservação dos Primatas Brasileiros (CPB), Praça Antenor Navarro 5. João Pessoa – PB, 58010-480, Brazil
E-mail: gbuss.icmbio@gmail.com

Butynski, Thomas M.

Sustainability Centre Eastern Africa, Lolldaiga Hills Research Programme, P. O. Box 149 Nanyuki 10400, Kenya
E-mail: tbutynski@aol.com

Carvalho, Rodrigo S.

Programa de Educação Ambiental (PREA) Av. Senhor dos Passos, 1155, Juiz de Fora, MG, Brasil - 36037-490
E-mail: rodrigosallesdecarvalho@gmail.com

Cervera, Laura

Universidad Laica Eloy Alfaro de Manabí, Manta, Ecuador
E-mail: laura.cervera24@gmail.com

Chaves, Óscar M.

Laboratório de Primatologia, Escola de Ciências, PUCRS, Av. Ipiranga, 6681 Prédio 12^a, Porto Alegre, RS 90619-900, Brazil
E-mail: ochaba@gmail.com

Chetry, Dilip

Aaranyak, 13 Toyab Ali Bye Lane, Bishnu Rabha Path, Beltola Tinali, PO: Beltola, Guwahati 781028, Assam, India
E-mail: dilip@aaranyak.org

Chetry, Rekha

Department of Zoology, Jawaharlal Nehru College, Boko, Kamrup, 781123, Assam, India
E-mail: chetryrekha@gmail.com

Cheyne, Susan M.

Department of Social Sciences, Oxford Brookes University, Oxford, UK
E-mail: s.cheyne@brookes.ac.uk

Cornejo, Fanny

ICTE, Stony Brook University, Circle Road, SBS Bldg N-221, Stony Brook, NY 11794-4364, USA
E-mail: fanny.cornejo@stonybrook.edu

CONTRIBUTORS' ADDRESSES

Davenport, Tim R. B.

Wildlife Conservation Society, P.O. Box 922,
Zanzibar, Tanzania
E-mail: tdavenport@wcs.org

de Azevedo, Renata B.

Instituto Chico Mendes de Conservação da
Biodiversidade/ICMBio, Centro Nacional de
Pesquisa e Conservação de Primatas Brasileiros/
CPB, João Pessoa/PB, Brazil
E-mail: renata.azevedo@icmbio.gov.br

De Gregorio, Chiara

Department of Life Sciences and Systems
Biology, Università degli Studi di Torino, Via
Verdi, 8 - 10124 Torino, P.I. 02099550010, C.F.
80088230018, Italy
E-mail: chiara.degregorio@unito.it

de Jong, Yvonne

Lolldaiga Hills Research Programme,
Kenya
Email: yvonne@wildsolutions.nl

de la Torre, Stella

University of Wisconsin, Madison, USA
E-mail: sdeltorre@usfq.edu.ec

De Luca, Daniela W.

Wildlife Conservation Society, PO Box 1475,
Mbeya, Tanzania
E-mail: ddeluca@wcs.org

Dempsey, Andrea

West African Primate Conservation Action, PO
Box GP1319, Accra, Ghana
E-mail: andrea.dempsey@wapca.org

Donati, Giuseppe

Department of Social Sciences, Oxford Brookes
University, Oxford, UK
E-mail: gdonati@brookes.ac.uk

Dreyer, William

GERP, Logement 34, Cité des Professeurs, Fort
Duchesnes, Antananarivo 101, Madagascar
E-mail: william.dreyer@mail.mcgill.ca

Dunn, Nicholas J.

Monkey World, out Hill, Longthorns, Wareham,
UK BH20 6HH
Email: Nic@monkeyworld.org

Fan, Pengfei

School of Life Sciences, Sun Yat-sen University,
Guangzhou, China
E-mail: fanpf1981@gmail.com

Farias, Izeni P.

Universidade Federal do Amazonas, Instituto de
Ciências Biológicas, Departamento de Biologia,
Manaus, Brazil
E-mail: izeni@evoamazon.net

Fei, Hanlan

School of Life Sciences, Sun Yat-sen University,
Guangzhou, China
E-mail: fanhj5@mail.sysu.edu.cn

Fransen, Sally J.

Programa de Educação Ambiental (PREA)
Av. Senhor dos Passos, 1155, Juiz de Fora, MG,
Brasil - 36037-490
Email: pheonix.07@hotmail.co.uk

Fredriksson, Gabriella

Institute of Systematics and Population Biology,
University of Amsterdam, Netherlands
E-mail: gabriella.fredriksson@gmail.com

Gamba, Marco

Università degli Studi di Torino
Via Verdi, 8 - 10124 Torino, Italy
E-mail: marco.gamba@unito.it

Giacoma, Cristina

Department of Life Sciences and Systems Biology,
Via Accademia Albertina 13, 10123 Torino, Italy
E-mail: cristina.giacoma@unito.it

Gonedelé Bi, Sery

Departement of Genetics, Cocody University,
Côte d'Ivoire
E-mail: gonedelebi@yahoo.fr

Gordo, Marcelo

Projeto Sauim-de-Coleira, Federal University of
Amazonas, Manaus, Brazil
E-mail: projetosauim@gmail.com

Hilser, Harry

Field Project Manager for Selamatkan Yaki,
Sulawesi, Indonesia
E-mail: harry@selamatkanyaki.com

Hirano, Zelinda B.

Cepesbi/ Projeto Bugio/ FURB (Universidade
Regional de Blumenau), Rua Rio de Janeiro 401,
bairro dos estados, Indaial-SC, CEP 89086793 Brazil
E-mail: zehirano@hotmail.com

Hrbek, Tomas

Laboratório de Evolução e Genética Animal—
Instituto de Ciências Biológicas, Universidade
Federal do Amazonas, Manaus, Brazil
E-mail: hrбек@evoamazon.net

Hunowu, Iwan

Wildlife Conservation Society, Northern Sulawesi,
Indonesia
E-mail: iwan6u@gmail.com

Igayara-Souza, Cláudia A.

Prefeitura Municipal de Guarulhos, Zoológico de
Guarulhos. Rua Dona Gloria Pagnoncelli, 344 Jd.
Rosa de França 07081-120 - Guarulhos, SP - Brazil
Email: claudiaigayara@uol.com.br

Ikemeh, Rachel Ashegbofe

SW/Niger Delta Forest Project, P.O. Box 3565,
900001, Garki, Abuja, Nigeria
E-mail: pi@swnigerdeltaforestproject.org

Jerusalinsky, Leandro

Instituto Chico Mendes de Conservação da
Biodiversidade (ICMBio), Centro Nacional de
Pesquisa e Conservação dos Primatas Brasileiros
(CPB), Praça Antenor Navarro 5. João Pessoa –
PB, 58010-480, Brazil
E-mail: ljerusalinsky@gmail.com

Johnson, Caspian L.

Bristol Zoological Society, c/o Bristol Zoo
Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: caspianjohnson@gmail.com

Kerhoas, Daphne

Bristol Zoological Society, c/o Bristol Zoo
Gardens, Clifton, Bristol BS8 3HA, UK
E-mail: dkerhoas@bristolzoo.org.uk

King, Juliet

Northern Rangelands Trust, Isiolo, Kenya
Email: julietking@nrt-kenya.org

Kivai, Stanislaus M.

Institute of Primate Research, P.O. Box 24481
Karen 00502, Nairobi, Kenya
E-mail: skivai@primateresearch.org

Koné, Inza

Swiss Center for Scientific Research in Côte
d'Ivoire, Yopougon, 01 BP 1303 Abidjan
E-mail: inza.kone@csrs.ci

Kühl, Hjalmar

Department of Primatology, **Max Planck Institute**
for Evolutionary Anthropology,
Deutscher Platz 6, D-04103 Leipzig, Germany
E-mail: kuehl@eva.mpg.de

Lacerda, Wagner R.

Biólogo - Mestre em Conservação e Manejo
de Recursos Naturais, Muriqui Instituto de
Biodiversidade, Cerne Consultoria Ambiental
E-mail: wr_lacerda@yahoo.com.br

Lagroteria, Diogo

DESMAN graduate, wildlife vet
E-mail: diogo.fauna@gmail.com

Leonard, Neahga

Cat Ba National Park, Cat Ba Island, Cat Hai
District, Hai Phong Province, Vietnam
E-mail: neahga.leonard@catbalangur.de

Lewis, Richard

Durrell Wildlife Conservation Trust, Lot II Y 49 J
Ampasani-malo, BP 8511 Antananarivo 101,
Madagascar
E-mail: richard.lewis@durrell.org

Louis, Jr., Edward E.

Director of Conservation Genetics, Omaha's
Henry Doorly Zoo and Aquarium, 3701 South 10th,
Omaha, NE 68107, USA
General Director, Madagascar Biodiversity
Partnership, NGO, VO 12 Bis A, Manakambahiny,
Antananarivo 101, Madagascar
E-mail: edlo@omahazoo.com

Loyola, Laura C.

Spatial Sciences Institute 3616 Trousdale Pkwy,
AHF B55 University of Southern California Los
Angeles, CA 90089
E-mail: loyola@usc.edu

Machaga, Sophy J.

Wildlife Conservation Society, PO Box 1475,
Mbeya, Tanzania
E-mail: smachaga@wcs.org

CONTRIBUTORS' ADDRESSES

Martinez, Jesus

Wildlife Conservation Society, Greater Madidi Landscape Conservation Program, Casilla 3-35181, San Miguel, La Paz, Bolivia and Wildlife Conservation Society, 185th Street and Southern Boulevard, Bronx, New York, 10460, U.S.A
E-mail: jmartinez@wcs.org

Matsuda Goodwin, Reiko

Dept. of Sociology and Anthropology, Fordham University, 441 E Fordham Road, Bronx, NY 10458-9993, USA
E-mail: reikogoodwin@gmail.com

McGraw, W. Scott

Department of Anthropology, The Ohio State University, 4064 Smith Laboratory, 174 West 18th Avenue, Columbus, OH 43210-1106, USA
E-mail: mcgraw.43@osu.edu

Meijaard, Erik

Borneo Futures
E-mail: emeijaard@gmail.com

Melo, Fabiano R.

Depto. Eng. Florestal – UFV, Campus Viçosa – Viçosa MG, 36.570-900, Brazil
E-mail: frmelo@ufv.br

Mendes, Sérgio L.

Instituto Nacional da Mata Atlântica - INMA/ MCTIC, Av. José Ruschi, Nº 4, Santa Teresa – ES, 29.650-000 Brazil
E-mail: direcao@inma.gov.br

Méndez-Carvajal, Pedro G.

Fundación Pro-Conservación de los Primates Panameños, Ciudad de Panamá, Panamá
E-mail: mendezp@fcprimatespanama.org

Miaretsoa, Longondraza

Dipartimento di Scienze della Vita e Biologia dei Sistemi, University of Torino, Turin, Italy
E-mail: miaretsoa.longondraza@unito.it

Montenegro, Mônica M. V.

Centro Nacional de Pesquisa e Conservação de Primatas Brasileiros, Instituto Chico Mendes de Conservação da Biodiversidade (CPB/ICMBio/ MMA), Brazil
Email: monica.montenegro@icmbio.gov.br

Mpunga, Noah E.

Wildlife Conservation Society, PO Box 1475, Mbeya, Tanzania
E-mail: nmpunga@wcs.org

Nadler, Tilo

Endangered Primate Rescue Center, Cuc Phuong National Park, Nho Quan District, Ninh Binh Province, Vietnam
E-mail: t.nadler@hust.edu.vn

Nary, Andrianjaka Rijaniaina Jean

Department de Biologie Animal, Faculté des Sciences, Université d'Antananarivo, B.P: 906, 101, Antananarivo, Madagascar
E-mail: naryandrianjaka@gmail.com

Nekaris, K. Anna I.

Nocturnal Primate Research Group, Oxford Brookes University, Oxford OX3 0BP, UK
E-mail: anekaris@brookes.ac.uk

Neves, Leonardo G.

Instituto de Estudos Socioambientais do Sul da Bahia (IESB) & Universidade Estadual de Santa Cruz (UESC), Ilhéus, Bahia, Brasil
E-mail: lgneves@yahoo.com

Nijman, Vincent

Nocturnal Primate Research Group, Oxford Brookes University, Oxford OX3 0BP, UK
E-mail: vnijman@brookes.ac.uk

Nixon, Stuart

North of England Zoological Society, Chester Zoo, Caughall Road, Upton-by-Chester, CH2 1LH, UK
E-mail: s.nixon@chesterzoo.org

Nowak, Matthew G.

PanEco Foundation, Wahid Hasyim No. 51/74, Medan 20154, Sumatera Utara, Indonesia
E-mail: nowak.mg@gmail.com

Oates, John F.

Hunter College of CUNY, New York, NY, USA and 34 Frampton Road, Hythe CT21 6JP, UK
E-mail: johnoates1@aol.com

Oklander, Luciana I.

Instituto de Biología Subtropical (IBS) y Universidad Nacional de Misiones (UNAM) Consejo Nacional de Investigaciones Científicas y Técnicas de Argentina (CONICET)
Email: lula_ok@yahoo.com.ar

Paciulli, Lisa M.

Biological Sciences, North Carolina State University, Hillsborough Street, Raleigh, NC 27605
E-mail: lisa_paciulli@ncsu.edu

Anna Pannick

Department of Biological Sciences, NC State University, 112 Derieux Place, Raleigh, NC 27695
E-mail: appannic@ncsu.edu

Passaro, Richard J.

Project Manager of the Cat Ba Langur Conservation Project from 2011-2014
E-mail: rickpassaro@yahoo.com

Patandung, Alfons

Wildlife Conservation Society, Indonesia.
E-mail: apatandung@gmail.com

Perkin, Andrew

C/o Tanzania Forest Conservation Group (TFCG), P.O. Box 23410, Dar es Salaam, Tanzania
E-mail: bwanakomba@yahoo.co.uk

Peterson, Colin

GERP (Groupe d'Etude et de Recherche sur les Primates de Madagascar), 34 Cité des Professeurs, Fort-Duchesne, Antananarivo, Madagascar
E-mail: colinrandpeterson@gmail.com

Phan Duy Thuc

Science Department, Cat Ba National Park, Cat Ba Island, Cat Hai District, Hai Phong Province, Vietnam
E-mail: pduythuc@gmail.com

Ponce, Gabriela

Wildlife Conservation Society (WCS), Casa No. 3, Avenida 15 de Marzo Flores, Peten, Guatemala, 17001, Guatemala.
E-mail: gponce@wcs.org

Port-Carvalho, Márcio

Seção de Animais Silvestres, Instituto Florestal de São Paulo, São Paulo, SP, Brazil
E-mail: marciopc@if.sp.gov.br

Portillo-Reyes, Héctor

Fundación de Ciencias para el Estudio y Conservación de la Biodiversidad (INCEBIO), Tegucigalpa, Honduras
E-mail: hectorportilloreyes@gmail.com

Pusparini, Wulan

Wildlife Conservation Society, Indonesia
E-mail: wpuspari@eco.umass.edu

Pozo Montuy, Gilberto

Conservación de la Biodiversidad del Usumacinta, A.C. Mexico. Gregorio Méndez 56, Col. Centro, Emiliano Zapata, Tabasco, Mex. C.P, 86990
E-mail: gilpozo@cobius.org

Raffel, Martina

Stiftung Artenschutz, Sentruper Str. 315, 48161 Münster, Germany
E-mail: martina.raffel@stiftung-artenschutz.de

Raharivololona, Brigitte M.

Anthropobiologie et Développement Durable, Faculte des Sciences, Université d'Antananarivo, PO Box 906, Antananarivo 101, Madagascar
E-mail: raharivololonabrigitte@yahoo.fr

Rahasia, Rivo

Program Studi Ilmu Kehutanan, Jurusan Budididaya Pertanian, Fakultas Pertanian Universitas Sam Ratulangi, Manado
Email: rivo.rahasia@gmail.com

Ramanamanjato, Jean-Baptiste

Tropical Biodiversity & Social Enterprise, Fort Dauphin, Madagascar
E-mail: jeanbaptiste.ramanamanjato@riotinto.com

Randimbiharirina, Doménico R.

Anthropobiologie et Développement Durable, Faculte des Sciences, Université d'Antananarivo, PO Box 906, Antananarivo 101, Madagascar
E-mail: rrogerdomnico84@gmail.com

Randrianarison, Rose Marie

LVDI International, California, USA
E-mail: lvdiusa@yahoo.com

Ralainasolo, Fidy

Madagascar Wildlife Conservation, Lake Alaotra, Madagascar
E-mail: Fidy.Ralainasolo@durrell.org

Ratsimbazafy, Jonah

General Director, Group D'Etude et and the Recherche Sur Les Primates de Madagascar (GERP), PB 779, Antananarivo 101, Madagascar
E-mail: jonah@gerp-mg.org

CONTRIBUTORS' ADDRESSES

Raveloarimalala, Lucile Mialisoa

Madagascar Wildlife Conservation, Lake Alaotra,
Madagascar
E-mail: rmialisoalucile@yahoo

Reibelt, Lena M.

Madagascar Wildlife Conservation, Lot 17420
bis Avaradrova Sud, 503 Ambatondrazaka,
Madagascar
E-mail: reibelt.lena@gmail.com

Ren, Guopeng

State Key Laboratory of Genetic Resources and
Evolution, Kunming Institute of Zoology,
Chinese Academy of Sciences, Kunming,
Yunnan, 650223 China
E-mail: zhu@mail.kiz.ac.cn

Rodríguez, Melissa E.

Asociación Territorios Vivos El Salvador (ATVES),
El Salvador. Villavicencio Plaza 2-19 Paseo
General Escalón y 99 Ave. Norte, Colonia
Escalón, San Salvador
E-mail: melissa.rodriguez@atves.org

Rodríguez-Beitia, Bonarge A.

Fundación Pro-Conservación de los Primates
Panameños (FCPP), PO Box: 0816-07905, Panama
Email: jardexp@hotmail.com

Rowe, Noel

Primate Conservation Inc. 1411 Shannock Road –
Charlestown, Rhode Island 02813-3726
E-mail: nrowe@primate.org

Röhe, Fábio

Department of Ecology, National Institute of
Amazonian Research, Manaus, Brazil
E-mail: fabiorohe@gmail.com

Rudran, Rasanayagam

Smithsonian's Conservation Biology Institute,
1500 Remount Road, Front Royal Virginia 22630, USA
E-mail: rudran@msn.com

Rylands, Anthony B.

Global Wildlife Conservation, c/o WeWork, 718
7th Street, N.W., Washington, DC 20001, USA
E-mail: arylands@globalwildlife.org

Salim, Agus

Hatfield Indonesia, Jakarta, Indonesia
E-mail: asalim@hatfieldgroup.com

Schrudde, Daniela

Project Manager of the Cat Ba Langur
Conservation Project from 2008-2011
E-mail: schrudde@hotmail.com

Sefczek, Timothy M.

The Ohio State University. 4005 Smith
Laboratory; 174 West 18th Avenue, Columbus,
Ohio 43210, USA
E-mail: sefczek.1@osu.edu

Shekelle, Myron

Department of Anthropology, Western
Washington University, Bellingham WA 98225, USA
E-mail: shekelle@tarsier.org

Silva, Alessandro A.

Programa de Educação Ambiental (PREA), Minas
Gerais, Brazil
E-mail: cassandroantunes@gmail.com

Solórzano, María Fernanda

Licenciada en Ciencias Biológicas y Ambientales,
Universidad Central del Ecuador (UCE), Ecuador.
Consultora Ambiental Independiente
Email: mfersolorzano@gmail.com

Sop, Tenekwetché

Max Planck Institute for Evolutionary
Anthropology Department of Primatology
Deutscher Platz 6 04103 Leipzig Germany
E-mail: tenekwetché_sop@eva.mpg.de

Stenke, Roswitha

Project Manager of the Cat Ba Langur
Conservation Project from 2000-2008
E-mail: rstenke@gmail.com

Supriatna, Jatna

Department of Biology Faculty of Mathematics
and Science (Conservation Biology), RCCC UI.
Lab Multidisiplin FMIPA Building, Kampus UI
Depok 16424 Indonesia
E-mail: jatna.supriatna@gmail.com

Ting, Nelson

Institute of Ecology and Evolution, University of
Oregon, USA
E-mail: nting@uoregon.edu

Torti, Valeria

Department of Life Sciences and Systems Biology,
University of Torino, Via Accademia Albertina 13,
10123 Torino, Italy
E-mail: valeria.torti@gmail.com

Valença-Montenegro, Mônica M.

Centro de Protecao de Primatas Brasileiros,
IBAMA, Brazil
E-mail: monica.montenegro@ibama.gov.br

Valente, Daria

Department of Life Sciences and Systems Biology,
University of Torino, Via Accademia Albertina 13,
10123 Torino, Italy
E-mail: daria.valente@unito.it

Vidal, Marcelo D.

Instituto Brasileiro do Meio Ambiente e dos
Recursos Naturais Renováveis – IBAMA, Brazil
E-mail: mderzi@provarzea.ibama.gov.br

Waeber, Patrick O.

Madagascar Wildlife Conservation, Lot 17420
bis Avaradrova Sud, 503 Ambatondrazaka,
Madagascar
E-mail: powaeber@gmail.com

Wallace, Robert

Wildlife Conservation Society, La Paz 3-35181,
Bolivia
E-mail: rwallace@wcs.org

Wiafe, Edward D.

Department of Environment and Natural
Resources, Presbyterian University College,
P.O.Box 393, Akropong-Akuapem, Ghana
E-mail: edward.wiafe@presbyuniversity.edu.gh

Wieczkowski, Julie

Department of Anthropology, Buffalo State
University, Buffalo, New York 14222, USA
E-mail: wieczkja@buffalostate.edu

Wich, Serge A.

Research Centre in Evolutionary Anthropology
and Palaeoecology, Liverpool John Moores
University, Liverpool, UK;
Institute for Biodiversity and Ecosystem
Dynamics, University of Amsterdam, Netherlands
E-mail: s.a.wich@ljmu.ac.uk

Wormell, Dominic

Durrell Wildlife Conservation Trust, Les Augrès
Manor, La Profonde Rue, Trinity, JE3 5BP, Jersey,
Channel Islands
E-mail: dominic.wormell@durrell.org

Zhang, Lu

School of Life Sciences, Sun Yat-Sen University,
Guangzhou, China
E-mail: yanlu@cloudmountain.cn

IUCN SSC PRIMATE SPECIALIST GROUP (PSG)

The **Chair** is *Russell A. Mittermeier*, Global Wildlife Conservation, Washington DC, USA, and the two **Deputy Chair** are *Anthony B. Rylands*, Global Wildlife Conservation, Washington DC, USA and *Christoph Schwitzer*, Bristol Zoological Society, Bristol Zoo Gardens, Bristol, UK

Vice Chairs: Section on Great Apes, *Dirck Byler*, Great Ape Conservation Director, Global Wildlife Conservation, Washington DC, USA, and *Serge A. Wich*, Liverpool John Moores University, UK

Vice Chair: Section on Small Apes, *Susan M. Cheyne*, Oxford Brookes University, Oxford, UK

Vice Chairs: Section on Human-Primate Interactions, *Siân Waters*, Department of Anthropology, Durham University, UK, and *Barbary Macaque Awareness and Conservation*, Tetouan, Morocco and *Susan M. Cheyne*, Oxford Brookes University, UK

There are **Regional Vice Chairs** for the principal areas where primates occur, as follows:

AFRICA SECTION

Rachel Ashegbofe Ikemeh, Niger Delta Forest Project, Nigeria
Inza Koné, Centre Suisse de Recherches Scientifiques, Côte d'Ivoire
David Osei, West African Primate Conservation Action (WAPCA), Ghana
Janette Wallis, University of Oklahoma, Norman, OK, USA

MADAGASCAR SECTION

Christoph Schwitzer, Bristol Zoological Society, Bristol, UK
Jonah Ratsimbazafy, Groupe d'étude et de recherche sur les primates (Gerp), Antananarivo, Madagascar
Steig Johnson, University of Calgary, Calgary, Canada

NEOTROPICAL SECTION

Mesoamerica

Liliana Cortés-Ortiz, Museum of Zoology and Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA

Andean Countries

Erwin Palacios, Conservación Internacional Colombia, Bogotá, Colombia
Eckhard W. Heymann, Deutsches Primatenzentrum, Göttingen, Germany
Fanny M. Cornejo, Stony Brook University, New York, USA
Stella de la Torre, San Francisco University of Quito, Quito, Ecuador
Diana C. Guzman Caro, Colombiana Primate Association, Bogotá, Colombia

Brazil and the Guianas

M Cecília M. Kierulff, Instituto Pri-Matas, Belo Horizonte, Minas Gerais, Brazil
Fabiano Rodrigues de Melo, Federal University of Viçosa, Viçosa, Minas Gerais, Brazil
Maurício Talebi, São Paulo Federal University, Diadema, São Paulo, Brazil
Leandro Jerusalinsky, Centro Nacional de Pesquisa and Conservação de Primatas Brasileiros, Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio), João Pessoa, Brazil

Southern Cone – Argentina, Bolivia, Paraguay, Uruguay

Martin Kowalewski, Estación Biológica Corrientes (EBCo), Corrientes, Argentina

ASIA SECTION

China

Baoguo Li, Northwest University, Xi'an, China

South-east Asia / Indochina

Jatna Supriatna, Conservation International Indonesia Program, Jakarta, Indonesia
Arif Setiawan, Faculty of Forestry, Gadjah Mada University, Yogyakarta, Indonesia
Christian Roos, Deutsches Primatenzentrum, Göttingen, Germany
Benjamin M. Rawson, World Wildlife Fund, Hanoi, Vietnam
Ramesh Boonratana, Mahidol University International College, Salaya, Nakhon Pathom, Thailand
Le Khac Quyet, Fauna & Flora International - Vietnam Conservation Support Programme, Hanoi, Vietnam
Duc Hoang Minh, Southern Institute of Ecology (SIE), Ho Chi Minh City, Vietnam

South Asia

Sanjay Molur, Wildlife Information Liaison Development, Coimbatore, Tamil Nadu, India
Dilip Chetry, Gibbon Conservation Centre, Assam, India

INTERNATIONAL PRIMATOLOGICAL SOCIETY (IPS)

The **President** is Karen Strier, University of Wisconsin-Madison, USA. The **Secretary General** is Júlio César Bicca-Marques, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil.

There are six Vice-Presidents: **Treasurer and Vice-President for Membership** – *Trudy Turner*, University of Wisconsin-Milwaukee, USA; **Vice-President for Communications** – Catherine Hobaiter, University of St Andrews, UK; **Vice-President for Conservation** – *Tatyana Humle*, University of Kent, UK; **Vice-President for Captive Care** – Stephen Ross, Lincoln Park Zoo, USA; **Vice-President for Education and Outreach** – Patricia Izar, University of São Paulo, Brazil; and **Vice President for Research** – Marina Cords, Columbia University, USA.

The **Species Survival Commission (SSC)** is one of six volunteer commissions of IUCN, a union of sovereign states, government agencies and non-governmental organizations. SSC's mission is to conserve biological diversity by developing and executing programmes to save, restore and wisely manage species and their habitats. Survival of the world's living primate species and subspecies is the principal mission of the IUCN SSC Primate Specialist Group (PSG), over 700 volunteer professionals who represent the front line in international primate conservation. The PSG website is www.primate-sg.org.

The **International Primatological Society (IPS)** was created to encourage all areas of non-human primatological scientific research, to facilitate cooperation among scientists of all nationalities engaged in primate research, and to promote the conservation of all primate species. The Society is organized exclusively for scientific, educational and charitable purposes. For more information about IPS, visit www.internationalprimatologicalsociety.org.

Global Wildlife Conservation (GWC) uses a flexible model to allow them to make a direct impact anywhere wildlife and ecosystems are most threatened. They work directly with local communities, indigenous groups, organizations, and governments to protect natural resources. This is critical for the long-term persistence of biodiversity and human societies on our only home – Earth. For more information about GWC, visit www.globalwildlife.org.

Bristol Zoological Society (BZS) runs Bristol Zoo Gardens, the Wild Place Project and the Institute of Conservation Science and Learning. BZS undertakes conservation action and conservation research in both the UK and around the world. Its mission is to save wildlife together through conservation action and engaging people with the natural world. For more information about BZS, visit www.bristolzoo.org.uk.

GOLDEN-HEADED LANGUR





ISBN 978-1-5272-4806-9



9 781527 248069 >