

COMMISSION E3

SOLAR IMPACT THROUGHOUT THE HELIOSPHERE

Impact solaire dans l'héliosphère

PRESIDENT
VICE-PRESIDENT
ADVISOR
ORGANIZING COMMITTEE

Nicole Vilmer
Claire Foullon
Ingrid Mann
Margit Haberreiter,
Olga Malandraki,
Masumi Shimojo,
Ilya G. Usoskin,

TRIENNIAL REPORT 2018-2021

1. Introduction

The IAU Commission E3 “Solar Impact through the Heliosphere” covers observational and theoretical aspects of the impacts of solar variability on the heliosphere, on the planetary magnetospheres and/or atmospheres as well as on the boundaries of the heliosphere with the interstellar medium. The primary goals are to investigate and understand the solar wind variability and its effect on the Earth (space weather), other planets and the boundaries of the heliosphere as well as the conditions under which energetic particles from the Sun and charged cosmic dust fill the heliosphere. Solar variability may occur on timescales of hours and days of the solar activity cycle and even on the lifetime of the Sun and the impact of this variability at different time scales controls space weather and space climate effects. Space weather activities have gained global visibility in the recent years and the topic has extended these recent years from fundamental research to applied research.

Key research topics for commission E3 are listed below:

1. Solar wind and its variability
2. Plasma and magnetic disturbances in the solar wind (magnetic clouds, Interplanetary shocks and coronal mass ejections)
3. Solar energetic particles in the inner heliosphere : their sources and propagation
4. Solar wind interaction with planets ; Space Weather
5. Extreme solar events and their influence on Earth and planets
6. Solar forcing on the heliosphere after the end of the Grand Solar Maximum
7. Multi-point observations in the inner heliosphere
8. Outer regions of the heliosphere and boundaries with the interstellar medium
9. Coordination between in-situ and remote-sensing observations (space and ground-based)

This range of scientific topics is particularly timely with the arrival of new data from a fleet of new spacecraft flying in the inner heliosphere (Parker Solar Probe, Solar Orbiter, Bepi Colombo). Synergies between multi-point in situ observations with these missions and previously launched missions (STEREO, WIND,...) and remote sensing observations from space missions (SDO, STEREO, Solar Orbiter, Hinode,...) and ground-based instruments should be strongly encouraged in the coming period. Future space missions

dedicated to solar and heliospheric studies should be launched in the near future by India (Aditya) and China (ASO-S), U.S. (PUNCH) and Japan (SOLAR-C-EUVST). Finally, NASA's Interstellar Mapping and Acceleration Probe (IMAP) looking outwards to observe the interaction of the solar wind with the interstellar medium should be launched in 2024.

During the past three years, ground-based radio observations from non-solar dedicated instruments have been largely used by the community to contribute to studies of the impact of solar activity on the heliosphere (Atacama Large millimeter/submillimeter array ALMA, the low Frequency Analyzer and Recorder (LOFAR) and Murchison Widefield Array (MWA). Together with the new observations from the Expanded Owens Valley Solar Array, these instruments pave the way for the use of observations from the future Square Kilometer Array (SKA) Observatory. The emerging field of space weather research will further develop in the coming years. The study of the influence of the recently discovered extreme solar events is a fast-evolving field of research which should be further developed. With the beginning of the new solar cycle which is largely predicted to be moderate, there is a further possibility to study and understand the solar forcing on the heliosphere in the case of a moderate level of solar activity.

2. Scientific activities and developments in the last 3 years

We present here a selection of a few advances achieved in the field in the last three years.

2.1. *Coronal/Heliospheric Physics and Space Weather with ground-based radio observatories*

In the last 3 years, ground based radio observations from non-solar dedicated instruments have led to many studies and results in the field of coronal and heliospheric physics.

At low frequencies, both LOFAR and MWA have started to be largely used by the heliospheric community. Brand-new results have been obtained with these instruments on the different radio emissions produced in the solar corona during flares and CMEs. Radio emissions produced by the propagation of electron beams (type III bursts) in the solar corona were used on one hand to deduce characteristics of the electron beams and of the density in the structures in which electrons propagate (Mann et al 2018; McCauley et al 2018; Reid 2020; Zhang et al 2020). Through the analysis of spectral fine structures and of their spatial evolution, new results were obtained on the scattering of the emission in the corona leading to new investigation of the turbulent nature of the corona (Chen et al 2018; Kuznetsov and Kontar 2019). Weak non-thermal radio emissions linked to nanoflare coronal heating theories have been analysed using MWA high sensitivity observations (Sharma et al 2018). Radio emissions associated with the propagation of shocks in the corona (type II bursts) were analysed together with EUV and white-light observations of Coronal Mass Ejections (CMEs) to track not only the propagation of shocks but also the conditions of CME-shock related particle acceleration (Chrysaphi et al 2018; Magdalenic et al 2020; Maguire et al 2021). A rare spatially resolved diagnostic of a CME magnetic field could also be achieved with MWA observations (Mondal et al 2020). Observations of the scintillation of the radio signal from pulsars with LOFAR have been used to derive properties of the solar wind (Tiburzi et al 2021). Finally, LOFAR observations allow also to infer characteristics of the ionospheric scintillations (Fallows et al 2020). These new studies demonstrate the larger use of LOFAR and MWA by the solar and heliospheric community. To go further and use e.g. LOFAR as a space-weather dedicated facility would require some upgrade of the system. This is the motivation behind

the LOFAR4SW projet of the EU which aims at designing a system upgrade such that the instrument can perform space weather observations (solar, heliospheric and ionospheric) (Carley et al 2020a). The ultimate goal will be to have a large-scale research facility in which completely simultaneous and independent observing modes provide continuous access to two research communities: radio astronomy and space weather research.

The Square Kilometer Array (SKA) Observatory was launched in February 2021. As for LOFAR, MWA, JVLA, the solar and heliospheric communities should take the new opportunities offered by SKA in terms of spatial, spectral, and temporal resolution and sensitivity to obtain major new insights in solar and heliospheric physics (e.g. Nindos et al 2019).

The scientific solar observations with the Atacama Large Millimeter/submillimeter Array (ALMA) started in December 2016. Due to the difficulty of the image synthesis, the achievements from the ALMA solar data had not appeared soon after the observations. However, many papers that are written based on the data obtained with ALMA are published in these three years. Since we can receive the millimeter/submillimeter-waves that propagate from the lower-middle chromospheres, and its images show us the thermal structures in the layers. The comparison studies between ALMA data and radiative MHD simulations are done actively for understanding the chromosphere, which is the root of the solar wind (Wedemeyer et al 2020; Eklund et al 2020; Martínez-Sykora et al 2020; Chintzoglou et al 2021)

In parallel to the use of observations from new radio telescopes developed for general astronomical use, recent advances in dedicated solar radio instrumentation have been performed to obtain observations with a broader frequency coverage with high spatial, spectral, and temporal resolution (Expanded Owens Valley Solar Array (EOVSA), Siberian Radio Heliograph (SRH), Mingantu Ultrawide Spectral Radiograph (MUSER). Such a progress in solar and heliospheric radio astronomy has led to the recent publication of research and review papers in a special issue of *Frontiers in Astronomy and Space Sciences* dedicated to Solar and Space Weather Radio Physics (Nindos 2020; Carley et al 2020b; Alissandrakis 2020; Reid 2020; Loukitcheva 2020; Vourlidas et al 2020; Kuroda et al 2020; Yan et al 2021; Klein 2021a,b).

2.2. Missions to the Sun

New solar and heliospheric missions have been launched in the last 3 years. The NASA mission Parker Solar Probe (PSP) was successfully launched in August 2018 and has already achieved 7 perihelia close to the Sun. This mission carries a set of in-situ instruments allowing to measure, close to the Sun, the parameters of the solar wind, the magnetic field, plasma waves and radio electromagnetic radiation, energetic particles, as well as an heliospheric imager. Already during the first perihelii, PSP returned a large amount of science data that led to a new picture of the source region of the solar wind. The first results were presented in *Nature* (2019) (Bale et al 2019; Howard et al 2019; Kasper et al 2019; McComas et al 2019) and a Special issue of the *Astrophysical Journal Supplement Series* was published in 2020 ([https://iopscience.iop.org/journal/0067-0049/page/Early Results from Parker Solar Probe](https://iopscience.iop.org/journal/0067-0049/page/Early%20Results%20from%20Parker%20Solar%20Probe)) with more than 50 science papers that provide more detailed results from the first two orbits. One of the surprising results is the existence of hundreds to a thousand intermittent reversals of the magnetic field accompanied by increases in the speed of the solar wind and reversal of the pitch angle of the suprathermal electron strahl indicating a kink of the magnetic field (switch-backs, jets, or spikes). A lot of papers have been published on their properties as well as on their origin and generation mechanisms. Plasma waves and micro-instabilities have been observed very close to the Sun leading to exceptional detections of electrostatic and electromagnetic waves,

electron holes, reconnection in current sheets, and type III radio bursts. The evolution of plasma turbulence between 0.17 and 1 AU has also been studied. A great number of small energetic particle events, small coronal mass ejections has been observed despite the fact that the first orbits were performed close to the solar minimum. Small plasma structures ejected from the Sun could be seen with the heliospheric imager on the PSP, as well as magnetic islands resulting from tearing-mode instability in the current sheet.

The ESA/NASA mission Solar Orbiter launched in February 2020 is the first mission combining in-situ and remote sensing observations of solar and heliospheric plasma together with an out of the ecliptic view of the phenomena at the end of the mission. A special issue of *Astronomy and Astrophysics* is in preparation to present the first results of the mission. Some of the new results are: observations of small impulsive energetic particle events within 1 AU despite the minimum phase of the solar cycle, radial dependence of CIR-associated suprathermal ions in the inner heliosphere (Mason, G. M. et al 2020, 2021; Gomez-Herrero, R. et al 2021; Allen, R. C. et al 2020)

2.3. *Advanced studies of extreme solar events*

An impressive progress has been achieved in studies of extreme solar events which form a severe hazard for the modern technological society. All severe solar energetic particle (SEP) events detected on Earth (ground-level enhancement, GLE) have been re-analyzed using a revised response function and an improved analysis method (e.g., Koldobskiy, S. et al 2021), making basis for a more precise analysis of such SEP events. The accuracy and reliability of reconstructions of the extreme SEP events for the past millennia has been greatly improved using terrestrial proxy data such as cosmogenic isotopes. The current state of the art has been summarized in a collective book by Miyake et al (2019) and further developed by an ISSI international team. Three extreme SEP events are known in the past: at 774 AD, 993 AD and 660 BC (Miyake et al 2012; Miyake et al 2013; O'Hare et al 2019), and two candidate events in 1052 and 1279 AD are still awaiting for confirmation (Brehm et al 2021). All these events are a factor 20–100 stronger than the strongest SEP event (23-Feb-1956) recorded during the instrumental era.

Another method to estimate the occurrence rate and severity of extreme SEP events is based on a projection of the statistic of super-flares recorded by the Kepler telescope on a large ensemble of sun-like stars (e.g., Notsu et al 2019). This statistic suggests an order-of-magnitude higher probabilities of super-flare occurrence on the Sun than that evaluated from the terrestrial solar proxies.

Thus, although the very fact of the super-flares (a factor of 100 or more stronger than those directly observed during the instrumental era) on the Sun is beyond any doubts, the exact occurrence probability and the severity of such events are not precisely known and need to be clarified in the future.

References

- Alissandrakis CE (2020) Structure of the solar atmosphere: a radio perspective. *Frontiers in Astronomy and Space Sciences* 7:74,
- Allen, R C, Mason, G M, Ho, G C, Rodríguez-Pacheco, J, Wimmer-Schweingruber, R F, Andrews, G B, Berger, L, Boden, S, Cernuda, I, Espinosa Lara, F, Freiherr von Forstner, J L, Gómez-Herrero, R, Hayes, J R, Kulkarni, S R, Lees, W J, Martin, C, Pacheco, D, al (2020) Suprathermal particles from corotating interaction regions during the first perihelion pass of solar orbiter. *A&A* , URL <https://doi.org/10.1051/0004-6361/202039870>
- Bale SD, Badman ST, Bonnell JW, Bowen TA, Burgess D, Case AW, Cattell CA, Chandran BDG, Chaston CC, Chen CHK, Drake JF, de Wit TD, Eastwood JP, Ergun RE, Farrell WM, Fong C, Goetz K, Goldstein M, Goodrich KA, Harvey PR, Horbury TS, Howes GG, Kasper JC, Kellogg PJ, Klimchuk JA, Korreck KE, Krasnoselskikh VV, Krucker S, Laker R,

- Larson DE, MacDowall RJ, Maksimovic M, Malaspina DM, Martinez-Oliveros J, McComas DJ, Meyer-Vernet N, Moncuquet M, Mozer FS, Phan TD, Pulupa M, Raouafi NE, Salem C, Stansby D, Stevens M, Szabo A, Velli M, Woolley T, Wygant JR (2019) Highly structured slow solar wind emerging from an equatorial coronal hole. *Nature* 576(7786):237–242,
- Brehm N, Bayliss A, Christl M, Synal H, Adolphi F, Beer J, Kromer B, Muscheler R, Solanki S, Usoskin I, Bleicher N, Bollhalder S, Tyers C, Wacker L (2021) Eleven-year solar cycles over the last millennium revealed by radiocarbon in tree rings. *Nature Geosci* 14:10–15,
- Carley EP, Baldovin C, Benthem P, Bisi MM, Fallows RA, Gallagher PT, Olberg M, Rothkaehl H, Vermeulen R, Vilmer N, Barnes D (2020a) Radio observatories and instrumentation used in space weather science and operations. *Journal of Space Weather and Space Climate* 10:7,
- Carley EP, Vilmer N, Vourlidas A (2020b) Radio observations of coronal mass ejection initiation and development in the low solar corona. *Frontiers in Astronomy and Space Sciences* 7:79,
- Chen X, Kontar EP, Yu S, Yan Y, Huang J, Tan B (2018) Fine Structures of Solar Radio Type III Bursts and Their Possible Relationship with Coronal Density Turbulence. *Astrophys Journ* 856(1):73, , 1801.07545
- Chintzoglou G, De Pontieu B, Martínez-Sykora J, Hansteen V, de la Cruz Rodríguez J, Szydlarski M, Jafarzadeh S, Wedemeyer S, Bastian TS, Sainz Dalda A (2021) ALMA and IRIS Observations of the Solar Chromosphere. I. An On-disk Type II Spicule. *Astrophys Journal* 906(2):82, , 2005.12717
- Chrysaphi N, Kontar EP, Holman GD, Temmer M (2018) CME-driven Shock and Type II Solar Radio Burst Band Splitting. *Astrophys Journ* 868(2):79, , 1810.08026
- Eklund H, Wedemeyer S, Szydlarski M, Jafarzadeh S, Guevara Gómez JC (2020) The Sun at millimeter wavelengths. II. Small-scale dynamic events in ALMA Band 3. *Astron and Astrophys* 644:A152, , 2010.06400
- Fallows RA, Forte B, Astin I, Allbrook T, Arnold A, Wood A, Dorrian G, Mevius M, Rothkaehl H, Matyjasiak B, Krankowski A, Anderson JM, Asgekar A, Avruch IM, Bentum M, Bisi MM, Butcher HR, Ciardi B, Dabrowski B, Damstra S, de Gasperin F, Duscha S, Eislöffel J, Franzen TMO, Garrett MA, Griebmeier JM, Gunst AW, Hoeft M, Hörandel JR, Iacobelli M, Intema HT, Koopmans LVE, Maat P, Mann G, Nelles A, Paas H, Pandey VN, Reich W, Rowlinson A, Ruiter M, Schwarz DJ, Serylak M, Shulevski A, Smirnov OM, Soida M, Steinmetz M, Thoudam S, Toribio MC, van Ardenne A, van Bemmelen IM, van der Wiel MHD, van Haarlem MP, Vermeulen RC, Vocks C, Wijers RAMJ, Wucknitz O, Zarka P, Zucca P (2020) A LOFAR observation of ionospheric scintillation from two simultaneous travelling ionospheric disturbances. *Journal of Space Weather and Space Climate* 10:10, , 2003.04013
- Gomez-Herrero, R, Pacheco, D, Kollhoff, A, Espinosa Lara, F, Freiherr von Forstner, J L, Dresing, N, Lario, D, Balmaceda, L, Krupar, V, Malandraki, O E, Aran, A, Bucik, R, Klassen, A, Klein, K-L, Cernuda, I, Eldrum, S, Reid, H, al (2021) First near-relativistic solar electron events observed by epd onboard solar orbiter. *A&A* , URL <https://doi.org/10.1051/0004-6361/202039883>
- Howard RA, Vourlidas A, Bothmer V, Colaninno RC, DeForest CE, Gallagher B, Hall JR, Hess P, Higginson AK, Korendyke CM, Kouloumvakos A, Lamy PL, Liewer PC, Linker J, Linton M, Penteado P, Plunkett SP, Poirier N, Raouafi NE, Rich N, Rochus P, Rouillard AP, Socker DG, Stenborg G, Thernisien AF, Viall NM (2019) Near-Sun observations of an F-corona decrease and K-corona fine structure. *Nature* 576(7786):232–236,
- Kasper JC, Bale SD, Belcher JW, Berthomier M, Case AW, Chandran BDG, Curtis DW, Gallagher D, Gary SP, Golub L, Halekas JS, Ho GC, Horbury TS, Hu Q, Huang J, Klein KG, Korreck KE, Larson DE, Livi R, Maruca B, Lavraud B, Louarn P, Maksimovic M, Martinovic M, McGinnis D, Pogorelov NV, Richardson JD, Skoug RM, Steinberg JT, Stevens ML, Szabo A, Velli M, Whittlesey PL, Wright KH, Zank GP, MacDowall RJ, McComas DJ, McNutt RL, Pulupa M, Raouafi NE, Schwadron NA (2019) Alfvénic velocity spikes and rotational flows in the near-Sun solar wind. *Nature* 576(7786):228–231,
- Klein KL (2021a) Radio astronomical tools for the study of solar energetic particles I. Correlations and diagnostics of impulsive acceleration and particle propagation. *Frontiers in Astronomy and Space Sciences* 7:105,
- Klein KL (2021b) Radio astronomical tools for the study of solar energetic particles II. Time-extended acceleration at subrelativistic and relativistic energies. *Frontiers in Astronomy and Space Sciences* 7:93,

- Koldobskiy, S, Raukunen, O, Vainio, R, Kovaltsov, G A, Usoskin, I (2021) New reconstruction of event-integrated spectra (spectral fluences) for major solar energetic particle events. *Astron Astrophys* 647:A132, , URL <https://doi.org/10.1051/0004-6361/202040058>
- Kuroda N, Fleishman GD, Gary DE, Nita GM, Chen B, Yu S (2020) Evolution of Flare-accelerated Electrons Quantified by Spatially Resolved Analysis. *Frontiers in Astronomy and Space Sciences* 7:22, , 2004.13155
- Kuznetsov AA, Kontar EP (2019) First imaging spectroscopy observations of solar drift pair bursts. *Astron and Astrophys* 631:L7, , 1910.09864
- Loukitcheva M (2020) Measuring magnetic field with Atacama Large Millimeter/Submillimeter Array. *Frontiers in Astronomy and Space Sciences* 7:45,
- Magdalenic J, Marqu e C, Fallows RA, Mann G, Vocks C, Zucca P, Dabrowski BP, Krankowski A, Melnik V (2020) Fine Structure of a Solar Type II Radio Burst Observed by LOFAR. *Astrophys Journ* 897(1):L15,
- Maguire CA, Carley EP, Zucca P, Vilmer N, Gallagher PT (2021) LOFAR Observations of a Jet-driven Piston Shock in the Low Solar Corona. *Astrophys Journ* 909(1):2, , 2101.05569
- Mann G, Breitling F, Vocks C, Aurass H, Steinmetz M, Strassmeier KG, Bisi MM, Fallows RA, Gallagher P, Kerdr on A, Mackinnon A, Magdalenic J, Rucker H, Anderson J, Asgekar A, Avruch IM, Bell ME, Bentum MJ, Bernardi G, Best P, Birzan L, Bonafede A, Broderick JW, Br uggen M, Butcher HR, Ciardi B, Corstanje A, de Gasperin F, de Geus E, Deller A, Duscha S, Eisl offel J, Engels D, Falcke H, Fender R, Ferrari C, Frieswijk W, Garrett MA, Griebmeier J, Gunst AW, van Haarlem M, Hassall TE, Heald G, Hessels JWT, Hoedt M, H orandel J, Horneffer A, Juette E, Karastergiou A, Klijn WFA, Kondratiev VI, Kramer M, Kuniyoshi M, Kuper G, Maat P, Markoff S, McFadden R, McKay-Bukowski D, McKean JP, Mulcahy DD, Munk H, Nelles A, Norden MJ, Orru E, Paas H, Pandey-Pommier M, Pandey VN, Pizzo R, Polatidis AG, Rafferty D, Reich W, R ottgering H, Scaife AMM, Schwarz DJ, Serylak M, Sluman J, Smirnov O, Stappers BW, Tagger M, Tang Y, Tasse C, ter Veen S, Thoudam S, Toribio MC, Vermeulen R, van Weeren RJ, Wise MW, Wucknitz O, Yatawatta S, Zarka P, Zensus JA (2018) Tracking of an electron beam through the solar corona with LOFAR. *Astron and Astrophys* 611:A57,
- Mart nez-Sykora J, De Pontieu B, de la Cruz Rodriguez J, Chintzoglou G (2020) The Formation Height of Millimeter-wavelength Emission in the Solar Chromosphere. *Astrophys Journal Letters* 891(1):L8, , 2001.10645
- Mason, G M, Ho, G C, Allen, R C, Rodr guez-Pacheco, J, Wimmer-Schweingruber, R F, Buc k, R, G omez-Herrero, R, Lario, D, von Forstner, J L F, Andrews, G B, Berger, L, Cernuda, I, Espinosa Lara, F, Lees, W J, Mart n, C, Pacheco, D, et al (2020) 3he-rich solar energetic particle events observed on the first perihelion pass of solar orbiter. *A&A* , URL <https://doi.org/10.1051/0004-6361/202039752>
- Mason, G M, Ho, G C, Allen, R C, Xu, Z G, Janitzek, N P, Freiherr von Forstner, J L, Kohlloff, A, Pacheco, D, Rodr guez-Pacheco, J, Wimmer-Schweingruber, R F, Bruce Andrews, G, Schlemm, C E, Seifert, H, Tyagi, K, Lees, W J, Hayes, J, G omez-Herrero, R, et al (2021) Quiet-time low energy ion spectra observed on solar orbiter during solar minimum. *A&A* , URL <https://doi.org/10.1051/0004-6361/202140540>
- McCaughey PI, Cairns IH, Morgan J (2018) Densities Probed by Coronal Type III Radio Burst Imaging. *Solar Phys* 293(10):132, , 1808.04989
- McComas DJ, Christian ER, Cohen CMS, Cummings AC, Davis AJ, Desai MI, Giacalone J, Hill ME, Joyce CJ, Krimigis SM, Labrador AW, Leske RA, Malandraki O, Matthaeus WH, McNutt RL, Mewaldt RA, Mitchell DG, Posner A, Rankin JS, Roelof EC, Schwadron NA, Stone EC, Szalay JR, Wiedenbeck ME, Bale SD, Kasper JC, Case AW, Korreck KE, MacDowall RJ, Pulupa M, Stevens ML, Rouillard AP (2019) Probing the energetic particle environment near the Sun. *Nature* 576:223–227,
- Miyake F, Nagaya K, Masuda K, Nakamura T (2012) A signature of cosmic-ray increase in ad 774-775 from tree rings in Japan. *Nature* 486:240–242,
- Miyake F, Masuda K, Nakamura T (2013) Another rapid event in the carbon-14 content of tree rings. *Nature Comm* 4:1748,
- Miyake F, Usoskin I, Poluianov S (eds) (2019) *Extreme Solar Particle Storms: The Hostile Sun*. IOP Publishing, Bristol, UK, , URL <http://dx.doi.org/10.1088/2514-3433/ab404a>
- Mondal S, Oberoi D, Vourlidis A (2020) Estimation of the Physical Parameters of a CME

- at High Coronal Heights Using Low-frequency Radio Observations. *Astrophys Journ* 893(1):28, , 1909.12041
- Nindos A (2020) Incoherent Solar Radio Emission. *Frontiers in Astronomy and Space Sciences* 7:57, , 2007.14888
- Notsu Y, Maehara H, Honda S, Hawley S, Davenport J, Namekata K, Notsu S, Ikuta K, Nogami D, Shibata K (2019) Do kepler superflare stars really include slowly rotating sun-like stars?—results using APO 3.5 m telescope spectroscopic observations and gaia-DR2 data. *Astrophys J* 876(1):58, , URL <https://doi.org/10.3847/2F1538-4357%2Fab14e6>
- O'Hare P, Mekhalidi F, Adolphi F, Raisbeck G, Aldahan A, Anderberg E, Beer J, Christl M, Fahrni S, Synal HA, Park J, Possnert G, Southon J, Bard E, Aster Team, Muscheler R (2019) Multiradionuclide evidence for an extreme solar proton event around 2,610 B.P. (660 BC). *Proc Nat Acad Sci* 116(13):5961–5966,
- Reid HAS (2020) A review of recent type III imaging spectroscopy. *Frontiers in Astronomy and Space Sciences* 7:56,
- Sharma R, Oberoi D, Arjunwadkar M (2018) Quantifying Weak Nonthermal Solar Radio Emission at Low Radio Frequencies. *Astrophys Journ* 852(2):69, , 1709.00878
- Tiburzi C, Shaifullah GM, Bassa CG, Zucca P, Verbiest JPW, Porayko NK, van der Wateren E, Fallows RA, Main RA, Janssen GH, Anderson JM, Bak Nielsen AS, Donner JY, Keane EF, Künsemöller J, Osłowski S, Griebmeier JM, Serylak M, Brügger M, Ciardi B, Dettmar RJ, Hoeft M, Kramer M, Mann G, Vocks C (2021) The impact of solar wind variability on pulsar timing. *Astron and Astrophys* 647:A84, , 2012.11726
- Vourlidas A, Carley EP, Vilmer N (2020) Radio Observations of Coronal Mass Ejections: Space Weather Aspects. *Frontiers in Astronomy and Space Sciences* 7:43,
- Wedemeyer S, Szydlarski M, Jafarzadeh S, Eklund H, Guevara Gomez JC, Bastian T, Fleck B, de la Cruz Rodriguez J, Rodger A, Carlsson M (2020) The Sun at millimeter wavelengths. I. Introduction to ALMA Band 3 observations. *Astron and Astrophys* 635:A71, , 2001.02185
- Yan Y, Chen Z, Wang W, Liu F, Geng L, Chen L, Tan C, Chen X, Su C, Tan B (2021) Mingantu spectral radiograph for solar and space weather studies. *Frontiers Astron Space Sci* 8:1
- Zhang P, Zucca P, Sridhar SS, Wang C, Bisi MM, Dabrowski B, Krankowski A, Mann G, Magdalenic J, Morosan DE, Vocks C (2020) Interferometric imaging with LOFAR remote baselines of the fine structures of a solar type-IIIb radio burst. *Astron and Astrophys* 639:A115, , 2005.09419