An aerial photograph of New York City at sunset. The sky is filled with golden, wispy clouds. In the background, the city skyline is visible, with the Empire State Building as the most prominent skyscraper. In the foreground, several multi-story brick buildings are shown. On the roof of one of these buildings, a group of people is gathered, and there is a rooftop garden with various plants and trees. The overall scene is bathed in the warm, golden light of the setting sun.

The New York City  
Community  
Air Survey

Neighborhood Air Quality  
2008 - 2016

# CONTRIBUTORS

Bureau of Environmental Surveillance and Policy  
Division of Environmental Health  
New York City Department of Health and Mental Hygiene

Queens College, City University of New York,  
Barry Commoner Center for Health and the Environment

# ACKNOWLEDGMENT

The Health Department is grateful to the following agencies for their logistical support and advice:

New York City Mayor's Office of Sustainability  
New York City Department of Transportation  
New York City Department of Parks and Recreation  
New York City Department of Environmental Protection  
New York State Department of Environmental Conservation

For more information about the Health Department's New York City Community Air Survey, visit [nyc.gov/health/nyccas](https://nyc.gov/health/nyccas).

For additional data and information on New York City's environment, sustainability and health outcomes, visit [nyc.gov/health/tracking](https://nyc.gov/health/tracking).

# Contents

Page

- 4 EXECUTIVE SUMMARY
- 5 BACKGROUND AND METHODS
- 9 RESULTS
- 21 APPENDICES

# EXECUTIVE SUMMARY

In 2007, as part of New York City's first long-term plan for environmental sustainability, the Health Department established the New York City Community Air Survey (NYCCAS), the largest ongoing urban air monitoring program of any U.S. city. NYCCAS data is used to inform local air pollution policies, to estimate exposures for health research, and to track changes in air quality over time. Beginning in 2015, the annual reporting of these results is mandated by Local Law 103.

This report:

- Provides a summary of the air monitoring program, site selection process, air quality monitoring and analysis methods, and descriptions of the pollutants measured
- Describes the trend in air pollutant levels from winter 2008-2009 through fall 2016 for fine particulate matter (PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>), nitric oxide (NO), black carbon (BC), wintertime sulfur dioxide (SO<sub>2</sub>), and summertime ozone (O<sub>3</sub>)
- Maps neighborhood air pollution levels by year and by season
- Identifies the sources that contribute to high levels of these pollutants in NYC neighborhoods

Major findings:

- Citywide, annual average PM<sub>2.5</sub>, NO<sub>2</sub>, NO, and BC levels have declined by 28%, 27%, 35%, and 24%, respectively, between the first year of monitoring (2009) and the most recent year (2016).
- The largest declines have been observed for SO<sub>2</sub> due to heating oil regulations. Wintertime average levels have declined by 95% over the eight-year period between the first winter of monitoring (2008-2009) and the most recent winter (2015-2016).

- Citywide, summertime average O<sub>3</sub> levels remained relatively stable across the eight years.
- Higher levels of PM<sub>2.5</sub>, NO<sub>2</sub>, NO, and BC continue to be observed in areas of higher traffic density, building density, and heat and hot water boiler density, and in industrial areas. O<sub>3</sub> levels remain higher in the outer boroughs, in areas that are downwind of high emissions density and in areas with fewer combustion emissions.

This report underscores the importance of emissions reduction efforts over the past decade and the continued need to reduce emissions citywide. Implementing new strategies and expanding existing measures within the City's sustainability plan, [OneNYC](#), and its roadmap to reduce greenhouse gas emissions, [80x50](#), will improve air quality and provide important public health benefits to all New Yorkers. These strategies and measures include converting the remaining residual oil boilers to ones that use cleaner heating fuels; transitioning to more efficient, less polluting light duty and heavy duty vehicles; reducing motor vehicle use by shifting to more sustainable modes of transportation; creating more efficient freight networks and expanding truck retrofit and replacement programs; and reducing fossil fuel combustion in buildings. Additionally, reducing emissions from other widely distributed sources of pollution, such as commercial charbroiling, will contribute to improved air quality in the future.

# BACKGROUND AND METHODS

**A**s part of NYC's first long-term plan for environmental sustainability, in 2007 the Health Department established the [New York City Community Air Survey](#) (NYCCAS), which is the largest ongoing urban air monitoring program of any U.S. city. The air quality monitoring network, which began collecting data in December 2008, is a collaboration between the Health Department and Queens College of the City University of New York and provides data to help inform the City's sustainability plan, [OneNYC](#). The objectives of NYCCAS are to:

- Measure air pollutants that affect public health across the city
- Identify local emission sources that impact neighborhood air quality
- Inform the public and city officials about air pollutant levels and clean air priorities
- Provide air pollution estimates for health studies

NYCCAS measures air pollutants that pose the most harm to the public's health. They include:

**Fine particles (PM<sub>2.5</sub>)** are tiny airborne solid and liquid particles less than 2.5 microns in diameter. PM<sub>2.5</sub> is the most harmful urban air pollutant, small enough to penetrate deep into the lungs and enter the bloodstream, worsening lung and heart disease, and leading to hospital admissions and premature deaths. PM<sub>2.5</sub> is also a human carcinogen.

PM<sub>2.5</sub> can either be directly emitted or formed in the atmosphere from other pollutants. Important local sources include fuel combustion in vehicles, boilers in buildings, power plants, construction equipment, marine vessels, and commercial cooking. PM<sub>2.5</sub> in NYC's air also comes from outside the city, from sources far upwind.

**Nitrogen dioxide (NO<sub>2</sub>) and nitric oxide (NO)** are part of a group of pollutants called “oxides of nitrogen” (NO<sub>x</sub>). Exposures to NO<sub>x</sub> are linked to increased emergency department visits and hospitalizations for respiratory conditions, particularly asthma. NO<sub>x</sub> also react with other compounds in the atmosphere to form PM<sub>2.5</sub> and O<sub>3</sub>. NO<sub>x</sub> is produced from a variety of combustion sources in NYC, including motor vehicles, buildings, marine vessels, and construction equipment.


**Sulfur dioxide (SO<sub>2</sub>)** in NYC is produced mainly from burning oils with high sulfur content, such as No. 4 and No. 6 oil (also known as residual fuel oil) or high sulfur No. 2 oil. Fuel oil in NYC is used mainly to heat buildings and for hot water, and some high-sulfur oil is also used to generate electric power and power marine vessels. SO<sub>2</sub> exposures can worsen lung diseases, causing hospitalizations and emergency department visits for asthma and other conditions. SO<sub>2</sub> also contributes to the formation of PM<sub>2.5</sub> in the atmosphere, resulting in exposures downwind of where it is emitted.

**Ozone (O<sub>3</sub>)**, at ground level, is formed through reactions in the atmosphere when NO<sub>x</sub> emissions combine with other airborne pollutants in the presence of sunlight. Therefore, measured O<sub>3</sub> concentrations are often highest downwind from high-emissions areas. In areas where there are high concentrations of fresh combustion emissions, NO<sub>x</sub> reacts with O<sub>3</sub> to reduce its concentrations. As a result, lower O<sub>3</sub> levels are observed near roadways, in city centers, and in other areas of high emissions density.

**Black carbon (BC)** is one component of PM<sub>2.5</sub> and is emitted from diesel exhaust and other sources, such as oil burning. Diesel exhaust particles, indicated by BC, can cause irritation of the breathing passages, leading to respiratory symptoms such as cough, or asthma exacerbation, and may increase the risk of cancer. BC pollution is also a contributor to global climate change.

NYCCAS air quality monitoring began in December 2008. Field teams sampled the air at 150 NYC locations per year during the first two years and at 60 to 100 locations per year in subsequent years (Figure 1). Samples are collected in all seasons for all pollutants, except O<sub>3</sub> and SO<sub>2</sub>, where samples are collected in the summer and winter seasons, respectively.

Figure 1: Monitoring locations.



The original 150 monitoring sites were selected to ensure that the ranges of traffic conditions, size and number of buildings, and land uses in NYC were adequately included while providing a balance in spatial coverage throughout the city. To do this, a digital map of the city was divided into a grid of more than 7,500 squares, each 300 by 300 meters, and each square was classified based on its traffic and building density. A random selection of squares was then drawn from this set, with high building and traffic density areas having an increased chance of selection as these areas are concentrated in a relatively small area of the city. This random site selection was used to locate 80% of the sampling sites. The remaining 20% of sites were selected in places with large remaining gaps in coverage from the random selection or near areas

of interest, such as high-traffic areas, transportation facilities, or major ongoing construction.

Each NYCCAS site is monitored for a two-week period in each season. The schedule of monitoring is assigned randomly so that the same number of sites across the city are monitored in each two-week period. In addition, "reference" sites - centrally located and away from nearby traffic and commercial or industrial activities - are monitored during every two-week period, year round. Data from these "reference" sites are used to adjust the measurements made at other sites for variation that occurs across the city over time, mainly due to weather conditions. For additional details on the selection methods for the original 150 sites, visit [NYCCAS First Winter Results](#), [NYCCAS Design and Implementation](#).

After the first two years of the study, the number of sites was reduced to between 60 and 100 sites, depending on the year, because of budget constraints and to free up resources to measure other pollutants and conduct additional air quality and health studies. The balance of source density and spatial density was preserved, through use of random selection methodologies similar to those described above. The patterns in air pollutant concentrations remained consistent year after year - areas of the city with higher concentrations tend to remain higher over time, while cleaner areas of the city remain cleaner - due to major emissions sources such as buildings and traffic remaining in fixed locations. Because of this, NYCCAS is able to track the geographic pattern of air quality over time with fewer locations than in the original design. Currently, routine NYCCAS air sampling occurs once per season at 60 of the original 150 sites, known as the "core" monitoring sites, and 30 additional locations which include 15 of the original 150 sites and 15 sites located in low income neighborhoods with previously lower monitoring density. The number of

Figure 2: NYCCAS team member deploys a monitor in the field.



reference sites was reduced from five to three after the first four years.

NYCCAS sampling is conducted using monitoring units mounted on lampposts 10 to 12 feet off the ground. The monitors include an air pump and filters to collect PM<sub>2.5</sub> while passive samplers mounted on the outside of unit absorb the gaseous pollutants NO<sub>x</sub>, SO<sub>2</sub>, and O<sub>3</sub>. Laboratory analysis of the filters and passive samplers determines the quantities of pollutants collected and their concentration in air is calculated. Quality control steps included confirming that the sampling pump was operating normally and collecting duplicate and unexposed samples for comparison with study samples.

NYCCAS data were analyzed using a “land-use regression” (LUR) model. LUR models estimate associations among pollution levels, average traffic, building emissions, land use, and other neighborhood factors around the monitoring sites. These associations were used to estimate the seasonal average air pollution levels at locations across the city, including locations where no measurements were taken. The LUR model is also used to assess sources that appear to contribute most to differences in pollution concentrations. For more details on the analysis methods, please see the technical appendices and scientific manuscripts available at [nyc.gov/health/nyccas](https://nyc.gov/health/nyccas). For details on the data sources used to create emissions indicators for the land-use regression modeling, please see Appendix 1.

The results of NYCCAS monitoring have been published in multiple public reports, scientific manuscripts, and annual online data updates. All reports and Health Department scientific studies are available on the NYCCAS website at [nyc.gov/health/nyccas](https://nyc.gov/health/nyccas). All neighborhood-level data and detailed neighborhood air quality reports are available for download through the [Health Department’s Environment & Health Data Portal](#).




# RESULTS:

## FINE PARTICULATE MATTER

At NYCCAS sites measured in each season for eight years:

- Seasonally adjusted street-level PM<sub>2.5</sub> concentrations declined by an average of 0.42 micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ) per year.
- Citywide, annual average levels at the 60 monitoring sites (locations measured across the eight years) declined by 28% between 2009 and 2016.
- In the most recent year (2016), seasonal average concentrations across NYCCAS monitoring sites ranged from 4.5 to 16.8  $\mu\text{g}/\text{m}^3$ .
- Across the eight-year period, higher levels were consistently seen at sites with higher nearby boiler and traffic emissions (Figure 3).

Figure 3: PM<sub>2.5</sub> levels at NYCCAS monitors, by density of nearby boilers emissions (A) and traffic emissions (B)<sup>1</sup>




<sup>1</sup>Boiler emissions represent estimated PM<sub>2.5</sub> emissions from all boiler types within 1,000 m. Traffic emissions were estimated as total traffic density, weighted by vehicle-specific PM<sub>2.5</sub> emissions rates from on-road vehicles within 100 m. High, Medium, and Low represent one third of sites ranked by source indicator density.

- PM<sub>2.5</sub> levels remained relatively higher throughout much of Manhattan, and in areas of higher traffic density, building density and industrial areas in the outer boroughs (Figure 4).

In the spatial model, the most important predictors of PM<sub>2.5</sub> concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
PM <sub>2.5</sub> emissions from heat and hot water boilers in buildings within 1,000 m	Combustion of heating oil and natural gas
Area of industrial land use within 1,000 m	Diesel exhaust particles from trucks idling and traveling through industrial areas. Industrial combustion equipment.
Traffic density, weighted by relative PM <sub>2.5</sub> emissions rates by vehicle type (car, truck, bus) within 250 m.	PM <sub>2.5</sub> emissions from all on-road motor vehicles based on vehicle miles and the relative emissions rates of different vehicle types.

Figure 4: PM<sub>2.5</sub> concentrations, 2016 annual average (left) and 2009-2016 annual averages (right)




# NITROGEN DIOXIDE

At NYCCAS sites measured in each season for eight years:

- Seasonally adjusted street-level NO<sub>2</sub> concentrations declined by an average of 1.1 parts per billion (ppb) per year.
- Citywide, annual average levels at the 60 monitoring sites (locations measured across the eight years) declined by 27% between 2009 and 2016.

- In the most recent year (2016), seasonal average concentrations across NYCCAS monitoring sites ranged from 5.8 to 44.5 ppb.
- Across the eight-year period, higher levels were consistently seen at sites with higher nearby building and traffic emissions (Figures 5-6).

Figure 5: NO<sub>2</sub> levels at NYCCAS monitors, by nearby building density (A) and traffic emissions (B)<sup>2</sup>




<sup>2</sup>Building density was estimated as total interior building area within 1,000 m of monitoring site. Density of nearby traffic emissions were estimated as total traffic density, weighted by vehicle-specific NO<sub>x</sub> emissions rates from on-road vehicles within 100 m. High, Medium, and Low represent one third of sites ranked by source indicator density.

In the spatial model, the most important predictors of NO<sub>2</sub> concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
Area of interior building space within 1,000 m	Combustion of heating oil and natural gas
Percent impervious surface within 100 m	Emissions of motor vehicles on paved roadways
Traffic density, weighted by relative NO <sub>x</sub> emissions rates and vehicle type (car, truck, bus) within 100 m	NO <sub>x</sub> emissions from all on-road motor vehicles based on vehicle miles and the relative emission rates of different vehicle types
Location on a bus route (compared to non-bus route locations)	Emissions from buses and other vehicles on busy roadways. Indicator of traffic congestion.

Figure 6: NO<sub>2</sub> concentrations, 2016 annual average (left) and 2009-2016 annual averages (right)




# NITRIC OXIDE

At NYCCAS sites measured in each season for eight years:

- Seasonally adjusted street-level NO concentrations declined by an average of 1.63 parts per billion (ppb) per year.
- Citywide, annual average levels at the 60 monitoring sites (locations measured across the eight years) declined by 35% between 2009 and 2016.

- In the most recent year (2016), seasonal average concentrations across NYCCAS monitoring sites ranged from 3.5 to 99.1 ppb.
- Across the eight-year period, higher levels were consistently seen at sites with higher nearby emissions from traffic sources and higher densities of nearby building boilers (Figures 7-8).

Figure 7: NO levels at NYCCAS monitors, by nearby NO<sub>x</sub> emissions from traffic (A) and nearby building boilers (B)<sup>3</sup>




<sup>3</sup> Density of nearby traffic emissions was estimated as total traffic density, weighted by vehicle-specific NO<sub>x</sub> emissions rates from on-road vehicles within 100 m. Nearby boiler density was estimated as the number of boilers within 250 m of monitoring sites, regardless of fuel type. High, Medium, and Low represent one third of sites ranked by source indicator density.

In the spatial model, the most important predictors of NO concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
Traffic density, weighted by relative NO <sub>x</sub> emissions rates and vehicle type (car, truck, bus) within 100 m	NO <sub>x</sub> emissions from all on-road motor vehicles based on vehicle miles and the relative emission rates of different vehicle types
Length of truck route within 50 m	Diesel exhaust
NO <sub>x</sub> emissions from heat and hot water boilers in buildings within 400 m, taking into account changes in building heating fuels over time.	Combustion of heating oil and natural gas
Number of building boilers within 250 m	Combustion of heating oil and natural gas

Figure 8: NO concentrations, 2016 annual average (left) and 2009-2016 annual averages (right)




# SULFUR DIOXIDE

At NYCCAS sites measured in each season for eight winters:

- In the most recent winter (2015-2016), 28% of sites measured SO<sub>2</sub> levels below the detection limit of the monitor (0.25 ppb).
- At the 72% of sites with levels above the detection limit, in winter 2015-2016, seasonally adjusted wintertime SO<sub>2</sub> concentrations declined by an average of 0.85 ppb per year between Winter 2008-2009 and Winter 2015-2016.
- At the 43 monitoring sites with levels above the detection limit in the most recent winter, citywide, wintertime average levels declined by 95% between winter 2008-2009 and winter 2015-2016.
- In the most recent winter (2015-2016), at the at the 43 monitoring sites with levels above the detection limit, seasonal average concentrations across NYCCAS monitoring sites ranged from 0.10 to 0.86 ppb.

Figure 9: SO<sub>2</sub> levels at NYCCAS monitors, by density of nearby residual oil boilers (A) and nearby nighttime population density (B)<sup>4</sup>




<sup>4</sup>Residual oil boiler density were estimated as number of Nos. 4 and 6 boilers within 1,000 m of monitoring sites. Nighttime population density was estimated as total nighttime population within 1,000 m of monitoring sites. High, Medium, and Low represent one third of sites ranked by source indicator density.

- Across the eight-year period, higher levels were consistently seen at sites in areas of high residual oil boiler density (Figures 9-10).

In the spatial model, the most important predictors of SO<sub>2</sub> concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
Time varying counts of boilers burning No. 4 and No. 6 oil within 1,000 m	Combustion of No. 4 and No. 6 heating oil, accounting for season-specific estimated counts of boilers.
Nighttime population within 1,000 m	Combustion of heating oil

Figure 10: SO<sub>2</sub> concentrations, 2015-2016 winter average (left) and 2009-2016 winter averages (right)






# BLACK CARBON

At NYCCAS sites measured in each season for eight years:

- Seasonally adjusted street-level BC concentrations declined by an average of 0.5 absorbance units (abs) per year.
- Citywide, annual average levels at the 60 monitoring sites (locations measured across the eight years) declined by 24% between 2009 and 2016.
- In the most recent year (2016), seasonal average concentrations across NYCCAS monitoring sites ranged from 0.4 to 2.3 abs.
- Across the eight-year period, higher levels were consistently seen at sites in industrial areas (reflecting increased truck traffic density) and in areas of high traffic emissions (Figures 11-12).

Figure 11: BC levels at NYCCAS monitors, by area of industrial tax lots (A) and nearby traffic emissions density (B)<sup>5</sup>




<sup>5</sup>Industrial land use area was estimated as the total area of industrial lots within 1,000 m. Density of nearby traffic emissions was estimated as traffic density, weighted by relative  $PM_{2.5}$  emissions rates by vehicle type, within 1,000 m. High, Medium, and Low represent one third of sites ranked by source indicator density.

In the spatial model, the most important predictors of BC concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
Area of industrial land use within 1,000 m	Diesel exhaust particles from trucks idling and traveling through industrial areas, industrial combustion equipment
Traffic density, weighted by relative PM <sub>2.5</sub> emissions rates and vehicle type (car, truck, bus) within 1,000 m	PM <sub>2.5</sub> emissions from all on-road motor vehicles based on vehicle miles and the relative emission rates of different vehicle types
Number of building boilers within 200 m	Combustion of heating oil and natural gas
Road length, weighted by traffic, within 50 m	Emissions from motor vehicles
Percent impervious road surface within 250 m	Emissions of motor vehicles on paved roadways

Figure 12: BC concentrations, 2016 annual average (left) and 2009-2016 annual averages (right)




# OZONE

At NYCCAS sites measured in each season:

- Since the first summer (2009), when relatively cool temperatures contributed to lower levels of O<sub>3</sub> citywide, summertime average levels varied minimally year to year (ranging from 29.0 ppb to 31.4 ppb between 2010 and 2016), without a consistent trend over this time period.
- In the most recent summer (2016), seasonal average O<sub>3</sub> varied from 21.8 to 43.5 ppb across the monitoring sites.
- Higher levels were consistently measured at sites with lower traffic density and lower NO<sub>2</sub> concentrations, reflecting less removal of ozone from the atmosphere in areas of fewer fresh combustion emissions (Figures 13-14).

Figure 13: O<sub>3</sub> levels at NYCCAS monitors, by nearby traffic density (A) and co-located NO<sub>2</sub> concentration (B)<sup>6</sup>




<sup>6</sup> Traffic density was estimated as annual average daily traffic (all types) within 1,000 m of monitoring sites. NO<sub>2</sub> concentrations are based on seasonal average monitored NO<sub>2</sub> levels at the same location. High, Medium, and Low represent one third of sites ranked by source indicator density.

In the spatial model, the most important predictors of O<sub>3</sub> concentrations were, in order of importance:

Indicator	Associated Sources and Interpretation
Level of NO <sub>2</sub> measured at the same location	Nitrogen oxides at elevated concentrations react with ground-level ozone and reduce levels
Tree cover within 50 m	Reduced levels through reactions of ozone with leaf surfaces

Figure 14: O<sub>3</sub> concentrations, 2016 summer average (left) and 2009-2016 summer averages (right)



# APPENDICES

1. DATA SOURCES FOR  
EMISSIONS INDICATORS

2. SEASONAL AVERAGE  
POLLUTANT MAPS

3. COMMUNITY DISTRICT  
AVERAGE POLLUTANT  
LEVELS



## APPENDIX 1

### DATA SOURCES FOR EMISSIONS INDICATORS





Source Category	Variables Examined (most calculated in buffers of 50 to 1,000 m)	Data Source
Cumulative Traffic Indicators	Unweighted and kernel-weighted traffic density	New York Metropolitan Transportation Council (NYMTC) traffic data, 2005; and U.S. Federal Highway Administration Highway Performance Monitoring System (HPMS) data, 2007
	Road density	Accident Location Information System (ALIS) road network data, 2008
	Kernel-weighted road density	ALIS network data
	Road density weighted by functional class	ALIS network; MPSI TrafficMetrix™ data, 1989-2006
	Road density kernel-weighted by functional class	ALIS network; MPSI TrafficMetrix™ data
	Traffic density weighted by relative emissions rates	NYMTC traffic data; emissions factors from Environmental Protection Agency's AP 42 database
	Number of signaled intersections	NYC Department of Transportation (DOT), 2008
Road-specific Measures	Average daily traffic (ADT) on nearest major road	NYMTC traffic data
	ADT/ Distance to nearest major road	NYMTC traffic data
	Location on a bus route	NYC DOT
	Distance to nearest road, by functional class	ALIS network; MPSI TrafficMetrix™ data





Source Category	Variables Examined (most calculated in buffers of 50 to 1,000 m)	Data Source
Truck/ Diesel-Related Measures	Unweighted traffic on designated truck routes	NYMTC traffic data
	Unweighted density of truck routes	NYMTC traffic data
	Kernel-weighted density of truck routes	NYMTC traffic data
	Distance to nearest truck route	NYMTC traffic data
	Trucks per day on nearest major road	NYMTC traffic data
Population Metrics	Census population density	U.S. Census Bureau 2000 data
	LandScan daytime, nighttime population density	Oak Ridge National Laboratory LandScan™ data, 2006
Built Space	Density of built space (building floor area)	NYC Department of City Planning Primary Land Use Tax Lot Output (PLUTO™) data, 2007
	Density of residential units	PLUTO™ data
	Total residential, factory, garage floor area	PLUTO™ data
	Estimated building boiler emissions for building heat and hot water	PLUTO™ data, EPA AP 42, NYC Department of Environmental Protection (NYC DEP) Registration and Certificate Permit Data, updated annually
	Area of commercial floor area	PLUTO™ data
Land Use	Area of industry and manufacturing	PLUTO™ data
	Area of heavy manufacturing	PLUTO™ data
	Area of gas stations	PLUTO™ data
	Area of tree cover	NYC Department of Parks and Recreation LiDAR data, 2010
	Percent impervious surface	United States Geological Survey, 2006
	Dominant land use type	PLUTO™ data





Source Category	Variables Examined (most calculated in buffers of 50 to 1,000 m)	Data Source
Permitted Emissions	Number of DEC permitted combustion sources	NYS Department of Environmental Conservation (DEC) permit data, 2005
	Number of DEP permitted combustion sources	NYC DEP permit data, 2008
	Number of DOB permitted boilers	NYC Department of Buildings (DOB) permit data, 2008
	Number of permitted combustion sources by fuel type (oil 2, 4, 6, natural gas)	DEP permit data, updated every 6 months
	Total BTU by fuel type (oil 2, 4, 6, natural gas)	DEP permit data
	Average BTU by fuel type (oil 2, 4, 6, natural gas)	DEP permit data
Transportation Facilities	Number of bus depots	NYC Department of Citywide Administrative Services (DCAS), 2008
	Minimum distance to bus depot, school bus depot	NYC DCAS; NYC Department of Education (DOE)
	Number of school bus depots	NYC DOE
	Number of school buses at nearest depot	NYC DOE
Distributed Facilities	Number of waste transfer stations	NYC Department of Sanitation inspections
	Minimum distance to waste transfer station, ferry terminal, water treatment facility	NYC DCAS
	Distance to nearest port, airport	NYC Office of Emergency Management







# APPENDIX 2 SEASONAL AVERAGE POLLUTANT MAPS, PM<sub>2.5</sub>, NO<sub>2</sub>, NO, BC









# APPENDIX 3 COMMUNITY DISTRICT AVERAGE POLLUTANT LEVELS

Table A3-1: Community district, annual average PM<sub>2.5</sub> and per-year decline in levels

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Manhattan	Midtown (CD5)	105	16.1	15.5	14.8	14.2	14.3	14.2	12.3	11.34	-0.61
Bronx	Fordham and University Heights (CD5)	205	12.1	11.4	11.7	10.5	10.1	10.1	9.8	8.34	-0.47
Bronx	Highbridge and Concourse (CD4)	204	12	11.2	11.6	10.4	10	10	9.7	8.25	-0.46
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	14.1	13.1	13.1	12.3	12.3	12.4	11.2	10.21	-0.46
Manhattan	Washington Heights and Inwood (CD12)	112	11.6	10.7	11.1	9.9	9.5	9.5	9.4	7.79	-0.46
Bronx	Kingsbridge Heights and Bedford (CD7)	207	11.5	10.7	11.1	9.8	9.4	9.5	9.3	7.78	-0.46
Manhattan	Upper East Side (CD8)	108	12.9	11.9	12.1	11.2	11.1	11.2	10.2	9.1	-0.45
Manhattan	Central Harlem (CD10)	110	11.6	10.6	11.1	9.9	9.6	9.7	9.3	7.89	-0.44
Manhattan	Morningside Heights and Hamilton Heights (CD9)	109	11.7	10.7	11.2	10.1	9.8	9.8	9.4	8.01	-0.44
Bronx	Morrisania and Crotona (CD3)	203	11.4	10.6	11.1	9.9	9.4	9.5	9.3	7.85	-0.44
Bronx	Belmont and East Tremont (CD6)	206	11.5	10.7	11.2	9.9	9.5	9.6	9.4	7.96	-0.43
Manhattan	Upper West Side (CD7)	107	12.2	11	11.5	10.4	10.3	10.4	9.8	8.49	-0.42
Manhattan	Financial District (CD1)	101	13.1	12.1	12.3	11.3	11.1	11.4	10.6	9.56	-0.42
Bronx	Riverdale and Fieldston (CD8)	208	11	10.1	10.7	9.3	8.9	9	9.2	7.49	-0.41
Bronx	Mott Haven and Melrose (CD1)	201	11.9	10.9	11.6	10.4	10	10.1	9.9	8.55	-0.40
Staten Island	Tottenville and Great Kills (CD3)	503	9.7	8.8	9.3	8.3	7.8	8.2	7.3	6.62	-0.39
Bronx	Parkchester and Soundview (CD9)	209	10.8	10	10.7	9.4	8.9	9.1	9	7.6	-0.39
Queens	Rego Park and Forest Hills (CD6)	406	10.6	9.8	10.1	9.1	8.7	8.9	8.5	7.5	-0.38
Manhattan	East Harlem (CD11)	111	11.5	10.4	11.1	10	9.7	9.8	9.5	8.17	-0.38
Manhattan	Clinton and Chelsea (CD4)	104	13.2	11.9	12.4	11.4	11.4	11.6	10.9	9.76	-0.37
Bronx	Morris Park and Bronxdale (CD11)	211	10.6	9.7	10.4	9.2	8.7	8.9	8.9	7.4	-0.37
Bronx	Hunts Point and Longwood (CD2)	202	11.6	10.7	11.5	10.3	9.8	10	9.8	8.49	-0.37
Manhattan	Greenwich Village and Soho (CD2)	102	12.8	11.6	12.1	11.1	10.9	11.2	10.6	9.53	-0.37
Queens	Hillcrest and Fresh Meadows (CD8)	408	10	9.3	9.7	8.6	8.2	8.4	8.2	7.01	-0.37
Brooklyn	Flatbush and Midwood (CD14)	314	10.3	9.7	10.1	8.9	8.6	8.9	8.4	7.45	-0.36
Queens	Kew Gardens and Woodhaven (CD9)	409	10.3	9.5	9.7	8.6	8.4	8.7	8.3	7.29	-0.36
Queens	Elmhurst and Corona (CD4)	404	10.7	9.8	10.3	9.2	8.9	9.2	8.7	7.71	-0.36
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	10.6	9.9	10.2	9.1	8.6	9	8.7	7.77	-0.36
Queens	Flushing and Whitestone (CD7)	407	10.2	9.4	10.1	8.9	8.4	8.7	8.5	7.26	-0.35
Queens	Jackson Heights (CD3)	403	10.3	9.2	9.9	8.8	8.4	8.7	8.3	7.27	-0.35

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Brooklyn	East Flatbush (CD17)	317	10.6	9.9	10.2	9.1	8.7	9.1	8.7	7.79	-0.35
Bronx	Williamsbridge and Baychester (CD12)	212	10.6	9.6	10.4	9	8.7	8.9	9.1	7.46	-0.35
Queens	South Ozone Park and Howard Beach (CD10)	410	9.8	9	9.2	8.1	8	8.2	7.9	6.85	-0.35
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	10.8	10	10.4	9.2	8.8	9.2	8.9	7.98	-0.35
Brooklyn	Borough Park (CD12)	312	10.3	9.6	10.1	8.9	8.5	8.9	8.5	7.49	-0.35
Brooklyn	Bensonhurst (CD11)	311	9.8	9.2	9.7	8.6	8.2	8.6	8	7.03	-0.35
Brooklyn	Brownsville (CD16)	316	10.8	10	10.3	9.1	8.8	9.2	8.9	7.97	-0.34
Queens	Jamaica and Hollis (CD12)	412	9.9	9.2	9.5	8.4	8.3	8.4	8.2	6.98	-0.34
Brooklyn	Coney Island (CD13)	313	9.4	8.8	9.4	8.2	8	8.3	7.6	6.65	-0.34
Brooklyn	Sheepshead Bay (CD15)	315	9.6	8.9	9.5	8.3	8.1	8.4	7.8	6.78	-0.34
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	10.2	9.4	10	8.8	8.4	8.8	8.4	7.39	-0.34
Staten Island	South Beach and Willowbrook (CD2)	502	9.8	8.8	9.5	8.3	8.1	8.5	7.8	6.89	-0.34
Queens	Bayside and Little Neck (CD11)	411	9.7	9	9.7	8.5	8.1	8.3	8.2	6.87	-0.34
Bronx	Throgs Neck and Co-op City (CD10)	210	10.5	9.5	10.4	9.1	8.7	8.9	9	7.51	-0.34
Queens	Queens Village (CD13)	413	9.4	8.7	9.2	8.1	7.9	8	7.9	6.53	-0.33
Brooklyn	East New York and Starrett City (CD5)	305	10.5	9.7	10	8.9	8.7	9	8.7	7.73	-0.33
Brooklyn	Bedford Stuyvesant (CD3)	303	10.8	9.9	10.4	9.2	8.8	9.3	9	8.09	-0.32
Brooklyn	Flatlands and Canarsie (CD18)	318	9.8	9.1	9.5	8.3	8.2	8.5	8.1	7.08	-0.32
Staten Island	St. George and Stapleton (CD1)	501	10	8.9	9.6	8.4	8.1	8.6	8.2	7.11	-0.32
Manhattan	Lower East Side and Chinatown (CD3)	103	11.8	10.6	11.3	10.1	9.9	10.3	9.9	8.9	-0.32
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	11.6	10.5	11.1	9.9	9.5	10	9.8	8.83	-0.32
Brooklyn	Bushwick (CD4)	304	10.8	9.8	10.3	9.2	8.8	9.3	9	8.07	-0.32
Queens	Ridgewood and Maspeth (CD5)	405	10.7	9.7	10.2	9.1	8.8	9.2	8.9	7.97	-0.31
Queens	Long Island City and Astoria (CD1)	401	10.7	9.4	10.4	9.2	8.9	9.2	9	7.82	-0.31
Brooklyn	Sunset Park (CD7)	307	11.1	10.3	10.9	9.7	9.2	9.7	9.6	8.52	-0.31
Queens	Rockaway and Broad Channel (CD14)	414	8.8	8	8.4	7.4	7.6	7.7	7.2	5.98	-0.31
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	11.4	10.4	11.1	9.9	9.4	9.9	9.9	8.86	-0.29
Queens	Woodside and Sunnyside (CD2)	402	11.9	10.6	11.5	10.3	10.1	10.5	10.3	9.27	-0.28
Brooklyn	Greenpoint and Williamsburg (CD1)	301	12	10.7	11.6	10.4	10.1	10.6	10.6	9.55	-0.25

Table A3-2: Community district, annual average NO<sub>2</sub> and per-year decline in levels

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Manhattan	Midtown (CD5)	105	46.8	42.1	41.8	40	38.7	37.1	32.9	31.64	-1.99
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	39.9	36.2	36.4	34.5	33.4	32.2	29.3	28.4	-1.53
Manhattan	Financial District (CD1)	101	36.6	33.4	33.5	31.1	30.3	28.9	26.7	26.27	-1.43
Manhattan	Clinton and Chelsea (CD4)	104	36.3	32.9	33.5	31.4	30.6	29.2	27.3	26.68	-1.30
Manhattan	Upper East Side (CD8)	108	34.5	31.2	31.7	29.6	28.7	27.8	25.8	24.91	-1.27
Manhattan	Greenwich Village and Soho (CD2)	102	34.8	31.8	32.3	30.1	29.3	28.1	26.3	25.94	-1.23
Manhattan	Morningside Heights and	109	31	27.6	28.1	25.5	25.2	23.6	22.9	22	-1.19
Manhattan	Central Harlem (CD10)	110	31.2	27.9	28.3	25.9	25.4	24	23.3	22.27	-1.18
Manhattan	Upper West Side (CD7)	107	31.9	28.6	29.3	27.1	26.5	25.1	23.9	23.15	-1.17
Bronx	Highbridge and Concourse	204	29.7	26.6	26.9	24.5	24	22.7	22.4	21.21	-1.11
Manhattan	Washington Heights and Lower East Side and	112	27.9	24.7	25	22.6	22.3	20.6	20.7	19.36	-1.11
Manhattan	Chinatown (CD3)	103	31.8	29.2	29.7	27.4	26.6	25.7	24.3	24	-1.09
Bronx	Fordham and University Heights (CD5)	205	29.6	26.5	26.8	24.6	23.9	22.7	22.7	21.18	-1.08
Manhattan	East Harlem (CD11)	111	29.8	27	27.6	25.2	24.7	23.6	22.9	22.09	-1.04
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	29.3	27.1	27.4	24.8	24.2	23.4	22.4	22.16	-1.02
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	27.8	25.9	26	23.4	22.8	22.3	21.3	20.74	-1.00
Bronx	Morrisania and Crotona (CD3)	203	27.7	25	25.3	23	22.3	21.6	21.4	20.07	-0.99
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	27	25.3	25.2	22.7	22.1	21.7	20.8	20.11	-0.97
Brooklyn	Bedford Stuyvesant (CD3)	303	27.6	25.7	25.9	23.5	22.8	22.2	21.3	20.79	-0.97
Bronx	Mott Haven and Melrose (CD1)	201	28.1	25.5	25.9	23.5	22.9	22.1	21.8	20.84	-0.97
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	27.7	25.9	26	23.3	22.7	22	21.4	21.17	-0.96
Bronx	Kingsbridge Heights and Bedford (CD7)	207	26.7	23.8	24.2	22.5	21.5	20.7	20.9	19.15	-0.94
Bronx	Belmont and East Tremont (CD6)	206	27.1	24.4	24.8	22.8	21.9	21.4	21.4	19.86	-0.91
Brooklyn	Sunset Park (CD7)	307	25.5	24	24	21.2	20.7	20.3	19.8	19.44	-0.89
Brooklyn	Brownsville (CD16)	316	25.6	24	24.2	21.9	21.4	20.9	20.1	19.29	-0.88
Bronx	Riverdale and Fieldston (CD8)	208	23.2	20.1	20.8	19.1	18.2	17.1	17.8	16.04	-0.88
Brooklyn	Bushwick (CD4)	304	25.9	24.2	24.7	22.4	21.7	21.2	20.3	19.82	-0.87
Brooklyn	East Flatbush (CD17)	317	25.1	23.6	23.6	21.3	20.7	20.5	19.8	18.94	-0.86
Brooklyn	Borough Park (CD12)	312	25.2	23.8	23.6	21.1	20.5	20.6	19.9	19.12	-0.85
Brooklyn	Flatbush and Midwood (CD14)	314	24.4	23	22.9	20.5	19.9	20	19.3	18.29	-0.84
Bronx	Hunts Point and Longwood (CD2)	202	26.1	23.9	24.3	22.1	21.3	21	20.9	19.79	-0.83
Brooklyn	Greenpoint and Williamsburg (CD1)	301	27	25.2	25.9	23.6	22.9	22.4	21.7	21.38	-0.81
Queens	Rego Park and Forest Hills (CD6)	406	25	23.5	24.1	21.8	21.1	20.9	20	19.31	-0.81
Queens	Elmhurst and Corona (CD4)	404	26.9	25.3	25.9	23.7	22.8	22.8	21.9	21.21	-0.80
Brooklyn	East New York and Starrett City (CD5)	305	23.4	22	22.4	20.2	19.8	19.3	18.7	17.83	-0.78
Queens	Kew Gardens and Woodhaven (CD9)	409	24.1	22.8	23.3	21	20.6	20.1	19.5	18.69	-0.77
Queens	Ridgewood and Maspeth (CD5)	405	24.1	22.6	23.3	21.1	20.4	20.1	19.4	18.83	-0.75
Bronx	Parkchester and Soundview (CD9)	209	24.6	22.6	23	21.1	19.9	20.2	20.1	18.81	-0.75
Queens	Long Island City and Astoria (CD1)	401	25.4	23.5	24.4	22.2	21.4	21.2	20.6	20.07	-0.74

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Queens	Jackson Heights (CD3)	403	25.6	24	24.7	22.6	21.6	21.7	21	20.32	-0.74
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	23.6	22.4	22.6	20.1	19.5	19.6	19.2	18.54	-0.73
Queens	Woodside and Sunnyside (CD2)	402	25.9	24.1	25.1	23	22.1	21.9	21.2	20.79	-0.72
Queens	South Ozone Park and Howard Beach (CD10)	410	22.2	21	21.5	19.3	19	18.6	18.2	17.21	-0.69
Queens	Hillcrest and Fresh Meadows (CD8)	408	22.4	21.3	21.7	19.4	18.7	18.8	18.3	17.63	-0.69
Brooklyn	Bensonhurst (CD11)	311	23	21.9	21.9	19.7	19	19.7	19.1	18.05	-0.67
Queens	Jamaica and Hollis (CD12)	412	22.3	21.3	21.6	19.4	19.1	18.9	18.5	17.63	-0.66
Bronx	Morris Park and Bronxdale (CD11)	211	23.4	21.2	21.8	20.4	19	19.4	19.6	17.91	-0.66
Brooklyn	Flatlands and Canarsie (CD18)	318	20.3	19.2	19.2	17.3	16.7	17.1	16.6	15.36	-0.65
Queens	Flushing and Whitestone (CD7)	407	22.7	21.5	21.9	19.8	18.7	19.4	19	18.2	-0.63
Staten Island	St. George and Stapleton (CD1)	501	20.7	19.3	20.5	18.1	17.6	17.1	16.9	16.86	-0.59
Bronx	Williamsbridge and Baychester (CD12)	212	22.1	19.7	20.6	19.5	18	18.3	19	16.93	-0.57
Brooklyn	Sheepshead Bay (CD15)	315	20.4	19.4	19.4	17.6	16.8	18	17.4	15.91	-0.55
Queens	Bayside and Little Neck (CD11)	411	20.5	19.6	19.9	17.9	16.8	17.8	17.6	16.78	-0.52
Queens	Queens Village (CD13)	413	19.9	19.1	19.3	17.4	16.9	17.3	17.2	16.21	-0.50
Bronx	Throgs Neck and Co-op City (CD10)	210	21.3	19.8	20.4	19	17.4	18.5	18.8	17.23	-0.49
Brooklyn	Coney Island (CD13)	313	18.4	17.6	17.7	16	15.1	16.6	16.1	14.64	-0.45
Staten Island	South Beach and Willowbrook (CD2)	502	16.5	15.2	16.3	14.5	13.8	14	13.6	13.36	-0.45
Staten Island	Tottenville and Great Kills (CD3)	503	14.7	13	13.3	12.4	11.4	12.5	11.8	11.44	-0.38
Queens	Rockaway and Broad Channel (CD14)	414	14.6	13.9	14.2	13	12.5	13.7	13.4	11.6	-0.30



Table A3-3: Community district, annual average NO and per-year decline in levels

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Manhattan	Midtown (CD5)	105	57.7	52.1	56	48.1	43.4	47.8	40.4	29.85	-3.37
Manhattan	Upper East Side (CD8)	108	51.8	46	50	42.2	37.7	41.3	34.2	27.23	-3.11
Manhattan	Upper West Side (CD7)	107	46.6	40.9	44.6	37.4	33.2	37.3	30.4	22.34	-2.96
Manhattan	Greenwich Village and Soho (CD2)	102	52.7	46.4	50.7	41.7	37.2	42.3	34.4	32.85	-2.72
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	49.3	44.7	48.2	41.2	37.4	42.3	36.4	27.43	-2.57
Manhattan	Clinton and Chelsea (CD4)	104	44.2	39.1	42.7	35.7	32.1	37.5	30.8	26.98	-2.16
Manhattan	Morningside Heights and Hamilton Heights (CD9)	109	34.1	29	31.9	26	23	27.5	22	16.92	-2.04
Bronx	Fordham and University Heights (CD5)	205	34.2	29.2	31.6	26.3	23.7	27.1	22.8	17.32	-1.98
Manhattan	Central Harlem (CD10)	110	33.6	28.3	31.4	25.3	22.4	26.9	21.4	17.22	-1.97
Manhattan	Washington Heights and Inwood (CD12)	112	32.3	27.2	29.6	24.4	21.7	25.5	20.7	16.36	-1.89
Manhattan	Lower East Side and Chinatown (CD3)	103	40.9	36	39.6	31.9	28.7	34.4	28.1	27	-1.85
Bronx	Kingsbridge Heights and Bedford (CD7)	207	31.1	26.1	28.4	23.3	20.8	23.6	20.1	15.84	-1.83
Bronx	Highbridge and Concourse (CD4)	204	32.3	27.7	30.2	24.9	22.2	26	21.8	16.75	-1.83
Manhattan	East Harlem (CD11)	111	33.4	28.9	31.9	26.2	23.5	28.3	23.4	18.56	-1.72
Manhattan	Financial District (CD1)	101	39.5	35.6	38.7	32.3	29.4	35.2	29.7	25.8	-1.65
Bronx	Belmont and East Tremont (CD6)	206	29.7	25	27.5	22.4	20.2	23.4	20	16.53	-1.57
Bronx	Morrisania and Crotona (CD3)	203	27.6	23.2	25.8	20.6	18.4	22.1	18.8	15.35	-1.44
Bronx	Mott Haven and Melrose (CD1)	201	28.4	24.3	26.9	21.8	19.5	23.9	19.9	16.34	-1.40
Bronx	Riverdale and Fieldston (CD8)	208	24.4	19.6	21.5	17.1	15.1	18.5	15	13.11	-1.35
Bronx	Hunts Point and Longwood (CD2)	202	26.1	22.2	24.8	19.8	17.7	21.4	18.4	15.69	-1.24
Queens	Rego Park and Forest Hills (CD6)	406	27.5	24.9	27.3	21.5	19.4	23.2	20.8	18.17	-1.19
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	28.3	25.3	27.9	21.6	18.9	24.1	20.6	19.9	-1.15
Bronx	Morris Park and Bronxdale (CD11)	211	23.7	19.8	22	17.4	15.8	18.4	16.5	14.21	-1.13
Queens	Long Island City and Astoria (CD1)	401	24.7	21.2	24	18.5	16.3	21.3	17.4	15.36	-1.13
Bronx	Parkchester and Soundview (CD9)	209	25.8	22.2	24.6	19.8	17.9	20.9	18.8	16.63	-1.12
Queens	Jackson Heights (CD3)	403	24.9	21.8	24.5	18.9	16.8	21.1	18.3	15.93	-1.10
Queens	Elmhurst and Corona (CD4)	404	26.6	23.8	26.4	20.6	18.5	22.9	20.1	17.82	-1.10
Queens	Woodside and Sunnyside (CD2)	402	27.1	23.9	26.6	20.9	18.7	24.3	20.2	17.98	-1.09
Queens	Hillcrest and Fresh Meadows (CD8)	408	24	21.8	24.3	18.9	17.1	19.2	18.4	16.11	-1.06
Bronx	Williamsbridge and Baychester (CD12)	212	23.1	18.9	20.9	16.6	15	17.6	15.9	14.13	-1.06
Brooklyn	Bushwick (CD4)	304	26.2	23.3	25.8	19.5	16.9	22.2	18.9	18.54	-1.06
Queens	Bayside and Little Neck (CD11)	411	22.6	20.5	23.2	18.1	16.5	17.3	17.4	15.08	-1.04
Queens	Flushing and Whitestone (CD7)	407	23.1	20.4	23	17.8	16	18.3	17.2	15.2	-1.04
Queens	Jamaica and Hollis (CD12)	412	23	21.3	23.7	18.5	16.8	18.3	17.8	15.64	-1.03
Queens	Queens Village (CD13)	413	21.3	19.9	22.4	17.6	16	16.2	16.7	14.39	-1.01
Queens	Kew Gardens and Woodhaven (CD9)	409	24.5	22.3	24.6	19	17	20.6	18.7	17.11	-1.00
Brooklyn	Greenpoint and Williamsburg (CD1)	301	27.6	24.5	27.2	21.1	18.6	24.6	20.5	20.06	-0.99

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	26.7	24.3	26.8	21.1	18.9	23.8	20.5	19.32	-0.97
Bronx	Throgs Neck and Co-op City (CD10)	210	24	20.7	23.1	18.4	16.9	19.1	18.2	16.14	-0.96
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	30	27	29.7	23.8	21.4	27.3	23.3	22.95	-0.92
Brooklyn	Flatbush and Midwood (CD14)	314	24.4	22.6	25.2	20.3	18.3	22.1	19.6	17.37	-0.90
Brooklyn	Bedford Stuyvesant (CD3)	303	25.7	22.9	25.4	19.4	16.9	22.4	19.5	19.03	-0.90
Queens	Ridgewood and Maspeth (CD5)	405	23.9	21.4	23.7	18	15.9	20.9	18.1	17.19	-0.88
Queens	South Ozone Park and Howard Beach (CD10)	410	22.4	20.8	22.8	17.7	15.9	19	17.5	16.1	-0.88
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	25.9	23.2	25.9	20.2	17.6	23.2	19.8	19.42	-0.87
Brooklyn	Brownsville (CD16)	316	24.4	22.2	24.4	18.8	16.4	21.2	18.9	18.3	-0.85
Brooklyn	East Flatbush (CD17)	317	24.3	22.4	24.7	19.5	17.5	21.9	19.4	18.17	-0.81
Brooklyn	East New York and Starrett City (CD5)	305	23.8	21.9	24	18.6	16.6	20.8	18.7	18.08	-0.81
Brooklyn	Sunset Park (CD7)	307	23.6	21.4	24.1	19	16.8	21.5	18.7	18.21	-0.73
Brooklyn	Borough Park (CD12)	312	21.6	19.9	22.6	17.9	15.9	19.8	17.5	16.14	-0.72
Queens	Rockaway and Broad Channel (CD14)	414	18.2	18	20.3	16.8	15.3	15.5	15.9	13.59	-0.70
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	21.1	19.4	22.3	17.9	15.9	19.5	17.5	15.66	-0.69
Brooklyn	Bensonhurst (CD11)	311	20	18.7	21.4	17.4	15.5	18.4	16.9	14.91	-0.66
Brooklyn	Sheepshead Bay (CD15)	315	19.2	18.5	21	17.3	15.6	17.8	17	14.65	-0.60
Brooklyn	Flatlands and Canarsie (CD18)	318	19.6	18.6	20.8	16.4	14.8	18.1	16.7	15.48	-0.57
Staten Island	St. George and Stapleton (CD1)	501	16.8	14.7	17.3	14.2	12.3	15.4	13.1	12.34	-0.56
Staten Island	Tottenville and Great Kills (CD3)	503	12.9	11.2	12.7	12.5	10.7	9.8	9.6	8.93	-0.55
Staten Island	South Beach and Willowbrook (CD2)	502	14.7	12.8	15.1	13.2	11.3	12.9	11.5	10.83	-0.50
Brooklyn	Coney Island (CD13)	313	17.3	16.6	19.2	16.1	14.4	16.1	15.6	13.72	-0.49

Table A3-4: Community district, wintertime average SO<sub>2</sub> and per-year decline in levels

Borough	Geography	ID	Winter Average 2008-2009	Winter Average 2009-2010	Winter Average 2010-2011	Winter Average 2011-2012	Winter Average 2012-2013	Winter Average 2013-2014	Winter Average 2014-2015	Winter Average 2015-2016	Slope (per year decline)
Bronx	Fordham and University Heights (CD5)	205	12.7	8.2	10.5	6.8	4.5	3.8	1.8	0.57	-1.66
Manhattan	Upper East Side (CD8)	108	12.1	8.5	10.4	6.3	4.1	3.8	1.5	0.45	-1.65
Bronx	Kingsbridge Heights and Bedford (CD7)	207	11.8	7.5	9.5	6.3	4	3.3	1.8	0.55	-1.53
Manhattan	Washington Heights and Inwood (CD12)	112	11.4	7.5	9.5	6.3	4.3	3.4	1.6	0.51	-1.50
Manhattan	Upper West Side (CD7)	107	11	7.6	9.4	6	4	3.5	1.4	0.44	-1.48
Manhattan	Midtown (CD5)	105	10.3	7	8.8	5.3	3.2	3.3	1.4	0.43	-1.38
Bronx	Highbridge and Concourse (CD4)	204	10.4	6.9	8.8	5.5	3.9	3.2	1.4	0.45	-1.38
Manhattan	Morningside Heights and Hamilton Heights (CD9)	109	9.3	6.3	7.9	5.1	3.5	2.9	1.2	0.39	-1.24
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	9.2	6.3	7.9	4.6	2.8	3	1.2	0.38	-1.24
Manhattan	Greenwich Village and Soho (CD2)	102	8.7	5.7	7.3	4.3	2.4	2.7	1.1	0.36	-1.16
Manhattan	Central Harlem (CD10)	110	8.2	5.6	7	4.4	3.1	2.7	1.1	0.36	-1.09
Bronx	Belmont and East Tremont (CD6)	206	8.2	5.3	6.8	4.3	2.8	2.5	1.3	0.38	-1.06
Manhattan	Clinton and Chelsea (CD4)	104	7.3	4.9	6.2	3.8	2.3	2.4	1	0.31	-0.97
Manhattan	Lower East Side and Chinatown (CD3)	103	7	4.6	5.9	3.3	1.8	2.2	0.9	0.29	-0.93
Manhattan	East Harlem (CD11)	111	7	4.8	6	3.7	2.5	2.4	1	0.3	-0.93
Bronx	Morrisania and Crotona (CD3)	203	7	4.7	6	3.6	2.4	2.2	1	0.32	-0.93
Bronx	Riverdale and Fieldston (CD8)	208	6.7	4.2	5.4	3.8	2.3	1.7	1	0.3	-0.87
Queens	Elmhurst and Corona (CD4)	404	6.3	4.5	5.6	2.8	1.8	2.4	0.9	0.3	-0.84
Queens	Jackson Heights (CD3)	403	6.1	4.3	5.4	2.8	1.8	2.3	0.9	0.28	-0.81
Bronx	Parkchester and Soundview (CD9)	209	5.9	3.9	4.9	2.9	1.9	2.1	1	0.29	-0.75
Bronx	Mott Haven and Melrose (CD1)	201	5.7	3.9	4.9	3	2.1	2	0.9	0.26	-0.75
Bronx	Morris Park and Bronxdale (CD11)	211	5.9	3.8	4.8	3	1.8	1.9	1.1	0.31	-0.74
Queens	Rego Park and Forest Hills (CD6)	406	5.3	3.8	4.7	2.3	1.3	1.8	0.7	0.26	-0.72
Bronx	Williamsbridge and Baychester (CD12)	212	5.8	3.6	4.5	3	1.7	1.7	1.2	0.31	-0.72
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	5	3.4	4.2	2	1.1	1.3	0.5	0.22	-0.69
Bronx	Hunts Point and Longwood (CD2)	202	5.1	3.5	4.4	2.5	1.7	1.8	0.8	0.25	-0.67
Queens	Long Island City and Astoria (CD1)	401	5	3.6	4.5	2.5	1.7	1.9	0.8	0.24	-0.67
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	4.8	3.2	4	2	1.1	1.3	0.5	0.21	-0.65
Queens	Woodside and Sunnyside (CD2)	402	4.8	3.4	4.3	2.3	1.5	1.7	0.7	0.22	-0.64
Brooklyn	Bushwick (CD4)	304	4.7	3.3	4.1	2	1.3	1.5	0.6	0.22	-0.64
Brooklyn	Flatbush and Midwood (CD14)	314	4.7	3.1	3.8	1.8	1	1.1	0.5	0.21	-0.63
Brooklyn	Bedford Stuyvesant (CD3)	303	4.7	3.2	4	2	1.2	1.4	0.6	0.21	-0.63
Manhattan	Financial District (CD1)	101	4.8	3	4	2.3	1.2	1.4	0.6	0.21	-0.63
Brooklyn	East Flatbush (CD17)	317	4.4	3	3.6	1.7	1	1.1	0.5	0.21	-0.60
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	4.4	2.9	3.7	1.9	1	1.2	0.5	0.19	-0.59
Brooklyn	Brownsville (CD16)	316	4.3	3.1	3.7	1.7	1.1	1.2	0.5	0.21	-0.59
Brooklyn	Borough Park (CD12)	312	4.4	2.8	3.6	1.7	0.9	1.1	0.5	0.2	-0.59
Brooklyn	Greenpoint and Williamsburg (CD1)	301	4.3	3	3.7	2	1.2	1.4	0.6	0.2	-0.58

Borough	Geography	ID	Winter Average 2008-2009	Winter Average 2009-2010	Winter Average 2010-2011	Winter Average 2011-2012	Winter Average 2012-2013	Winter Average 2013-2014	Winter Average 2014-2015	Winter Average 2015-2016	Slope (per year decline)
Queens	Kew Gardens and Woodhaven (CD9)	409	4.2	3.1	3.7	1.8	1.1	1.4	0.5	0.24	-0.58
Queens	Flushing and Whitestone (CD7)	407	4.5	3	3.9	2	1.2	1.9	0.7	0.26	-0.57
Queens	Ridgewood and Maspeth (CD5)	405	4.1	2.9	3.6	1.8	1.1	1.4	0.6	0.21	-0.55
Bronx	Throgs Neck and Co-op City (CD10)	210	4.4	2.8	3.6	2.1	1.2	1.6	0.9	0.26	-0.54
Queens	Hillcrest and Fresh Meadows (CD8)	408	4	2.8	3.4	1.7	0.9	1.5	0.5	0.26	-0.53
Brooklyn	East New York and Starrett City (CD5)	305	3.8	2.8	3.3	1.5	1	1.1	0.5	0.21	-0.52
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	3.9	2.5	3.2	1.6	0.8	1.1	0.4	0.17	-0.52
Brooklyn	Sunset Park (CD7)	307	3.8	2.3	3.1	1.5	0.8	1	0.4	0.17	-0.50
Brooklyn	Bensonhurst (CD11)	311	3.6	2.4	2.9	1.3	0.8	1	0.5	0.18	-0.47
Queens	South Ozone Park and Howard Beach (CD10)	410	3.4	2.5	2.9	1.4	0.8	1.1	0.4	0.22	-0.46
Queens	Bayside and Little Neck (CD11)	411	3.7	2.4	3.1	1.6	0.8	1.5	0.6	0.27	-0.46
Queens	Jamaica and Hollis (CD12)	412	3.4	2.5	2.9	1.4	0.7	1.2	0.4	0.25	-0.46
Brooklyn	Sheepshead Bay (CD15)	315	3.3	2.3	2.7	1.2	0.7	0.8	0.4	0.18	-0.45
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	3.4	2.1	2.8	1.3	0.7	1	0.4	0.16	-0.44
Brooklyn	Flatlands and Canarsie (CD18)	318	3.2	2.3	2.7	1.2	0.8	0.9	0.4	0.18	-0.43
Queens	Queens Village (CD13)	413	3.1	2.2	2.5	1.3	0.6	1.1	0.4	0.26	-0.40
Brooklyn	Coney Island (CD13)	313	2.9	2	2.4	1	0.6	0.8	0.4	0.17	-0.38
Queens	Rockaway and Broad Channel (CD14)	414	2.4	1.9	1.9	0.9	0.5	0.6	0.3	0.2	-0.33
Staten Island	St. George and Stapleton (CD1)	501	2.6	1.4	2.1	1	0.5	1	0.3	0.13	-0.32
Staten Island	South Beach and Willowbrook (CD2)	502	2.2	1.2	1.8	0.8	0.4	0.9	0.3	0.12	-0.26
Staten Island	Tottenville and Great Kills (CD3)	503	1.8	1	1.5	0.6	0.4	0.7	0.2	0.11	-0.22

Table A3-5: Community district, annual average BC and per-year decline in levels

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Bronx	Morrisania and Crotona (CD3)	203	1.4	1.4	1.4	1.2	1.2	1.2	1.1	0.97	-0.06
Bronx	Riverdale and Fieldston (CD8)	208	1.3	1.2	1.2	1.1	1.1	1	1	0.81	-0.06
Bronx	Highbridge and Concourse (CD4)	204	1.5	1.5	1.5	1.3	1.4	1.3	1.2	1.07	-0.06
Bronx	Parkchester and Soundview (CD9)	209	1.3	1.3	1.3	1.1	1.1	1.1	1	0.9	-0.06
Manhattan	Washington Heights and Inwood (CD12)	112	1.4	1.4	1.4	1.3	1.3	1.2	1.1	1	-0.06
Bronx	Belmont and East Tremont (CD6)	206	1.4	1.4	1.4	1.3	1.3	1.2	1.1	1.01	-0.06
Bronx	Mott Haven and Melrose (CD1)	201	1.5	1.5	1.5	1.4	1.4	1.3	1.2	1.12	-0.06
Manhattan	Morningside Heights and Hamilton Heights (CD9)	109	1.4	1.4	1.4	1.3	1.3	1.2	1.1	1.03	-0.06
Manhattan	Central Harlem (CD10)	110	1.5	1.4	1.5	1.3	1.3	1.3	1.2	1.09	-0.05
Bronx	Hunts Point and Longwood (CD2)	202	1.4	1.5	1.5	1.3	1.3	1.3	1.2	1.08	-0.05
Bronx	Kingsbridge Heights and Bedford (CD7)	207	1.3	1.3	1.3	1.2	1.2	1.1	1.1	0.92	-0.05
Bronx	Williamsbridge and Baychester (CD12)	212	1.2	1.2	1.2	1.1	1.1	1	1	0.82	-0.05
Bronx	Fordham and University Heights (CD5)	205	1.5	1.5	1.5	1.4	1.4	1.3	1.3	1.12	-0.05
Bronx	Morris Park and Bronxdale (CD11)	211	1.2	1.2	1.2	1.1	1.1	1	1	0.83	-0.05
Bronx	Throgs Neck and Co-op City (CD10)	210	1.2	1.2	1.2	1.1	1.1	1	1	0.88	-0.05
Queens	Long Island City and Astoria (CD1)	401	1.2	1.2	1.2	1.1	1.1	1	1	0.88	-0.05
Queens	Flushing and Whitestone (CD7)	407	1.1	1.1	1.1	1	1	0.9	0.9	0.8	-0.04
Queens	Woodside and Sunnyside (CD2)	402	1.5	1.5	1.5	1.4	1.4	1.3	1.3	1.21	-0.04
Manhattan	Upper West Side (CD7)	107	1.5	1.4	1.5	1.3	1.3	1.3	1.3	1.14	-0.04
Queens	Elmhurst and Corona (CD4)	404	1.2	1.1	1.1	1	1	1	0.9	0.87	-0.04
Queens	Rego Park and Forest Hills (CD6)	406	1.1	1.1	1.1	1	0.9	0.9	0.9	0.83	-0.04
Brooklyn	East New York and Starrett City (CD5)	305	1.1	1.1	1	1	0.9	0.8	0.9	0.83	-0.04
Brooklyn	Coney Island (CD13)	313	0.8	0.9	0.9	0.8	0.7	0.6	0.6	0.65	-0.04
Brooklyn	East Flatbush (CD17)	317	1	1.1	1	1	0.9	0.8	0.8	0.81	-0.04
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	1.3	1.3	1.3	1.2	1.1	1.1	1.1	1.04	-0.04
Brooklyn	Borough Park (CD12)	312	1	1	1	0.9	0.8	0.8	0.8	0.74	-0.04
Queens	Jackson Heights (CD3)	403	1.1	1	1.1	1	0.9	0.9	0.9	0.77	-0.04
Brooklyn	Brownsville (CD16)	316	1.1	1.1	1.1	1	0.9	0.9	0.9	0.85	-0.04
Brooklyn	Flatbush and Midwood (CD14)	314	1	1	1	0.9	0.8	0.8	0.8	0.75	-0.04
Manhattan	East Harlem (CD11)	111	1.4	1.4	1.4	1.3	1.3	1.3	1.2	1.1	-0.04
Staten Island	St. George and Stapleton (CD1)	501	1	0.9	0.9	0.9	0.8	0.8	0.7	0.72	-0.04
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	1.1	1.1	1.1	1	0.9	0.9	0.9	0.87	-0.04
Queens	Kew Gardens and Woodhaven (CD9)	409	1	1	1	0.9	0.8	0.8	0.8	0.77	-0.04
Brooklyn	Bedford Stuyvesant (CD3)	303	1.1	1.1	1	1	0.9	0.9	0.9	0.83	-0.04
Queens	Ridgewood and Maspeth (CD5)	405	1.1	1.1	1.1	1	1	0.9	0.9	0.86	-0.04
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	1.3	1.3	1.3	1.2	1.2	1.1	1.1	1.06	-0.04

Borough	Geography	ID	Annual Average 2009	Annual Average 2010	Annual Average 2011	Annual Average 2012	Annual Average 2013	Annual Average 2014	Annual Average 2015	Annual Average 2016	Slope (per year decline)
Queens	South Ozone Park and Howard Beach (CD10)	410	0.9	0.9	0.9	0.8	0.7	0.7	0.7	0.68	-0.04
Brooklyn	Bushwick (CD4)	304	1.1	1.1	1.1	1	1	0.9	0.9	0.87	-0.04
Queens	Hillcrest and Fresh Meadows (CD8)	408	1	1	1	0.9	0.9	0.8	0.8	0.77	-0.04
Manhattan	Upper East Side (CD8)	108	1.6	1.6	1.6	1.5	1.5	1.5	1.4	1.33	-0.04
Brooklyn	Flatlands and Canarsie (CD18)	318	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.68	-0.04
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	1	1	1	0.9	0.9	0.8	0.8	0.78	-0.04
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	1.7	1.6	1.6	1.5	1.5	1.5	1.5	1.37	-0.04
Brooklyn	Sunset Park (CD7)	307	1.2	1.2	1.2	1.1	1.1	1	1	0.99	-0.04
Manhattan	Financial District (CD1)	101	1.4	1.4	1.4	1.3	1.3	1.3	1.2	1.15	-0.04
Brooklyn	Bensonhurst (CD11)	311	0.9	0.9	0.9	0.8	0.8	0.7	0.7	0.7	-0.04
Manhattan	Greenwich Village and Soho (CD2)	102	1.8	1.7	1.8	1.6	1.6	1.6	1.6	1.53	-0.04
Manhattan	Lower East Side and Chinatown (CD3)	103	1.6	1.5	1.5	1.4	1.4	1.4	1.4	1.29	-0.04
Manhattan	Midtown (CD5)	105	1.9	1.8	1.9	1.8	1.7	1.8	1.7	1.61	-0.03
Brooklyn	Sheepshead Bay (CD15)	315	0.8	0.9	0.9	0.8	0.7	0.6	0.7	0.67	-0.03
Queens	Bayside and Little Neck (CD11)	411	1	1	1	0.9	0.9	0.9	0.8	0.77	-0.03
Staten Island	South Beach and Willowbrook (CD2)	502	0.9	0.9	0.9	0.9	0.8	0.8	0.7	0.69	-0.03
Brooklyn	Greenpoint and Williamsburg (CD1)	301	1.5	1.4	1.4	1.4	1.3	1.3	1.3	1.22	-0.03
Manhattan	Clinton and Chelsea (CD4)	104	1.7	1.6	1.6	1.6	1.5	1.6	1.5	1.39	-0.03
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	1	1	1	1	0.9	0.9	0.8	0.81	-0.03
Staten Island	Tottenville and Great Kills (CD3)	503	0.8	0.8	0.8	0.8	0.7	0.7	0.6	0.63	-0.03
Queens	Queens Village (CD13)	413	0.8	0.9	0.9	0.8	0.7	0.7	0.7	0.68	-0.03
Queens	Jamaica and Hollis (CD12)	412	0.9	1	0.9	0.9	0.8	0.8	0.8	0.75	-0.03
Queens	Rockaway and Broad Channel (CD14)	414	0.6	0.7	0.7	0.6	0.6	0.5	0.5	0.56	-0.02

Table A3-6: Community district, summertime average O<sub>3</sub> and per-year change in levels

Borough	Geography	ID	Summer Average 2009	Summer Average 2010	Summer Average 2011	Summer Average 2012	Summer Average 2013	Summer Average 2014	Summer Average 2015	Summer Average 2016	Slope (per year decline)
Queens	Rockaway and Broad Channel (CD14)	414	33.7	39.1	39.1	40.4	36.9	36.3	37.5	38.26	0.14
Brooklyn	Coney Island (CD13)	313	30.6	37.4	37.5	37.9	34.6	33.2	35.8	36.33	0.19
Bronx	Throgs Neck and Co-op City (CD10)	210	27.8	35	34.9	36	32.6	32.5	31.7	34	0.19
Bronx	Williamsbridge and Baychester (CD12)	212	26.6	33.8	33.2	35	31.6	31.4	29.8	33.19	0.21
Brooklyn	Sheepshead Bay (CD15)	315	29.3	36.2	36.2	36.8	33.6	32.4	34.8	35.37	0.25
Bronx	Morris Park and Bronxdale (CD11)	211	26.1	33.7	33.5	35	31.6	31.4	30.5	33.21	0.29
Brooklyn	Flatlands and Canarsie (CD18)	318	28.9	36.1	35.9	36.6	33.6	33.1	34.7	35.71	0.35
Brooklyn	Bensonhurst (CD11)	311	27.4	34.9	34.5	34.9	32	31.5	33.4	34.51	0.36
Queens	Flushing and Whitestone (CD7)	407	26.3	34	34	34.5	31.5	32	31.9	33.48	0.37
Queens	Bayside and Little Neck (CD11)	411	26.3	33.6	33.1	33.9	30.8	32.3	31.4	33.06	0.37
Queens	Woodside and Sunnyside (CD2)	402	26.4	33.3	32.8	33.8	30.7	30.9	31.6	33.37	0.37
Queens	Long Island City and Astoria (CD1)	401	26.3	33.5	33.3	34.5	31.2	31.1	31.8	33.44	0.38
Bronx	Parkchester and Soundview (CD9)	209	25.6	33.5	33.6	34.9	31.5	31.3	31.2	33.25	0.38
Staten Island	South Beach and Willowbrook (CD2)	502	27	35.2	34.1	33.7	31.3	31.6	32.5	35.11	0.40
Staten Island	Tottenville and Great Kills (CD3)	503	27.6	36.5	35.9	35.3	32.6	31.9	34.1	36.19	0.40
Queens	Queens Village (CD13)	413	27.8	34.4	33.7	35	31.9	33.7	32.6	34.43	0.41
Queens	Jackson Heights (CD3)	403	25.5	33.1	33.1	33.7	30.7	30.8	31.5	32.97	0.41
Queens	Ridgewood and Maspeth (CD5)	405	26.1	33.7	33.3	33.7	31.1	31.5	32.1	33.76	0.45
Queens	Elmhurst and Corona (CD4)	404	24.9	32.7	32.6	32.9	30.2	30.5	31.2	32.75	0.46
Bronx	Hunts Point and Longwood (CD2)	202	25.5	33.6	33.8	35.3	31.9	31.6	31.8	33.74	0.46
Brooklyn	Bay Ridge and Dyker Heights (CD10)	310	25.6	33.4	32.5	33	30.3	30.7	31.8	33.51	0.47
Brooklyn	Greenpoint and Williamsburg (CD1)	301	25.4	32.5	31.6	32.7	29.8	30.5	31	32.99	0.47
Bronx	Riverdale and Fieldston (CD8)	208	23.7	32	31.3	33.8	30.4	30.5	28.8	32.52	0.48
Queens	South Ozone Park and Howard Beach (CD10)	410	27.2	34.6	34.2	34.8	32.3	33	33.1	34.86	0.48
Staten Island	St. George and Stapleton (CD1)	501	24.9	33	31.5	31.5	29.2	30.3	30.4	33.4	0.48
Brooklyn	East New York and Starrett City (CD5)	305	26.7	34.3	33.9	34.4	31.9	32.3	32.9	34.57	0.49
Brooklyn	Flatbush and Midwood (CD14)	314	25.6	33.4	32.8	33.5	30.8	30.7	32.3	33.56	0.49
Queens	Hillcrest and Fresh Meadows (CD8)	408	25.2	32.9	32.5	33	30.4	31.7	31.3	32.98	0.49
Bronx	Belmont and East Tremont (CD6)	206	23.7	32.1	31.9	33.8	30.5	30.3	29.8	32.5	0.50
Bronx	Kingsbridge Heights and Bedford (CD7)	207	23.6	31.9	31.4	33.5	30.3	30.3	29.2	32.46	0.50
Brooklyn	Borough Park (CD12)	312	25.3	33.2	32.4	33.1	30.4	30.6	32	33.45	0.51
Queens	Rego Park and Forest Hills (CD6)	406	24.5	32.5	32.2	32.5	30	30.6	31.1	32.7	0.51
Queens	Jamaica and Hollis (CD12)	412	26.4	33.6	33	33.9	31.3	32.8	32.2	34.03	0.51
Brooklyn	East Flatbush (CD17)	317	25.7	33.6	33	33.6	31	31.2	32.4	33.88	0.52
Brooklyn	Bushwick (CD4)	304	25.3	33	32.3	32.9	30.4	31	31.7	33.47	0.53
Queens	Kew Gardens and Woodhaven (CD9)	409	25.4	33.2	32.8	33.2	30.8	31.7	31.8	33.58	0.53

Borough	Geography	ID	Summer Average 2009	Summer Average 2010	Summer Average 2011	Summer Average 2012	Summer Average 2013	Summer Average 2014	Summer Average 2015	Summer Average 2016	Slope (per year decline)
Brooklyn	Brownsville (CD16)	316	25.4	33.3	32.6	33.2	30.7	31.2	32.1	33.74	0.54
Bronx	Mott Haven and Melrose (CD1)	201	24	32.1	32.1	34	30.6	30.5	30.8	32.81	0.56
Bronx	Morrisania and Crotona (CD3)	203	23.4	31.9	31.9	33.7	30.4	30.3	30.2	32.58	0.57
Brooklyn	Sunset Park (CD7)	307	24.5	32.6	31.4	32.2	29.6	30.6	31.2	33.19	0.58
Brooklyn	Bedford Stuyvesant (CD3)	303	23.9	31.8	30.8	31.6	29.1	29.9	30.6	32.55	0.59
Manhattan	East Harlem (CD11)	111	22.8	30.6	30.3	32.3	28.9	29	29.6	31.66	0.59
Brooklyn	South Crown Heights and Lefferts Gardens (CD9)	309	24.2	32.3	31.4	32.1	29.6	30.3	31.2	32.98	0.60
Brooklyn	Crown Heights and Prospect Heights (CD8)	308	23.6	31.7	30.7	31.5	29	29.8	30.7	32.54	0.62
Brooklyn	Park Slope and Carroll Gardens (CD6)	306	23.4	31.3	29.9	31	28.4	29.7	30	32.27	0.62
Bronx	Fordham and University Heights (CD5)	205	22.2	31	30.7	32.9	29.7	29.7	29.1	32.06	0.63
Brooklyn	Fort Greene and Brooklyn Heights (CD2)	302	22.7	30.5	29.1	30.2	27.6	28.9	29.4	31.63	0.64
Manhattan	Lower East Side and Chinatown (CD3)	103	21.6	28.8	27.4	28.8	26.1	27.3	28	30.41	0.65
Manhattan	Upper West Side (CD7)	107	21.4	28.7	27.9	30.2	26.9	27.4	28.1	30.55	0.67
Bronx	Highbridge and Concourse (CD4)	204	22.3	31	30.8	33	29.7	29.8	29.7	32.2	0.67
Manhattan	Washington Heights and Inwood (CD12)	112	21.9	30.7	30.3	32.8	29.5	29.7	28.9	32.01	0.67
Manhattan	Central Harlem (CD10)	110	21.6	29.8	29.5	31.8	28.5	28.6	29.1	31.42	0.71
Manhattan	Upper East Side (CD8)	108	20.8	28.1	27.4	29.2	26.2	26.6	28	30.12	0.71
Manhattan	Morningside Heights and Hamilton Heights (CD9)	109	21.3	29.6	29.2	31.6	28.3	28.5	28.8	31.29	0.72
Manhattan	Greenwich Village and Soho (CD2)	102	20.5	27.7	26.1	27.7	25.1	26.5	27.2	29.87	0.73
Manhattan	Clinton and Chelsea (CD4)	104	20.3	27.3	25.9	27.9	25.1	26.1	27.1	29.76	0.75
Manhattan	Stuyvesant Town and Turtle Bay (CD6)	106	17.8	25.3	24	25.5	23.1	24.1	25.8	28.23	0.87
Manhattan	Financial District (CD1)	101	19.1	27	25.2	26.6	24.4	26.1	27.1	29.81	0.90
Manhattan	Midtown (CD5)	105	14.4	22.5	21	22.5	20.7	21.8	24.3	26.92	1.16



